Published in final edited form as:

Vis Neurosci. 2019 January ; 36: E006. doi:10.1017/S095252381900004X.

Synaptic inhibition tunes contrast computation in the retina

Nicholas W. Oesch^{1,2} and Jeffrey S. Diamond¹

¹Synaptic Physiology Section, National Institute of Neurological Disorders and Stroke, National Institutes of Health, 35 Convent Drive, Bethesda, MD 20892-3701

²Present address: University of California, San Diego, Department of Psychology, Department of Ophthalmology, 9500 Gilman Drive MC#0109, La Jolla, CA 92093-0109

Abstract

Inhibition shapes activity and signal processing in neural networks through numerous mechanisms mediated by many different cell types. Here, we examined how one type of GABAergic interneuron in the retina, the A17 amacrine cell, influences visual information processing. Our results suggest that A17s, which make reciprocal feedback inhibitory synapses onto rod bipolar cell (RBC) synaptic terminals, extend the luminance range over which RBC synapses compute temporal contrast and enhance the reliability of contrast signals over this range. Inhibition from other amacrine cells does not influence these computational features. Although A17-mediated feedback is mediated by both GABA_A and GABA_C receptors, the latter plays the primary role in extending the range of contrast computation. These results identify specific functions for an inhibitory interneuron subtype, as well as specific synaptic receptors, in a behaviorally relevant neural computation.

Keywords

Amacrine cell; Weber contrast; ribbon synapse; feedback inhibition

Introduction

Interneurons in the brain exhibit remarkable morphological, molecular and physiological diversity, suggesting that different interneuron subtypes may perform specific computational tasks. Consistent with this idea, interneuron subtypes target distinct populations of neurons and specific subcellular compartments within neurons, including functionally presynaptic compartments such as the distal axon and synaptic terminal. The computational impact of presynaptic inhibition is complex and depends on factors including the local chloride reversal potential, activity patterns and release probability in the target axon (Engelman & MacDermott, 2004; Kullmann *et al.*, 2005). In the retina, previous work has shown that presynaptic inhibition extends the dynamic range of light-evoked transmission from rod and cone bipolar cells (Euler & Masland, 2000; Sagdullaev *et al.*, 2006).

Here, we examined the impact of presynaptic inhibition by a single interneuron subtype on visual signaling in the rod pathway, which mediates night vision. Rod bipolar cells (RBCs) receive synaptic signals from rod photoreceptors and encode luminance (ϕ) in their membrane potential (V_{RBC} ; Berntson & Taylor, 2000; Euler & Masland, 2000; Oesch & Diamond, 2011). RBC synapses continuously release neurotransmitter and encode V_{RBC} (and luminance) in the frequency of synaptic vesicle release. Moreover, a step increase in luminance from one level (ϕ_1) to another, brighter level (ϕ_2) (temporal contrast) elicits a transient bout of vesicle release that corresponds to the change between luminance levels. This transient release is proportional to Weber contrast ($C_w = [\phi_2 - \phi_1]/\phi_1$) (Oesch & Diamond, 2011), a metric reflecting the psychophysical sensitivity to changes in sensory stimuli (Weber, 1834; Aguilar & Stiles, 1954; Whittle, 1994).

Synaptic release from RBCs can be recorded as an excitatory postsynaptic current (EPSC) in postsynaptic AII amacrine cells. In response to a sustained stimulus, this EPSC consists of two kinetically distinct phases: transient and sustained. The size of the transient phase of the EPSC is influenced by the size of the readily releasable pool (RRP), the synaptic vesicles docked at the presynaptic membrane that are primed for immediate release. At RBC synapses, the RRP size reflects a dynamic balance between vesicle release from and replenishment to presynaptic release sites and varies with V_{RBC} . (Holt *et al.*, 2004; Jarsky *et al.*, 2011; Oesch & Diamond, 2011; Graydon *et al.*, 2014): As V_{RBC} becomes more depolarized with increased luminance, RRP size decreases. Consequently, RRP size constitutes a physical (inverse) readout of luminance (Figure 1A), and the transient bout of vesicle release corresponds to the change in RRP size between the two luminance levels. Because V_{RBC} encodes luminance in roughly a log-linear fashion (Oesch & Diamond, 2011), the extent to which the readout of this computation (i.e., the contrast gain) remains consistent over the physiological luminance range depends on the linearity of the $\log(\phi_1)$ -RRP relationship (Figure 1A,B).

RBC synaptic terminals receive inhibitory synaptic input from numerous subtypes of GABAergic and glycinergic amacrine cells (Dowling & Boycott, 1966; Sterling & Lampson, 1986; Eggers & Lukasiewicz, 2006a, b), including reciprocal GABAergic feedback inhibition from A17 amacrine cells (Nelson & Kolb, 1985; Dong & Hare, 2003). GABAergic inhibition has been shown to expand the dynamic range of RBC signaling (Euler & Masland, 2000), but a specific role for A17s has not been investigated. Past work has suggested that A17-mediated inhibition may mediate center surround inhibition in the rod pathway, (Nelson & Kolb, 1985; Bloomfield, 1992; Völgyi *et al.*, 2002; Zhang *et al.*, 2002), enhance and sharpen responses near visual threshold (Dong & Werblin, 1998; Dong & Hare, 2003; Grimes *et al.*, 2010), or improve the signal-to-noise characteristics of single-photon responses (Grimes *et al.*, 2015). Here, we examined how presynaptic inhibition influences the dynamic range and consistency of contrast computation at the RBC terminal over a range of background luminance (ϕ_1). We recorded light-evoked EPSCs from AIIIs in acutely prepared slices of dark-adapted rat retina and pharmacologically manipulated different components of presynaptic inhibition. Rather than universally decreasing release, the presence of GABAergic inhibition actually increased the transient EPSC evoked by a step increase in luminance over most of the background luminance range, enabling RBC synapses to compute Weber contrast over a wider range of luminance conditions. This effect

appears to be mediated primarily by feedback inhibition from A17s: blocking inhibition from other GABAergic and glycinergic amacrine cells subtly modulated the gain of the RRP vs ϕ_1 relationship but did not affect the dynamic range of Weber contrast computation. We also found that GABA_C receptors, located on RBC terminals and activated by GABA released from A17s (Grimes *et al.*, 2009; Grimes *et al.*, 2015), mediate most of the A17's inhibitory effects on contrast computation.

Materials and Methods

Slice and Recording Procedures

All animal procedures were conducted in accordance with NIH guidelines, as approved by the National Institute of Neurological Disorders and Stroke Animal Care and Use Committee (ASP 1220). Acute slices (200 μ m thick) were prepared from retina isolated from Sprague-Dawley rats (p18-p22) as previously described (Singer & Diamond, 2003; Oesch & Diamond, 2011). Rats were dark-adapted for 2–3 h prior to dissection and all subsequent procedures were performed in darkness or under IR illumination (940 nm). Retinas were isolated, and all subsequent procedures and recordings were performed, in Ames Medium (United States Biological) equilibrated with 95% O₂/5% CO₂. For physiological recordings, retinal slices were superfused with equilibrated Ames media at 4–5 mL/min at 35–37°C. Pharmacological agents were added to the superfusate at the specified final concentrations: 0.5 μ M TTX (Ascent Scientific), 5 μ M SR-95531 (Ascent Scientific), 3 μ M strychnine (Sigma), and 50 μ M TPMPA (Tocris). RBCs and AIIs were targeted visually using IR (940 nm) DIC or Dodt contrast video microscopy. Whole-cell voltage-clamp recordings were made with pipettes (4–6 M Ω) filled with solution containing (in mM) 90 CsCH₃SO₃, 20 TEA-Cl, 10 HEPES, 10 EGTA, 10 phosphocreatine disodium salt hydrate, 4 Mg-ATP, 0.4 Na-GTP. Recordings were obtained using an Axopatch 700B amplifier (Molecular Devices), low-pass Bessel filtered at 4 kHz and digitized at 20–50 kHz with an Instrutech ITC-18 A/D board (HEKA Elektronik) controlled by custom acquisition software written in IgorPro (WaveMetrics). Traces were down-sampled to 10 kHz and digitally filtered using a binomial smoothing algorithm (IgorPro) to approximate a Gaussian filter with a characteristic cutoff frequency of 2 kHz.

Light Stimulation

Light stimuli were generated with a 525-nm LED and directed through a 1.0 NA 40 X water-immersion objective (Zeiss) to uniformly illuminate the 0.45 mm-diameter field of view. LED intensity controlled by pulse width modulation of the LED (500 Hz; 0.1% resolution) using a custom-built constant current power supply controlled digitally via the ITC18 and acquisition software and further attenuated with neutral density filters in a motorized filter wheel (ThorLabs), also controlled by the acquisition software. Photon flux was measured at the focal plane of the microscope objective using a DR-2000 radiometer (Gamma Scientific) and was converted to photoisomerizations (r^*)/rod assuming a collecting area of 0.5 μ m (Field and Rieke, 2002). Light stimuli were presented at 15-second intervals and each set of light stimuli was presented 3 times.

Data Analysis

Unless indicated otherwise, all data are reported as mean $\pm$ SD. Statistical comparisons were made with *t*-tests ($\alpha=0.05$) and, in the case of multiple (*n*) comparisons, Bonferroni correction ($\alpha=0.05/n$) was applied.

To measure transient release components for either light-evoked or paired recordings, we integrated release over the transient component (0.5 sec) and subtracted the baseline offset measured immediately preceding the transient response to isolate the transient change in release. (Measuring the peak current yielded qualitatively similar results.) For comparison, we normalized all transient and sustained responses to the maximal response. To determine threshold (10% RRP depletion), saturation (90% RRP depletion) and dynamic range, we fit the integrated release from the second transient in the depletion protocol to the Hill Function:

$$y = base + \frac{max - base}{\left(1 + \left(\frac{x_{half}}{x}\right)^{rate}\right)},$$

using the IgorPro “CurveFit” routine. We then rescaled the y-axis between 0 and 1 and measured the value x-value where y crossed 0.1 ($\phi_{saturation}$) and 0.9 ($\phi_{threshold}$). The dynamic range was calculated as the difference between $\phi_{saturation}$ and $\phi_{threshold}$. To compare how “linear” RRP depletion was relative to log luminance, we used IgorPro’s CurveFit routine to fit the data to function $(m - \log(\phi)) + b$. The chi-square was measured for each fit against the data, and pairwise comparisons were made using the two-tailed *t*-test. Contrast gain distributions were fit with a sigmoid function for the purposes of qualitative comparison:

$$y = base + \left[\frac{max}{1 + \exp\left(\frac{x_{1/2} - x}{rate}\right)} \right].$$

Results

All amacrine cells receive glutamatergic input from RBCs via AMPA receptors that provide a linear, high-fidelity readout of RBC synaptic release (Heidelberger & Matthews, 1992; Heidelberger *et al.*, 1994; von Gersdorff & Matthews, 1997; Hartveit, 1999; Singer & Diamond, 2003; Singer *et al.*, 2004; Oesch & Diamond, 2011). Our previous work demonstrated that the steady-state size of the RRP is a key substrate for computing contrast signals at the RBC (Oesch & Diamond, 2011), but we did not examine how inhibitory inputs to the RBC terminal influence contrast coding. To address this, we first recorded light-evoked changes in whole-cell currents in AII in dark-adapted acute rat retinal slices with synaptic inhibition intact (Figure 2). To elicit primarily rod-driven responses, light stimulus levels were kept well below the threshold for cone activation (<100 rhodopsin photoisomerizations per rod per s [$r^*/rod/s$]; Nikonov *et al.*, 2006; Ke *et al.*, 2014)).

The full-field luminance was stepped from complete darkness to different luminance (ϕ_1) values (0.08 – 72 $r^*/rod/s$) for 1 s, enough time for RBC release and RRP size to settle to

new steady-state levels (Figure 2A). The luminance was then stepped to a higher level (ϕ_2 ; 72 r*/rod/s; Figure 2A) to evoke release of the remaining RRP (Oesch & Diamond, 2011). Light-evoked EPSCs were recorded from the AII amacrine cells and the transient EPSC size was quantified by integrating the transient component of the EPSC (after subtracting the preceding sustained EPSC baseline). The sum of the two transient responses was relatively constant over the entire range of ϕ_1 levels (squares, Figure 2B) and equivalent to the transient response evoked by a step from darkness directly to ϕ_2 , indicating that the sum reflects release of the entire RRP (i.e., the total number of vesicles available in darkness). Thus, the transient EPSC evoked by ϕ_1 reflects the change in RRP size between complete darkness and ϕ_1 , and the transient EPSC evoked by the second step to ϕ_2 reflects the RRP size at each ϕ_1 (Oesch & Diamond, 2011). Light steps were brief and slices were kept in complete darkness between trials to minimize slower adaptation processes in the rod pathway (Grimes *et al.*, 2014; Ke *et al.*, 2014); accordingly, we saw little change in sensitivity over the course of an experiment.

To test the effects of inhibition on light-evoked EPSCs in AII, this light stimulation protocol was repeated after the slice was superfused with extracellular solution containing SR95531 (5 μ M), TPMPA (50 μ M), and strychnine (3 μ M) to block GABA_A, GABA_C and glycine receptors, respectively (Figure 2A, gray traces). In the absence of inhibition, transient responses to ϕ_2 decreased more steeply as a function of ϕ_1 . To quantify this, we measured the luminance values where RRP occupancy fell below 90% and 10% of maximum occupancy from the Hill fit to the data. In the presence of inhibitory blockers, luminance corresponding to 10% maximum occupancy decreased significantly compared to control (2.25 ± 1.1 r*/rod/sec vs. 15.6 ± 6.4 r*/rod/sec, $n=9$; $p=2.3 \times 10^{-4}$, paired *t*-test; Figure 2B,C). Therefore, blocking inhibition reduced the dynamic range (luminance range between the 10% and 90% maximal occupancy levels) over which RBC synapses could encode luminance changes, from 39.8 ± 3.1 dB in control conditions to 25.6 ± 4.9 dB with inhibition blocked ($n=9$, $p=3.9 \times 10^{-5}$, paired *t*-test; see Methods), with no significant change in the size of the RRP ($n=9$, $p=0.75$, paired *t*-test; Figure 2D). These results are analogous to those reported in ON ganglion cells of the mouse retina (Sagdullaev *et al.*, 2006) and RBCs in the rat retina (Euler & Masland, 2000).

By expanding the dynamic range of the transient responses to ϕ_2 , inhibition may enhance the range over which RBCs report contrast consistently (Figure 1). To test this hypothesis, we repeated the two-step stimulus paradigm but varied luminance steps so that the Weber contrast between ϕ_1 and ϕ_2 was nearly the same (Weber contrast = 1.97 ± 0.073) in each trial across a wide range of ϕ_1 values (0.1 – 50 r*/rod/s; Figure 3A). Under control conditions, transient release evoked by the step from ϕ_1 to ϕ_2 remained constant up to $\phi_1 = 10$ r*/rod/s, matching the constant Weber contrast in the stimulus (Figure 3B); contrast was not encoded above 10 r*/rod/s, presumably because at these luminance levels the RRP was nearly completely depleted (Figure 2B) and further luminance increases could not elicit transient release. Blocking inhibition during constant contrast stimulation reduced the ϕ_1 range over which Weber contrast was encoded (Figure 3B; see Methods), as predicted from the observed changes in dynamic range (Figure 2D). These results demonstrate that synaptic inhibition increases the range of background luminance over which Weber contrast is signaled.

Given the diversity of inhibitory signaling in the retina, we next examined how specific types of inhibition influenced contrast signaling. RBC axon terminals receive at least three pharmacologically distinguishable classes of inhibitory inputs. Here we refer to two of the three classes as “lateral” inhibition, because they are mediated by amacrine cells that are driven by other bipolar cells (Eggers *et al.*, 2007; Chávez *et al.*, 2010). Lateral inhibition from glycinergic amacrine cells is blocked completely by the glycine receptor antagonist strychnine, whereas lateral inhibition from GABAergic amacrine cells is eliminated by blocking Na_v channels with tetrodotoxin (TTX; Chávez & Diamond, 2008). This allows us to isolate the third component, reciprocal GABAergic feedback inhibition from A17 amacrine cells, which inhibits the same RBC terminal from which it receives glutamatergic excitation; reciprocal inhibition persists in the presence of TTX but is eliminated by GABA receptor antagonists (Hartveit, 1999; Chávez *et al.*, 2006).

We examined the effect of lateral inhibition on the relationship between RRP and luminance by repeating this stimulation protocol, first in control solution and then in the presence of strychnine (3 μM) and TTX (0.5 μM) (Figure 4A). Blocking lateral inhibition did not significantly change the slope of the $\log(\phi_1)$ vs. $Q_{\text{transient}}$ relationship ($p=0.53$, paired *t*-test; Figure 4B), and we did not detect a significant change in the dynamic range (34.5 ± 3.9 vs 37.7 ± 2.3 dB, $n=6$, $p=0.07$, paired *t*-test). Accordingly, in experiments in which ϕ_2 was varied to maintain constant contrast (as in Figure 3), blocking lateral inhibition did not change the range over which contrast was encoded reliably (Figure 4C,D).

Previous work has demonstrated that GABA release from A17s can activate both GABA_A and GABA_C receptors on RBC terminals (Pan & Lipton, 1995; Hartveit, 1999; Euler & Masland, 2000; Singer & Diamond, 2003; Chávez *et al.*, 2006; Chávez *et al.*, 2010; Grimes *et al.*, 2015). In ON cone bipolar cells, GABA_C receptor-mediated feedback enhances the dynamic range of synaptic signaling (Sagdullaev *et al.*, 2006). GABAergic inhibition has been shown to expand the dynamic range of RBC responses (Euler & Masland, 2000), but a specific role for GABA_C receptors was not examined. Here, blocking GABA_C receptors with TPMPA significantly reduced the range over which RBC synapses encoded luminance (from 38.1 ± 5.1 to 29.2 ± 4.6 dB, $n=12$, $p=6.8 \times 10^{-5}$, paired *t*-test), similar to the effects blocking inhibition entirely (Figure 3) and analogous to effects reported in the cone pathway (Sagdullaev *et al.*, 2006). Accordingly, TPMPA also reduced the range over which RBC transient responses reliably encoded contrast (Figure 5D). These results show that GABA_C receptors play a major role in mediating the A17's modulation of contrast gain in RBC synapses.

Discussion

Here we show that lateral and reciprocal presynaptic inhibition, mediated by pharmacologically separable populations of interneurons, play distinct roles in modulating synaptic release and contrast/luminance coding at RBC synapses in rat retina. We show that reciprocal but not lateral feedback synapses onto RBC terminals equalize the contrast gain across a range of background luminance. Furthermore, we demonstrate that presynaptic GABA_C receptors on RBC terminals, activated by reciprocal feedback from A17 amacrine cells, play the primary role in extending the range of contrast computation.

Computational roles of presynaptic inhibition

These results indicate that reciprocal synaptic inhibition from the A17 amacrine cell expands the range of luminance over which RBC synapses encode Weber contrast. In addition, reciprocal feedback inhibition onto RBC synaptic terminals maintains the contrast responses (contrast gain) at a consistent level across a range of luminance. Ganglion cell recordings and psychophysical tests of luminance perception both demonstrate that contrast gain is largely constant across a wide range of background luminance (Enroth-Cugell & Shapley, 1973), suggesting that this function may be involved in a fundamental optimization of the contrast signal to meet downstream demands in the visual system. This work contributes to our understanding of the mechanisms that may underlie this optimized contrast signaling.

The A17 amacrine cell's role traditionally has been thought to mediate center surround inhibition (Nelson & Kolb, 1985; Bloomfield, 1992; Völgyi *et al.*, 2002; Zhang *et al.*, 2002); more recent evidence suggests, however, that A17 output synapses operate as independent reciprocal feedback processors (Grimes *et al.*, 2010), contrary to their proposed role in integrating surround information. A subsequent study suggested a role for A17-mediated feedback in resolving weak stimuli near the visual threshold (Grimes *et al.*, 2015). Earlier work indicated that reciprocal GABAergic feedback shapes the time course of bipolar cell release in salamander and rabbit retina (Dong & Werblin, 1998; Dong & Hare, 2003). The present study elaborates on this hypothesis, demonstrating how A17-mediated reciprocal inhibition can help compute changes in luminance over time (i.e., temporal contrast) in a perceptually relevant context.

Though our experimental results point to a clear computational task achieved by A17-mediated reciprocal feedback, they do not identify a particular role for the lateral inhibition provided by other GABAergic and glycinergic amacrine cells in the rod pathway. Blocking lateral inhibition exerted only a very minor effect on synaptic release from the RBC synapse and did not alter the shape, tuning, or operating range of the RRP-luminance relationship. Given the prominence of lateral inhibitory synapses on the RBC terminal (Sterling & Lampson, 1986), it seems likely that lateral inhibition influences a different computation that was not engaged in the experiments performed here. One additional possibility is that relevant lateral connections were severed during the slicing procedure, perhaps leading to a bias toward more local circuitry. Future experiments examining feedback inhibition in the wholemount retina may address this issue.

Computing Weber contrast as gain normalization

Gain normalization, a fundamental computational strategy employed in all sensory systems and many other neural circuits, enables a particular perceptual quality or stimulus feature to be represented equally (normalized) across a wide range of input strength (Carandini & Heeger, 2012; Hong & Wilson, 2013). The Weber contrast computation in RBC terminals is a straightforward example of this strategy, because the signal representing a perceptually relevant feature (in this case the luminance change in space-time), is normalized to the intensity (luminance) of the background upon which the signal occurs, resulting in a perceptually equal response to a percent change contrast stimulus independent of background lighting conditions (Shapley & Enroth-Cugell, 1984).

Inhibition has been implicated in gain normalization elsewhere in the brain, but universal rules (if any) for inhibition's role remain incompletely resolved (Mitchell & Silver, 2003; Wilson *et al.*, 2012; Kepecs & Fishell, 2014; Roux & Buzsaki, 2015). Interestingly, the reciprocal inhibition described here does not strictly perform subtraction or division, two computations most typically associated with inhibition and neural adaptation, but rather a combination of the two. In the RBC-A17 circuit, the dynamics of reciprocal inhibition seems specifically tuned to boost or suppresses contrast signals as needed to establish more consistent contrast gain over the operating range for the RBC ribbon synapse.

Distinct roles for different components of feedback inhibition

While it is well established that different inhibitory interneurons perform distinct tasks, the work presented here demonstrates that different inhibitory receptors within the same synapse also may play distinct roles, possibly increasing the computational capacity of a single synapse. At synapses between ON cone bipolar cells and ganglion cells, activation of presynaptic GABA_C receptors reduces the extent of multivesicular release and selectively limits activation of perisynaptic NMDA receptors on ganglion cell dendrites (see also Sagdullaev *et al.*, 2006). Although both A17s and AII amacrine cells express functional NMDA receptors (Veruki *et al.*, 2018), their postsynaptic responses are mediated exclusively by AMPA receptors (Singer & Diamond, 2003; Chávez *et al.*, 2006; Veruki *et al.*, 2018). Because blocking the GABA_C receptor-mediated component of reciprocal inhibition causes effects that are similar to those observed when all reciprocal feedback inhibition is blocked (cf. Figures 2D and 5D), it appears that GABA_C receptors are the primary effectors that modulate contrast gain at the RBC synapse. Effects of GABA_A receptor blockade alone are difficult to interpret due to GABA_A receptor-mediated inhibition between amacrine cells (Eggers & Lukasiewicz, 2011). GABA_A and GABA_C receptors may underlie distinct parallel functions at RBC synapses, but our stimulus protocols may not have engaged those mediated by GABA_A receptors. Further experiments are required to fully understand the roles for these receptors in RBC signaling.

Acknowledgments

We thank Dr. Wei Li for helpful comments on the manuscript. This work was supported by the NINDS Intramural Research Program (1ZIANS003039 to JSD).

References

- Aguilar M & Stiles W. (1954). Saturation of the Rod Mechanism of the Retina at High Levels of Stimulation. *Optica Acta: International Journal of Optics* 1, 59–65.
- Berntson A & Taylor WR. (2000). Response characteristics and receptive field widths of on-bipolar cells in the mouse retina. *J Physiol* 524 Pt 3, 879–889. [PubMed: 10790165]
- Bloomfield SA. (1992). Relationship between receptive and dendritic field size of amacrine cells in the rabbit retina. *J Neurophysiol* 68, 711–725. [PubMed: 1432044]
- Carandini M & Heeger DJ. (2012). Normalization as a canonical neural computation. *Nature Reviews Neuroscience* 13, 51–62.
- Chávez AE & Diamond JS. (2008). Diverse mechanisms underlie glycinergic feedback transmission onto rod bipolar cells in rat retina. *J Neurosci* 28, 7919–7928. [PubMed: 18667624]
- Chávez AE, Grimes WN & Diamond JS. (2010). Mechanisms underlying lateral GABAergic feedback onto rod bipolar cells in rat retina. *J Neurosci* 30, 2330–2339. [PubMed: 20147559]

- Chávez AE, Singer JH & Diamond JS. (2006). Fast neurotransmitter release triggered by Ca influx through AMPA-type glutamate receptors. *Nature* 443, 705–708. [PubMed: 17036006]
- Dong CJ & Hare WA. (2003). Temporal modulation of scotopic visual signals by A17 amacrine cells in mammalian retina in vivo. *J Neurophysiol* 89, 2159–2166. [PubMed: 12686583]
- Dong CJ & Werblin FS. (1998). Temporal contrast enhancement via GABAC feedback at bipolar terminals in the tiger salamander retina. *J Neurophysiol* 79, 2171–2180. [PubMed: 9535976]
- Dowling JE & Boycott BB. (1966). Organization of the primate retina: electron microscopy. *Proc R Soc Lond B Biol Sci* 166, 80–111. [PubMed: 4382694]
- Eggers ED & Lukasiewicz PD. (2006a). GABA(A), GABA(C) and glycine receptor-mediated inhibition differentially affects light-evoked signalling from mouse retinal rod bipolar cells. *J Physiol* 572, 215–225. [PubMed: 16439422]
- Eggers ED & Lukasiewicz PD. (2006b). Receptor and transmitter release properties set the time course of retinal inhibition. *J Neurosci* 26, 9413–9425. [PubMed: 16971525]
- Eggers ED & Lukasiewicz PD. (2011). Multiple pathways of inhibition shape bipolar cell responses in the retina. *Vis Neurosci*, 1–14. [PubMed: 21348136]
- Eggers ED, McCall MA & Lukasiewicz PD. (2007). Presynaptic inhibition differentially shapes transmission in distinct circuits in the mouse retina. *J Physiol* 582, 569–582. [PubMed: 17463042]
- Engelman HS & MacDermott AB. (2004). Presynaptic ionotropic receptors and control of transmitter release. *Nat Rev Neurosci* 5, 135–145. [PubMed: 14735116]
- Enroth-Cugell C & Shapley RM. (1973). Adaptation and dynamics of cat retinal ganglion cells. *J Physiol* 233, 271–309. [PubMed: 4747229]
- Euler T & Masland RH. (2000). Light-evoked responses of bipolar cells in a mammalian retina. *J Neurophysiol* 83, 1817–1829. [PubMed: 10758094]
- Graydon CW, Zhang J, Oesch NW, Sousa AA, Leapman RD & Diamond JS. (2014). Passive diffusion as a mechanism underlying ribbon synapse vesicle release and resupply. *J Neurosci* 34, 8948–8962. [PubMed: 24990916]
- Grimes WN, Li W, Chavez AE & Diamond JS. (2009). BK channels modulate pre- and postsynaptic signaling at reciprocal synapses in retina. *Nat Neurosci* 12, 585–592. [PubMed: 19363492]
- Grimes WN, Schwartz GW & Rieke F. (2014). The synaptic and circuit mechanisms underlying a change in spatial encoding in the retina. *Neuron* 82, 460–473. [PubMed: 24742466]
- Grimes WN, Zhang J, Graydon CW, Kachar B & Diamond JS. (2010). Retinal parallel processors: more than 100 independent microcircuits operate within a single interneuron. *Neuron* 65, 873–885. [PubMed: 20346762]
- Grimes WN, Zhang J, Tian H, Graydon CW, Hoon M, Rieke F & Diamond JS. (2015). Complex inhibitory microcircuitry regulates retinal signaling near visual threshold. *J Neurophysiol* 114, 341–353. [PubMed: 25972578]
- Hartveit E (1999). Reciprocal synaptic interactions between rod bipolar cells and amacrine cells in the rat retina. *J Neurophysiol* 81, 2923–2936. [PubMed: 10368409]
- Heidelberger R, Heinemann C, Neher E & Matthews G. (1994). Calcium dependence of the rate of exocytosis in a synaptic terminal. *Nature* 371, 513–515. [PubMed: 7935764]
- Heidelberger R & Matthews G. (1992). Calcium influx and calcium current in single synaptic terminals of goldfish retinal bipolar neurons. *J Physiol* 447, 235–256. [PubMed: 1317429]
- Holt M, Cooke A, Neef A & Lagnado L. (2004). High mobility of vesicles supports continuous exocytosis at a ribbon synapse. *Curr Biol* 14, 173–183. [PubMed: 14761649]
- Hong EJ & Wilson RI. (2013). Olfactory Neuroscience: Normalization Is the Norm. *Current Biology* 23, R1091–R1093. [PubMed: 24355783]
- Jarsky T, Cembrowski M, Logan SM, Kath WL, Rieke H, Demb JB & Singer JH. (2011). A synaptic mechanism for retinal adaptation to luminance and contrast. *J Neurosci* 31, 11003–11015. [PubMed: 21795549]
- Ke JB, Wang YV, Borghuis BG, Cembrowski MS, Rieke H, Kath WL, Demb JB & Singer JH. (2014). Adaptation to background light enables contrast coding at rod bipolar cell synapses. *Neuron* 81, 388–401. [PubMed: 24373883]

- Kepecs A & Fishell G. (2014). Interneuron cell types are fit to function. *Nature* 505, 318–326. [PubMed: 24429630]
- Kullmann DM, Ruiz A, Rusakov DM, Scott R, Semyanov A & Walker MC. (2005). Presynaptic, extrasynaptic and axonal GABAA receptors in the CNS: where and why? *Prog Biophys Mol Biol* 87, 33–46. [PubMed: 15471589]
- Mitchell SJ & Silver RA. (2003). Shunting inhibition modulates neuronal gain during synaptic excitation. *Neuron* 38, 433–445. [PubMed: 12741990]
- Nelson R & Kolb H. (1985). A17: a broad-field amacrine cell in the rod system of the cat retina. *J Neurophysiol* 54, 592–614. [PubMed: 4045539]
- Nikonov SS, Kholodenko R, Lem J & Pugh EN Jr., (2006). Physiological features of the S-and M-cone photoreceptors of wild-type mice from single-cell recordings. *J Gen Physiol* 127, 359–374. [PubMed: 16567464]
- Oesch NW & Diamond JS. (2011). Ribbon synapses compute temporal contrast and encode luminance in retinal rod bipolar cells. *Nat Neurosci* 14, 1555–1561. [PubMed: 22019730]
- Pan ZH & Lipton SA. (1995). Multiple GABA receptor subtypes mediate inhibition of calcium influx at rat retinal bipolar cell terminals. *J Neurosci* 15, 2668–2679. [PubMed: 7722621]
- Roux L & Buzsaki G. (2015). Tasks for inhibitory interneurons in intact brain circuits. *Neuropharmacology* 88, 10–23. [PubMed: 25239808]
- Sagdullaev BT, McCall MA & Lukasiewicz PD. (2006). Presynaptic inhibition modulates spillover, creating distinct dynamic response ranges of sensory output. *Neuron* 50, 923–935. [PubMed: 16772173]
- Shapley RM & Enroth-Cugell C. (1984). Visual adaptation and retinal gain controls. *Prog Ret Eye Res* 3, 263–346.
- Singer JH & Diamond JS. (2003). Sustained Ca²⁺ entry elicits transient postsynaptic currents at a retinal ribbon synapse. *J Neurosci* 23, 10923–10933. [PubMed: 14645488]
- Singer JH, Lassoova L, Vardi N & Diamond JS. (2004). Coordinated multivesicular release at a mammalian ribbon synapse. *Nat Neurosci* 7, 826–833. [PubMed: 15235608]
- Sterling P & Lampson LA. (1986). Molecular specificity of defined types of amacrine synapse in cat retina. *J Neurosci* 6, 1314–1324. [PubMed: 2423657]
- Veruki ML, Zhou Y, Castilho A, Morgans CW & Hartveit E. (2018). Extrasynaptic NMDA receptors on rod pathway amacrine cells: molecular composition, activation, and signaling. *J Neurosci*
- Völgyi B, Xin D & Bloomfield SA. (2002). Feedback inhibition in the inner plexiform layer underlies the surround-mediated responses of AII amacrine cells in the mammalian retina. *J Physiol* 539, 603–614. [PubMed: 11882691]
- von Gersdorff H & Matthews G. (1997). Depletion and replenishment of vesicle pools at a ribbon-type synaptic terminal. *J Neurosci* 17, 1919–1927. [PubMed: 9045721]
- Weber E (1834). *De Pulsu, resorptione, auditu et tactu. Annotationes anatomicae et physiologicae* Koehler CF, Leipzig.
- Whittle P (1994). *The Psychophysics of Contrast Brightness*. In *Lightness, Brightness, and Transparency*, ed. Gilchrist A Lawrence Erlbaum Associates, Hillsdale, NJ.
- Wilson NR, Runyan CA, Wang FL & Sur M. (2012). Division and subtraction by distinct cortical inhibitory networks in vivo. *Nature* 488, 343–348. [PubMed: 22878717]
- Zhang J, Li W, Trexler EB & Massey SC. (2002). Confocal analysis of reciprocal feedback at rod bipolar terminals in the rabbit retina. *J Neurosci* 22, 10871–10882. [PubMed: 12486181]

Figure 1 |. Logistic functions determine response gain

A, Theoretical intensity response functions plotted over the working range of luminance signaling in RBCs. The change in response level from ϕ_1 to ϕ_2 is shown on the y-axis for a perfectly linear-log function ($y = (m - \log(\phi)) + b$; black) and for a nonlinear-log Hill function (gray). B, Plot of contrast gain over the range of ϕ_1 values predicted from the difference in response levels determined by the two curves shown in panel A.

Figure 2 | Inhibition alters the relationship between ϕ_1 and RRP size

A, *Top*, intensity of the light stimulus over time for a family of light stimuli to measure RRP depletion. *Bottom*, average responses across 10 cells for control conditions (black traces) and in the presences of 0.5 μM TTX, 3 μM Strychnine, 5 μM SR95531, and 50 μM TPMPA (gray traces) to block all inhibition. Traces are presented in order of increasing ϕ_1 intensity from the top down. Dashed box denotes time window for calculating charge transfer. **B**, Average charge transfer of the transient peak in response to ϕ_1 (red) and ϕ_2 (blue), plotted against ϕ_1 and normalized to the ϕ_1 step to 72 $r^*/\text{rod/sec}$, for control responses. Dashed lines are the Hill equation fits to the data. Black circles show the sum of the responses to ϕ_1 and ϕ_2 at each ϕ_1 . Error bars indicate standard deviation. **C**, Average charge transfer of the transient peak in response to ϕ_1 (red) and ϕ_2 (blue), normalized to the ϕ_1 step to 72 $r^*/\text{rod/sec}$, during inhibitory block. Solid lines show Hill equation fits to the data. Error bars

indicate standard deviation. D , Average (solid lines) and standard error (shading), for Hill fits to each cell in control conditions (black) and during inhibitory block (gray; $n=10$).

Figure 3 | Inhibition alters contrast gain across a range of ϕ_1

A, Top, intensity of the light stimulus over time for a family of light stimuli to measure contrast gain. Bottom, average responses across 11 cells for control conditions (black traces) and in the presence of 0.5 μM TTX, 3 μM strychnine, 5 μM SR95531, and 50 μM TPMPA (gray traces) to block all inhibition. *B*, Average charge transfer of the transient peak in response ϕ_2 in control (black symbols) and during inhibitory blockade (gray symbols) plotted against the ϕ_1 and normalized to the maximal response in control. Solid lines show sigmoid fits for qualitative comparisons between the two distributions. Dashed line plots the actual Weber contrast between ϕ_1 and ϕ_2 for each pair, plotted against ϕ_1 and scaled for comparison to transient EPSCs. Error bars indicate standard deviation. Asterisks indicate p values that reached significance following Bonferroni correction (i.e., $\alpha=0.01$; **: $p<0.01$, ***: $p<0.001$).

Figure 4 | Lateral inhibition does not influence ϕ_1 -RRP relationship or contrast gain

A, Top, intensity of the light stimulus over time for a family of light stimuli to measure RRP depletion. Bottom, average responses across 6 cells for control conditions (black traces) and in the presence of 0.5 μ M TTX and 3 μ M strychnine (gray traces) to block lateral inhibition. Traces are presented in order of increasing ϕ_1 intensity from the top down. **B**, Average charge transfer of the transient peak in response to ϕ_2 , for control responses (black symbols), and during block of lateral inhibition with 0.5 μ M TTX and 3 μ M strychnine (gray symbols) plotted against ϕ_1 and normalized to the ϕ_1 step to 72 $r^*/rod/sec$ in control. Error bars indicate standard deviation. Solid lines are Hill fits to the data. (n=6). **C**, Top, intensity of the light stimulus over time for a family of light stimuli to measure contrast gain. Bottom, average responses across 5 cells for control conditions (black traces) and in the presence of 0.5 μ M TTX and 3 μ M strychnine (gray traces) to block all inhibition. **D**, Average charge transfer of the transient peak in response to ϕ_1 , for control responses (black symbols), and

during block of lateral inhibition with 0.5 μM TTX and 3 μM strychnine (gray symbols) plotted against ϕ_1 and normalized to the maximal response. Error bars indicate standard deviation (n=5). Solid lines show sigmoid fits for qualitative comparisons between the two distributions.

Figure 5 | GABA_C receptors are responsible for shaping contrast gain

A, Top, intensity of the light stimulus over time for a family of light stimuli to measure RRP depletion. Bottom, average responses across 12 cells for control conditions (0.5 μM TTX, 3 μM strychnine; black traces) and following addition of 50 μM TPMPA (gray traces) to block GABA_C receptors. **B**, Average charge transfer of the transient peak in response ϕ_2 in control (black symbols) and during GABA_C blockade (gray symbols), plotted against the ϕ_1 and normalized to the maximal response in control. Error bars indicate standard deviation ($n=12$). **C**, Top, intensity of the light stimulus over time for a family of light stimuli to measure contrast gain. Bottom, average responses across 12 cells for control conditions (0.5 μM TTX, 3 μM strychnine; black traces) with the addition of 50 μM TPMPA (gray traces) to block GABA_C receptors. **D**, Average charge transfer of the transient peak in response ϕ_2 in control (black symbols) and during GABA_C blockade (gray symbols), plotted against the ϕ_1 and normalized to the maximal response in control. Error bars indicate standard deviation

(n=12). Asterisks indicate p values that reached significance following Bonferroni correction (i.e., $\alpha=0.01$; **: $p<0.01$). Solid lines show sigmoid fits for qualitative comparisons between the two distributions.