Published in final edited form as:

Pharmacol Biochem Behav. 2019 April ; 179: 55–62. doi:10.1016/j.pbb.2019.02.003.

Evaluation of SYA16263 as a new potential antipsychotic agent without catalepsy.

Barbara A. Bricker^a, Kwame Peprah^a, Hye J. Kang^{b,c}, and S. Y. Ablordeppey^{a,*}

^aDivision of Basic Pharmaceutical Sciences, Florida A&M University, College of Pharmacy and Pharmaceutical Sciences, Tallahassee, FL 32307, USA

^bDepartment of Pharmacology, School of Medicine, University of North Carolina at Chapel Hill, Chapel Hill, NC 27599-7365, USA

^cNational Institute of Mental Health Psychoactive Drug Screening Program (NIMH PDSP), School of Medicine, University of North Carolina at Chapel Hill, Chapel Hill, NC 27599-7365, USA

Abstract

SYA16263 exhibited moderate radioligand binding affinity at the D₂ receptor and produced inhibition of apomorphine-induced climbing behavior in mice with an ED₅₀ value of 3.88 mg/kg IP, predicting potential antipsychotic effects in humans.

Analysis of plasma and brains from rats injected IP with SYA16263 over the course of 24 hours revealed a log [brain]/[plasma] (log BB) at C_{max} observed equal to 1.08, indicating that SYA16263 enters the brain and is predicted to cross the blood brain barrier (BBB) readily.

When tested in animal behavior tests for catalepsy, SYA16263 did not produce catalepsy at doses up to 19 times the apomorphine ED₅₀ value predicting little or no extra-pyramidal (EPS) side effects in humans. This is similar to aripiprazole, which is associated with a low incidence of EPS in humans, but unlike haloperidol which is known to cause severe EPS in humans.

Functional activities for SYA16263 show that it acts as a D₂ agonist at both the Gi and β-arrestin pathways, similar to, but better than aripiprazole, which could account for the absence of the catalepsy observed.

Taken together, the receptor binding profile, the functional status, the animal behavioral tests and the log BB value, all provide evidence for further pre-clinical testing of SYA16263 as a potential antipsychotic agent with an interesting profile and a unique mechanism of action resulting in no EPS even up to 19 times the ED₅₀ value.

*Corresponding author.

Author Information

BAB contributed to experimental design, collected, analyzed and interpreted the in vivo studies data and the writing of the manuscript. K.P. synthesized the compounds tested in this manuscript. H.D.K. contributed to the functional assays under the supervision of the Director of the NIMH PDSP, Prof Bryan L Roth. SYA originated the ideas expressed in the manuscript, contributed to the experimental design, analyzed and interpreted the data, wrote and edited the manuscript.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Graphical Abstract

Keywords

Antipsychotic; D₂ receptor; β -arrestin pathway; Catalepsy; Aripiprazole; Apomorphine-induced climbing

1. Introduction

Schizophrenia affects about 1% of the world's population (Saha et al., 2005) and here in the US, the estimate is 1.1% (NIMH, 2017). The symptoms of schizophrenia are three-fold: a) positive b) negative and c) cognitive symptoms. The positive symptoms often involve hallucinations, delusions, thought disorders and movement disorders. The negative symptoms include disruptions to normal emotions and behaviors, poverty of speech and a lack of the ability to plan and sustain activities. Cognitive symptoms are subtle and may involve trouble focusing, poor executive functioning and problems with working memory. Schizophrenia is also associated with suicidal ideation and 9-13% of patients eventually take their own lives (Inskip et al., 1998; Meltzer, 1999). While estimates vary, up to 1/3 of patients are treatment resistant (TRS) (Essock et al., 1996; Lehman et al., 2004) and current drugs are either ineffective or they produce serious adverse side effects which lead to patient non-compliance. These findings underscore the need to seek novel treatment options for patients with schizophrenia.

First generation antipsychotic (FGA) drugs such as haloperidol presumably act primarily by blocking dopamine D₂ receptors (D₂RS) (Besnard et al., 2012; Kapur and Mamo, 2003; Wang, 2018). The efficacy of the FGAs is however hampered by their capacity to induce Parkinsonism-like side effects that become very debilitating with long-term use (Correll et al., 2004; Girgis et al., 2011; Tapp, 2008). The second generation antipsychotic (SGA) drugs are efficacious against the positive symptoms and may also attenuate some of the negative symptoms (Boter et al., 2009; Tandon et al., 2008). In addition to their interaction at the D₂R, each interacts with additional receptors including 5HT_{1A}, 5HT_{2A}, 5HT₇, alpha (α), and H₁ receptors which may be responsible for the subtle differences in their clinical profiles. While the EPS side effects of the SGAs are somehow muted, they produce weight gain and metabolic disorders which limit their widespread utility (Lieberman et al., 2005), (Boehm et al., 2004; Buckley et al., 2005; Wong et al., 2010).

Many CNS disorders are treated with therapeutic agents that directly or indirectly target G protein-coupled receptors (GPCR). In the canonical or conventional model of GPCR signaling, trimeric G protein signaling is initiated at the plasma membrane and is terminated by β -arrestins, which prevent coupling between receptors and G-proteins, and promote receptor internalization (Watari et al., 2014). The internalization process contributes further to the mechanism of G-protein signal desensitization by promoting receptor lysosomal degradation (Delom and Fessart, 2011; Mills, 2007). Studies using biochemical, pharmacological, biophysical, and mice studies have now revealed that several GPCRs, including the D_2 receptor subtype (D_2R), do not always follow this conventional signaling and desensitization paradigm. In fact, it is now widely accepted that GPCRs can signal not only through G protein activation but also through β -arrestins to scaffold intracellular signaling molecules with distinct pharmacological properties that could result in favorable therapies. Work by Caron and others (Urs et al., 2014; Urs et al., 2012) has now provided additional validation of the physiological role of this signaling pathway, using genetic approaches such as the deletion of GSK3 β from D_2 receptor-expressing neurons, to study the effects of antipsychotics downstream of the D_2 receptor, confirming the role of the β -arrestin pathway. Thus, agonists and antagonists that selectively discriminate between these signaling mechanisms, through the stabilization of an associated GPCR conformation, might become novel therapeutic tools that can lead to obtaining compounds with reduced side effects.

It has previously been shown that the D_2R is the main target of clinically effective antipsychotics and D_2R engages an Akt/GSK3 signaling pathway via the scaffolding of a β -arrestin2/Akt/PP2A/GSK3 complex (Beaulieu et al., 2005). Therefore, a strategy to activate β -arrestin2 selectively and spare the G-protein dependent signaling has been proposed and validated as a novel approach to the development of new antipsychotic agents. There is also now a preponderance of evidence in rodent models indicating that activation of 5HT $_{1A}R$ produces several effects including prevention of extrapyramidal symptoms (EPS) induced by dopamine D_2R blockade, enhancement of dopaminergic neurotransmission in the frontal cortex and inhibition of NMDA receptor antagonist-induced cognitive and social interaction deficits (Newman-Tancredi, 2010). Furthermore, blocking the 5HT $_7R$ is suggested to improve cognitive impairments in animals (Bonaventure et al., 2011). Lurasidone, a more recently introduced antipsychotic drug with favorable cognitive outcomes, activates 5-HT $_{1A}R$ and blocks 5-HT $_7R$ (Huang et al., 2012). Therefore, we have hypothesized that an antipsychotic agent which activates β -arrestinergic signaling through the D_2R , activates 5HT $_{1A}R$ [$K_i < 10$ nM] and blocks 5HT $_7R$ [$K_i < 10$ nM] should constitute an effective antipsychotic agent. It is equally important that the compounds avoid H_1R , 5HT $_{2C}R$, 5HT $_{2B}R$ and not activate 5HT $_{2A}R$. Weight gain and the metabolic syndrome are associated with binding to H_1R and 5HT $_{2C}R$ (Kroeze et al., 2003; Matsui-Sakata et al., 2005; Opgen-Rhein et al., 2010) while induction of valvular heart disease is associated with activating the 5HT $_{2B}R$ (Rothman et al., 2000; Setola et al., 2003; Setola and Roth, 2005). While often targeted in antipsychotic drug development, activating the 5HT $_{2A}R$ could produce hallucinogenic effects (Glennon et al., 1982; Glennon et al., 1980; Rasmussen et al., 1986; Smith et al., 1998).

The present study investigates the potential of SYA16263 to serve as an antipsychotic agent using the apomorphine-induced climbing behavior test in mice. The results are then compared to another potential antipsychotic, SYA013, which does not activate the β -arrestin pathway at the D₂R, the typical antipsychotic haloperidol, and the β -arrestin pathway activator aripiprazole. In addition, the potential of these compounds to induce catalepsy in rats, an animal behavioral test with predictive ability for possible extra-pyramidal side effects in humans, was investigated, and the results compared and discussed in light of their CNS receptor binding profiles.

2. Materials and methods

2.1. Animals

Reversal of apomorphine-induced climbing behavior experiments were performed using male, albino, Swiss-Webster mice (21 - 29 g), (5-7 weeks old). Catalepsy experiments were carried out on male Sprague-Dawley rats (107-168 g), (5-6 weeks old). All animals were from Harlan Laboratories, Inc. Animals were housed in the Florida A & M University Animal Care facility which is fully AAALAC accredited, and operates with a 12 h light/dark cycle and controlled temperature (24±2 °C). Animals were given free access to food and water and at least 5 days to adjust before experiments were begun. Animals were fasted the night before each experiment. All experimental procedures were performed in accordance with protocols approved by the Florida A & M University Institutional Animal Care and Use Committee.

2.2. Drugs and chemicals

All chemicals were purchased from Sigma-Aldrich (St. Louis, MO, USA) unless otherwise stated. SYA16263 (CLog P = 4.36) and SYA013 (CLog P = 5.15) were synthesized and characterized by us (Ablordepey et al., 2008; Peprah et al., 2012) at Florida A & M University with CHN values within 0.4% of theoretical values. The compounds were dissolved in filtered (0.22 μ) 1% lactic acid vehicle for all of the animal studies. Haloperidol and apomorphine hydrochloride hemihydrate (APO), were obtained from Sigma-Aldrich. Haloperidol and aripiprazole were dissolved in filtered (0.22 μ) 1% lactic acid. APO was dissolved in HPLC grade water, followed by ascorbic acid (0.1% w/w) dissolution followed by sodium chloride (0.9% w/w) on the morning of the experiments in an amber vial. Lactic acid (ACROS), ascorbic acid, phosphate buffered saline (PBS) (Fisher) and sodium chloride (Fisher) were ACS reagent grade). The d-amphetamine hemisulfate salt was from Sigma-Aldrich, and the water used to make solutions was HPLC grade. Diethyl ether was from Fluka, residue analysis grade. The acetic acid, water, methanol, and acetonitrile were HPLC grade. The sodium sulfate was analytical reagent grade. Doses are expressed as the free base for all compounds except SYA013 which is the HC1 salt, and were given in a volume of 10 mL/kg by intraperitoneal (ip) injection, except for apomorphine, which was given by subcutaneous (sc) injection. The HPLC internal standard DS-49 (CLog P = 5.17) used for calibration was also synthesized at Florida A & M University (Sikazwe et al., 2004) with CHN values within 0.4% of theoretical values as determined by CHN analysis.

The CLog P values in this report were calculated using ChemDraw Ultra, version 11.0.1 obtained from CambridgeSoft.

2.3. Animal behavioral experiments

All behavioral experiments were conducted under standard temperatures (68-79 °F) and humidity (30-70%) in the animal care facility.

2.3.1. Reversal of apomorphine-induced climbing—A modified climbing test (Needham et al., 1996; Protais et al., 1976) was used with Swiss-Webster mice to predict potential antipsychotic activity. Inhibition of the apomorphine-induced climbing indicates antipsychotic properties. Five mice per dose were injected first with haloperidol, SYA16263, or vehicle and returned to their home cage, then thirty minutes later injected with 1.5 mg/kg (as free base) apomorphine, and placed in cylindrical wire cages (12 cm diameter, 14 cm height) and observed for climbing behavior at 10 and 20 minutes after the apomorphine injection. Climbing behavior was assessed as follows: For all 4 paws up on the cage wall, not on the floor, a score of 2 was assigned (0% inhibition). For 3 or 2 paws on the cage wall, a score of 1 was assigned. For 1 or 0 paws on the cage wall, a score of 0 was assigned (100% inhibition). Scores were expressed as mean percent climbing inhibition, and plotted in Fig. 4. Prism 5.03, GraphPad software, Inc., non-linear regression software was used to calculate ED₅₀S.

2.3.2. Catalepsy bar test—To assess the potential for extrapyramidal side effects (EPS) in humans, the catalepsy bar test was used with rats (Kleven et al., 2005) utilizing a semi-automated instrument from Med Associates, Inc., St. Albans, VT (Bricker et al., 2014). We have used a second test of catalepsy that is reported to be sensitive to the anticataleptic actions of 5HT_{1A} receptor agonists, the crossed-legs position (CLP) test (Depoortere et al., 2007; Kleven et al., 2005). A righting test (McCreary et al., 2007; Reeve et al., 1992) was performed following the two catalepsy tests to check for sedation or other effects.

Rats were injected ip, then 60 minutes later, tested in the CLP test for 30 s, followed by the bar test for 30 s, followed by the righting test, then returned to their home cage. This was repeated at 3 and 6 min from the start of the first trial. Five rats were used for each dose. The mean of the 3 trials was used as the catalepsy response for each test for each rat (Depoortere et al., 2007; Kleven et al., 2005).

2.3.3. Catalepsy crossed-legs position (CLP) test—Rats were placed on the stainless steel floor of the catalepsy test chamber, the abdomen toward the floor, and the hind paws brought forward and the front paws backward so that the ipsilateral hind paws could hold onto the top of the front paws and the time (measured with a stopwatch) the rat stayed in this position with either one or both sets of paws was recorded up to 30 s (Depoortere et al., 2007; Kleven et al., 2005).

2.3.4. Righting test—Rats were placed gently on their back and observed immediately following the bar and CLP tests. If the rat did not stay in this position, and flipped over without assistance it was scored as “righted” (McCreary et al., 2007; Reeve et al., 1992).

2.4. Statistical analyses

All statistical analyses were performed using Prism 5.03, GraphPad Software Inc. All error bars are standard error of the mean (SEM). One-way ANOVA with Dunnett's post tests were used, with *, **, and *** indicating $p < 0.05$, 0.01, and 0.001 respectively compared with the vehicle group.

2.5. Receptor binding and GloSensor cAMP functional assays

Protocols for D_2 , and other receptor radioligand binding assays and GloSensor cAMP functional assays for G_i - or G_s -coupled GPCRs are available at the National Institute of Mental Health Psychoactive Drug Screening Program Web site, verified 12-7-18: <https://pdspdb.unc.edu/pdspWeb/content/PDSP%20Protocols%20II%202013-03-28.pdf>.

2.6. β -Arrestin Tango assay

Recruitment of β -arrestin to agonist stimulated D_2R or $5-HT_{1A}$ were performed using the "Tango"-type assay described in Barnea (Barnea et al., 2008), with modifications, to evaluate SYA16263 and SYA16264. Briefly, HTLA cells stably expressing β -arrestin-TEV protease and a tetracycline transactivator-driven luciferase were plated in 15-cm dishes in DMEM containing 10% FBS and transfected with D_2 or $5-HT_{1A}$ Tango construct (PMID25895059). The next day the cells were plated and challenged with the reference agonist or test ligand (dose-responses with 16 concentrations). After 18 h, the medium was removed and replaced with BriteGlo reagent (Promega), and luminescence per well was read using a TriLux plate reader. Data were normalized to vehicle (0%) and quipirole (100%) controls and regressed using the sigmoidal dose-response function built into GraphPad Prism 5.0. The results are reported in Table 2.

2.7. Pharmacokinetic analyses

2.7.1. Sample preparation for HPLC-PDA—Plasma and brain tissues were stored at -20°C until ready for extraction, and then prepared and extracted as described in (Sampson et al., 2014; Shimokawa et al., 2005). Briefly, 0.5 mL rat plasma plus internal standard was made basic with 100 μL 2M NaOH and extracted with diethyl ether. Supernatant was dried under nitrogen and then reconstituted in 200 μL of mobile phase. Centrifugation at 10,000 g afforded a supernatant for HPLC injection. Frozen rat brains were homogenized with 2 parts phosphate buffered saline (PBS), sonified, and internal standard added to 1 mL (~1 gram) of homogenate which was then extracted with acetonitrile, centrifuged, and the supernatant evaporated under nitrogen. Residue was mixed with water and sodium hydroxide, extracted with diethyl ether, dried under nitrogen, and reconstituted with 200 μL of mobile phase. After centrifugation at 10,000 g, the extract was injected into the HPLC.

2.7.2. HPLC conditions—As reported in (Sampson et al., 2014; Shimokawa et al., 2005), reversed phase HPLC with PDA detection at 254 nm was used to analyze the extracts. The mobile phase was 55% methanol 45% (0.01M sodium sulfate in 4% v/v acetic acid) at a flowrate of 1 mL/min, and run time of 15 minutes. Retention time and PDA spectra were used to identify analytes.

2.7.3. Determination of SYA16263 levels in plasma and brain samples—A single bolus dose of 50 mg/kg SYA16263 ($12.9 \times \text{ED}_{50}$) was given by IP injection to an average of 4 rats per time point group, and blood and brains collected at 5, 10, 15 and 30 min, 1hr, 2hr, 3hr, 6hr, 9hr, and 24 hr. For control purposes, vehicle was injected into 4 rats as well. The internal standard method was used to correct for % recoveries. Calibration lines were calculated using linear regression from 7-8 points, and had coefficients of determination of 0.9988 or better.

3. Results and Discussion

3.1. Biochemistry studies

3.1.1. Evaluation of chemical stability in plasma—The chemical stability of SYA16263 in rat plasma incubated at 37°C was evaluated prior to the in vivo studies in order to ascertain that the drug remained intact during the experiments.

HPLC analysis indicated no change in concentration over a 24 hour period for a 0.71 mg/mL preparation. One-way ANOVA indicated no significant difference [$F(2,6)=0.4361$; $p>0.05$] between concentration means at 0, 2 and 24h time points (Fig. 1).

3.1.2. Brain uptake of SYA16263: Plasma and brain concentrations of SYA16263— $\text{Log} [\text{brain}]/[\text{plasma}]$ (Log BB) at C_{max} is a predictor of the degree of drug incorporation into the brain (Palenicek et al., 2011) and a value > 0.3 predicts compounds will readily cross the blood brain barrier (BBB) (Olsen et al., 2008; Vilar et al., 2010). To evaluate the brain permeability of SYA16263, a time profile of the plasma and brain concentrations was conducted after IV injections of 50.0 mg/kg using previous methods (Bricker et al., 2012). As shown in Fig. 2, SYA16263 enters the brain quickly with $t_{\text{max}} = 10$ min. In addition, the brain levels significantly exceeded those in the plasma with $\text{log BB} = 1.08$ (log BB of haloperidol is 1.34) indicating a high degree of brain permeability.

3.2. Neurochemistry studies I of the CNS: Radioligand receptor binding studies

3.2.1. Prediction and comparison of the potential of compounds to exhibit antipsychotic properties in animal behavioral studies: Binding of analogs at CNS receptors—We have previously synthesized SYA16263 and several haloperidol analogs and screened them at CNS receptors including D_2R , $5\text{HT}_{1\text{A}}$, $5\text{HT}_{2\text{A}}$, $5\text{HT}_{2\text{C}}$, 5-HT_7 and H_1 receptors (Peprah et al., 2012) (Table 1) due to their relevance in antipsychotic drug discovery. The structures of key compounds discussed in this article are presented in Fig. 3. The compounds were found to bind to the D_2R with moderate affinity and with varying affinities at the other receptors. Of particular note is the high affinity of SYA16263 at the $5\text{HT}_{1\text{A}}\text{R}$ which may suggest a potential to overcome EPS and to contribute to the treatment of the negative and cognitive symptoms of schizophrenia (Bardin et al., 2006; Broekkamp et al., 1988; Christoffersen and Meltzer, 1998; Diaz-Mataix et al., 2005; Erhart et al., 2006; Goff et al., 1991; Ichikawa and Meltzer, 2000; Invernizzi et al., 1988; Kapur and Remington, 1996; Kleven et al., 2005; Leucht et al., 1999; Lieberman et al., 1989; Moss et al., 1993; Rollema et al., 2000; Rollema et al., 1997; Sprouse et al., 1999). In addition to its affinity to D_2R and $5\text{HT}_{1\text{A}}\text{R}$, SYA16263 also binds with high affinity to the D_4R , and moderately to

D₃R and 5HT₇R and does not bind significantly to 5HT_{2C}R or M₁R-M₅R. These observations are noteworthy since these latter two receptors are associated with weight gain (Kroeze et al., 2003; Matsui-Sakata et al., 2005; Opgen-Rhein et al., 2010) and cognitive deficits (Anagnostaras et al., 2003; Seeger et al., 2004) respectively. SYA16263 also binds about 100-fold more poorly to H₁R compared to clozapine, suggesting a lower propensity to induce sedation and/or weight gain.

3.3. Behavioral studies

3.3.1. Evaluation of haloperidol, SYA16263 and SYA013 in an in vivo mouse model of schizophrenia, reversal of apomorphine-induced climbing—Based on its binding profile in comparison to clozapine (Table 1), SYA16263 was selected for evaluation in a mouse model of the positive symptoms of schizophrenia (Needham et al., 1996; Protais et al., 1976) and the results are reported in Fig. 4. SYA16263 exhibited dose dependent antagonism to apomorphine climbing behavior in mice, with higher doses producing greater inhibition [$F(6, 28)=24.53$; $p<0.0001$]. The results were also compared to those of haloperidol [$F(6, 28)=13.09$; $p<0.0001$] and SYA013 [$F(7, 42)=6.495$; $p<0.0001$] which were previously reported by our lab (Ablordeppey et al., 2008; Sampson et al., 2014). The results suggest that SYA16263 is predicted to produce antipsychotic-like effect as expected, by its inhibition of APO-induced climbing behavior ($ED_{50}=3.88$ mg/kg, 95% CI 3.03–4.97, $n=5$ mice per dose).

3.3.2. Evaluation of SYA16263, haloperidol, SYA013, and aripiprazole for induction of catalepsy in rats—Most FGAs produce undesirable EPS in humans which manifests as catalepsy in animals. This occurs in addition to their antipsychotic actions, presumably as a result of blockade of D₂R. However, compounds such as aripiprazole do not produce significant catalepsy, presumably as a result of their activation rather than blockade of the D₂R (Bardin et al., 2006; Broekkamp et al., 1988; Christoffersen and Meltzer, 1998; Goff et al., 1991; Invernizzi et al., 1988; Kleven et al., 2005; Liebman et al., 1989; Moss et al., 1993; Rollema et al., 2000; Rollema et al., 1997).

Thus, an evaluation of the potential of an agent to induce catalepsy in animals is a required step in the development of new antipsychotic agents. When SYA16263 was evaluated in rats for catalepsy induction, there was an absence of catalepsy (< 20 seconds on bar) even up to 19 times its ED_{50} value (75 mg/kg), suggesting it may have a unique mechanism of action. The results are shown in Fig. 5 for SYA16263 [$F(6, 32)=2.736$; $p<0.05$] along with the standards, haloperidol [$F(6, 45)=66.37$; $p<0.0001$], SYA013 [$F(3, 49)=0.9125$; $p>0.05$], and aripiprazole [$F(2, 29)=21.09$; $p<0.0001$]. All animals righted themselves following each test.

We have also evaluated the potential of SYA 16263 to induce catalepsy in the CLP catalepsy test to confirm the absence of catalepsy. As shown in Fig. 6, while catalepsy produced by haloperidol [$F(4, 20)=32.25$; $p<0.0001$] and aripiprazole [$F(2, 12)=11.30$; $p<0.01$] showed a dose dependent trend of catalepsy in this test, SYA 16263 failed to elicit any signs of catalepsy up to 19 times its ED_{50} value. All animals righted themselves following each test. It is worthy to note that the aripiprazole catalepsy result is consistent with a meta-analysis

(Bernagie et al., 2016) indicating a mean EPS incidence of 17.1 % (95 % CI 0.128–0.223) in pediatric and adolescent patients.

3.4. Neurochemistry studies II of the CNS

3.4.1. Evaluation of functional selectivity of SYA16263 and SYA013 compared to aripiprazole—To ascertain whether SYA16263 acts as an agonist or antagonist at the D₂R, a functional assay was carried out on SYA16263 and SYA013 with aripiprazole as the positive control given that functionally selective ligands promoting β -arrestin recruitment to D₂R may produce antipsychotic-like properties without catalepsy (Allen et al., 2011; Chen et al., 2012). Using the D₂R β -arrestin Tango Assay (Barnea et al., 2008; Kroeze et al., 2015) the effects of the compounds on signaling were obtained and are reported in Tables 2 and 3.

3.4.2. β -Arrestin Tango assay at D₂R and absence of catalepsy—Recruitment of β -arrestin-2 to agonist stimulated D₂R was performed using a “Tango” assay (Barnea et al., 2008; Kroeze et al., 2015) to evaluate SYA16263 and SYA013. The results show that SYA16263 acts as an agonist at both the G protein dependent and independent β -arrestin signaling pathways. Interestingly, its functional profile at the D₂R mimics that of aripiprazole albeit superior in its efficacy. For example, while it has only a moderate binding affinity to D₂R [K_i = 124 nM], SYA16263 is less potent but more efficacious as an agonist [EC_{50} = 5.7 nM; E_{max} = 64.4%] than aripiprazole [EC_{50} = 1.0 nM; E_{max} = 51.0%] in activating G_i pathway at the D₂R. Also, while less potent, it is more efficacious in promoting β -arrestin-2 recruitment to D₂R, [EC_{50} = 10.5 nM; E_{max} = 92%] than aripiprazole [EC_{50} = 4.0 nM; E_{max} = 62%] (Table 2). These results are consistent with the absence of catalepsy observed with SYA16263. On the other hand, SYA013 neither showed significant agonist or antagonist properties at the D₂R and thus, the results have not provided an explanation as to why SYA013 produces antipsychotic-like activity and has limited catalepsy. Further experiments to evaluate the functional behavior of these compounds at other D₂-like and 5-HT receptors are planned to elucidate the basis of these observations.

3.4.3. Functional selectivity at the 5-HT_{1A} receptor—The results of functional evaluation of SYA16263 at 5-HT_{1A}R showed that it does not activate G protein coupled 5HT_{1A}R signaling [EC_{50} = 0; E_{max} = -23.7] and instead, shows extreme bias in β -arrestin recruitment to the 5HT_{1A}R [IC_{50} = 149 nM; E_{max} = 54.4%, Table 3]. The implication of this unique signaling is under further investigation.

CONCLUSION

In summary, we have identified 1-(4-(4-fluorophenyl)butyl)-4-(pyridin-2-yl)piperazine or SYA16263 as a D₂R ligand with an ED₅₀ value of 3.88 mg/kg, in reversing APO-induced climbing behavior in mice. Evaluation of catalepsy induction indicated that SYA16263 does not produce catalepsy even at greater than 19 times its ED₅₀ value. Functional analysis revealed SYA16263 activates both G-protein and β -arrestin signaling at the D₂R, thus suggesting that SYA16263 has a signaling profile similar to that of aripiprazole at the D₂R. However, SYA16263 binds with a higher affinity at the 5HT_{1A}R than at the D₂R and

functional evaluation at the 5-HT_{1A}R demonstrated SYA16263 selectively activates β -arrestin recruitment to the 5HT_{1A}R and spares any significant interaction at the G-Protein dependent signaling pathway. The studies herein support this unique mechanism of action, making SYA16263 a potential antipsychotic to be further studied.

Acknowledgements

The authors gratefully acknowledge the financial support of the National Institute of General Medical Studies (NIGMS) grant #5SC1GM116724-03. We also acknowledge the financial support from Anxiolytech toward the ongoing in vivo studies. We thank the NIMH Psychoactive Drug Screening Program for carrying out the screening and functional evaluations reported herein. This work was supported in part by the Pharmaceutical Research Center NIH/NCRR 1 C06-RR12512-01 Grant.

References

- Ablordeppey SY, Altundas R, Bricker B, Zhu XY, Kumar EV, Jackson T, Khan A, Roth BL, 2008 Identification of a butyrophenone analog as a potential atypical antipsychotic agent: 4-[4-(4-chlorophenyl)-1,4-diazepan-1-yl]-1-(4-fluorophenyl)butan-1-one. *Bioorg. Med. Chem.* 16(15), 7291–7301. [PubMed: 18595716]
- Allen JA, Yost JM, Setola V, Chen X, Sassano MF, Chen M, Peterson S, Yadav PN, Huang XP, Feng B, Jensen NH, Che X, Bai X, Frye SV, Wetsel WC, Caron MG, Javitch JA, Roth BL, Jin J, 2011 Discovery of beta-arrestin-biased dopamine D2 ligands for probing signal transduction pathways essential for antipsychotic efficacy. *Proc. Natl. Acad. Sci. U. S. A* 108(45), 18488–18493. [PubMed: 22025698]
- Anagnostaras SG, Murphy GG, Hamilton SE, Mitchell SL, Rahnama NP, Nathanson NM, Silva AJ, 2003 Selective cognitive dysfunction in acetylcholine M1 muscarinic receptor mutant mice. *Nat. Neurosci* 6(1), 51–58. [PubMed: 12483218]
- Bardin L, Kleven MS, Barret-Grevoz C, Depoortere R, Newman-Tancredi A, 2006 Antipsychotic-like vs cataleptogenic actions in mice of novel antipsychotics having D2 antagonist and 5-HT_{1A} agonist properties. *Neuropsychopharmacology* 31(9), 1869–1879. [PubMed: 16237379]
- Barnea G, Strapps W, Herrada G, Berman Y, Ong J, Kloss B, Axel R, Lee KJ, 2008 The genetic design of signaling cascades to record receptor activation. *Proc. Natl. Acad. Sci. U. S. A* 105(1), 64–69. [PubMed: 18165312]
- Beaulieu JM, Sotnikova TD, Marion S, Lefkowitz RJ, Gainetdinov RR, Caron MG, 2005 An Akt/beta-arrestin 2/PP2A signaling complex mediates dopaminergic neurotransmission and behavior. *Cell* 122(2), 261–273. [PubMed: 16051150]
- Besnard J, Ruda GF, Setola V, Abecassis K, Rodriguiz RM, Huang XP, Norval S, Sassano MF, Shin AI, Webster LA, Simeons FR, Stojanovski L, Prat A, Seidah NG, Constam DB, Bickerton GR, Read KD, Wetsel WC, Gilbert IH, Roth BL, Hopkins AL, 2012 Automated design of ligands to polypharmacological profiles. *Nature* 492(7428), 215–220. [PubMed: 23235874]
- Boehm G, Racoosin JA, Laughren TP, Katz R, 2004 Consensus development conference on antipsychotic drugs and obesity and diabetes: response to consensus statement. *Diabetes Care* 27(8), 2088–2089; author reply 2089–2090.
- Bonaventure P, Aluisio L, Shoblock J, Boggs JD, Fraser IC, Lord B, Lovenberg TW, Galici R, 2011 Pharmacological blockade of serotonin 5-HT(7) receptor reverses working memory deficits in rats by normalizing cortical glutamate neurotransmission. *PLoS One* 6(6), e20210. [PubMed: 21701689]
- Boter H, Peuskens J, Libiger J, Fleischhacker WW, Davidson M, Galderisi S, Kahn RS, group E.s., 2009 Effectiveness of antipsychotics in first-episode schizophrenia and schizophreniform disorder on response and remission: an open randomized clinical trial (EUFEST). *Schizophr. Res* 115(2–3), 97–103. [PubMed: 19819114]
- Bricker B, Jackson T, Boateng B, Zhu XY, Ablordeppey SY, 2012 Evaluation of the behavioral and pharmacokinetic profile of SYA013, a homopiperazine analog of haloperidol in rats. *Pharmacol. Biochem. Behav* 102(2), 294–301. [PubMed: 22588199]

- Bricker B, Sampson D, Ablordeppey SY, 2014 Evaluation of the potential of antipsychotic agents to induce catalepsy in rats: assessment of a new, commercially available, semi-automated instrument. *Pharmacol. Biochem. Behav* 120, 109–116. [PubMed: 24565832]
- Broekkamp CL, Oosterloo SK, Berendsen HH, van Delft AM, 1988 Effect of metergoline, fenfluramine, and 8-OHDPAT on catalepsy induced by haloperidol or morphine. *Naunyn Schmiedeberg Arch. Pharmacol.* 338(2), 191–195. [PubMed: 2972934]
- Buckley PF, Miller DD, Singer B, Arena J, Stirewalt EM, 2005 Clinicians' recognition of the metabolic adverse effects of antipsychotic medications. *Schizophr. Res* 79(2-3), 281–288. [PubMed: 15964743]
- Chen X, Sassano MF, Zheng L, Setola V, Chen M, Bai X, Frye SV, Wetsel WC, Roth BL, Jin J, 2012 Structure-functional selectivity relationship studies of beta-arrestin-biased dopamine D(2) receptor agonists. *J. Med. Chem* 55(16), 7141–7153. [PubMed: 22845053]
- Christoffersen CL, Meltzer LT, 1998 Reversal of haloperidol-induced extrapyramidal side effects in cebus monkeys by 8-hydroxy-2-(di-n-propylamino)tetralin and its enantiomers. *Neuropsychopharmacology* 18(5), 399–402. [PubMed: 9536454]
- Correll CU, Leucht S, Kane JM, 2004 Lower risk for tardive dyskinesia associated with second-generation antipsychotics: a systematic review of 1-year studies. *Am. J. Psychiatry* 161(3), 414–425. [PubMed: 14992963]
- Delom F, Fessart D, 2011 Role of Phosphorylation in the Control of Clathrin-Mediated Internalization of GPCR. *Int. J. Cell Biol* 2011, 246954. [PubMed: 21765832]
- Depoortere R, Bardin L, Auclair AL, Kleven MS, Prinssen E, Colpaert F, Vacher B, Newman-Tancredi A, 2007 F15063, a compound with D2/D3 antagonist, 5-HT 1A agonist and D4 partial agonist properties. II. Activity in models of positive symptoms of schizophrenia. *Br. J. Pharmacol* 151(2), 253–265. [PubMed: 17375086]
- Diaz-Mataix L, Scorza MC, Bortolozzi A, Toth M, Celada P, Artigas F, 2005 Involvement of 5-HT1A receptors in prefrontal cortex in the modulation of dopaminergic activity: role in atypical antipsychotic action. *J. Neurosci* 25(47), 10831–10843. [PubMed: 16306396]
- Erhart SM, Marder SR, Carpenter WT, 2006 Treatment of schizophrenia negative symptoms: future prospects. *Schizophr. Bull* 32(2), 234–237. [PubMed: 16492797]
- Essock SM, Hargreaves WA, Dohm FA, Goethe J, Carver L, Hipshman L, 1996 Clozapine eligibility among state hospital patients. *Schizophr. Bull* 22(1), 15–25. [PubMed: 8685657]
- Girgis RR, Merrill DB, Vorel SR, Edward K, Kimberly P, You M, Pikalov A, Whitehead R, Lieberman JA, 2011 Aripiprazole versus haloperidol treatment in early-stage schizophrenia. *J. Psychiatr. Res* 45(6), 756–762. [PubMed: 20937506]
- Glennon RA, Young R, Benington F, Morin RD, 1982 Behavioral and serotonin receptor properties of 4-substituted derivatives of the hallucinogen 1-(2,5-dimethoxyphenyl)-2-aminopropane. *J. Med. Chem* 25(10), 1163–1168. [PubMed: 7143352]
- Glennon RA, Young R, Rosecrans JA, Kallman MJ, 1980 Hallucinogenic agents as discriminative stimuli: a correlation with serotonin receptor affinities. *Psychopharmacology (Berl.)* 68(2), 155–158. [PubMed: 6776558]
- Goff DC, Midha KK, Brotman AW, McCormick S, Waites M, Amico ET, 1991 An open trial of buspirone added to neuroleptics in schizophrenic patients. *J. Clin. Psychopharmacol* 11(3), 193–197. [PubMed: 2066458]
- Huang M, Horiguchi M, Felix AR, Meltzer HY, 2012 5-HT1A and 5-HT7 receptors contribute to lurasidone-induced dopamine efflux. *Neuroreport* 23(7), 436–440. [PubMed: 22415605]
- Ichikawa J, Meltzer HY, 2000 The effect of serotonin(1A) receptor agonism on antipsychotic drug-induced dopamine release in rat striatum and nucleus accumbens. *Brain Res.* 858(2), 252–263. [PubMed: 10708676]
- Inskip HM, Harris EC, Barraclough B, 1998 Lifetime risk of suicide for affective disorder, alcoholism and schizophrenia. *Br. J. Psychiatry* 172, 35–37. [PubMed: 9534829]
- Invernizzi RW, Cervo L, Samanin R, 1988 8-Hydroxy-2-(di-n-propylamino) tetralin, a selective serotonin1A receptor agonist, blocks haloperidol-induced catalepsy by an action on raphe nuclei medianus and dorsalis. *Neuropharmacology* 27(5), 515–518. [PubMed: 2969083]

- Kapur S, Mamo D, 2003 Half a century of antipsychotics and still a central role for dopamine D2 receptors. *Prog. Neuropsychopharmacol. Biol. Psychiatry* 27(7), 1081–1090. [PubMed: 14642968]
- Kapur S, Remington G, 1996 Serotonin-dopamine interaction and its relevance to schizophrenia. *Am. J. Psychiatry* 153(4), 466–476. [PubMed: 8599393]
- Kleven MS, Barret-Grevoz C, Bruins Slot L, Newman-Tancredi A, 2005 Novel antipsychotic agents with 5-HT(1A) agonist properties: role of 5-HT(1A) receptor activation in attenuation of catalepsy induction in rats. *Neuropharmacology* 49(2), 135–143.
- Kroeze WK, Hufeisen SJ, Popadak BA, Renock SM, Steinberg S, Ernsberger P, Jayathilake K, Meltzer HY, Roth BL, 2003 H1-histamine receptor affinity predicts short-term weight gain for typical and atypical antipsychotic drugs. *Neuropsychopharmacology* 28(3), 519–526. [PubMed: 12629531]
- Kroeze WK, Sassano MF, Huang XP, Lansu K, McCorvy JD, Giguere PM, Sciaky N, Roth BL, 2015 PRESTO-TANGO as an open-source resource for interrogation of the druggable human GPCRome. *Nat. Struct. Mol. Biol* 22(5), 362–369. [PubMed: 25895059]
- Lehman AF, Lieberman JA, Dixon LB, McGlashan TH, Miller AL, Perkins DO, Kreyenbuhl J, American Psychiatric A., Steering Committee on Practice, G., 2004. Practice guideline for the treatment of patients with schizophrenia, second edition. *Am. J. Psychiatry* 161(2 Suppl), 1–56.
- Leucht S, Pitschel-Walz G, Abraham D, Kissling W, 1999 Efficacy and extrapyramidal side-effects of the new antipsychotics olanzapine, quetiapine, risperidone, and sertindole compared to conventional antipsychotics and placebo. A meta-analysis of randomized controlled trials. *Schizophr. Res.* 35(1), 51–68. [PubMed: 9988841]
- Lieberman JA, Stroup TS, McEvoy JP, Swartz MS, Rosenheck RA, Perkins DO, Keefe RS, Davis SM, Davis CE, Lebowitz BD, Severe J, Hsiao JK, Clinical Antipsychotic Trials of Intervention Effectiveness, I., 2005 Effectiveness of antipsychotic drugs in patients with chronic schizophrenia. *N. Engl. J. Med* 353(12), 1209–1223. [PubMed: 16172203]
- Liebman JM, Gerhardt SC, Gerber R, 1989 Effects of 5-HT1A agonists and 5-HT2 antagonists on haloperidol-induced dyskinesias in squirrel monkeys: no evidence for reciprocal 5-HT-dopamine interaction. *Psychopharmacology (Berl.)* 97(4), 456–461. [PubMed: 2524857]
- Matsui-Sakata A, Ohtani H, Sawada Y, 2005 Receptor occupancy-based analysis of the contributions of various receptors to antipsychotics-induced weight gain and diabetes mellitus. *Drug Metab. Pharmacokinet.* 20(5), 368–378. [PubMed: 16272755]
- McCreary AC, Glennon JC, Ashby CR Jr., Meltzer HY, Li Z, Reinders JH, Hesselink MB, Long SK, Herremans AH, van Stuijvenberg H, Feenstra RW, Kruse CG, 2007 SLV313 (1-(2,3-dihydrobenzo[1,4]dioxin-5-yl)-4-[5-(4-fluoro-phenyl)-pyridin-3-ylmethyl]-piperazine monohydrochloride): a novel dopamine D2 receptor antagonist and 5-HT1A receptor agonist potential antipsychotic drug. *Neuropsychopharmacology* 32(1), 78–94. [PubMed: 16710314]
- Meltzer HY, 1999 Suicide and schizophrenia: clozapine and the InterSePT study. *International Clozaril/Leponex Suicide Prevention Trial. J. Clin. Psychiatry* 60 Suppl 12, 47–50.
- Mills IG, 2007 The interplay between clathrin-coated vesicles and cell signalling. *Semin. Cell Dev. Biol.* 18(4), 459–470. [PubMed: 17692542]
- Moss LE, Neppe VM, Drevets WC, 1993 Buspirone in the treatment of tardive dyskinesia. *J. Clin. Psychopharmacol.* 13(3), 204–209. [PubMed: 8102622]
- Needham PL, Atkinson J, Skill MJ, Heal DJ, 1996 Zotepine: preclinical tests predict antipsychotic efficacy and an atypical profile. *Psychopharmacol. Bull* 32(1), 123–128. [PubMed: 8927661]
- Newman-Tancredi A, 2010 The importance of 5-HT1A receptor agonism in antipsychotic drug action: rationale and perspectives. *Curr Opin Investig Drugs* 11(7), 802–812.
- Olsen CK, Brennum LT, Kreilgaard M, 2008 Using pharmacokinetic-pharmacodynamic modelling as a tool for prediction of therapeutic effective plasma levels of antipsychotics. *Eur. J. Pharmacol* 584(2-3), 318–327. [PubMed: 18325493]
- Opgen-Rhein C, Brandl EJ, Muller DJ, Neuhaus AH, Tiwari AK, Sander T, Dettling M, 2010 Association of HTR2C, but not LEP or INSIG2, genes with antipsychotic-induced weight gain in a German sample. *Pharmacogenomics* 11(6), 773–780. [PubMed: 20504252]
- Palenicek T, Balikova M, Rohanova M, Novak T, Horacek J, Fujakova M, Hoschl C, 2011 Behavioral, hyperthermic and pharmacokinetic profile of para-methoxymethamphetamine (PMMA) in rats. *Pharmacol. Biochem. Behav* 98(1), 130–139. [PubMed: 21167195]

- Peprah K, Zhu XY, Eyunni SV, Setola V, Roth BL, Ablordeppey SY, 2012 Multi-receptor drug design: Haloperidol as a scaffold for the design and synthesis of atypical antipsychotic agents. *Bioorg. Med. Chem* 20(3), 1291–1297. [PubMed: 22245230]
- Protais P, Costentin J, Schwartz JC, 1976 Climbing behavior induced by apomorphine in mice: a simple test for the study of dopamine receptors in striatum. *Psychopharmacology (Berl.)* 50(1), 1–6. [PubMed: 827755]
- Rasmussen K, Glennon RA, Aghajanian GK, 1986 Phenethylamine hallucinogens in the locus coeruleus: potency of action correlates with rank order of 5-HT₂ binding affinity. *Eur. J. Pharmacol* 132(1), 79–82. [PubMed: 3816969]
- Reeve B, Dingwall B, Darlington CL, Scott SJ, Sansom AJ, Smith PF, 1992 Simple device for quantifying drug effects on the righting reflex. *Pharmacol. Biochem. Behav* 42(1), 183–185. [PubMed: 1528940]
- Rollema H, Lu Y, Schmidt AW, Sprouse JS, Zorn SH, 2000 5-HT_{1A} receptor activation contributes to ziprasidone-induced dopamine release in the rat prefrontal cortex. *Biol. Psychiatry* 48(3), 229–237. [PubMed: 10924666]
- Rollema H, Lu Y, Schmidt AW, Zorn SH, 1997 Clozapine increases dopamine release in prefrontal cortex by 5-HT_{1A} receptor activation. *Eur. J. Pharmacol* 338(2), R3–5. [PubMed: 9456005]
- Rothman RB, Baumann MH, Savage JE, Rauser L, McBride A, Hufeisen SJ, Roth BL, 2000 Evidence for possible involvement of 5-HT_{2B} receptors in the cardiac valvulopathy associated with fenfluramine and other serotonergic medications. *Circulation* 102(23), 2836–2841. [PubMed: 11104741]
- Saha S, Chant D, Welham J, McGrath J, 2005 A systematic review of the prevalence of schizophrenia. *PLoS Med.* 2(5), e141. [PubMed: 15916472]
- Sampson D, Bricker B, Zhu XY, Peprah K, Lamango NS, Setola V, Roth BL, Ablordeppey SY, 2014 Further evaluation of the tropane analogs of haloperidol. *Bioorg. Med. Chem. Lett.* 24(17), 4294–4297. [PubMed: 25070422]
- Schmidt AW, Lebel LA, Howard HR Jr., Zorn SH, 2001 Ziprasidone: a novel antipsychotic agent with a unique human receptor binding profile. *Eur. J. Pharmacol* 425(3), 197–201. [PubMed: 11513838]
- Seeger T, Fedorova I, Zheng F, Miyakawa T, Koustova E, Gomeza J, Basile AS, Alzheimer C, Wess J, 2004 M₂ muscarinic acetylcholine receptor knock-out mice show deficits in behavioral flexibility, working memory, and hippocampal plasticity. *J. Neurosci* 24(45), 10117–10127. [PubMed: 15537882]
- Setola V, Hufeisen SJ, Grande-Allen KJ, Vesely I, Glennon RA, Blough B, Rothman RB, Roth BL, 2003 3,4-methylenedioxymethamphetamine (MDMA, “Ecstasy”) induces fenfluramine-like proliferative actions on human cardiac valvular interstitial cells in vitro. *Mol. Pharmacol* 63(6), 1223–1229. [PubMed: 12761331]
- Setola V, Roth BL, 2005 Screening the receptorome reveals molecular targets responsible for drug-induced side effects: focus on ‘fen-phen’. *Expert Opin. Drug Metab. Toxicol.* 1(3), 377–387. [PubMed: 16863450]
- Shapiro DA, Renock S, Arrington E, Chiodo LA, Liu LX, Sibley DR, Roth BL, Mailman R, 2003 Aripiprazole, a novel atypical antipsychotic drug with a unique and robust pharmacology. *Neuropsychopharmacology* 28(8), 1400–1411. [PubMed: 12784105]
- Shimokawa Y, Akiyama H, Kashiya E, Koga T, Miyamoto G, 2005 High performance liquid chromatographic methods for the determination of aripiprazole with ultraviolet detection in rat plasma and brain: application to the pharmacokinetic study. *J. Chromatogr. B Analyt. Technol. Biomed. Life Sci.* 821(1), 8–14.
- Sikazwe DM, Li S, Mardenborough L, Cody V, Roth BL, Ablordeppey SY, 2004 Haloperidol: towards further understanding of the structural contributions of its pharmacophoric elements at D₂-like receptors. *Bioorg. Med. Chem. Lett.* 14(23), 5739–5742. [PubMed: 15501032]
- Smith RL, Canton H, Barrett RJ, Sanders-Bush E, 1998 Agonist properties of N,N-dimethyltryptamine at serotonin 5-HT_{2A} and 5-HT_{2C} receptors. *Pharmacol. Biochem. Behav* 61(3), 323–330. [PubMed: 9768567]

- Sprouse JS, Reynolds LS, Braselton JP, Rollema H, Zorn SH, 1999 Comparison of the novel antipsychotic ziprasidone with clozapine and olanzapine: inhibition of dorsal raphe cell firing and the role of 5-HT_{1A} receptor activation. *Neuropsychopharmacology* 21(5), 622–631. [PubMed: 10516958]
- Tandon R, Belmaker RH, Gattaz WF, Lopez-Ibor JJ Jr., Okasha A, Singh B, Stein DJ, Olie JP, Fleischhacker WW, Moeller HJ, Section of Pharmacopsychiatry WPA, 2008 World Psychiatric Association Pharmacopsychiatry Section statement on comparative effectiveness of antipsychotics in the treatment of schizophrenia. *Schizophr. Res.* 100(1-3), 20–38. [PubMed: 18243663]
- Tapp A, Wood A, Kilzieh N, Martin L, Kennedy A, Raskind M, 2008 P.3.c.029 Incidence of tardive dyskinesia in patients receiving first or second generation antipsychotic therapy. *Eur. Neuropsychopharmacology* 18(4), S420.
- Urs NM, Nicholls PJ, Caron MG, 2014 Integrated approaches to understanding antipsychotic drug action at GPCRs. *Curr. Opin. Cell Biol.* 27, 56–62. [PubMed: 24680431]
- Urs NM, Snyder JC, Jacobsen JP, Peterson SM, Caron MG, 2012 Deletion of GSK3 β in D₂R-expressing neurons reveals distinct roles for beta-arrestin signaling in antipsychotic and lithium action. *Proc. Natl. Acad. Sci. U. S. A* 109(50), 20732–20737. [PubMed: 23188793]
- Vilar S, Chakrabarti M, Costanzi S, 2010 Prediction of passive blood-brain partitioning: straightforward and effective classification models based on in silico derived physicochemical descriptors. *J. Mol. Graph. Model* 28(8), 899–903. [PubMed: 20427217]
- Wang S, Che T, Levit A, Shoichet BK, Wacker D.k and Roth BL 2018 Structure of the D₂ dopamine receptor bound to the atypical antipsychotic drug risperidone. *Nature Accelerated Article Preview* 10.1038/nature25758 (2018).
- Watari K, Nakaya M, Kurose H, 2014 Multiple functions of G protein-coupled receptor kinases. *J. Mol. Signal* 9(1), 1–9. [PubMed: 24597858]
- Wong EH, Tarazi FI, Shahid M, 2010 The effectiveness of multi-target agents in schizophrenia and mood disorders: Relevance of receptor signature to clinical action. *Pharmacol. Ther* 126(2), 173–185. [PubMed: 20171983]

Web References

- (NIMH, 2017) https://www.nimh.nih.gov/health/statistics/prevalence/file_148221.pdf Last accessed 12-7-2018.
- 2). https://www.ema.europa.eu/documents/scientific-discussion/abilify-epar-scientific-discussion_en.pdf. Last accessed 1-11-2019.
- 3). <https://www.rxlist.com/clozaril-drug.htm#description>. Last accessed 1-11-2019.

Fig. 1.
Chemical stability of SYA16263 in plasma. No significant difference in means (one-way ANOVA), $n = 3$ was observed. Error bars are SEM.

Fig. 2.
Time profile of plasma and brain concentrations of SYA16263 in a rat, n=3-4 rats/group.
Error bars are SEM.

Figure 3.
Structures of key agents discussed in this article

Inhibition of Apomorphine-Induced Climbing Behavior

Fig. 4. SYA16263, haloperidol, and SYA013 inhibited APO-induced climbing behavior in mice. N=5 mice per dose. Error bars are SEM. One-way ANOVA with Dunnett's post tests. *, **, and *** indicate $p < 0.05$, 0.01 , and 0.001 respectively compared with the vehicle group

Fig. 5.

Haloperidol produced catalepsy in rats at clinical doses while SYA16263, SYA013, and aripiprazole did not (<20s on bar). In fact, no catalepsy was observed up to 75 mg/kg SYA16263, or at 19 X the ED₅₀ value of SYA16263. N=5 mice per dose. Error bars are SEM. One-way ANOVA with Dunnett's post tests. *, **, and *** indicate p<0.05, 0.01, and 0.001 respectively compared with the vehicle group.

Fig. 6.

Haloperidol produced catalepsy in the CLP test (>20 sec position maintained) in rats while aripiprazole and SYA16263 did not. A mean of zero seconds was observed for vehicle, 0.24 mg/kg haloperidol, and all doses of SYA16263 up to 19 times the ED₅₀ value. n= 5-9 rats/dose. Error bars are SEM. One-way ANOVA with Dunnett's post tests. *, **, and *** indicate p<0.05, 0.01, and 0.001 respectively compared with the vehicle group.

Table 1

*Binding affinities [Mean K_i (nM) n=3/4 experiments] of SYA16263, SYA013, and several standard antipsychotics at relevant CNS receptors.

Compound	5HT _{1A}	5HT _{2A}	5HT ₇	D ₁	D ₂	D ₃	D ₄	D ₅	5HT _{2C}	H ₁	M ₁
^a <i>Aripiprazole</i>	5.6	8.7	10.3	1960	3.3	9.7	510	2590	76	25.1	6780
^b <i>Haloperidol</i>	3600	120	1100	120	1.4	2.5	3.3	147 ^c	4700	440	1600
^b <i>Clozapine</i>	140	8.9	66	290	130	240	54.0	454 ^d	17.0	1.8	1.8
<i>SYA013</i>	117	23.6	78.0	163	43.3	159	6.6	NA	1425	189	872
<i>SYA16263</i>	1.1	50.0	90.0	238	124	86	3.5	1451	>10K	167	MPA

* All standard errors are within 20% of the mean values indicated. Each result represents the mean of at least two determinations, each in triplicate. Data for aripiprazole, haloperidol and clozapine were obtained from ^aref (Shapiro et al., 2003), ^bref (Schmidt et al., 2001), ^cref (Tyler et al., 2017), and ^dref (Web ref 3). MPA = Missed primary assay threshold of 50% inhibition at 10 μ M. NA = not available.

Table 2Radioligand binding affinity (K_i) and functional activities at D₂ receptor.

$(EC_{50}/IC_{50}, \text{ nM}; E_{\text{max}} \%)^a$					
CMPD	D ₂ Binding (K_i nM)	D ₂ Agonist (G_i -cAMP)	D ₂ Antagonist	D ₂ Tango Agonist	D ₂ Tango Antagonist
^b Aripiprazole	(3.3,3.9)	1.0; 51.0	0.0	4.0; 62	0.0
SYA16263	124	5.7; 64.4	0.0	10.5; 92	0.0; -3.8
SYA013	43.3	0.0;16.4	>10K; 49.6	0.0	ND

^aResults from NIMH PDSP.^bref (Chen et al., 2012; Shapiro et al., 2003). ND = Not determined. Quinpirole is the positive controls for D₂R. Tango assays are for β -arrestin-2 recruitment.

Table 3Radioligand binding (K_i , nM) and functional activities at 5-HT_{1A}R.

(EC ₅₀ /IC ₅₀ , nM; E _{max} %) ^a					
CMPD	5-HT _{1A} R Binding (K_i nM)	Agonist G _i -cAMP	Antagonist	Tango Agonist	Tango Antagonist
Methysergide	25.0	ND	ND	ND	ND
SYA16263	1.1	0.0; -23.7	0.0	149; 54.4	0.0
SYA013	117.4	0.0; 4.4	0.0; 6.5	ND	ND

^aResults from NIMH PDSP.^bref (Chen et al., 2012; Shapiro et al., 2003). ND = Not determined. Methysergide is the positive control for 5-HT_{1A}R. Tango assays are for β -arrestin-2 recruitment.