Published in final edited form as:

Anesthesiology. 2019 July ; 131(1): 7–9. doi:10.1097/ALN.0000000000002710.

Breathing Life into Pulmonary Physiology: How Age and Body Weight Impair Oxygenation

Jeffrey Thiboutot, MD MHS¹ and Robert H. Brown, MD MPH^{1,2,3,4}

¹Department of Medicine, Division of Pulmonary and Critical Care Medicine, Johns Hopkins University, Baltimore, MD.

²Department of Anesthesiology and Critical Care Medicine, Johns Hopkins University, Baltimore, MD.

³Department of Environmental Health and Engineering, Johns Hopkins University, Baltimore, MD

⁴Department of Radiology, Johns Hopkins University, Baltimore, MD

In the words of the famous H. L. Mencken, one of the most influential writers of the last century, “For every complex problem there is an answer that is clear, simple, and wrong.” Impaired arterial oxygenation is a complex problem. While one solution to impaired oxygenation is to increase the inspired oxygen, that is not necessarily the correct solution. As we parse out the various factors that cause worsening oxygenation, we get closer to understanding the reasons and thus closer to more accurate solutions.

As advances in anesthetic and surgical techniques have progressed over the past few decades, we continue to perform more surgeries on patients who previously were thought to be at too high a risk. Two determinates of worsening outcomes for patients undergoing anesthesia are old age and obesity. Traditionally, increased age is commonly associated with increasing co-morbidities, and obesity is associated with obstructive sleep apnea and atelectasis, all of which portend to poorer anesthesia outcomes. Beyond these superficial assessments, there is much to be learned about what is happening physiologically during anesthesia as it relates to these determinates. In the current issue of the journal, Hedenstierna and colleagues studied the pathophysiological effects of increased age and body weight on impaired oxygenation during anesthesia.¹

Caring for patients with obesity and morbid obesity has become a regular part of everyday practice. It is estimated that worldwide, by 2025, global obesity will surpass 18% of men and 21% of women², with even greater rates in high-income countries. Similarly, it is expected that by 2030, the number of people worldwide aged 60 and over will rise by nearly 56% to nearly 1.6 billion people³. With advances in areas such as transplant, oncologic, laparoscopic, and robotic surgeries, more procedures are safely being performed in the elderly^{4–6}. As these obese and elderly patients will respond differently to our approaches

Corresponding Author: Jeffrey Thiboutot, MD MHS, Johns Hopkins University, Department of Medicine, Division of Pulmonary and Critical Care Medicine, 5th Floor, 1830 E. Monument St., Baltimore, MD 21205, Jthibou1@jhmi.edu.

Conflicts of Interest: The authors declare no competing interests

and treatments, a better understanding of the physiologic alterations induced by aging and weight is needed to best care for these populations.

The healthy lung comprises a heterogeneous distribution of various proportions of ventilation (V) and perfusion (Q). This is commonly assessed by their ratio V/Q, which is highest at the apices and decreases towards the bases. Shunt is an abnormal physiologic phenomenon that is at the extreme end of the V/Q spectrum (no V, all Q). Both areas of low V/Q and of shunt can lead to oxygen impairment. Until recently, there have been limited studies of the effects of anesthesia on V/Q and oxygen impairment. One method to assess ventilation and V/Q defects is the multiple inert gas elimination technique (MIGET). MIGET uses infused inert gases with different solubilities and measures systemic and pulmonary artery concentrations as well as mixed expired concentrations of these inert gases to calculate V/Q.⁷

In this issue, Hedenstierna et. al. capitalized on MIGET technology to assess if V/Q differences exist in healthy patients (American Society of Anesthesiologist, ASA, class 1) undergoing anesthesia.¹ MIGET, Computed Tomography (CT) scans of the chest, and PaO₂ measurements were performed in healthy patients across varying ages and body weights both in the awake and anesthetized states to assess the drivers of impaired oxygenation under anesthesia. In this study, increased oxygen impairment was associated with older age and increased BMI in the anesthetized patients. MIGET revealed that areas of low V/Q increased with age, but had no association with BMI. Shunt on the other hand, followed a quadratic distribution with a peak at age 45 years, and increased linearly with BMI. Areas of low V/Q became a more important drivers of impaired oxygenation than shunt in the elderly adults. The explanation for the increased shunt with high body weights is fairly intuitive (and confirmed by CT findings). Higher body weights caused increased areas of atelectasis in the dependent portions of the lungs, through which blood is shunted, leading to impaired oxygenation. In addition, in the supine position, the size of these dependent portions of the lungs increase, further worsening atelectasis and shunt. However, what is most interesting was their finding that the peak effects of shunt occurred at age 45 years. The authors hypothesized that these areas of low V/Q may actually be preventing atelectasis. If one has existing areas of low V/Q prior to anesthesia, during pre-oxygenation with higher FiO₂ (0.8–1), the nitrogen existing at the alveoli will not have enough time to be washed out. Then, following the induction of anesthesia, when FiO₂ is reduced, in areas of low ventilation, the oxygen will still be absorbed but the larger amount of remaining nitrogen will act to stent open the alveoli. This theory has been supported by previous studies^{8–10}.

The work did have several limitations. This was a retrospective study that used previously acquired data from eight studies performed at two sites with slightly differing population demographics, general anesthetic choices, and respiratory settings. Also, the studies were only performed on healthy (ASA class 1) patients. Whether co-morbidities will affect these findings remains to be determined. In addition, only non-obese (BMI < 30 kg/m²) subjects were studied. While there are no obvious reasons why the relationship between weight and impaired oxygenation should be different in obese (or morbidly obese) individuals compared to simply overweight ones, further studies including these populations should be performed. It is currently leap of faith, albeit small, to extrapolate these findings to obese patients.

Another limitation is that while MIGET measurements can give an overall estimate of V/Q, it has no spatial resolution, so the sites of mismatch within the lungs cannot be identified. Unfortunately, this technique is not amenable to clinical practice and will remain as only a research tool.

So how do these findings affect clinical practice? One size may not fit all. While recruitment maneuvers may still be useful in overweight (and obese) patients to reduce atelectasis, recruitment maneuvers in elderly patients may not be helpful, as their oxygenation impairment may be driven by low V/Q, not atelectasis (shunt). As suggested by the authors, in the elderly patients where low V/Q will be the cause of the impaired oxygenation, one should consider increasing lung volume, such as through the use of PEEP to stabilize the airways, and decreasing expiratory flow rates to keep airways open. While these findings are provocative and may lead to alternative interventions for different groups of patients with impaired oxygenation, they should be verified in the populations of interest, ideally through randomized controlled trials.

Acknowledgments

Funding Statement: Research reported in this publication was supported by the National Heart, Lung, and Blood Institute of the National Institutes of Health under Award Numbers RO1HL121788 and F32HL144121. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

References

1. Hedenstierna G, Tokics L, Scarramuzzo G, Rothen HU, Edmark L, Ohrvik J. Oxygenation Impairment During Anesthesia: Influence of Age and Body Weight. *Anesthesiology*. 2019;00(00).
2. Trends in adult body-mass index in 200 countries from 1975 to 2014: a pooled analysis of 1698 population-based measurement studies with 19.2 million participants. *Lancet* (London, England). 2016;387(10026):1377–1396.
3. United Nations DoEaSA, Population Division. World Population Ageing 2015. ST/ESA/SER.A/390. 2015.
4. Ceccarelli G, Andolfi E, Biancafarina A, et al. Robot-assisted surgery in elderly and very elderly population: our experience in oncologic and general surgery with literature review. *Aging clinical and experimental research*. 2017;29(Suppl 1):55–63. [PubMed: 27905087]
5. Segall L, Nistor I, Pascual J, et al. Criteria for and Appropriateness of Renal Transplantation in Elderly Patients With End-Stage Renal Disease: A Literature Review and Position Statement on Behalf of the European Renal Association-European Dialysis and Transplant Association Descartes Working Group and European Renal Best Practice. *Transplantation*. 2016;100(10):e55–65. [PubMed: 27472096]
6. Otake A, Sasase A, Suzuki A, et al. Assessment of age-specific safety of laparoscopic surgery in elderly patients with ovarian tumors. *The journal of obstetrics and gynaecology research*. 2016;42(3):297–301. [PubMed: 26628082]
7. Mélot C Contribution of multiple inert gas elimination technique to pulmonary medicine. 5. Ventilation-perfusion relationships in acute respiratory failure. *Thorax*. 1994;49(12):1251–1258. [PubMed: 7878564]
8. Edmark L, Kostova-Aherdan K, Enlund M, Hedenstierna G. Optimal oxygen concentration during induction of general anesthesia. *Anesthesiology*. 2003;98(1):28–33. [PubMed: 12502975]
9. Sum Ping SJ, Makary LF, Van Hal MD. Factors influencing oxygen store during denitrogenation in the healthy patient. *Journal of clinical anesthesia*. 2009;21(3):183–189. [PubMed: 19464611]
10. Rothen HU, Sporre B, Engberg G, Wegenius G, Reber A, Hedenstierna G. Prevention of atelectasis during general anaesthesia. *Lancet* (London, England). 1995;345(8962):1387–1391.