Received:
22 January 2019

Revised:
12 March 2019

Accepted:
13 March 2019

<https://doi.org/10.1259/bjr.20190078>

Cite this article as:

Gaur P, Gedroyc W, Hill P. ADPKD—what the radiologist should know. *Br J Radiol* 2019; **92**: 20190078.

REVIEW ARTICLE

ADPKD—what the radiologist should know

¹PRITIKA GAUR, BSc, MBBS, ¹WLADYSLAW GEDROYC, MBBS, MRCP, FRCR and ²PETER HILL, MBBS, BSc, FRCP

¹Department of Radiology, Imperial College Healthcare NHS Trust, London, United Kingdom

²Renal Medicine, Imperial College Healthcare NHS Trust, London, United Kingdom

Address correspondence to: Dr. Pritika Gaur

E-mail: p.gaur@nhs.net

ABSTRACT:

Autosomal dominant polycystic kidney disease (ADPKD) is the most common renal genetic disorder and a leading cause of end stage renal failure (ESRF) affecting over 12 million people worldwide. Whilst the mainstay of diagnosis has historically favoured the imaging domain, the progression of disease was until very recently thought to be best monitored via biochemical analysis, *i.e.* measurement of estimated glomerular filtration rate. Imaging modalities such as sonography, CT and MRI have more recently proven to be key in monitoring disease progression. As much as half of the renal parenchyma can be lost with no real derangement in renal function. Tolvaptan, a vasopressin antagonist has been shown to slow disease progression and preserve renal function. Here we discuss at length the pathogenesis of ADPKD, the various diagnostic challenges surrounding its evaluation, new treatment options and monitoring of disease progression via serial imaging. We also propose monitoring of the efficacy of Tolvaptan at slowing the rate of deterioration in renal function in patients with ADPKD through MRI guided volumetric analysis of the kidneys.

INTRODUCTION

ADPKD is the commonest renal genetic disorder affecting almost 60,000 patients in the United Kingdom alone and over 12 million worldwide.¹ It is characterized by progressive cystic dilatation of renal tubules causing architectural distortion, increase in kidney size and decline in renal function often culminating in end stage renal disease. Most inherited polycystic kidney disease is acquired in an autosomal dominant fashion (ADPKD) and usually presents symptomatically in adult life. A rarer disease—autosomal recessive PKD presents much earlier during the antenatal or neonatal period. We focus our attention on ADPKD in our review.

GENETICS AND PATHOGENESIS

Affected individuals have a 50% risk of passing it on to their offspring. Although almost one in four cases of ADPKD present with no known family history; a large proportion of these are due to undiagnosed cases in family members and roughly 6% are as a result of *de novo* mutation²

Gene mutations affecting PKD1 (on the long arm of chromosome 16³ and PKD2 (located on the long arm of chromosome 4) affect 85 and 15% of cases respectively.^{4,5} These mutations cause dysfunction of transmembrane proteins in the primary cilia of renal tubular epithelial cells, namely polycystin 1 and 2. The underlying mutation primarily

affects the tubular cells which primarily which will proliferate continuously resulting in an increased number of cells; there is resultant dilatation of the tubular segments and eventual bulging of cysts into the interstitium. Eventually there is separation of the cyst from the tubule becoming an autonomous cyst. Abnormal ciliary function correlates with disproportionately increased levels of intracellular calcium and disruption of local homeostasis via upregulation of cAMP pathways in renal collecting ducts and distal nephrons, leads to expanding cysts filling with fluid.⁶ Usually, this is triggered by arginine vasopressin which promotes cyst cell proliferation and cyst fluid secretion.⁷

Cysts are originally attributed to 1–2% of all nephrons within various segments of the kidney, however as they grow in number and size they cause apoptosis of proximal tubule cells, obstruct adequate flow of urine, and compress adjacent tubular vasculature thereby activating the renin-angiotensin system. This leads to damaged renal parenchyma, urinary tract infections and hypertension respectively⁸. It has been hypothesized that injured tubules and expanding cysts release inflammatory cytokines causing fibrosis and eventual progression to ESRF

PRESENTATION

Case presentation is usually in early adult life between the ages of 20–30 years usually either as part of screening due

Figure 1. Unilateral renal sonographic appearance of a 44-year-old female, known to have PCKD. Presence of simple cysts with post cystic acoustic enhancement and nephromegaly is noted. PCKD, polycystic kidney disease.

to knowledge of the mutation within a family member, vague abdominal discomfort or incidentally as part of sonographic evaluation of the abdomen for another cause. Once the cysts exponentially increase in size and number, symptoms of ADPKD and its complications may present. These may include poor urinary concentration, infection, intracystic bleeding, haematuria, nephrolithiasis, systemic hypertension and eventually renal insufficiency.⁸

Extra renal manifestations of ADPKD can involve presence of cysts within other organs such as the liver, pancreas, spleen, seminal vesicle cysts and arachnoid cysts within the brain. Higher instances of vascular connective tissue like disorders have also been noted within the spectrum of ADPKD including presence of vascular aneurysms (aortic and intracranial), abdominal herniation, mitral valve insufficiency secondary to prolapse, diverticulitis and genitourinary cysts. Further, links to metal health disorders such as depression have been published.^{9,10} These extrarenal complications although occurring with low frequency must be considered by the radiologist whilst evaluating relevant imaging studies. Figures attached of extra renal complications, ICA aneurysm, diverticulitis.

DIAGNOSIS

There are various methods of diagnosing ADPKD, including sonographic evaluation (Figure 1) and genetic testing. Genetic testing is however not frequently employed in the diagnosis of ADPKD, the main reason being the associated cost and labour requirement. This is currently only being used in exceptional circumstances for research purposes and in those patients in whom ADPKD must be ruled out with absolute certainty (living kidney donor).¹¹ Conflicting evidence is demonstrated with regard to confident exclusion of disease based upon the imaging modality utilised and age exclusion criteria. According to Pei et al, ultrasound exclusion is unreliable below the age of 14, whereas the detection of 2 or fewer cysts below the age of 40 rules out PKD.¹² A recent conference discussing controversies surrounding PKD diagnosis with a view to improving global

Table 1.

Age (years)	Number of Cysts	Modality
15–39	three or more (in total)	US
40–59	two in each kidney	US
>60	four in each kidney	US
16–40	More than 10 (in both kidneys)	MRI

outcomes; the notion of ultrasound confidently excluding the diagnosis in those aged below 40 was deemed suboptimal.¹³

Family screening of at risk relatives involves sonographic evaluation of the kidneys, again using the criteria described above.

Regardless of these controversies the abdominal ultrasound is a reliable, repeatable, sensitive and biologically safe method of diagnosis; and currently remains the preferred method of diagnosis in those with a positive family history of the disease.

Ravine and Demetriou devised specific criteria for the diagnosis of Type 1 and Type 2 ADPKD respectively. In clinical practice however, we are not necessarily equipped to differentiate between the two and hence an integrated approach based on a positive family history and including MR criteria was devised by Pei et al.^{12,14}

Pei et Criteria for diagnosis of ADPKD (Table 1).

DIFFERENTIAL DIAGNOSES

There are other diseases which cause cystic change or produce renal lesions which may create diagnostic confusion. The radiological manifestations of these must be considered for accurate diagnoses. Listed in the table below are some important differential diagnoses with their radiological appearances.¹⁵ All of the diseases below may show cortical renal cysts (Table 2).

COMPLICATIONS AND RENAL MANIFESTATIONS OF ADPKD

Patients with ADPKD are imaged not only to make the initial diagnosis, but also to monitor disease progression due to the high risk of renal complications. These include complicated cysts, nephrolithiasis, perinephric haemorrhage and cyst rupture. Rarely, renal cell carcinoma may also present itself on a background of ADPKD.

NEPHROLITHIASIS

This is more common in patients with ADPKD compared to the background general population, occurring in around 20% of patients with ADPKD.¹⁶ Therefore this is an important differential to bear in mind in patients with ADPKD presenting with flank pain. This can be due to metabolic disturbances within the collecting system promoting formation of calculi. Additionally kidney size is a good indicator of likelihood of developing renal stones, with larger kidney size correlating with an increased likelihood of stone formation.

Table 2. Differential diagnosis of cystic kidney disease and their imaging manifestations

Differential diagnosis	Radiological features	Additional features	Examples
Tuberous sclerosis: gene mutation affecting TSC1 and TSC2.	<p>Renal angiomyolipomas: type of benign renal lesion composed of vascular, smooth muscle and fat.</p> <p>Ultrasound: hyperechoic lesions in the renal cortex.</p> <p>CT: Macroscopic fat within lesions. Albeit some lipid poor AMLs.</p> <p>MRI : fat saturated sequences with in and out of Phase imaging demonstrate fat rich lesions.</p>	<p>Neurological: Cortical and subcortical tubers</p> <p>Abdominal: Renal AML</p> <p>Thoracic: Lymphangioleiomyomatosis (Figure 2)</p> <p>Musculoskeletal: bony sclerosis</p>

(Continued)

Table 2. (Continued)

Differential diagnosis	Radiological features	Additional features	Examples
<p>Von Hippel Lindau: mutation of VHL tumour suppressor gene on chromosome 3.</p> <p>Development of numerous benign tumours</p>	<p>Renal cell carcinomas, usually are bilateral in combination with bilateral renal cysts.</p>	<p>Neurological: CNS hemangioblastomas</p> <p>Abdominopelvic: liver cysts, pheochromocytomas, pancreatic cysts,</p>	<p>Case courtesy of A.Prof Frank Gaillard, rID: 4314 demonstrating a cerebellar hemangioma</p> <p>Case courtesy of Dr Mohammad Taghi Niknejad, rID: 6,1889</p>

(Continued)

Table 2. (Continued)

Differential diagnosis	Radiological features	Additional features	Examples
congenital cystic Kidney Disease: mutation in the PKHD1 gene (polycystic kidney and hepatic disease) location on chromosome 6p.	Bilateral renal involvement, from a young age unlike ADPKD. The cysts rarely exceed 1–2 cm in diameter.		<p>Case courtesy of Dr Hani Salam, rID: 9635</p>
Medullary sponge disease	Ultrasound: echogenic medullary pyramids X-ray and CT: Clusters of pyramidal medullary calcification	Lesions can be bilateral

(Continued)

Table 2. (Continued)

Differential diagnosis	Radiological features	Additional features	Examples
Acquired multicystic kidney disease This occurs in patients who have end stage renal failure and usually are on dialysis. No other history of cystic renal disease.	Multiple bilateral small mostly cortical, but can be renal medullary cysts The kidneys are generally decreased in size	Hyperplastic cysts have been considered a source of renal cell carcinoma	<p>Case courtesy of Dr Mostafa Mahmoud El-Feky, rID: 5,4566</p>
Local renal cystic disease	Renal cysts involve segments of the kidney or the entire kidney but are <i>always</i> unilateral. The cysts are always separated by renal parenchyma.		<p>Case courtesy of Dr Mohammad Taghi Niknejad, rID: 6,0730</p>

AML, angiomyolipoma; CNS, central nervous system; VHL, von Hippel-Lindau.

Figure 2. CT appearances of PCKD in axial section. PCKD, polycystic kidney disease.

Imaging features may be confusing especially on ultrasound examination with presence of multiple background renal cysts, some of which may contain calcification. Diagnostic uncertainty with renal stones may occur since all calcifications cast a posterior acoustic enhancement. Therefore, the best imaging modality is high resolution CT, which detects stones and calcifications. Composition of calculi usually includes uric acid or calcium oxalate.

COMPLEX CYSTS

Simple cysts are of low density on CT and hypoechoic on ultrasound, whilst on MRI they demonstrate high signal intensity on T_2 weighted imaging, and low signal intensity on T_1 weighted imaging. Infection, haemorrhage or formation of renal cell carcinoma may be the cause of developing complex cysts. They may be hyperechoic on ultrasound, show hyperdensity on CT with calcification, and return low signal on T_1 weighted MRI with variable signal intensity on T_2 weighted MR imaging (Figure 3c). These do not enhance post gadolinium administration^{17 18}

Angiographic evaluation may be required in cyst rupture which causes a disproportionate amount of bleeding, with consideration of embolization and/or nephrectomy if active extravasation is demonstrated acutely.¹⁹ Intracystic haemorrhage usually

Figure 3. (A) MRI T_2 weighted images (B). Liver cysts in PCKD. (C). T1 MRI sequence showing mixed signal intensity in the cysts representing various stages of haemorrhage into cysts; one of the complications of ADPKD. A, B and C Left to Right. ADPKD, autosomal dominant polycystic kidney disease; PCKD, polycystic kidney disease.

causes haematuria and flank pain, with the imaging features of complex cysts described above. If however, there is a direct communication of the cyst with the renal pelvis this may result in painless frank haematuria.

Intracystic infection, pyelonephritis or renal abscess normally present with fever, pain and evidence of a high signal T_1 weighted cyst on MRI which does not enhance on contrast administration. Since infection and haemorrhage into cysts can demonstrate similar features on MRI, the clinical picture must guide management. If renal obstruction is suspected sonographic evaluation or CT should be considered. ^{18}F FDG-PET studies can be used to detect avidity of infected kidney cysts more accurately than MRI,²⁰ and also support CT guided needle aspirations in cases of recurrent cyst formation.

RENAL CELL CARCINOMA

Although renal cell carcinoma has been reported in patients with ADPKD, there has been no firm causal link associated with the disease. There are Bosniak criteria which classify cyst characteristics into those which require intervention versus those that are likely benign. (Table 3 below).

Cysts which show suspicious imaging features such as irregular thickened walls and internal separations can be considered for repeat interval imaging, although MRI can be extremely useful in delineating benign from malignant lesions. Cysts which enhance post-gadolinium administration (unlike most benign complex cysts) will require biopsy and even surgical intervention. As was discussed earlier, complex cysts due to infection and bleeding on the whole do not demonstrate post-contrast administration enhancement. It is sometimes possible though for complicated haemorrhage and infected cysts to demonstrate enhancement which would probably require histological sampling or surgical resection as a neoplasm cannot be ruled out.²²

TREATMENT

Polycystic kidney disease lends itself to further complications during the course of the illness; cyst infections and cyst haemorrhage being relatively common occurrences. Often, imaging-based diagnosis of these is difficult due to the high proteinaceous content of cysts and variable signals on MR sequences. (Figure 3 above shows mixed signal intensity arising from cysts). Furthermore, general complications of chronic renal failure—hypertension, anaemia and hyperphosphataemia are seen with disease progression which are treated with targeted medication acting directly at the end-organ damage sites.

More recently, a novel therapy with a highly selective vasopressin antagonist drug namely Tolvaptan has been trialled and successfully used in adults with Stage 1–3 ADPKD. In a patient with ADPKD, arginine vasopressin (anti-diuretic hormone) induced production of cyclic-AMP in renal collecting ducts and distal nephrons, promotes retention of urine and causes cyst cell proliferation and cyst fluid secretion. Antagonism of the vasopressin-2 receptor by Tolvaptan on the contrary results in reduced urine osmolality and increased water secretion with decreased cyst cell proliferation. Tolvaptan has stringent commencement

Table 3. Summary of Bosniak cyst criteria ²¹

Category	Description	Management
IA	Simple cyst. No septations, no calcifications, water attenuation, no enhancement	Nil
IIA	May contain hairline thin septa, fine calcification in septa, short uniformly high attenuation lesions (3 cm), no measurable enhancement	Nil
IIB	Multiple hairline thin septa, minimal thickening of wall or septa, calcification which may be thick or nodular, no measurable enhancement, 3 cm	Repeat imaging
III	Thickened irregular or thickened walls, measureable enhancement	Surgical for histology or resection
IV	All criteria of III, but also distinct soft tissue component to the cyst which shows enhancement	Surgical resection

criteria in accordance with the NICE guidelines, with treatment only possible in adults over the age of 18 or with evidence of rapidly progressive disease defined by continuous fall in estimated glomerular filtration rate (eGFR) >5 ml/min in 1 year and more than 5% increase renal volume per year. The TEMPO trial 3:4 showcases evidence that 3 years of total treatment reduced growth in total kidney volume and reduced rate of renal function decline when compared to placebo; the mean increase in total kidney volume per year was 2.8% for patients on Tolvaptan when compared to 5.5% for patients on a placebo. The rate of formation of cysts is reduced and lower event rates such as cyst haemorrhage and infection have been reported. Some of the common side-effects been reported are those that are of the normal physiological disturbance caused by the pharmacodynamic properties of a V2 receptor antagonist causing polyuria, nocturia, polydipsia and thirst. Other idiosyncratic side-effects include derangement of liver enzymes and therefore liver function tests.

MONITORING DISEASE PROGRESSION

It has been postulated that there is no biochemical evidence of renal function decline (eGFR) despite ongoing cyst proliferation and loss of over 50% of functioning renal parenchyma; making renal function measurements unreliable at early stages of the disease.¹⁵ Kidney and cyst volume and size measurements have been shown to be better predictors of progression of disease in the monitoring of ADPKD.²³

Sonographic evaluation has been a front line aid in diagnosis and evaluation for complications of ADPKD. It is fast, biologically safe and repeatable. However, being an operator-dependant imaging modality it gives rise to interuser variability with low reproducibility; of particular importance in serial kidney measurements. Here we evaluate the use of various imaging modalities with an emphasis on MRI.

Information for almost 30 years of follow up of total kidney volume from 2355 patients has been collated by the PKD Outcomes Consortium including observational datasets from various sources, *i.e.* Consortium for Radiologic Imaging Studies (CRISP).^{24,25} These demonstrate use of total kidney volume as a baseline prognostic biomarker decline in eGFR. Further, changes in total kidney volume (TKV) is shown to be a useful marker in

monitoring disease progression with image-based kidney and/or cyst volume measurement.

The Mayo Clinic Model is an online risk calculator which risk stratifies patients into various classes A–E with estimated annual kidney growth rates based on height and age adjusted total kidney volumes :

Class 1A < 1.5%

Class 1B 1.5–3.0%

Class 1C 3.0–4.5%

Class 1 D 4.5–6.0%

Class 1 E > 6.0% per year.

Patients in Class 1C–1E are predicted to be at risk of rapid disease progression, and accord to the Mayo criteria, taking into account the age and eGFR, Tolvaptan has been recommended for these patients. The role of radiologist therefore in accurately measuring renal volumes in a repeatable manner is paramount.

IMAGING MODALITIES AND TECHNIQUES USED FOR VOLUME MEASUREMENT

Ultrasound

The use of ultrasound for diagnostic purposes has been discussed above. This modality of imaging has several drawbacks affecting the accuracy of TKV estimation due to interoperator variability. CRISP data suggest that kidneys with >17 cm length would be unreliably assessed on sonographic evaluation²⁶. Other imaging modalities such as CT and MRI would be better suited to volume measurement.

CT

This is an accurate method of TKV estimation although requires patient exposure to ionising radiation and nephrotoxic iodinated contrast. It has been proven to be of similar accuracy as MRI in volume measurement.

MRI

A number of studies have been conducted which show use of gadolinium contrast T1 sequences for the assessment of renal volume through the means of manual stereology. Nephrogenic systemic fibrosis was shown to affect patients with pre-existing impaired renal function secondary to gadolinium administration. The newer macrocyclic gadolinium compound used in modern centres has not been linked with NSF,²⁷ however recent recommendations following CRISP data are as follows; acquisition of T_2 weighted fast spin echo images with fat saturation and without contrast administration as ideal for volume measurement. Several methods of measurement techniques have been described, requiring a range of interpreter time impacting technique selection for the busy radiologist.

The following methods have been described for volume measurement which can be utilized in kidney volume measurement.

Stereology for CT/MRI

Grid points are defined corresponding to regions covering the entire organ in various cross-sectional slices; volume calculation is performed by summation of products of the resulting areas

Ellipsoid formula for US/MRI/CT

This is when KV is a measure of AP diameter x transaxial diameter x craniocaudal length x $\pi/6$. This formula is 87% accurate when used in MRI/CT studies, however, only demonstrates 21% accuracy in sonographic evaluation. Studies have shown although this shows good correlation with actual volume measurements, in itself this is probably not an accurate enough technique and shows great interobserver variability. This is seen in Figure 4 [Figure 4](#)

Further volume measurement strategies such as the **Mid-slice, Semi-automated and Automated** techniques have been described. These provide good accuracy, the gold-standard remains manual planimetry providing 100% accuracy.

We will now focus on the preferred volumetric analysis which we use at our centre and has been shown to be the gold standard.

Figure 4. Ellipsoid Method

Figure 5. Cavalieri's Principle

Manual planimetry or the modified Cavalieri method for CT/MRI

[Figure 5](#) Diagrammatic representation of how volumes of various slices through an object can be calculated using the Cavalieri Principle.

The kidney volume can be estimated using the Cavalieri principle of volume estimation via measurement of consecutive sections. All PACS systems allow for "lesion" segmentation which is a platform for calculating the volume of the kidneys through tracing an outline onto each cross-sectional image of the kidney, the most accurate but a little more time consuming. [Figure 6](#) demonstrates the same kidney which was calculated to be 1681 cm³ as compared to 2772 cm³ via the ellipsoid formula demonstrated above.

This method is based on the mathematical principle set out by Bonaventura Cavalieri; The cut surface areas of these sections are calculated and summated to provide the total cut surface area. Multiplication with the mean slice thickness will then estimate the object volume. This is demonstrated by the equation $V = t \times \sum A_{cm}^3$

Studies have reported little interuser variability and good agreement in repeat measures for the modified Cavalieri technique when used for volume measurement of brain volume²⁸ and fibroid measurement²⁹. It is important for the region of interest

Figure 6. Manual Planimetry on MRI using manual segmentation

to have clear boundaries such as the kidney; as a failure to distinguish the boundaries of interested structures may contribute to bias.

As the gold standard method of volume measurement, we recommend using this technique for monitoring disease progression in ADPKD.

DISCUSSION

As radiologists, we have a large role in imaging renal and extra-renal complications of this disease and in distinguishing it from other cystic diseases which may have immense consequences for future patient management. Furthermore, an understanding of the role of imaging in ADPKD is important in optimal modern management of these patients through what may be decades of managing the condition. The development of new drugs which slow rate of deterioration of ADPKD changes the radiologists' role here from a bystander to an active participant in helping to monitor the efficacy of novel therapy. It is crucial to understand that the lack of regenerative capacity of nephrons renders the use of eGFR, and as such other biochemical markers, non-viable in monitoring efficacy of treatment; as the eGFR will almost never

improve due to the irreversible inflammatory fibrotic damage to renal nephrons³⁰. As established in our section relating to monitoring disease progression, the more reliable upcoming marker, especially in the early stages of the disease is TKV. Currently an online tool for use in selection of patients for clinical trials, mayo criteria, risk stratifies patients into various classes which can subsequently be used for determining commencement of treatment with Tolvaptan.

Irazabal et al demonstrated that measurement of TKV via the ellipsoid method shows good correlation with stereology method, the accuracy is a mere 21%.³¹ This reaffirms that although adequate for prediction of disease progression, ultimately other methods such as manual planimetry should be sought.⁽²⁵⁾

Total kidney volume is one of the few criteria is one of the very few FDA-certified biomarkers for clinical trial use which makes it all the more imperative for us to accurately calculate it. MRI and CT are in current use for most clinical trials related to volume estimation in ADPKD. Out of the various methods of volume estimation, the highest inter- and intraobserver agreement rates were estimated in planimetry and stereology methods, both

being more reproducible for the expert operator and the former for the beginner operator.³² Of interest, these methods show more accuracy on CT in comparison to MRI due to slice thickness and utilisation of the Cavalieri method; however, this would not be feasible due to the chronic nature of ADPKD and serial imaging requirements over decades and the radiation hazards carried by CT. MRI is therefore the most accurate method practically applicable to imaging requests, with the time requirement being the biggest factor to impact on MRI reporting adding on average at least 30 additional minutes to produce accurate reproducibility.³³ We must evaluate the additional value this carries in enhancing the radiological report, considering volume estimation is useful in such patients particularly around decision making on Tolvaptan commencement.³⁴ A request from the renal physician specifying volume measurement after patient selection would lend huge help in guiding radiologists to prioritise

the imaging requiring additional reporting time, with a pathway similar to vasculature evaluation for potential kidney donors—these can then be assigned subspecialists within Radiology to take on responsibility for TKV estimation reporting in addition to producing a standard MRI report.

This will be invaluable for renal physicians in selecting patients for therapy commencement enabling risk stratification through kidney growth per year in combination with eGFR. It will allow new opportunities to continually build our database for TKV as a prognostic indicator. In addition, data can be assimilated on the efficacy of Tolvaptan as a viable treatment outside clinical trials by demonstrating rate of reduction in kidney growth per year, opening further avenues into investigating the role of vasopressin antagonists in various patients with their differing comorbidities and how this plays into everyday clinical practice.

REFERENCES

- Gabow PA. Autosomal dominant polycystic kidney disease. *N Engl J Med Overseas Ed* 1993; **329**: 332–42. doi: <https://doi.org/10.1056/NEJM199307293290508>
- Ong ACM, Devuyt O, Knebelmann B, Walz G, for Inherited EE. Autosomal dominant polycystic kidney disease: the changing face of clinical management. *The Lancet* 2015; **385**: 1993–2002. doi: [https://doi.org/10.1016/S0140-6736\(15\)60907-2](https://doi.org/10.1016/S0140-6736(15)60907-2)
- Ward C, Peral B, Hughes J, Thomas S, Gamble V, MacCarthy A, et al. The polycystic kidney disease 1 gene encodes a 14 kb transcript and lies within a duplicated region on chromosome 16. The European polycystic kidney disease Consortium. *Cell* 1994; **77**: 881–94.
- Torres VE, Harris PC, Pirson Y. Autosomal dominant polycystic kidney disease. *The Lancet* 2007; **369**: 1287–301. doi: [https://doi.org/10.1016/S0140-6736\(07\)60601-1](https://doi.org/10.1016/S0140-6736(07)60601-1)
- Mochizuki T, Wu G, Hayashi T, Xenophontos SL, Veldhuisen B, Saris JJ, et al. PKD2, a gene for polycystic kidney disease that encodes an integral membrane protein. *Science* 1996; **272**: 1339–42. doi: <https://doi.org/10.1126/science.272.5266.1339>
- Cowley BD. Calcium, cyclic AMP, and MAP kinases: dysregulation in polycystic kidney disease. *Kidney Int* 2008; **73**: 251–3. doi: <https://doi.org/10.1038/sj.ki.5002695>
- Chapman AB, Johnson A, Gabow PA, Schrier RW. The renin-angiotensin-aldosterone system and autosomal dominant polycystic kidney disease. *N Engl J Med* 1990; **323**: 1091–6. doi: <https://doi.org/10.1056/NEJM199010183231602>
- Galarreta CI, Grantham JJ, Forbes MS, Maser RL, Wallace DP, Chevalier RL. Tubular obstruction leads to progressive proximal tubular injury and atubular glomeruli in polycystic kidney disease. *Am J Pathol* 2014; **184**: 1957–66. doi: <https://doi.org/10.1016/j.ajpath.2014.03.007>
- Grantham JJ, Torres VE. The importance of total kidney volume in evaluating progression of polycystic kidney disease. *Nat Rev Nephrol* 2016; **12**: 667–77. doi: <https://doi.org/10.1038/nrneph.2016.135>
- Luciano RL, Dahl NK. Extra-renal manifestations of autosomal dominant polycystic kidney disease (ADPKD): considerations for routine screening and management. *Nephrology Dialysis Transplantation* 2014; **29**: 247–54. doi: <https://doi.org/10.1093/ndt/gft437>
- Rossetti S, Strmecki L, Gamble V, Burton S, Sneddon V, Peral B, et al. Mutation analysis of the entire PKD1 gene: genetic and diagnostic implications. *Am J Hum Genet* 2001; **68**: 46–63. doi: <https://doi.org/10.1086/316939>
- Barua M, Pei Y. Diagnosis of autosomal-dominant polycystic kidney disease: an integrated approach. *Semin Nephrol* 2010; **30**(No. 4): 356–65. doi: <https://doi.org/10.1016/j.semnephrol.2010.06.002>
- Chapman AB, Devuyt O, Eckardt K-U, Gansevoort RT, Harris T, Horie S, et al. Autosomal-dominant polycystic kidney disease (ADPKD): Executive summary from a kidney disease: improving global outcomes (KDIGO) controversies conference. *Kidney Int* 2015; **88**: 17–27. doi: <https://doi.org/10.1038/ki.2015.59>
- Pei Y, Obaji J, Dupuis A, Paterson AD, Magistroni R, Dicks E, et al. Unified criteria for ultrasonographic diagnosis of ADPKD. *J Am Soc Nephrol* 2009; **20**: 205–12. doi: <https://doi.org/10.1681/ASN.2008050507>
- Chapman AB, Guay-Woodford LM, Grantham JJ, Torres VE, Bae KT, Baumgarten DA, et al. Renal structure in early autosomal-dominant polycystic kidney disease (ADPKD): the Consortium for radiologic imaging studies of polycystic kidney disease (CRISP) cohort. *Kidney Int* 2003; **64**: 1035–45. doi: <https://doi.org/10.1046/j.1523-1755.2003.00185.x>
- Rahbari-Oskouei F, Mittal A, Mittal P, Chapman A. Renal relevant radiology: radiologic imaging in autosomal dominant polycystic kidney disease. *Clin J Am Soc Nephrol* 2014; **9**: 406–15. doi: <https://doi.org/10.2215/CJN.08940813>
- Gabow PA. Autosomal dominant polycystic kidney disease. *N Engl J Med* 1993; **329**: 332–42. doi: <https://doi.org/10.1056/NEJM199307293290508>
- Bisceglia M, Galliani CA, Senger C, Stallone C, Sessa A: renal cystic diseases: a review. *Adv Anat Pathol* 2006; **13**: 26–56.
- Bleeker-Rovers CP, de Sévaux RG, van Hamersvelt HW, Corstens FH, Oyen WJ. Diagnosis of renal and hepatic cyst infections..
- By 18-F-fluorodeoxyglucose positron emission tomography in autosomal dominant polycystic kidney disease. *Am J Kidney Dis* 2003; **41**: E18–E21.
- Bosniak MA. The current radiological approach to renal cysts. *Radiology* 1986; **158**: 1–10PubMed. doi: <https://doi.org/10.1148/radiology.158.1.3510019>
- Keith DS, Torres VE, King BF, Zinck H, Farrow GM: renal cell carcinoma in

- autosomal dominant polycystic kidney disease. *J Am Soc Nephrol* 1994; **4**: 1661–9.
23. Grantham JJ. Clinical practice. Autosomal dominant polycystic kidney disease. *N Engl J Med* 2008; **359**: 1477–85. doi: <https://doi.org/10.1056/NEJMcp0804458>
 24. Perrone RD, Neville J, Chapman AB, Gitomer BY, Miskulin DC, Torres VE, et al. Therapeutic area data standards for autosomal dominant polycystic kidney disease: a report from the polycystic kidney disease outcomes Consortium (PKDOC). *Am J Kidney Dis* 2015; **66**: 583–90. doi: <https://doi.org/10.1053/j.ajkd.2015.04.044>
 25. Perrone RD, Mouksassi M-S, Romero K, Czerwiec FS, Chapman AB, Gitomer BY, et al. Total Kidney Volume Is a Prognostic Biomarker of Renal Function Decline and Progression to End-Stage Renal Disease in Patients With Autosomal Dominant Polycystic Kidney Disease. *Kidney International Reports* 2017; **2**: 442–50. doi: <https://doi.org/10.1016/j.ekir.2017.01.003>
 26. Bhutani H, Smith V, Rahbari-Oskouei F, Mittal A, Grantham JJ, Torres VE, et al. A comparison of ultrasound and magnetic resonance imaging shows that kidney length predicts chronic kidney disease in autosomal dominant polycystic kidney disease. *Kidney Int* 2015; **88**: 146–51. doi: <https://doi.org/10.1038/ki.2015.71>
 27. Reiter T, Ritter O, Prince MR, Nordbeck P, Wanner C, Nagel E, et al. Minimizing risk of nephrogenic systemic fibrosis in cardiovascular magnetic resonance. *J Cardiovasc Magn Reson* 2012; **14**: 31. doi: <https://doi.org/10.1186/1532-429X-14-31>
 28. Sahin B, Elfaki A. Estimation of the volume and volume fraction of brain and brain structures on radiological images. *NeuroQuantology* 2011; **10**. doi: <https://doi.org/10.14704/nq.2012.10.1.429>
 29. Quinn SD, Vedelago J, Kashef E, Gedroyc W, Regan L. Measurement of uterine fibroid volume: a comparative accuracy and validation of methods study. *European Journal of Obstetrics & Gynecology and Reproductive Biology* 2013; **171**: 161–5. doi: <https://doi.org/10.1016/j.ejogrb.2013.08.036>
 30. Ibrahim HN, Foley R, Tan L, Rogers T, Bailey RF, Guo H, et al. Long-term consequences of kidney donation. *N Engl J Med* 2009; **360**: 459–69. doi: <https://doi.org/10.1056/NEJMoa0804883>
 31. Irazabal MV, Rangel LJ, Bergstralh EJ, Osborn SL, Harmon AJ, Sundsbak JL, et al. Imaging classification of autosomal dominant polycystic kidney disease: a simple model for selecting patients for clinical trials. *J Am Soc Nephrol* 2015; **26**: 160–72. doi: <https://doi.org/10.1681/ASN.2013101138>
 32. Magistroni R, Corsi C, Martí T, Torra R. A review of the imaging techniques for measuring kidney and cyst volume in establishing autosomal dominant polycystic kidney disease progression. *Am J Nephrol* 2018; **48**: 67–78. doi: <https://doi.org/10.1159/000491022>
 33. Tangri N, Hougen I, Alam A, Perrone R, McFarlane P, Pei Y. Total kidney volume as a biomarker of disease progression in autosomal dominant polycystic kidney disease. *Canadian Journal of Kidney Health and Disease* 2017; **4**: 205435811769335. doi: <https://doi.org/10.1177/2054358117693355>
 34. Kistler AD, Poster D, Krauer F, Weishaupt D, Raina S, Senn O, et al. Increases in kidney volume in autosomal dominant polycystic kidney disease can be detected within 6 months. *Kidney Int* 2009; **75**: 235–41. doi: <https://doi.org/10.1038/ki.2008.558>