Received:
09 August 2018

Revised:
07 January 2019

Accepted:
27 February 2019

<https://doi.org/10.1259/bjr.20180707>

Cite this article as:
Hell MM, Achenbach S. CT support of cardiac structural interventions. *Br J Radiol* 2019; **92**: 20180707.

REVIEW ARTICLE

CT support of cardiac structural interventions

MICHAELA M HELL, MD and STEPHAN ACHENBACH, MD

Department of Cardiology, Faculty of Medicine, Friedrich-Alexander-Universität Erlangen-Nürnberg (FAU), Erlangen, Germany

Address correspondence to: Dr Michaela M Hell
E-mail: michaela.hell@uk-erlangen.de

Abstract:

Due to its high temporal and isotropic spatial resolution, CT has become firmly established for pre-procedural imaging in the context of structural heart disease interventions. CT allows to very exactly measure dimensions of the target structure, CT can provide information regarding the access route and, as a very valuable addition, volumetric CT data sets can be used

to identify fluoroscopic projection angulations to optimally visualize the target structure and place devices. This review provides an overview of current methods and applications of pre-interventional CT to support adult cardiac interventions including transcatheter aortic valve implantation, percutaneous mitral valve intervention, left atrial appendage occlusion and paravalvular leak closure.

INTRODUCTION

The most frequent indication of cardiac CT imaging is to detect and rule out coronary artery stenosis.¹⁻³ Intravenous contrast is injected for the purpose of coronary lumen visualization and the procedure is referred to as “coronary CT angiography” (CTA). In a broader sense, “cardiac CTA” refers to contrast-enhanced visualization of the morphology and structure of the entire heart and adjacent great vessels. In fact, due to its high spatial resolution which is well below 1 mm in all orientations, contrast-enhanced cardiac CT imaging has emerged as a valuable tool for morphological cardiac imaging in the past years.⁴ With their close to isotropic spatial resolution, modern CT scanners allow the reconstruction of data sets in any desired plane without losing the ability to visualize small structures and perform exact measurements.

In this context, pre-procedural CT imaging for transcatheter aortic valve implantation (TAVI) is the most frequent indication, but there are numerous other interventions which potentially benefit from cardiac CTA and are discussed in the current review: these include percutaneous interventions of the mitral, tricuspid and pulmonary valves, left atrial appendage occlusion, the interventional closure of paravalvular leaks related to prosthetic valve replacement as well as percutaneous revascularization of coronary chronic total occlusions. For an interventional cardiologist, there are three major aspects for which CT imaging can provide valuable information when planning structural cardiac interventions: precise information regarding the dimensions of the target structure, information on optimal

fluoroscopic projections to provide orthogonal visualization of the target structure without foreshortening, and assessment of the vascular access route. For user convenience, several software solutions are offered for pre-procedural CT data analysis in structural cardiac interventions. They have been reviewed in recent publications and are not part of the current review.⁵

Although cardiac CT is a robust modality that can provide a large amount of information for pre-procedural planning, a potential drawback is the need for intravenous contrast enhancement. Patients may occasionally have contraindications for iodinated contrast. Radiation exposure is a further potential concern and efforts must be undertaken to keep radiation dose within reasonable limits. However, many patients under consideration for structural heart interventions – particularly, TAVI – are of relatively high age and radiation exposure will be of limited relevance.

General aspects of CT image acquisition for structural heart disease

Specific acquisition parameters depend on the CT system that is used, but some general principles should be taken into consideration:

- To achieve high spatial resolution, multi slice scanner systems should be used that provide a reconstructed slice width of 0.6 to 0.75 mm.
- Minimization of cardiac motion artefacts is achieved by ECG-synchronized imaging, either through retrospective gating or prospective triggering. Diastolic imaging is commonly applied, timed at approximately 70%

of the cardiac cycle. However, in patients with high heart rates and irregular rhythms, systolic imaging may be the preferred option (e.g. 250–350 ms after the R-wave peak). If necessary, the entire cardiac cycle can be covered to provide both systolic and diastolic image data, albeit at the expense of increased radiation exposure. The possibility to pre-treat patients with beta-blockers for rate control depends on the patient's underlying cardiac disease and clinical condition. While usually considered mandatory for coronary imaging, heart rate control does not carry the same relevance when CT is performed in the context of structural heart interventions since image quality does not have to be as high as for coronary artery visualization.

- Respiration must be suspended during acquisition to avoid severe motion artefacts. Patients must therefore be thoroughly instructed regarding breathhold commands prior to imaging.
- The volume of iodinated contrast should be minimized, common doses are 50–100 ml. Typically, a peripheral venous contrast injection rate of at least 4 ml s⁻¹ should be applied to achieve adequate intravascular contrast (consider 3 ml s⁻¹ in patients with renal insufficiency). For pre-procedural TAVI imaging, protocols with substantially lower amounts of contrast agent have been published.^{6,7}

Transcatheter aortic valve implantation

Transcatheter aortic valve implantation has emerged as treatment of choice for symptomatic severe aortic stenosis in patients with a high operative risk.⁸ As opposed to open heart surgery, direct visualization of the target area is not available during the TAVI procedure, making pre-procedural imaging crucial for the success of the intervention. CT is the preferred imaging method in this context.⁹ CT imaging should include evaluation of aortic root and aortic annulus dimensions, assessment of aortic valve structure and calcifications, determination of fluoroscopic viewing angles exactly orthogonal to the aortic annulus as well as analysis of the vascular access route. Measurement of the aortic valve anatomical orifice area is feasible by CT. However, echocardiography remains the preferred image modality to establish the severity of aortic valve stenosis.

While imaging of the heart and aortic root requires ECG-synchronized imaging, the aorta and ilio-femoral arteries can be imaged in a non-gated mode. Frequently, two separate data sets are acquired: A gated data set of the heart and a non-gated, much more rapid acquisition to cover the vascular access route. Splitting acquisition into two different data sets can reduce radiation exposure and contrast volume (due to faster volume coverage of the non-gated scan). When sizing the aortic annulus, dynamic changes during the cardiac cycle have to be taken into account. The short-axis diameter, area, and perimeter have been shown to exhibit a small, but significant increase in size during systole compared with diastole.^{10,11} Therefore, diastolic compared with systolic measurements can result in undersizing TAVI prostheses in patients with large conformational changes.

The aortic annulus is a virtual, ovally shaped structure formed by connecting the three most caudal insertion points of the aortic valve cusps (so-called “hinge points”).¹² Importantly, it is

the diameter of the aortic annulus and not the aortic root diameter that dictates the size of the prosthesis, irrespectively of the inserted prosthesis type.¹³ Incorrect sizing of the aortic valve prosthesis can lead to severe complications. Prosthesis undersizing (<10% greater than the CT-measured aortic annulus) carries an increased risk of poor hemodynamics, paravalvular leak or prosthesis embolization.^{14,15} Oversizing, on the other hand, can lead to incomplete expansion of the prosthesis (leading to both valvular and paravalvular regurgitation) or, more likely, to annulus rupture.¹⁶ In order to accurately assess aortic annulus size, a stepwise approach is typically applied. First, double-oblique multiplanar reconstructions are created in a plane that is exactly aligned with the aortic annulus (Figure 1A,B,C).¹⁷ Once the aortic annulus plane has been obtained, three different methods for determining the aortic annulus diameter have been proposed (Figure 1D,E,F): (1) calculation of the mean diameter from the long and short diameters of the oval aortic annulus, (2) planimetry of the aortic annulus area and (3) measurement of the circumference of the aortic annulus. In the latter two cases, the mean diameter is calculated under the assumption that after implantation of a prosthesis, the annulus will assume full circularity while maintaining the identical area and circumference. Especially for area measurements, this assumption is mathematically incorrect, but not to an extent that this would influence clinical practice. Interobserver comparisons have demonstrated a moderate to excellent agreement for manual assessment of the aortic annulus size.^{18,19} The use of automated algorithms is feasible in daily routine with accurate and reproducible results.^{20–25} Infrequently, patients may have annulus sizes for which no suitable prostheses are currently available (e.g. diameters less than 18 mm or greater than 31 mm).

The extent of calcification of the aortic annulus and commissures as well as of the left ventricular outflow tract (LVOT) has been associated with the risk of post-procedural paravalvular aortic regurgitation which, in turn, has been shown to significantly increase mortality in the short and long-term course.^{26,27} Marwan et al reported a significantly higher rate of post-procedural paravalvular regurgitation in patients with annular calcification when using a balloon-expandable prosthesis.²⁸ Reduced aortic annular flexibility, prosthesis undersizing and a shallow position of the prosthesis frame in the LVOT have also been identified as factors increasing the rate of post-procedural paravalvular regurgitation.^{29,30}

Coronary ostial occlusion is a severe complication after TAVI. It has an incidence of less than 1% but is frequently fatal.³¹ A short distance between the aortic annulus and the coronary ostia and a shallow sinus of Valsalva have been shown to be associated with coronary obstruction (Figure 2).³² Risk is further increased by the presence of bulky calcifications on the aortic cusps.³³ Using appropriately oriented multiplanar reconstructions, the distance from the aortic annular plane to the left and the right coronary ostium and the lengths of the native leaflets can easily be measured.³⁴ There is no minimum distance from the annular plane to the coronary ostia which could be considered “safe”. Much rather, the presence of large calcifications on the aortic valve cusps, the depth of the sinus of Valsalva, the cusp length

Figure 1. Determination of aortic annulus dimensions in CT. (A,B) Creation of a double-oblique plane of the aortic annulus (white bold line) containing the lowest insertion points of the right, left, and noncoronary aortic valve cusps (C, arrows). Ao, aorta; LV, left ventricle. Three different approaches to determine aortic annulus size have been suggested: (D) From the long and short diameters, the mean diameter can be calculated. (E) From the annulus area, the diameter can be deduced under the assumption that this area changes to a circle when a valve is implanted. (F) From the circumference, the diameter can be derived assuming that the circumference will stay constant during the implantation and that the annulus will achieve a perfectly circular shape after prosthesis implantation.

Figure 2. Assessment of native valve leaflet length and coronary ostium height. (A) Routinely, the length of the native valve leaflet and the height of the left and right coronary ostia are measured to predict possible ostial occlusion during prosthesis implantation. LM, left main coronary artery; RCA, right coronary artery. (B) Aortic root of a 82-year-old female patient showing a shallow left sinus of Valsalva, a native left leaflet length of 13 mm and an annulus-ostium distance of 12 mm, portending a moderate risk of ostial occlusion. (C) Subtotal stenosis of the left main coronary artery ostium after transcatheter implantation of an aortic valve prosthesis (arrow). (D) Left main coronary artery ostium after implantation of a stent (arrow).

Figure 3. Determination of a suitable fluoroscopic angulation. (A) Double-oblique plane at the level of the aortic valve commissures. The white line corresponds to the angulation of an image plane that is orthogonal both to the aortic annulus plane and to the commissural line between the left and noncoronary aortic valve. R, right coronary cusp; N, noncoronary cusp; L, left coronary cusp. (B) The predicted angulation (LAO 2°/caudal 7°) was used during the TAVI procedure as C-arm angulation for the first aortogram. (C) Final angiogram with the implanted TAVI prosthesis.

and the ostium-annulus distance on the one hand, versus the operative risk on the other hand have to be considered for the final decision to undergo TAVI in the individual patient.

Of critical importance for the catheter-based implantation of bioprosthetic aortic valves is the identification of a fluoroscopic projection that provides an exactly orthogonal visualization of the aortic valve plane. Suitable views can be determined intra procedurally by repeated angiograms but this may be time-consuming and requires repeated contrast injections.³⁵ CT data sets can be used to identify suitable projections that provide an orthogonal view on the aortic valve plane with a significant reduction in contrast agent volume required during the TAVI procedure compared to a conventional approach (Figure 3).^{36–39} The slightly different position of the patient between CT acquisition and the TAVI procedure causes no noticeable problem.

Percutaneous mitral valve interventions

Mitral regurgitation is the second most common valvular disease in adult patients.⁸ Approximately 20% of patients with heart failure and 12% of patients with previous myocardial infarction suffer from at least moderate mitral regurgitation.^{40,41} While the aortic valve anatomy is well suited for the percutaneous implantation of tubular prostheses, the mitral valve has a complex saddle-shaped structure with a non-planar annulus, an anterior and posterior peak, a complex subvalvular apparatus and highly variable anatomy.⁴² The mitral valve further lacks a circular, fibrous annular structure to support a prosthesis and is in close proximity to the LVOT. Simply transferring the TAVI implantation technique for treating mitral regurgitation is not an option, so that several percutaneous repair devices have been developed. The MitraClip® device is the most frequently used method.^{43,44} Based on the principle of surgical edge-to-edge mitral valve repair, the MitraClip® device clamps and converges the anterior and posterior leaflets thereby reducing the grade of regurgitation. However, effectiveness is inferior to surgical mitral valve repair.⁴⁵ Recently, supported by advanced

imaging guidance, the technique of transcatheter mitral valve implantation (TMVI) has been further developed.^{46–50} Given the complex problems of mitral valve interventions listed above, pre-procedural CT imaging for TMVI must address anatomical assessment and measurement of the mitral valvular and sub valvular apparatus, assessment of annular calcification, prediction of LVOT obstruction, investigation of mitral regurgitation aetiology and prediction of an orthogonal fluoroscopic angulation.

Assessment of the mitral valvular and sub valvular apparatus using 2D imaging has been shown to over simplify the complex 3D geometry. Hence, 3D segmentation of the mitral annulus should be favored.⁴² Measurement of leaflet lengths and angulations using thin-section multiplanar reconstruction images has been demonstrated to provide good correlation with 3D transthoracic echocardiography.⁵¹ To best account for the needs of TMVI, a D-shaped model of the mitral annulus involving truncation of the saddle-shaped mitral annular contour at a virtual interconnecting line between both fibrous trigones (the trigone-to-trigone distance) has been introduced.⁵² From this model, the annular area and perimeter can readily be derived.

A relevant procedural complication of transcatheter-based implantation of a prosthetic valve in mitral position is the obstruction of the LVOT due to the elongation of the outflow tract by the implanted device, previously described as “neo-LVOT”.⁵³ Patient-specific, CT-based virtual, stereolithographic device simulation as well as 3D-printing approaches have been suggested to allow pre-procedural identification of patients which are at an increased anatomical risk for a narrow neo-LVOT.⁵⁴ Identification of mitral valve prolapse has been shown to be highly accurate in comparison to traditional echocardiographic imaging.⁵⁵ Further characteristics of mitral valve prolapse, including myxomatous and degenerative leaflets and disjunction between the posterior mitral valve leaflet insertion and the atrioventricular junction, can also be identified.

Figure 4. Caseous Mitral Annular Calcification. Caseous mitral annular calcification (arrows). (A) Density within the liquefied core is typically similar to contrast enhanced blood. (B) Hence, an additional non-enhanced acquisition is highly useful to identify this condition.

Degenerative mitral annular calcification, commonly found in the elderly population, is most frequently located at the posterior rim but its extent can vary from mild localized to exuberant circumferential involvement.⁵⁶ A rare variation of mitral annular calcification is caseous annular calcification, most frequently found along the posterior rim. Caseous mitral annular calcification is typically a large-volume, tumor-like lesion which can be easily detected by CT (while difficult to judge by echocardiography). Cases of spontaneous regression of the caseous calcification have been reported.⁵⁷ The density within caseous mitral calcification can be similar to the contrast-enhanced blood pool, so that combined visualization with contrast-enhanced and non-contrast images may be required (Figure 4).^{57,58} The identification of caseous annular calcification is of importance as it may interfere with the apposition of self-expandable TMVI systems.⁴²

Similar to TAVI, pre-procedural CT allows the prediction of suitable C-arm angulations orthogonal to the mitral annulus for the TMVI procedure. Due to a relatively vertical orientation of the mitral annulus, very cranio-caudal angulations have been described to be necessary.⁵⁶

Transcatheter valve interventions of the right heart
With the success of transcatheter therapies for aortic and mitral valve disease, percutaneous treatments have subsequently been developed for severe tricuspid regurgitation.⁵⁹ Tricuspid valve anatomy and its relationships with surrounding structures, the aetiology and severity of regurgitation, right ventricular anatomy and function are key aspects to assess in patients considered for transcatheter tricuspid valve therapy. CT has been shown to accurately assess tricuspid annulus geometry and dimensions.⁶⁰ For measurements of the right ventricle, an excellent correlation with cardiac MRI has been demonstrated.⁶¹ Depending on the type of interventional device used, specific anatomical aspects have to be assessed preprocedurally to minimize complication

risk. Implantation of transcatheter valves into the caval veins, so-called heterotopic transcatheter valve implantation, requires the measurement of caval vein diameters at the cavoatrial junction for device sizing and the assessment of the distance between the inferior cavoatrial junction to the first hepatic vein in order to avoid vein obstruction by the device.⁵⁹ For transcatheter tricuspid annuloplasty, evaluation of the course of the right coronary artery along the atrioventricular groove and the distance to the tricuspid annulus is crucial to avoid impingement during implantation of the suture pledgets and anchoring systems.⁶² Preprocedural CT imaging for implantation of a spacer device, designed to reduce the leaflet coaptation gap caused by tricuspid annular dilatation and leaflet tethering, includes the measurement of the tricuspid valve annulus and right ventricular dimensions, the distance from the tricuspid annulus plane to the right ventricular septal free wall groove as well as the luminal diameters of the left subclavian and axillary veins.⁵⁹ When transcatheter valve implantation into a surgically placed ring or prosthesis is considered, the valve size can be selected based on the sewing ring's nominal size or on the measurements of the reconstructed CT cross-sectional views of the tricuspid valve prosthesis or ring.⁵⁹

Whereas transcatheter pulmonary valve replacement has revolutionized pediatric cardiology by using a catheter-based rather than surgical approach in many congenital heart disease patients, it plays only a minor role in adult cardiology. As radiation may be an issue for this predominately young patient cohort, echocardiography and cardiac magnetic resonance tomography are the common imaging modalities used for procedure planning.⁶³ Valve anatomy, right ventricular outflow tract dimensions and the distance of the coronary arteries from the outflow tract are important parameters to be assessed preprocedurally.⁶⁴

Left atrial appendage occlusion

The left atrial appendage (LAA) is the origin of thrombus formation in about 90% of patients who develop atrial thrombi in the

context of atrial fibrillation.⁶⁵ LAA morphology is highly variable. Four main configurations have been described (“chicken wing”, “windsock”, “cauliflower”, and “cactus”), with the “chicken wing” morphology reported to have the lowest risk for thrombus formation.⁶⁶ Occlusion of the LAA provides effective stroke prevention and is clinically used in patients with a high bleeding risk on anticoagulation therapy.⁶⁵ Several percutaneous LAA closure devices have been developed (with the WATCHMAN™ device the only one to have FDA approval).^{67,68} Correct sizing of the LAA is crucial for successful and safe device implantation. Serious complications have been reported in 4% of cases and include hemodynamically relevant pericardial effusion, severe bleeding, device embolization or stroke.⁶⁹ Pre-procedural imaging is commonly limited to transesophageal echocardiography but some operators also use CT. CT analysis, in this case, includes reporting of LAA ostial diameters and LAA depth, LAA morphology including the number of lobes, as well as the assessment of the anatomic relation to surrounding structures (pulmonary veins, mitral annulus, left circumflex artery) for possible interference with the deployed occluder. LAA dimensions are highly dependent on the patient’s volume status. It is therefore recommended to hydrate the patient before imaging (usually 500–1000 ml saline solution) in order to achieve comparable filling conditions to the implantation procedure (left atrial pressure >12 mmHg).⁷⁰ As LAA occlusion devices are delivered transvenously and through the interatrial septum, identification of atrial septal aneurysms and of septal lipomatous hypertrophy is important. The presence of thrombi in the LAA must be reported.^{71,72} It should be noted that filling of the LAA with contrast is often incomplete, potentially mimicking the presence of a thrombus. A delayed second acquisition 60 sec after the initial contrast bolus may therefore be useful in order to identify thrombi. The method has been reported to have 100% sensitivity and 99% specificity for LAA thrombus detection (Figure 5).⁷³

In order to measure LAA dimensions using two-dimensional (2D) CT images, oblique views are superior over orthogonal views in the horizontal, coronal, and sagittal plane, since they better account for the typically elliptical shape of the LAA orifice.⁶⁶ While 3-dimensional CT reconstructions may offer a more comprehensive visualization of the LAA, superiority in sizing over 2D oblique imaging could not be demonstrated.⁶⁶ The LAA orifice is defined as the plane connecting the pulmonary vein ridge superiorly to the inferior junction of the left atrium and the LAA at the level of the circumflex artery.⁷⁴ An important aspect to take into consideration is the fact that the “landing zone” of the occlusion device varies between different occluder types^{72,74}. The WATCHMAN™ device is placed approximately 10–20 mm distal to the LAA ostium (Figure 6). As the sheath-constrained length of the WATCHMAN™ device is very similar to its unconstrained diameter once deployed, the length of the appendage along a clear line of sight from the proposed landing zone should be at least as long as the diameter of the device selected.⁷² The AMPLATZER Amulet device, consisting of a self-expanding nitinol mesh forming a lobe and disk, is implanted with the lobe anchoring approximately 10 mm inside of the LAA orifice.⁷⁵

Periprocedural guidance by transesophageal echocardiography is common. However, one has to be aware that underestimation of the LAA ostium diameter by 2D-transesophageal echocardiography compared to CT-based measurements can easily occur if the plane is not exactly aligned with the largest diameter of the LAA orifice.⁷⁶

Paravalvular leak occlusion

Paravalvular leaks are a common complication of surgical (incidence of relevant leak: 2–18%) and interventional prosthetic valve replacement (11–12%) and are associated with increased mortality.^{77–81} Aortic paravalvular leaks are most commonly

Figure 5. Detection of left atrial appendage thrombus by CT angiography. (A) First pass CT angiogram demonstrating a filling defect (white arrow) in the left atrial appendage suggestive of a thrombus. (B) Delayed phase CT angiogram acquired 60 sec after contrast application confirms the presence of a thrombus (white arrow).

Figure 6. Measurement of left atrial appendage ostium. (A) The plane of the LAA orifice is formed by the pulmonary vein ridge superiorly to the inferior junction of the left atrium and the LAA at the level of the circumflex artery. In a transaxial plane, the left circumflex coronary artery is identified (white arrow). Perpendicular to the left circumflex (white line), an oblique section is created which produces a vertical long-axis view of the heart (B) with the left circumflex coronary artery (white arrow) displayed in cross-section below the left atrial appendage (LAA) ostium. (C) Depending on the applied device system, the landing zone of the occluder varies. For a WATCHMAN™ occluder, the plane of implantation is identified approximately 10–20 mm (*, here 15 mm) from the ostium (dashed line) and measured between the left circumflex coronary artery and the edge of the landing zone (blue line). (D) Short axis of the plane of implantation for measurement of minimum (20 mm) and maximum (25 mm) diameter predicting a 27 mm WATCHMAN™ occluder. (E) Imaging of the implanted 27 mm WATCHMAN™ occluder.

located at the noncoronary cusp, while mitral paravalvular leaks are typically found in an anterolateral/posteromedial location.⁸² The lesser degree of invasiveness and convincing success rates favor a transcatheter approach for closure, with surgical repair being reserved for excessively large paravalvular leaks, contraindications to transcatheter closure, and unsuccessful procedures.^{83–85} Echocardiography is used to initially identify the presence and hemodynamic severity of paravalvular leaks but it can be difficult to assess the exact location, size and course of the sometimes serpiginous leaks. CT angiography can provide detailed information on the exact location and morphological characteristics of paravalvular leaks including size, course and shape as well as the presence of multiple communications.⁸⁶ Since metal components of the valve prosthesis can cause beam hardening artifacts, it is crucial to minimize motion artefacts during acquisition. It is therefore recommended to maximize image quality by using beta-blockers for heart rate control and to revert to systolic acquisition if heart rates remain >65 beats/min.^{87,88}

An acquisition covering the entire R-R-interval can demonstrate valve dehiscence with excess valve motion, to allow assessment of mechanical valve leaflet motion, and to reduce the risk for inaccurate assessment of a potential paravalvular leak tract.⁸⁶ For differentiation between contrast continuity and pledgets placed during surgery, an additional non-contrast scan should usually be considered. When assessing the image, adequate windowing is essential, a very soft window has been suggested to minimize beam hardening in prosthetic valves.⁸⁹ Based on our experience, we start with the set up *window 1500 HU and center 400 HU* with subsequent windowing adjustments (Figure 7A). The CT orientation for identifying the location of a paravalvular leak is demonstrated in Figure 7B–H. To report the location of the paravalvular leak, it has been recommended to use the simple triangulation method based on anatomic relationships of the atrial septum, LAA, and aortic valve with anatomically correct terminology (anterior vs posterior, superior vs inferior and lateral vs

Figure 7. (A) Paravalvular leak (arrow) in a patient with a prosthetic aortic valve. Windowing is set to window 1500 HU and center 400 HU to minimize beam hardening effects. (B-H) A 53-year-old male patient with a paravalvular leak 8 months after bioprosthetic aortic valve replacement. (B,C,D) Pre-procedural CT angiography shows the presence of a paravalvular leak (white arrow) between the aortic valve prosthesis (Edwards CE Perimount 23 mm) and the noncoronary cusp. (E) CT-derived measurement in a double oblique plane revealed a 0.48×0.73 cm leak. The purple line corresponds to the desired plane in (F) which is orthogonal to the prosthetic valve plane (RAO 14°/caudal 6°) and was used during the implantation procedure for C-arm angulation (G). (G) Invasive angiogram of the implanted 7 mm occluder (Ductus arteriosus occluder Hyperion PDA-I 07) (white arrow) in RAO 14° and caudal 6° angulation. (H) Device enhancement imaging of the implanted occluder (white arrow).

medial).^{86,90} Furthermore, assessment of the shape and size of the leak may be helpful for device selection.

Percutaneous revascularization of coronary chronic total occlusions

A coronary total occlusion (CTO) is defined as a coronary artery with complete interruption of the antegrade blood flow exceeding 3 months duration as proven by prior angiography or estimated from the clinical course.⁹¹ Percutaneous revascularization of CTOs is recommended in patients with angina resistant to medical therapy or with a large area of documented ischaemia in the territory of the occluded vessel.⁹² However, it is a technically challenging procedure with only moderate success rates which is complicated by the fact that visualization of the anatomic and morphological features of the occluded segment in invasive angiography is incomplete.⁹³ For this reason, coronary CTA can be particularly advantageous adjunct to invasive angiography for the visualization of long and tortuous CTO lesions, the detection of multiple occlusion sites and ostial occlusions and when the distal CTO segment cannot be visualized in invasive angiography.^{94–97} Scores such as the CT-RECTOR score permit prediction of procedural success rates. The CT-RECTOR score includes 6 CTA-based and clinical parameters: the presence of multiple occlusions, a blunt stump, severe calcification, bending of the occluded segment, a CTO age ≥ 12 months, and previously

failed PCI,⁹⁸ assigning one point for each parameter and adding them per CTO lesion. The likelihood of successful guidewire crossing within 30 min ranges from 95% for a CT-RECTOR score of 0 to 22% for a score of 3.

Furthermore, the CTO intervention itself may be facilitated by coronary CTA. Higher CTO intervention success rates in comparison to non-CTA guidance have been reported when CTA datasets were directly imported to the catheterization laboratory with the possibility of automatic alignment according to the angulation of the C-arm.⁹⁹ CT guidance by real-time automated co-registration of coronary CTA centreline and calcification onto live fluoroscopic images has been shown to be feasible.¹⁰⁰

CONCLUSION

In summary, CT is an increasingly important imaging tool for the expanding field of structural cardiac disease interventions. Current indications for pre-procedural CT imaging in cardiac interventional procedures include transcatheter aortic valve implantation, percutaneous interventions of the mitral, tricuspid and pulmonary valves, LAA occlusion, paravalvular leak occlusion as well as percutaneous revascularization of coronary chronic total occlusions. The most important information CT can provide concerns the access route, measurement of dimensions of the target structure and identification of an optimal viewing angulation.

REFERENCES

- Montalescot G, Sechtem U, Achenbach S, Andreotti F, Arden C, Budaj A, et al. 2013 ESC guidelines on the management of stable coronary artery disease: the task Force on the management of stable coronary artery disease of the European Society of cardiology. *Eur Heart J* 2013; **34**: 2949–3003. doi: <https://doi.org/10.1093/eurheartj/ehz296>
- Roffi M, Patrono C, Collet JP, Mueller C, Valgimigli M, Andreotti F, et al. ESC guidelines for the management of acute coronary syndromes in patients presenting without persistent ST-segment elevation. Task Force for the management of acute coronary syndromes in patients presenting without persistent ST-segment elevation of the European Society of cardiology (ESC). *G Ital Cardiol* 2016; **17**: 831–72.
- Al-Mallah MH, Aljizeeri A, Villines TC, Srichai MB, Alsaleek A. Cardiac computed tomography in current cardiology guidelines. *J Cardiovasc Comput Tomogr* 2015; **9**: 514–23. doi: <https://doi.org/10.1016/j.jcct.2015.09.003>
- Schoenhagen P, Numburi U, Halliburton SS, Aulbach P, von Roden M, Desai MY, et al. Three-dimensional imaging in the context of minimally invasive and transcatheter cardiovascular interventions using multi-detector computed tomography: from pre-operative planning to intra-operative guidance. *Eur Heart J* 2010; **31**: 2727–40. doi: <https://doi.org/10.1093/eurheartj/ehq302>
- Hell M, Marwan M, Gaede L, Achenbach S. Software innovations in computed tomography for structural heart disease interventions. *EuroIntervention* 2016; **12** Suppl X(Suppl X): X68–X74. doi: <https://doi.org/10.4244/EIJV12SXA13>
- Wuest W, Anders K, Schuhbaeck A, May MS, Gauss S, Marwan M, et al. Dual source multidetector CT-angiography before transcatheter aortic valve implantation (TAVI) using a high-pitch spiral acquisition mode. *Eur Radiol* 2012; **22**: 51–8. doi: <https://doi.org/10.1007/s00330-011-2233-0>
- Kok M, Turek J, Muhl C, Reinartz SD, Gohmann RF, Nijssen EC, et al. Low contrast media volume in pre-TAVI CT examinations. *Eur Radiol* 2016; **26**: 2426–35. doi: <https://doi.org/10.1007/s00330-015-4080-x>
- Baumgartner H, Falk V, Bax JJ, De Bonis M, Hamm C, Holm PJ, et al. 2017 ESC/EACTS guidelines for the management of valvular heart disease. *Eur Heart J* 2017; **38**: 2739–91. doi: <https://doi.org/10.1093/eurheartj/ehx391>
- Achenbach S, Delgado V, Hausleiter J, Schoenhagen P, Min JK, Leipsic JA. SCCT expert consensus document on computed tomography imaging before transcatheter aortic valve implantation (TAVI)/transcatheter aortic valve replacement (TAVR). *J Cardiovasc Comput Tomogr* 2012; **6**: 366–80. doi: <https://doi.org/10.1016/j.jcct.2012.11.002>
- Suchá D, Tuncay V, Prakken NHJ, Leiner T, van Ooijen PMA, Oudkerk M, et al. Does the aortic annulus undergo conformational change throughout the cardiac cycle? A systematic review. *Eur Heart J Cardiovasc Imaging* 2015; **16**: jev210–7. doi: <https://doi.org/10.1093/ehjci/jev210>
- Murphy DT, Blanke P, Alaamri S, Naoum C, Rubinshtein R, Pache G, et al. Dynamism of the aortic annulus: effect of diastolic versus systolic CT annular measurements on device selection in transcatheter aortic valve replacement (TAVR). *J Cardiovasc Comput Tomogr* 2016; **10**: 37–43. doi: <https://doi.org/10.1016/j.jcct.2015.07.008>
- Piazza N, de Jaegere P, Schultz C, Becker AE, Serruys PW, Anderson RH. Anatomy of the aortic valvar complex and its implications for transcatheter implantation of the aortic valve. *Circ Cardiovasc Interv* 2008; **1**: 74–81. doi: <https://doi.org/10.1161/CIRCINTERVENTIONS.108.780858>
- Zamorano JL, Badano LP, Bruce C, Chan K-L, Gonçalves A, Hahn RT, et al. EAE/ASE recommendations for the use of echocardiography in new transcatheter interventions for valvular heart disease. *Eur Heart J* 2011; **32**: 2189–214. doi: <https://doi.org/10.1093/eurheartj/ehz259>
- Willson AB, Webb JG, Labounty TM, Achenbach S, Moss R, Wheeler M, et al. 3-dimensional aortic annular assessment by multidetector computed tomography predicts moderate or severe paravalvular regurgitation after transcatheter aortic valve replacement: a multicenter retrospective analysis. *J Am Coll Cardiol* 2012; **59**: 1287–94. doi: <https://doi.org/10.1016/j.jacc.2011.12.015>
- Bax JJ, Delgado V, Bapat V, Baumgartner H, Collet JP, Erbel R, et al. Open issues in transcatheter aortic valve implantation. Part 1: patient selection and treatment strategy for transcatheter aortic valve implantation. *Eur Heart J* 2014; **35**: 2627–38. doi: <https://doi.org/10.1093/eurheartj/ehu256>
- Barbanti M, Yang T-H, Rodès Cabau J, Tamburino C, Wood DA, Jilaihawi H, et al. Anatomical and procedural features associated with aortic root rupture during balloon-expandable transcatheter aortic valve replacement. *Circulation* 2013; **128**: 244–53. doi: <https://doi.org/10.1161/CIRCULATIONAHA.113.002947>
- Achenbach S, Schuhbaeck A, Min JK, Leipsic J. Determination of the aortic annulus plane in CT imaging—a step-by-step approach. *JACC Cardiovasc Imaging* 2013; **6**: 275–8. doi: <https://doi.org/10.1016/j.jcmg.2012.06.015>
- Schuhbaeck A, Achenbach S, Pflederer T, Marwan M, Schmid J, Nef H, et al. Reproducibility of aortic annulus measurements by computed tomography. *Eur Radiol* 2014; **24**: 1878–88. doi: <https://doi.org/10.1007/s00330-014-3199-5>
- Schmidkonz C, Marwan M, Klinghammer L, Mitschke M, Schuhbaeck A, Arnold M, et al. Interobserver variability of CT angiography for evaluation of aortic annulus dimensions prior to transcatheter aortic valve implantation (TAVI). *Eur J Radiol* 2014; **83**: 1672–8. doi: <https://doi.org/10.1016/j.ejrad.2014.06.001>
- Gopal A, Grayburn PA, Mack M, Chacon I, Kim R, Montenegro D, et al. Noncontrast 3D CMR imaging for aortic valve annulus sizing in TAVR. *JACC Cardiovasc Imaging* 2015; **8**: 375–8. doi: <https://doi.org/10.1016/j.jcmg.2014.11.011>
- Stortecky S, Heg D, Gloekler S, Wenaweser P, Windecker S, Buellesfeld L. Accuracy and reproducibility of aortic annulus sizing using a dedicated three-dimensional computed tomography reconstruction tool in patients evaluated for transcatheter aortic valve replacement. *EuroIntervention* 2014; **10**: 339–46. doi: <https://doi.org/10.4244/EIJV10I3A59>
- Watanabe Y, Morice M-C, Bouvier E, Leong T, Hayashida K, Lefèvre T, et al. Automated 3-dimensional aortic annular assessment by multidetector computed tomography in transcatheter aortic valve implantation. *JACC Cardiovasc Interv* 2013; **6**: 955–64. doi: <https://doi.org/10.1016/j.jcin.2013.05.008>
- Van Linden A, Kempfert J, Blumenstein J, Möllmann H, Kim W-K, Alkaya S, et al. Manual versus automatic detection of aortic annulus plane in a computed tomography scan for transcatheter aortic valve implantation screening. *Eur J Cardiothorac Surg* 2014; **46**: 207–12 discussion 12. doi: <https://doi.org/10.1093/ejcts/ezt600>
- Lou J, Obuchowski NA, Krishnaswamy A, Popovic Z, Flamm SD, Kapadia SR, et al. Manual, semiautomated, and fully

- automated measurement of the aortic annulus for planning of transcatheter aortic valve replacement (TAVR/TAVI): analysis of interchangeability. *J Cardiovasc Comput Tomogr* 2015; **9**: 42–9. doi: <https://doi.org/10.1016/j.jcct.2014.11.003>
25. Blanke P, Spira EM, Ionasec R, Meinel FG, Ebersberger U, Scheuering M, et al. Semiautomated quantification of aortic annulus dimensions on cardiac CT for TAVR. *JACC Cardiovasc Imaging* 2014; **7**: 320–2. doi: <https://doi.org/10.1016/j.jcmg.2013.11.005>
 26. Van Belle E, Juthier F, Susen S, Vincentelli A, Lung B, Dallongeville J, et al. Postprocedural aortic regurgitation in balloon-expandable and self-expandable transcatheter aortic valve replacement procedures: analysis of predictors and impact on long-term mortality: insights from the FRANCE2 registry. *Circulation* 2014; **129**: 1415–27. doi: <https://doi.org/10.1161/CIRCULATIONAHA.113.002677>
 27. Jilaihawi H, Makkar RR, Kashif M, Okuyama K, Chakravarty T, Shiota T, et al. A revised methodology for aortic-valvar complex calcium quantification for transcatheter aortic valve implantation. *Eur Heart J Cardiovasc Imaging* 2014; **15**: 1324–32. doi: <https://doi.org/10.1093/ehjci/jeu162>
 28. Marwan M, Achenbach S, Ensminger SM, Pflederer T, Ropers D, Ludwig J, et al. CT predictors of post-procedural aortic regurgitation in patients referred for transcatheter aortic valve implantation: an analysis of 105 patients. *Int J Cardiovasc Imaging* 2013; **29**: 1191–8. doi: <https://doi.org/10.1007/s10554-013-0197-7>
 29. Watanabe Y, Lefèvre T, Arai T, Hayashida K, Bouvier E, Hovasse T, et al. Can we predict postprocedural paravalvular leak after Edwards SAPIEN transcatheter aortic valve implantation? *Catheter Cardiovasc Interv* 2015; **86**: 144–51. doi: <https://doi.org/10.1002/ccd.25665>
 30. Masri A, Schoenhagen P, Svensson L, Kapadia SR, Griffin BP, Tuzcu EM, et al. Dynamic characterization of aortic annulus geometry and morphology with multimodality imaging: predictive value for aortic regurgitation after transcatheter aortic valve replacement. *J Thorac Cardiovasc Surg* 2014; **147**: 1847–54. doi: <https://doi.org/10.1016/j.jtcvs.2013.05.047>
 31. Gnreux P, Head SJ, Van Mieghem NM, Kodali S, Kirtane AJ, Xu K, et al. Clinical outcomes after transcatheter aortic valve replacement using valve academic research Consortium definitions: a weighted meta-analysis of 3,519 patients from 16 studies. *J Am Coll Cardiol* 2012; **59**: 2317–26. doi: <https://doi.org/10.1016/j.jacc.2012.02.022>
 32. Ribeiro HB, Webb JG, Makkar RR, Cohen MG, Kapadia SR, Kodali S, et al. Predictive factors, management, and clinical outcomes of coronary obstruction following transcatheter aortic valve implantation: insights from a large multicenter registry. *J Am Coll Cardiol* 2013; **62**: 1552–62. doi: <https://doi.org/10.1016/j.jacc.2013.07.040>
 33. Barbanti M. Avoiding Coronary Occlusion and Root Rupture in TAVI - The Role of Pre-procedural Imaging and Prosthesis Selection. *Interv Cardiol* 2015; **10**: 94–7. doi: <https://doi.org/10.15420/icr.2015.10.2.94>
 34. Tops LF, Wood DA, Delgado V, Schuijff JD, Mayo JR, Pasupati S, et al. Noninvasive evaluation of the aortic root with multislice computed tomography implications for transcatheter aortic valve replacement. *JACC Cardiovasc Imaging* 2008; **1**: 321–30. doi: <https://doi.org/10.1016/j.jcmg.2007.12.006>
 35. Kasel AM, Cassese S, Leber AW, von Scheidt W, Kastrati A. Fluoroscopy-guided aortic root imaging for TAVR: "follow the right cusp" rule. *JACC Cardiovasc Imaging* 2013; **6**: 274–5. doi: <https://doi.org/10.1016/j.jcmg.2012.06.014>
 36. Gurvitch R, Wood DA, Leipsic J, Tay E, Johnson M, Ye J, et al. Multislice computed tomography for prediction of optimal angiographic deployment projections during transcatheter aortic valve implantation. *JACC Cardiovasc Interv* 2010; **3**: 1157–65. doi: <https://doi.org/10.1016/j.jcin.2010.09.010>
 37. Arnold M, Achenbach S, Pfeiffer I, Ensminger S, Marwan M, Einhaus F, et al. A method to determine suitable fluoroscopic projections for transcatheter aortic valve implantation by computed tomography. *J Cardiovasc Comput Tomogr* 2012; **6**: 422–8. doi: <https://doi.org/10.1016/j.jcct.2012.10.008>
 38. Hell MM, Biburger L, Marwan M, Schuhbaeck A, Achenbach S, Lell M, et al. Prediction of fluoroscopic angulations for transcatheter aortic valve implantation by CT angiography: influence on procedural parameters. *Eur Heart J Cardiovasc Imaging* 2017; **18**: 906–14. doi: <https://doi.org/10.1093/ehjci/jeu144>
 39. Samim M, Stella PR, Agostoni P, Kluin J, Ramjankhan F, Budde RPJ, et al. Automated 3D analysis of pre-procedural MDCT to predict annulus plane angulation and C-arm positioning: benefit on procedural outcome in patients referred for TAVR. *JACC Cardiovasc Imaging* 2013; **6**: 238–48. doi: <https://doi.org/10.1016/j.jcmg.2012.12.004>
 40. Robbins JD, Maniar PB, Cotts W, Parker MA, Bonow RO, Gheorghiadu M. Prevalence and severity of mitral regurgitation in chronic systolic heart failure. *Am J Cardiol* 2003; **91**: 360–2. doi: [https://doi.org/10.1016/S0002-9149\(02\)03172-7](https://doi.org/10.1016/S0002-9149(02)03172-7)
 41. Bursi F, Enriquez-Sarano M, Nkomo VT, Jacobsen SJ, Weston SA, Meverden RA, et al. Heart failure and death after myocardial infarction in the community: the emerging role of mitral regurgitation. *Circulation* 2005; **111**: 295–301. doi: <https://doi.org/10.1161/01.CIR.0000151097.30779.04>
 42. Blanke P, Naoom C, Webb J, Dvir D, Hahn RT, Grayburn P, et al. Multimodality imaging in the context of transcatheter mitral valve replacement: establishing consensus among modalities and disciplines. *JACC Cardiovasc Imaging* 2015; **8**: 1191–208. doi: <https://doi.org/10.1016/j.jcmg.2015.08.004>
 43. Feldman T, Wasserman HS, Herrmann HC, Gray W, Block PC, Whitlow P, et al. Percutaneous mitral valve repair using the edge-to-edge technique: six-month results of the Everest phase I clinical trial. *J Am Coll Cardiol* 2005; **46**: 2134–40. doi: <https://doi.org/10.1016/j.jacc.2005.07.065>
 44. Feldman T, Kar S, Rinaldi M, Fail P, Hermiller J, Smalling R, et al. Percutaneous mitral repair with the MitraClip system: safety and midterm durability in the initial Everest (endovascular valve Edge-to-Edge repair study) cohort. *J Am Coll Cardiol* 2009; **54**: 686–94. doi: <https://doi.org/10.1016/j.jacc.2009.03.077>
 45. Magruder JT, Crawford TC, Grimm JC, Fredi JL, Shah AS. Managing mitral regurgitation: focus on the MitraClip device. *Med Devices* 2016; **9**: 53–60. doi: <https://doi.org/10.2147/MDER.S86645>
 46. Bapat V, Buellesfeld L, Peterson MD, Hancock J, Reineke D, Buller C, et al. Transcatheter mitral valve implantation (TMVI) using the Edwards foris device. *EuroIntervention* 2014; **10 Suppl U**(Suppl U): U120–U128. doi: <https://doi.org/10.4244/EIJV10SUA18>
 47. Herrmann HC, Maisano F. Transcatheter therapy of mitral regurgitation. *Circulation* 2014; **130**: 1712–22. doi: <https://doi.org/10.1161/CIRCULATIONAHA.114.009881>
 48. Moat N, Duncan A, Lindsay A, Quarto C, Blanke P, Leipsic J, et al. Transcatheter mitral valve replacement for the treatment of mitral regurgitation: in-hospital outcomes of an apically tethered device. *J Am Coll Cardiol* 2015; **65**: 2352–3. doi: <https://doi.org/10.1016/j.jacc.2015.01.066>

49. Cheung A, Webb J, Verheye S, Moss R, Boone R, Leipsic J, et al. Short-term results of transapical transcatheter mitral valve implantation for mitral regurgitation. *J Am Coll Cardiol* 2014; **64**: 1814–9. doi: <https://doi.org/10.1016/j.jacc.2014.06.1208>
50. Sondergaard L, Brooks M, Ihlemann N, Jonsson A, Holme S, Tang M, et al. Transcatheter mitral valve implantation via transapical approach: an early experience. *Eur J Cardiothorac Surg* 2015; **48**: 873–8discussion 77–8. doi: <https://doi.org/10.1093/ejcts/ezu546>
51. Shanks M, Delgado V, Ng ACT, van der Kley F, Schuijff JD, Boersma E, et al. Mitral valve morphology assessment: three-dimensional transesophageal echocardiography versus computed tomography. *Ann Thorac Surg* 2010; **90**: 1922–9. doi: <https://doi.org/10.1016/j.athoracsur.2010.06.116>
52. Blanke P, Dvir D, Cheung A, Ye J, Levine RA, Precious B, et al. A simplified D-shaped model of the mitral annulus to facilitate CT-based sizing before transcatheter mitral valve implantation. *J Cardiovasc Comput Tomogr* 2014; **8**: 459–67. doi: <https://doi.org/10.1016/j.jcct.2014.09.009>
53. Blanke P, Naoum C, Dvir D, Bapat V, Ong K, Muller D, et al. Predicting LVOT Obstruction in Transcatheter Mitral Valve Implantation: Concept of the Neo-LVOT. *JACC Cardiovasc Imaging* 2017; **10**: 482–5. doi: <https://doi.org/10.1016/j.jcmg.2016.01.005>
54. Kohli K, Wei ZA, Yoganathan AP, Oshinski JN, Leipsic J, Blanke P. Transcatheter mitral valve planning and the Neo-LVOT: utilization of virtual simulation models and 3D printing. *Curr Treat Options Cardiovasc Med* 2018; **20**: 99. doi: <https://doi.org/10.1007/s11936-018-0694-z>
55. Feuchtner GM, Alkadhhi H, Karlo C, Sarwar A, Meier A, Dichtl W, et al. Cardiac CT angiography for the diagnosis of mitral valve prolapse: comparison with echocardiography. *Radiology* 2010; **254**: 374–83. doi: <https://doi.org/10.1148/radiol.2541090393>
56. Grover R, Ohana M, Arepalli CD, Sellers SL, Mooney J, Kueh S-H, et al. Role of MDCT imaging in planning mitral valve intervention. *Curr Cardiol Rep* 2018; **20**: 16. doi: <https://doi.org/10.1007/s11886-018-0960-4>
57. Deluca G, Correale M, Ieva R, Del Salvatore B, Gramenzi S, Di Biase M. The incidence and clinical course of caseous calcification of the mitral annulus: a prospective echocardiographic study. *J Am Soc Echocardiogr* 2008; **21**: 828–33. doi: <https://doi.org/10.1016/j.echo.2007.12.004>
58. Plank F, Al-Hassan D, Nguyen G, Raju R, Wheeler M, Thompson C, et al. Caseous calcification of the mitral annulus. *Cardiovasc Diagn Ther* 2013; **3**: E1–3. doi: <https://doi.org/10.3978/j.issn.2223-3652.2013.06.01>
59. Prihadi EA, Delgado V, Hahn RT, Leipsic J, Min JK, Bax JJ. Imaging Needs in Novel Transcatheter Tricuspid Valve Interventions. *JACC Cardiovasc Imaging* 2018; **11**: 736–54. doi: <https://doi.org/10.1016/j.jcmg.2017.10.029>
60. van Rosendaal PJ, Joyce E, Katsanos S, Debonnaire P, Kamperidis V, van der Kley F, et al. Tricuspid valve remodelling in functional tricuspid regurgitation: multidetector row computed tomography insights. *Eur Heart J Cardiovasc Imaging* 2016; **17**: 96–105. doi: <https://doi.org/10.1093/ehjci/jev140>
61. Maffei E, Messalli G, Martini C, Nieman K, Catalano O, Rossi A, et al. Left and right ventricle assessment with cardiac CT: validation study vs. cardiac Mr. *Eur Radiol* 2012; **22**: 1041–9. doi: <https://doi.org/10.1007/s00330-011-2345-6>
62. van Rosendaal PJ, Delgado V, Bax JJ. The tricuspid valve and the right heart: anatomical, pathological and imaging specifications. *EuroIntervention* 2015; **11** Suppl W(Suppl W): W123–W127. doi: <https://doi.org/10.4244/EIJV11SWA35>
63. Renapurkar RD, El-Sherief AH, Prieto L, Kapadia SR, Schoenhagen P. Transcatheter structural cardiac intervention: a radiology perspective. *AJR Am J Roentgenol* 2015; **204**: W648–W662. doi: <https://doi.org/10.2214/AJR.14.12571>
64. Holoshitz N, Hijazi ZM. Transcatheter pulmonary valve replacement: valves, techniques of implantation and outcomes. *Interv Cardiol* 2013; **5**: 465–77N. doi: <https://doi.org/10.2217/ica.13.36>
65. Kirchhof P, Benussi S, Kotecha D, Ahlsson A, Atar D, Casadei B, et al. 2016 ESC guidelines for the management of atrial fibrillation developed in collaboration with EACTS. *Eur Heart J* 2016; **37**: 2893–9622016. doi: <https://doi.org/10.1093/eurheartj/ehw210>
66. Wang Y, Di Biase L, Horton RP, Nguyen T, Morhanty P, Natale A. Left atrial appendage studied by computed tomography to help planning for appendage closure device placement. *J Cardiovasc Electrophysiol* 2010; **21**: 973–82. doi: <https://doi.org/10.1111/j.1540-8167.2010.01814.x>
67. Price MJ, Holmes DR. Mechanical closure devices for atrial fibrillation. *Trends Cardiovasc Med* 2014; **24**: 225–31. doi: <https://doi.org/10.1016/j.tcm.2014.06.001>
68. Meier B, Blaauw Y, Khattab AA, Lewalter T, Sievert H, Tondo C, et al. EHRA/EAPCI expert consensus statement on catheter-based left atrial appendage occlusion. *EuroIntervention* 2015; **10**: 1109–25. doi: https://doi.org/10.4244/EIJY14M09_18
69. Holmes DR, Kar S, Price MJ, Whisenant B, Sievert H, Doshi SK, et al. Prospective randomized evaluation of the Watchman left atrial appendage closure device in patients with atrial fibrillation versus long-term warfarin therapy: the prevail trial. *J Am Coll Cardiol* 2014; **64**: 1–1210.1016/j.jacc.2014.04.029. doi: <https://doi.org/10.1016/j.jacc.2014.04.029>
70. Spencer RJ, DeJong P, Fahmy P, Lempereur M, Tsang MYC, Gin KG, et al. Changes in left atrial appendage dimensions following volume loading during percutaneous left atrial appendage closure. *JACC Cardiovasc Interv* 2015; **8**: 1935–41. doi: <https://doi.org/10.1016/j.jcin.2015.07.035>
71. Kwong Y, Troupis J. Cardiac CT imaging in the context of left atrial appendage occlusion. *J Cardiovasc Comput Tomogr* 2015; **9**: 13–18. doi: <https://doi.org/10.1016/j.jcct.2014.11.005>
72. Ismail TF, Panikker S, Markides V, Foran JP, Padley S, Rubens MB, et al. CT imaging for left atrial appendage closure: a review and pictorial essay. *J Cardiovasc Comput Tomogr* 2015; **9**: 89–102. doi: <https://doi.org/10.1016/j.jcct.2015.01.011>
73. Romero J, Husain SA, Kelesidis I, Sanz J, Medina HM, Garcia MJ. Detection of left atrial appendage thrombus by cardiac computed tomography in patients with atrial fibrillation: a meta-analysis. *Circ Cardiovasc Imaging* 2013; **6**: 185–94. doi: <https://doi.org/10.1161/CIRCIMAGING.112.000153>
74. Wang DD, Eng M, Kupsky D, Myers E, Forbes M, Rahman M, et al. Application of 3-Dimensional Computed Tomographic Image Guidance to WATCHMAN Implantation and Impact on Early Operator Learning Curve: Single-Center Experience. *JACC Cardiovasc Interv* 2016; **9**: 2329–40. doi: <https://doi.org/10.1016/j.jcin.2016.07.038>
75. Saw J, Lempereur M. Percutaneous left atrial appendage closure: procedural techniques and outcomes. *JACC Cardiovasc Interv* 2014; **7**: 1205–20. doi: <https://doi.org/10.1016/j.jcin.2014.05.026>
76. Clemente A, Avoglierio F, Berti S, Paradossi U, Jamagidze G, Rezzaghi M, et al. Multimodality imaging in preoperative assessment of left atrial appendage transcatheter occlusion with the Amplatzer cardiac plug. *Eur Heart J Cardiovasc Imaging*

- 2015; **16**: 1276–87. doi: <https://doi.org/10.1093/ehjci/jev097>
77. Ionescu A, Fraser AG, Butchart EG. Prevalence and clinical significance of incidental paraprosthetic valvar regurgitation: a prospective study using transoesophageal echocardiography. *Heart* 2003; **89**: 1316–21. doi: <https://doi.org/10.1136/heart.89.11.1316>
 78. Nishimura RA, Otto CM, Bonow RO, Carabello BA, Erwin JP, Guyton RA, et al. 2014 AHA/ACC guideline for the management of patients with valvular heart disease: a report of the American College of Cardiology/American Heart Association Task Force on practice guidelines. *J Am Coll Cardiol* 2014; **63**: e57–1853rd. doi: <https://doi.org/10.1016/j.jacc.2014.02.536>
 79. Hwang HY, Choi J-W, Kim H-K, Kim K-H, Kim K-B, Ahn H. Paravalvular leak after mitral valve replacement: 20-year follow-up. *Ann Thorac Surg* 2015; **100**: 1347–52. doi: <https://doi.org/10.1016/j.athoracsur.2015.03.104>
 80. Smith CR, Leon MB, Mack MJ, Miller DC, Moses JW, Svensson LG, et al. Transcatheter versus surgical aortic-valve replacement in high-risk patients. *N Engl J Med Overseas Ed* 2011; **364**: 2187–98. doi: <https://doi.org/10.1056/NEJMoa1103510>
 81. Leon MB, Smith CR, Mack M, Miller DC, Moses JW, Svensson LG, et al. Transcatheter aortic-valve implantation for aortic stenosis in patients who cannot undergo surgery. *N Engl J Med Overseas Ed* 2010; **363**: 1597–607. doi: <https://doi.org/10.1056/NEJMoa1008232>
 82. Ruiz CE, Hahn RT, Berrebi A, Borer JS, Cutlip DE, Fontana G, et al. Clinical trial principles and endpoint definitions for Paravalvular leaks in surgical prosthesis: an expert statement. *J Am Coll Cardiol* 2017; **69**: 2067–87. doi: <https://doi.org/10.1016/j.jacc.2017.02.038>
 83. Koo HJ, Lee JY, Kim GH, Kang J-W, Kim Y-H, Kim D-H, et al. Paravalvular leakage in patients with prosthetic heart valves: cardiac computed tomography findings and clinical features. *Eur Heart J Cardiovasc Imaging* 2018; **19**: 1419–27. doi: <https://doi.org/10.1093/ehjci/jex341>
 84. Goel K, Eleid MF. Paravalvular leak in structural heart disease. *Curr Cardiol Rep* 2018; **20**: 18. doi: <https://doi.org/10.1007/s11886-018-0959-x>
 85. Ruiz CE, Jelnin V, Kronzon I, Dudy Y, Del Valle-Fernandez R, Einhorn BN, et al. Clinical outcomes in patients undergoing percutaneous closure of periprosthetic paravalvular leaks. *J Am Coll Cardiol* 2011; **58**: 2210–7. doi: <https://doi.org/10.1016/j.jacc.2011.03.074>
 86. Lesser JR, Han BK, Newell M, Schwartz RS, Pedersen W, Sorajja P. Use of cardiac CT angiography to assist in the diagnosis and treatment of aortic prosthetic paravalvular leak: a practical guide. *J Cardiovasc Comput Tomogr* 2015; **9**: 159–64. doi: <https://doi.org/10.1016/j.jcct.2014.11.001>
 87. Mahabadi AA, Achenbach S, Burgstahler C, Dill T, Fischbach R, Knez A, et al. Safety, efficacy, and indications of beta-adrenergic receptor blockade to reduce heart rate prior to coronary CT angiography. *Radiology* 2010; **257**: 614–23. doi: <https://doi.org/10.1148/radiol.10100140>
 88. Leipsic J, LaBounty TM, Ajlan AM, Earls JP, Strovski E, Madden M, et al. A prospective randomized trial comparing image quality, study Interpretability, and radiation dose of narrow acquisition window with widened acquisition window protocols in prospectively ECG-triggered coronary computed tomography angiography. *J Cardiovasc Comput Tomogr* 2013; **7**: 18–24. doi: <https://doi.org/10.1016/j.jcct.2013.01.001>
 89. O'Neill AC, Martos R, Murtagh G, Ryan ER, McCreery C, Keane D, et al. Practical tips and tricks for assessing prosthetic valves and detecting paravalvular regurgitation using cardiac CT. *J Cardiovasc Comput Tomogr* 2014; **8**: 323–7. doi: <https://doi.org/10.1016/j.jcct.2014.07.001>
 90. Rihal CS, Sorajja P, Booker JD, Hagler DJ, Cabalka AK. Principles of percutaneous paravalvular leak closure. *JACC Cardiovasc Interv* 2012; **5**: 121–30. doi: <https://doi.org/10.1016/j.jcin.2011.11.007>
 91. Sianos G, Werner GS, Galassi AR, Papafaklis MI, Escaned J, Hildick-Smith D, et al. Recanalisation of chronic total coronary occlusions: 2012 consensus document from the EuroCTO Club. *EuroIntervention* 2012; **8**: 139–45. doi: <https://doi.org/10.4244/EIJV8I1A21>
 92. Neumann FJ, Sousa-Uva M, Ahlsson A, Alfonso F, Banning AP, Benedetto U, et al. ESC/EACTS guidelines on myocardial revascularization. *Eur Heart J* 2018; **2018**.
 93. Grantham JA, Marso SP, Spertus J, House J, Holmes DR, Rutherford BD. Chronic total occlusion angioplasty in the United States. *JACC Cardiovasc Interv* 2009; **2**: 479–86. doi: <https://doi.org/10.1016/j.jcin.2009.02.008>
 94. Magro M, Schultz C, Simsek C, Garcia-Garcia HM, Regar E, Nieman K, et al. Computed tomography as a tool for percutaneous coronary intervention of chronic total occlusions. *EuroIntervention* 2010; **6 Suppl G** (6 Suppl): G123–31G:G123–31. doi: <https://doi.org/10.4244/EIJV4I5A102>
 95. Opolski MP, Achenbach S. CT Angiography for Revascularization of CTO: Crossing the Borders of Diagnosis and Treatment. *JACC Cardiovasc Imaging* 2015; **8**: 846–58. doi: <https://doi.org/10.1016/j.jcmg.2015.05.001>
 96. García-García HM, van Mieghem CAG, Gonzalo N, Meijboom WB, Weustink AC, Onuma Y, et al. Computed tomography in total coronary occlusions (CTTO registry): radiation exposure and predictors of successful percutaneous intervention. *EuroIntervention* 2009; **4**: 607–16. doi: <https://doi.org/10.4244/EIJV4I5A102>
 97. Opolski MP, Kepka C, Achenbach S, Juraszynski Z, Pregowski J, Kruk M, et al. Coronary computed tomographic angiography for prediction of procedural and intermediate outcome of bypass grafting to left anterior descending artery occlusion with failed visualization on conventional angiography. *Am J Cardiol* 2012; **109**: 1722–8. doi: <https://doi.org/10.1016/j.amjcard.2012.02.015>
 98. Opolski MP, Achenbach S, Schuhbäck A, Rolf A, Möllmann H, Nef H, et al. Coronary computed tomographic prediction rule for time-efficient guidewire crossing through chronic total occlusion: insights from the CT-RECTOR multicenter registry (computed tomography Registry of chronic total occlusion revascularization). *JACC Cardiovasc Interv* 2015; **8**: 257–67. doi: <https://doi.org/10.1016/j.jcin.2014.07.031>
 99. Rolf A, Werner GS, Schuhbäck A, Rixe J, Möllmann H, Nef HM, et al. Preprocedural coronary CT angiography significantly improves success rates of PCI for chronic total occlusion. *Int J Cardiovasc Imaging* 2013; **29**: 1819–27. doi: <https://doi.org/10.1007/s10554-013-0258-y>
 100. Ghoshhajra BB, Takx RAP, Stone LL, Girard EE, Brilakis ES, Lombardi WL, et al. Real-time fusion of coronary CT angiography with X-ray fluoroscopy during chronic total occlusion PCI. *Eur Radiol* 2017; **27**: 2464–73. doi: <https://doi.org/10.1007/s00330-016-4599-5>