Published in final edited form as:

Am J Obstet Gynecol. 2019 July ; 221(1): 41.e1–41.e7. doi:10.1016/j.ajog.2019.03.004.

Pelvic organ prolapse as a function of levator ani avulsion, hiatus size, and strength

Victoria L. HANDA, MD MHS,

Johns Hopkins School of Medicine, Baltimore, MD

Jennifer ROEM, MS,

Johns Hopkins Bloomberg School of Public Health, Baltimore, MD

Joan L. BLOMQUIST, MD,

Greater Baltimore Medical Center, Baltimore, MD

Hans Peter DIETZ, MD PhD, and

University of Sydney, Sydney, New South Wales, Australia

Alvaro MUÑOZ, PhD

Johns Hopkins Bloomberg School of Public Health, Baltimore, MD

STRUCTURED ABSTRACT

BACKGROUND: Obstetrical levator ani muscle avulsion is detected after 10–30% of vaginal deliveries and is associated with pelvic organ prolapse later in life. However, the mechanism by which levator avulsion may contribute to prolapse is unknown.

OBJECTIVES: This study investigated the extent by which size of the levator hiatus and pelvic muscle weakness may explain the association between levator avulsion and pelvic organ prolapse.

STUDY DESIGN: This was a supplementary study of a longitudinal cohort of parous women enrolled 5–10 years after first delivery and assessed annually for prolapse (defined as descent beyond the hymen) for up to 9 annual visits. For this sub-study, vaginally parous participants were assessed for levator avulsion using 3dimensional transperineal ultrasound. Ultrasound was performed at a median interval of 11 years from delivery. Ultrasound volumes were also used to measure levator hiatus area with Valsalva. Pelvic muscle strength was measured with perineometry. Women with and without pelvic organ prolapse were compared for levator avulsion, levator hiatus area, and pelvic muscle strength, using multivariable logistic regression yielding a measure of mediation. Bootstrap methods were used to calculate the confidence interval corresponding to the measure of mediation by hiatus area and pelvic muscle strength.

Corresponding author: Victoria L. Handa, MD MHS; 4940 Eastern Ave, Room A121; Baltimore, MD 21224-2780; (410) 550-0336; F (410) 550-0196; vhanda1@jhmi.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Presented at the 39th Annual Scientific Meeting of the American Urogynecologic Society, Chicago, IL, October 9-13, 2018

RESULTS: Prolapse was identified in 109/429 (25%) and was significantly associated with levator avulsion (odds ratio (OR) 4.17, 95% confidence interval (CI) 2.28, 7.31). Prolapse was also associated with levator hiatus area (OR 1.52 per 5cm², 95%CI 1.34, 1.73) and inversely with muscle strength (OR 0.87 per 5cm H₂O, 95%CI 0.81, 0.94). In a multivariable logistic model including levator avulsion, levator hiatus area, and strength, the association between levator avulsion and prolapse was substantially attenuated and indeed was no longer statistically significant (OR 1.75, 95%CI 0.91, 3.39). Hiatus area and strength mediated 61% (95% CI 34%, 106%) of the association between avulsion and prolapse. Furthermore, since the 95% confidence interval for this estimate contained 100%, it cannot be ruled out that the two markers fully mediate the effect of avulsion on prolapse.

CONCLUSIONS: The strong association between pelvic organ prolapse and levator avulsion can be explained to a large extent by a larger levator hiatus and weaker pelvic muscles after levator avulsion.

UNSTRUCTURED ABSTRACT

Obstetrical levator ani muscle avulsion is detected after 10–30% of vaginal deliveries and is associated with pelvic organ prolapse later in life. However, the mechanism by which levator avulsion may contribute to prolapse is unknown. This study investigated the extent by which size of the levator hiatus and pelvic muscle weakness may explain the association between levator avulsion and pelvic organ prolapse. This was a supplementary study of a longitudinal cohort of parous women enrolled 5–10 years after first delivery and assessed annually for prolapse (defined as descent beyond the hymen) for up to 9 annual visits. For this sub-study, vaginally parous participants were assessed for levator avulsion using 3dimensional transperineal ultrasound. Ultrasound was performed at a median interval of 11 years from delivery. Ultrasound volumes were also used to measure levator hiatus area with Valsalva. Pelvic muscle strength was measured with perineometry. Women with and without pelvic organ prolapse were compared for levator avulsion, levator hiatus area, and pelvic muscle strength, using multivariable logistic regression yielding a measure of mediation. Bootstrap methods were used to calculate the confidence interval corresponding to the measure of mediation by hiatus area and pelvic muscle strength. Prolapse was identified in 109/429 (25%) and was significantly associated with levator avulsion (odds ratio (OR) 4.17, 95% confidence interval (CI) 2.28, 7.31). Prolapse was also associated with levator hiatus area (OR 1.52 per 5cm², 95%CI 1.34, 1.73) and inversely with muscle strength (OR 0.87 per 5cm H₂O, 95%CI 0.81, 0.94). In a multivariable logistic model including levator avulsion, levator hiatus area, and strength, the association between levator avulsion and prolapse was substantially attenuated and indeed was no longer statistically significant (OR 1.75, 95%CI 0.91, 3.39). Hiatus area and strength mediated 61% (95% CI 34%, 106%) of the association between avulsion and prolapse. Furthermore, since the 95% confidence interval for this estimate contained 100%, it cannot be ruled out that the two markers fully mediate the effect of avulsion on prolapse. In conclusion, the strong association between pelvic organ prolapse and levator avulsion can be explained to a large extent by a larger levator hiatus and weaker pelvic muscles after levator avulsion.

Keywords

levator ani muscle avulsion; pelvic muscle strength; pelvic organ prolapse

INTRODUCTION

The probability that a woman will develop pelvic organ prolapse is significantly increased by vaginal childbirth [1]. Specifically, prior research suggests a 5-fold increase in the odds of pelvic organ prolapse 5–10 years after vaginal birth versus cesarean [2].

The reasons for this association are unclear, although one hypothesized mechanism involves obstetrical injury to the levator ani muscles. Recent studies have explored the relationship between obstetrical levator injury, levator hiatus size, levator strength, and prolapse [3–10]. Levator ani muscle avulsion, in which the levator muscle is detached from its origin at the pubis, is observed after 10–30% of vaginal deliveries [3,4,5] and is significantly associated with the development of prolapse later in life [6,7,8]. However, the biological mechanism for this strong association has not been explained. Levator avulsion appears to lead to a significantly larger levator hiatus and significantly poorer muscle strength [9]. It is unknown whether these changes in levator structure and function could explain the development of pelvic organ prolapse after vaginal childbirth.

The objective of this research was to determine the relative contributions of levator avulsion, the size of the levator hiatus, and levator contraction strength to pelvic organ prolapse after vaginal childbirth. Our first hypothesis was that levator hiatus area and pelvic muscle weakness are each independent and significant risk factors for pelvic organ prolapse. Our second hypothesis was that these changes in muscle structure and function explain the association between levator avulsion and prolapse. More specifically, we hypothesized that after controlling for levator weakness and hiatus size, the residual direct effect of levator avulsion on pelvic organ prolapse would be reduced. By addressing these hypotheses in a cohort of vaginally parous women, our goal was to provide insights into the mechanistic relationship between vaginal childbirth and pelvic organ prolapse.

MATERIALS AND METHODS

Data for this research were acquired in the setting of a longitudinal cohort study of parous women [2,10]. The methods for the parent study have been described previously [2,10]. Briefly, parous women were enrolled 5–10 years from first delivery and followed annually for the incidence of pelvic floor disorders. From May 2015 to April 2017, each vaginally parous study participant was offered participation in this supplementary study [8,9].

This supplementary study included the measurement of pelvic muscle strength and assessment of the levator ani muscle using 3dimensional transperineal ultrasound. Each woman was offered participation at one of her regularly scheduled study visits. Pelvic floor muscle strength (PFMS) was measured during voluntary pelvic muscle contraction with the Peritron perineometer [11, 12]. PFMS was defined as the average increase in pressure generated over two contraction attempts, both obtained after the participant had been coached in the contraction technique and after an investigator confirmed correct technique via palpation. Women reporting latex allergy did not participate in this aspect of the research.

To assess the levator ani muscle anatomy, ultrasound volumes were captured as cine loops using a GE Voluson s6 system, with a RAB2–6-RS convex transducer (General Electric Healthcare, Chicago, IL), held in the sagittal plane on the perineum. Volumes were obtained for each participant at rest, with maximum Valsalva and with pelvic floor muscle contraction. Levator avulsion was identified on tomographic images [13], obtained during maximum contraction. A complete levator ani avulsion was diagnosed if volumes demonstrated a complete separation between the levator muscle and the pubis ramus for at least three contiguous “slices”: at the plane of minimal hiatal dimension and for at least 5 mm above that level [13]. All suspected avulsions were confirmed independently by two investigators. We also measured the area of the levator hiatus with maximal Valsalva (in the plane of minimal hiatal dimension [14]), using the area tool provided by GE 4Dview software (General Electric Healthcare, Chicago, IL).

Prolapse was assessed annually in this cohort using the Pelvic Organ Prolapse Quantification (POPQ) examination [15]. Prolapse was defined as descent of the cervix or any vaginal segment beyond the hymen. In addition, participants were asked about prior surgery for pelvic organ prolapse. At the time of the ultrasound assessment, we calculated the cumulative incidence, defined as the proportion of women who had met criteria for prolapse (at any time prior to or at the time of the study ultrasound), either on physical examination or by a report of surgery for treatment of prolapse.

The database of the parent study provided additional participant data, including primary race, age (at the time of ultrasound), and body mass index (kg/m^2) measured at the time of ultrasound. Obstetrical data included maternal age at first vaginal delivery, any forceps-assisted birth, any deliveries with second stage of labor greater than 120 minutes, any deliveries with birth weight greater than 4 kilograms, and history of obstetrical anal sphincter laceration. These obstetrical data considered all deliveries for each woman up until the time of the ultrasound.

In addition, since pelvic muscle strength was characterized in this study, participants were asked whether they had been in a prior supervised program of pelvic muscle exercises.

The analytic plan for this study was influenced by prior work in this cohort demonstrating that levator avulsion is associated with a significantly larger levator hiatus and significantly poorer muscle strength (as suggested by lower peak perineometry pressures) [9]. A logistic regression model was developed to estimate the odds of avulsion as a function of levator hiatus area and PFMS. This model was then used to calculate each woman’s propensity of avulsion, based on her hiatus area and PFMS. Specifically, the log odds of avulsion was calculated for each woman and the interquartile range was calculated for the study population. Among women diagnosed with avulsion in the study population, according to criteria established by Tukey [16], outliers were defined as those with a log odds of avulsion below the first quartile by more than 1.5 times the interquartile range. These participants were considered outliers and were excluded from the primary statistical analyses, as their inclusion would be expected to distort statistical models. However, sensitivity analysis was also performed with the inclusion of the outliers to assess their influence on the study findings.

For the primary statistical analysis, the outcome of interest was pelvic organ prolapse; the exposures of interest were presence of levator avulsion, levator hiatus size, and PFMS. Logistic regression was employed to estimate the odds of pelvic organ prolapse as a function of each of the exposures. Specifically, the logarithm of the odds of pelvic organ prolapse was modeled using the following regression equation:

$$\alpha_0 + \alpha_1 \text{avulsion} + \alpha_2(\text{hiatus size} - 25)/5 + \alpha_3(\text{PFMS} - 30)/5$$

Given this regression equation, the crude univariable association of avulsion with prolapse is captured by α_1^* , defined as the value of α_1 resulting from imposing $\alpha_2 = 0$ and $\alpha_3 = 0$ (i.e., univariate association of avulsion and prolapse).

Our second hypothesis was that these changes in muscle structure and function explain the association between levator avulsion and prolapse. To assess the indirect effect of avulsion on prolapse (i.e., mediated by hiatus area and PFMS), we calculated $(\alpha_1^* - \alpha_1)/\alpha_1^*$. This statistic can be read as the percentage of the effect of avulsion on prolapse that is explained by (mediated via) the hiatus area and PFMS. To calculate confidence intervals for this statistic, we considered that estimates of α_1 and α_1^* are two statistics from regressions on the same data set. Thus, they are subjected to statistical dependencies for which a bootstrap approach offers a way to calculate standard errors. Specifically, to calculate a 95% confidence interval for $(\alpha_1^* - \alpha_1)/\alpha_1^*$ we produced 1000 bootstrap samples (i.e., sampling N times with replacement from the N participants of the total study population) and carried out the two logistic regressions (with and without the hiatus area and PFMS) to arrive to 1000 estimates of $(\alpha_1^* - \alpha_1)/\alpha_1^*$ so that the 2.5th and 97.5th percentiles define the 95% confidence interval.

In an additional statistical model, the population was stratified by levator avulsion in order to determine the role of levator hiatus and PFMS on prolapse in women with the same levator avulsion status. In the two resultant populations, we carried out two logistic regression models, considering prolapse as a function of levator hiatus area and PFMS.

Inferences and statistical comparisons were carried out based on maximum likelihood methods using SAS version 9.4 (Cary, NC).

RESULTS

Of 550 eligible women, 10 (2%) declined to participate in this supplemental study, 86 (16%) did not return for a visit during the study interval, 1 had uninterpretable ultrasound volumes, and 21 (4%) did not undergo perineometry (15 due to latex allergy, 6 for other reasons). Thus, 432 vaginally parous women contributed data for this research. At the time of these ultrasound studies, the median age was 43 years. Parity was equal to one for 22%, equal to 2 for 57% and equal to three or more for 21% of participants. Only 7 of 432 participants (2%) reported participation in a program of supervised exercises

Of these 432 women, 112 (25%) had prolapse, including seven with a history of prior surgery for prolapse (but no prolapse on examination) and 105 with prolapse beyond the hymen on physical examination (six of whom also reported a history of surgery for prolapse). Comparing women with and without prolapse (Table 1), there were no statistically significant differences in race, body mass index, macrosomia or prolonged second stage. However, compared to women without prolapse, those with prolapse were older, and were more likely to have experienced forceps delivery.

Levator avulsion was diagnosed among 64 of 432 (15%) women. Comparing women with versus those without levator avulsion (Figure 1), women with avulsion (black closed circles) had larger hiatus area (medians: 34.57 cm² versus 25.03 cm²; $p < 0.001$) and lower perineometry measures (medians: 13.5 cmH₂O versus 28.5 cmH₂O; $p < 0.001$) when compared to women without levator avulsion (gray open circles). The exceptions were the three cases shown in black closed diamonds. Although these three women had ultrasound findings that met diagnostic criteria for avulsion, their hiatus area and PFMS measures were very different from the rest of those with avulsion. This led to the question of whether these three cases should be included in further analyses. Using the method for identifying outliers established by Tukey [16], these three participants were confirmed as outliers. More specifically, using their levator hiatus and PFMS measures to predict their probability of avulsion, using $\{1 + \exp(3.52 - 0.56(\text{hiatus area} - 25)/5 + 0.61(\text{PFMS} - 30)/5)\}^{-1}$, we calculated that the probabilities these three women had an avulsion were 0.35%, 0.43% and 0.31%, respectively. Thus, assuming accurate measurement of hiatus area and PFMS, it was very unlikely these women had levator avulsion. Their data were excluded from further analysis, leaving 429 women for analysis, of whom 61 had levator avulsion and 109 had prolapse.

Among the remaining 429 women, prolapse was significantly associated with levator hiatus size (Figure 2, panel A). Across five ordinal categories of hiatus area size (from $<20\text{ cm}^2$ to 35 cm^2), the odds of prolapse increased more than 7-fold: the relative odds (OR) for prolapse among women with hiatus area 35 cm^2 versus $<20\text{ cm}^2$ was 7.22 (95% confidence interval (CI) 3.40, 15.32). Moreover, in an analysis that considered levator hiatus area as a continuous variable, the relative odds for prolapse was found to increase by 50% per 5cm² increase in hiatus size (OR 1.52, 95% CI 1.34, 1.73).

Prolapse also increased as PFMS decreased (Figure 2, panel B). For women with peak pressure $35\text{ cm H}_2\text{O}$, the odds of prolapse were halved (OR 0.45; 95% CI 0.26, 0.77) in comparison with women who had peak pressure $<20\text{ cmH}_2\text{O}$. Considering PFMS as a continuous variable, for every 5cmH₂O increase in peak pressure, the odds for prolapse was reduced by 13% (OR 0.87, 95% CI 0.81, 0.94).

In univariable logistic regression, the relative odds for prolapse associated with levator avulsion was 4.17 (95% CI 2.28, 7.31). Given the association between avulsion, levator hiatus size and PFMS depicted in Figure 1, multivariable logistic regression was next used to investigate whether levator hiatus area and pelvic muscle weakness were each independent and significant risk factors for pelvic organ prolapse (Table 2). In a multivariable model with

these three variables, the association between levator avulsion and prolapse was substantially attenuated and indeed was no longer statistically significant (OR 1.75, 95%CI 0.91, 3.39).

For this multivariable logistic regression model, we estimated that the percent of the association between avulsion and prolapse (OR= 4.17) mediated via hiatus area and PFMS was 61% ($= 100 * (\log(4.17) - \log(1.75))/\log(4.17)$). The bootstrap-based 95% confidence interval for this estimate was from 34% to 106%. By the lower bound of the confidence interval being above zero, this confirms there is significant mediation. By the upper bound being above 100%, the possibility that hiatus area and PFMS fully mediate the avulsion-prolapse association cannot be ruled out. We also considered separate models with hiatus area and PFMS as individual mediators of avulsion, which resulted in partial mediation of 38% (95%CI 22%, 68%) and 19% (95%CI 2%, 40%), respectively. Therefore, while hiatus area and muscle strength separately mediate a portion of the association between prolapse and avulsion, when considered together they may completely mediate the avulsion-prolapse association.

Table 2 also demonstrates the results of stratified multivariable regression models, in which women with and without levator avulsion were considered separately. Among 368 women without levator avulsion, the association between prolapse and both hiatus area and PFMS were statistically significant. Furthermore, the association between levator hiatus area and prolapse was strongest for women with levator avulsion. Specifically, among 61 women with levator avulsion, the odds of prolapse was more than doubled for each 5cm² increase in levator hiatus area (OR 2.43, 95%CI 1.51, 3.91).

Finally, in a sensitivity analysis that considered inclusion of the three outliers in the multivariable regression model, the impact of these three cases demonstrated their undue influence. When these three cases are included in the multivariable model, the OR for avulsion was increased (from 1.75 without them) to 2.27, and was statistically significant (95%CI 1.21, 4.29). In addition, the association between prolapse and PFMS increased towards the null (OR=0.93), being only marginally significant (95%CI 0.86, 1.00).

COMMENT

Our results confirm a strong association between prolapse and levator avulsion, previously reported by Dietz [7] and Delancey [6]. In addition, the present study extends those findings by demonstrating significant associations between prolapse and two characteristics seen among women with levator avulsion [9]: increased levator hiatus area and decreased muscle strength (as measured by the vaginal pressure generated during perineometry). With respect to levator hiatus size, for every 5cm² increase in area, the odds of prolapse was increased by 50% (OR 1.52, 95%CI 1.34, 1.73). This association is similar to an observation in prior study of care-seeking women, in whom hiatal area was associated with stage 2 support [17]. In addition, we found that PFMS was inversely and significantly associated with the odds of prolapse (OR 0.87, 95% CI 0.81, 0.94).

The results of the present study also suggests that the association between levator avulsion and prolapse might be explained, to a large extent (and possibly fully), by the increase in the

levator hiatus size and the reduction in PFMS seen among women with levator avulsion. This is an important finding, which leads us to speculate that levator hiatus size and pelvic muscle weakness may be the mechanism by which levator avulsion is linked to prolapse later in life. This hypothesis would be consistent with biomechanical models that suggest that levator impairment, and more specifically hiatus size, is a critical contributor to anterior vaginal prolapse [18]. This could suggest possible targets for prolapse prevention among vaginally parous women. However, the residual association between prolapse and levator avulsion in the multivariable model (OR 1.75), while not statistically significant, cannot be completely discounted. This might suggest some additional mechanism by which levator avulsion may be associated with prolapse.

An additional contribution of this study is the use of each woman's levator hiatus area and PFMS measures to validate her diagnosis of avulsion. Using an established method for identifying avulsion [13], 64 women were initially diagnosed with avulsion and this diagnosis was confirmed independently by two readers. However, using the levator hiatus area and PFMS, 3/64 (5%) women diagnosed with avulsion were considered unlikely to have had avulsion and were excluded from our analysis. This additional validation of the avulsion diagnosis, building on the rigorous use of validated criteria, is a strength of this study. Such validation could improve the diagnostic accuracy of the ultrasound diagnosis of avulsion for further research studies and could provide additional valuable diagnostic criteria for avulsion as the potential clinical value of this diagnosis is established.

Additional strengths of this study were the use of a rigorous definition of prolapse, as well as the masking of observers (reducing the potential for bias). Also, the sample size provided adequate power to address the proposed aims of this study.

Women in this study were relatively young (mid 40's) at the time of this study and our results might not be generalizable to older women. The focus on relatively young women was intentional, as the impact of childbirth is more likely to be manifested in women with the onset of prolapse at a relatively young age. Indeed, although the peak hazard for prolapse likely occurs more than 20 years after delivery, data from this study cohort suggest that the strong association between childbirth and the incidence of prolapse is sustained over time [19]. Nevertheless, it is possible that different mechanisms might contribute to prolapse that develops later in life.

A potential weakness of this study was perineometry may not accurately reflect pelvic muscle strength. While perineometry has a strong correlation with more traditional measures of levator strength, such as Brink score, and has demonstrated high inter- and intra-rater reliability [20], this technique may be influenced by a Valsalva effort during pelvic muscle contraction. An additional weakness of this study is that levator hiatus size and PFMS were assessed many years after childbirth. An argument for a causal relationship with prolapse would be stronger if these characteristics had been assessed prior to the development of prolapse. While we acknowledge this limitation, there is some reason to believe that PFMS, as assessed with perineometry, is relatively constant over time. For example, research in this population found that PFMS is essentially constant over 4 years of observation [21]. Thus, we believe that PFMS measured 11 years after childbirth is a reasonable approximation of

strength a decade before. Less is known about changes in levator hiatus area over time. In one study of 101 vaginally parous women, Chan and colleagues found an increase in mean levator hiatus area between 1 year and 3–5 years postpartum (14.24 cm² versus 16.71 cm², $p < 0.005$) [22]. Furthermore, they found that the increase in hiatal area was greater among 28 women with levator avulsion versus those without avulsion. These findings, together with the results of the present study, argue for further longitudinal studies of levator biometry and pelvic organ support over time.

The findings of the present study provide important clues about the sequelae of levator avulsion and the pathophysiology of prolapse among parous women. These results suggest that hiatus size and muscle strength may be important markers to identify women at greatest risk to develop prolapse. Further studies are needed to determine whether interventions to reduce hiatus size or increase muscle strength would be effective prevention strategies to reduce the incidence of prolapse later in life.

Disclosure:

Funded by the *Eunice Kennedy Shriver* National Institute of Child Health and Human Development (R01HD082070 and R01HD056275); VLH received an honorarium from UpToDate for published material on a related topic.

REFERENCES

1. Hallock JL, Handa VL. The Epidemiology of Pelvic Floor Disorders and Childbirth: An Update. *Obstet Gynecol Clin North Am*. 2016;43:1–13. [PubMed: 26880504]
2. Handa VL, Blomquist JL, Knoepp LR, Hoskey KA, McDermott KC, Muñoz A. Pelvic floor disorders 5–10 years after vaginal or cesarean childbirth. *Obstet Gynecol*. 2011;118:777–84 [PubMed: 21897313]
3. DeLancey JO, Kearney R, Chou Q, et al. The appearance of levator ani muscle abnormalities in magnetic resonance images after vaginal delivery. *Obstet Gynecol* 2003;101:46–53. [PubMed: 12517644]
4. Dietz HP, Lanzarone V. Levator trauma after vaginal delivery. *Obstet Gynecol* 2005;106:707–12. [PubMed: 16199625]
5. Caudwell-Hall J, Kamisan Atan I, Martin A, et al. Intrapartum predictors of maternal levator ani injury. *Acta Obstet Gynecol Scand* 2017;96:426–431. [PubMed: 28117880]
6. DeLancey JO, Morgan DM, Fenner DE, et al. Comparison of levator ani muscle defects and function in women with and without pelvic organ prolapse. *Obstet Gynecol* 2007;109:295–302. [PubMed: 17267827]
7. Dietz HP, Simpson JM. Levator trauma is associated with pelvic organ prolapse. *BJOG*. 2008;115:979–84. [PubMed: 18503571]
8. Handa VL, Blomquist JL, Roem J, Muñoz A, Dietz HP. Pelvic Floor Disorders After Obstetric Avulsion of the Levator Ani Muscle. *Female Pelvic Med Reconstr Surg*. 2019; 25:3–7. [PubMed: 30285979]
9. Handa VL, Blomquist JL, Roem J, Muñoz A, Dietz HP. Levator Morphology and Strength After Obstetric Avulsion of the Levator Ani Muscle. *Female Pelvic Med Reconstr Surg*. 2018 9 27 [Epub ahead of print]
10. Handa VL, Blomquist JL, Roem J, Muñoz A. Longitudinal study of quantitative changes in pelvic organ support among parous women. *Am J Obstet Gynecol*. 2018;218:320.e1–320.e7 [PubMed: 29277626]
11. Friedman S, Blomquist JL, Nugent JM, et al. Pelvic muscle strength after childbirth. *Obstet Gynecol* 2012;120:1021–8 [PubMed: 23090518]

12. Middlekauff ML, Egger MJ, Nygaard IE, Shaw JM. The impact of acute and chronic strenuous exercise on pelvic floor muscle strength and support in nulliparous healthy women. *Am J Obstet Gynecol.* 2016 9;215(3):316.e1–7 [PubMed: 26899909]
13. Dietz H, Bernardo M, Kirby A, et al. Minimal criteria for the diagnosis of avulsion of the puborectalis muscle by tomographic ultrasound *Int Urogynecol J* 2011;22:699–704 [PubMed: 21107811]
14. Dietz HP, Shek C, De Leon J, et al. Ballooning of the levator hiatus. *Ultrasound Obstet Gynecol.* 2008;31:676–80. [PubMed: 18470963]
15. Bump RC, Mattiasson A, Bo K, et al. The standardization of terminology of female pelvic organ prolapse and pelvic floor dysfunction. *Am J Obstet Gynecol* 1996;175:10–7. [PubMed: 8694033]
16. Tukey JW. *Exploratory Data Analysis.* 1977 Addison-Wesley, Reading, MA.
17. Dietz HP, Franco AVM, Shek KL, et al. Avulsion injury and levator hiatal ballooning: two independent risk factors for prolapse? An observational study. *Acta Obstet Gynecol Scand* 2012, 91:211–4. [PubMed: 22050558]
18. Chen L, Ashton-Miller JA, DeLancey JO. A 3D finite element model of anterior vaginal wall support to evaluate mechanisms underlying cystocele formation. *J Biomech.* 2009;42(10):1371–7. [PubMed: 19481208]
19. Blomquist JL, Muñoz A, Carroll M, Handa VL. Association of Delivery Mode With Pelvic Floor Disorders After Childbirth. *JAMA.* 2018 12 18;320(23):2438–2447. [PubMed: 30561480]
20. Hundley AF, Wu JM, Visco AG. A comparison of perineometer to brink score for assessment of pelvic floor muscle strength. *Am J Obstet Gynecol.* 2005 5;192(5):1583–91. [PubMed: 15902162]
21. Myer ENB, Roem JL, Lovejoy DA, et al. Longitudinal changes in pelvic floor muscle strength among parous women. *Am J Obstet Gynecol.* 2018 6 11 pii: S0002–9378(18)30490–3. doi: 10.1016/j.ajog.2018.06.003. [Epub ahead of print]
22. Chan SSC, Cheung RYK, Lee LL, et al. Longitudinal pelvic floor biometry: which factors affect it? *Ultrasound Obstet Gynecol.* 2018;51:246–252. doi: 10.1002/uog.17446. [PubMed: 28236365]

CONDENSATION:

The association between levator ani avulsion and pelvic organ prolapse may be fully mediated by increased levator hiatus area and reduced muscle strength.

AJOG AT A GLANCE:

A. This study was conducted to clarify mechanisms by which obstetrical levator avulsion may lead to pelvic organ prolapse later in life.

B. The key findings are that two characteristics associated with levator avulsion, increased levator hiatus area and decreased muscle strength, are strongly associated with prolapse and may explain the association between levator avulsion and prolapse.

C. While it has previously been shown that levator avulsion is strongly associated with prolapse, these results also suggests that this association might be explained, to a large extent (and possibly fully), by the increase in the levator hiatus size and the reduction in pelvic muscle strength seen among women with levator avulsion.

Figure 1:

Levator hiatus area (in cm²) and PFMS (mean peak pressure during voluntary contraction, in cmH₂O), by levator avulsion. Data for women with levator avulsion shown in black closed circles. Three outliers (with levator avulsion) are shown in black closed diamonds.

Figure 2:

Percentage of women with pelvic organ prolapse, as a function of (A) levator hiatus area at maximum Valsalva, measured on 3-dimensional transperineal ultrasound; (B) PFMS (mean peak pressure at maximum voluntary contraction), measured via perineometry.

Table 1:Characteristics of 432 vaginally parous women, by prolapse status^a

Characteristic ^b	No prolapse (n=320)	Prolapse (n=112)	P value
Age at ultrasound, years	42.8 [39.5, 46.7]	45.0 [41.0, 48.3]	0.008
Years from first vaginal birth to ultrasound visit	10.5 [9.3, 13.3]	12.3 [9.6, 14.6]	0.006
Black race (vs. nonblack)	12% (37)	8% (9)	0.298
Body mass index (kg/m ²) at ultrasound	25.6 [22.7, 29.3]	25.9 [22.8, 30.5]	0.362
Any ^c vaginal delivery with macrosomia (>4 kg)	14% (45)	17% (19)	0.457
Any ^c vaginal delivery with 2 nd stage >2hr	28% (89)	32% (36)	0.384
Any ^c obstetric anal sphincter laceration	15% (48)	22% (25)	0.075
Any ^c vacuum-assisted vaginal delivery	11% (35)	14% (16)	0.345
Any ^c forceps-assisted vaginal delivery	10% (33)	25% (28)	<.001

^a Prolapse was defined as any examination in which Ba, C, or Bp was beyond the hymen or history of surgery for prolapse^b Categorical variables reported as percent (n); continuous variables reported as median [interquartile range].^c “Any” refers to an occurrence across all deliveries.

Associations between pelvic organ prolapse and levator avulsion, levator hiatus area, and pelvic muscle strength (n=429).

Table 2:

	Univariable analyses	Multivariable analysis	Stratified by avulsion	
			No avulsion (n=368)	Avulsion (n=61)
Levator avulsion	4.17 (2.38, 7.31)	1.75 (0.91, 3.39)		
Levator hiatus area (per 5cm ²)	1.52 (1.34, 1.73)	1.46 (1.28, 1.67)	1.36 (1.18, 1.57)	2.43 (1.51, 3.91)
Pelvic muscle strength (per 5cm H ₂ O)	0.87 (0.81, 0.94)	0.90 (0.83, 0.98)	0.91 (0.84, 0.99)	0.85 (0.57, 1.24)

Statistically significant results showed in bold font.