Published in final edited form as:

J Cogn Neurosci. 2019 February ; 31(2): 299–313. doi:10.1162/jocn_a_01353.

Grasping neurons in the ventral premotor cortex of macaques are modulated by social goals

Gino Coudé¹, Giulia Toschi⁴, Fabrizia Festante³, Marco Bimbi², James Bonaiuto¹, and Pier Francesco Ferrari^{1,2}

¹Institut des sciences cognitives Marc Jeannerod, CNRS; Université Lyon 1

²Department of Medicine and Surgery, Neuroscience Unit, Università di Parma

³Department of Developmental Neuroscience, IRCCS Stella Maris Foundation, Pisa

⁴Department of Biomedical and Neuromotor Sciences, Università di Bologna

Abstract

While it is established that F5 neurons can distinguish between non-social goals such as bringing food to the mouth for eating or placing it in a container, it is not clear whether they discriminate between social and non-social goals. Here, we recorded single-unit activity in ventral premotor cortex (PMv) of two female macaques and used a simple reach-to-grasp motor task in which a monkey grasped an object with a precision grip in three conditions which only differed in terms of their final goal, that is, a subsequent motor act that was either social (placing in the experimenter's hand or *Hand* condition) or non-social (placing in a container or bringing to the mouth for eating, *Container* or *Mouth* condition). We found that during the execution of the grasping motor act, the response of a sizable proportion of F5 motor neurons was modulated by the final goal of the action, with some having a preference for the social goal condition. Our results reveal that the representation of goal-directed actions in PMv is influenced by contextual information not only extracted from physical cues, but also from cues endowed with biological or social value. Our study suggests that the activity of grasping neurons in the premotor cortex is modulated by social context.

INTRODUCTION

Naturalistic actions involve a hierarchy of goals, with subgoals acting in the service of an overarching goal (Rizzolatti et al., 2014). For example, the overarching goal of “eating” involves the subgoal of “taking possession”. Distinct motor acts like reaching, grasping and biting are each endowed with a specific motor goal. For instance, “grasping” implies having the motor goal of taking possession of an object. When planning and executing an action such as “grasping and eating a piece of food”, an individual must use the overarching goal, “eating” in this example, to select an appropriate sequence of subgoals and the motor acts to achieve them.

In normal daily life, animals and humans do not just act independently from their conspecifics, but are often required to perform actions directed to or jointly with others (Sebanz et al., 2006). The motor system, therefore, must express a high degree of versatility in order to activate similar motor programs but aimed at different goals, both non-social and social (e.g. placing an object into a container or placing an object into the hand of another individual). Social goals involve coordination of an action not only with the external environment, but also with other individuals who are not static environmental features, but have their own intentions, motives, and internal states.

The ventral premotor area F5 is a core region for coding the goal of hand motor acts such as grasping, tearing, and holding (Rizzolatti et al., 2009). The neurons of F5 have the crucial property of coding goals independently of the sequence of movements or the effector used to achieve them (Rizzolatti et al., 1988; Umiltà et al., 2008), thus displaying a high level of motor abstraction (Rizzolatti et al., 2002). In area F5 and the inferior parietal lobule (IPL), the activation of specific neurons during the execution of a goal-directed motor act (grasping to place or grasping to eat), is modulated by the final goal of the action (eating or placing) (Fogassi et al., 2005; Bonini et al., 2011). In these studies, the monkey was trained to execute two motor acts: grasp a piece of food and bring it to the mouth, or grasp an object and place it into a container (as well as to observe the experimenter performing the same two motor acts). A large proportion of motor neurons in F5 and IPL responded differentially to the same act (grasping) according to the final goal of the action in which the act was embedded. The ability of the motor system to encode the same action according to the final goal likely derives from the presence of contextual cues provided by the container in the grasp-to-place condition (i.e. the presence of a container inside which the object was subsequently placed) or by the type of object to grasp (i.e. a piece of food) in the grasp-to-eat condition.

While it is established that F5 neurons can distinguish between non-social goals such as bringing food to the mouth for eating and placing in a container, it is not clear whether they also code social goals. Here we devised a simple reach-to-grasp motor task in which a monkey grasped an object with a precision grip in three conditions which only differed in term of their final goal, that is, a subsequent motor act that is either social (placing in the experimenter's hand or "hand" condition) or non-social (placing in a container or bringing to the mouth for eating, "container" or "mouth" condition). We found that during performance of the grasping motor act, the response of a sizable proportion of F5 motor neurons was modulated by the final goal, with some having a preference for the social goal condition.

EXPERIMENTAL PROCEDURES

Animals and surgical procedures

Two captive-born and individually housed adult female rhesus macaques (*Macaca mulatta*) served as subjects (M1 and M2). The animal handling, as well as surgical and experimental procedures, complied with the European guidelines (86/609/EEC 2003/65/EC Directives and 2010/63/EU) and Italian laws (D.L. 116–92, D.L. 26–2014) in force on the care and use of laboratory animals, and were approved by the Veterinarian Animal Care and Use Committee of the University of Parma (Prot. 78/12 17/07/2012) and authorized by the Italian Health

Ministry (D.M. 294/2012-C, 11/12/2012). The monkeys were housed and handled in strict accordance with the recommendations of the Weatherall Report about good animal practice. The well-being and health conditions of the monkeys were constantly monitored by the institutional veterinary doctor of the University of Parma.

A titanium head post (Crist Instrument, Hagerstown, MD, USA) was surgically implanted on the skull using titanium screws. A cilux recording chamber (18X18 mm, Alpha-Omega, Germany) was stereotaxically implanted and secured with dental cement. For both procedures, each animal was deeply anaesthetized with ketamine hydrochloride (5 mg/kg i.m.) and medetomidine hydrochloride (0.1 mg/kg i.m.) and its heart rate, temperature and respiration were carefully monitored and kept within physiological range. Pain medication was routinely given after surgery: (Dexamethasone, 2mg/kg, every 12 hours, from 1 day before to 3 days after surgery; Ketoprofen, 5mg/kg, every 12 hours for 3 days following surgery)

Monkey task

The basic task is illustrated in figure 1A. The monkey was seated facing a table (60X60 cm) onto which a metallic cube was placed along the monkey body midline, at 13 cm from monkey's hand starting position. The monkey had to reach and grasp the object and then place it in a small container located 10 cm to the left of the grasping location. At the beginning of each trial the monkey had to keep her right hand on a handle attached to the table for at least 1000 ms (Fig. 1A-I), after which a transparent barrier was removed to give the "go" signal and the monkey was required to grasp the object (Fig. 1A-II) and place it in the container (Fig. 1A-III). A juice reward (and a solid food reward) was delivered after 500–1000 ms if the monkey correctly executed the trial (Fig. 1A-IV). The task was run in one or more blocks of five trials each until a minimum total of 12 trials per condition was reached. Any trial in which the grasping action was not properly executed was aborted and no reward was delivered. Note that the reward consisted of juice (0.14 ml) and a piece of apple (1cm³) for each trial and was the same for all three conditions (See Experimental conditions below). This reward schedule was used because in the *Mouth* condition (see below) the monkey had to grasp a piece of food, bring it to the mouth, and eat it. Therefore, in the other two conditions we decided to introduce the same type of reward to avoid possible differential neuronal activity according to the type of reward.

Experimental conditions

The three experimental conditions are illustrated in figure 1B. The first part of the task was the same in each condition: the monkey was required to release the handle and grasp an object or a piece of food with a precision grip. The experimental conditions differed only in the second part of the task in which the monkey had to: Fig. 1B-I) place the object in a container (*Container* condition); Fig. 1B-II) place the object in the hand of the experimenter (*Hand* condition); or Fig. 1B-III) bring the object (a food morsel) to the mouth and eat it (*Mouth* condition). Since the grasped object was a metallic cube in the *Container* and *Hand* conditions, but a cubic piece of apple in the *Mouth* condition, we had to balance the reward value across all conditions. Therefore, in addition to the juice reward delivered in all conditions, the same pieces of apple used in the *Mouth* condition were given to the monkey

after the correct completion of the trials in the *Container* and *Hand* conditions. The conditions were executed in randomized blocks of 5 trials.

Data recording

Neuronal activity was recorded with a linear multi-site electrode (16-channel, 250 micrometers spacing; U-probe, Plexon inc., Dallas, TX, USA) and digitalized at 40kHz using the Omniplex 16-channel recording system (Plexon). Task control (including handle holding, hand-object contact detection and reward delivery) was computer-controlled through a customized Labview© program. Contact-detection panels (Crist Instrument, Hagerstown, MD, USA) were used to record the exact moments in which the monkey's hand released the handle and touched the target. The latter event was used to align the neuronal activity recorded for every trial in all conditions.

We collected usable motor neuron data in 15 sessions from Monkey 1 and in 17 sessions from Monkey 2. Neuronal activity was recorded from area F5 of the ventral premotor cortex (see below for details). During the insertion of the electrode shaft in the brain we ensured that the topmost electrode was positioned under the dura, but remained outside of the cortex, so that it could be used as a reference channel. Of the 15 available recording channels, two were dedicated to eye movement control for another task. The remaining 13 channels were used for single-unit recording. The multi-electrode was lowered in the brain through the intact dura and the general properties of the neurons were tested. When motor properties for grasping were found, we waited for approximately one hour to allow the neural activity to stabilize and then started to record single units using the behavioral protocol described above.

Preliminary testing of neuronal activity

Before proceeding with the neural testing using our behavioral protocol, single- and multiunit activity were systematically tested for visuomotor properties to identify recording sites in area F5 endowed with hand grasping motor activity (Rozzi et al., 2008; see Maranesi et al., 2012). Briefly, we required the monkey to grasp food items in various conditions (i.e. with eyes closed, or without flexing the wrist, elbow or shoulder) enabling us to disentangle neuronal activity related to visual stimulation, reaching or grasping objects. Also, to exclude the possible presence of mouth-related responses, we tested changes in neural activity related to the delivery of small pieces of food directly into the mouth while the monkey's eyes were closed.

Data analysis

Neuronal spike waveforms were classified offline into units using a commercial spike-sorting software (OfflineSorter, Plexon, USA). The procedure we used for spike sorting was as follows: We set the detection threshold at -2% of the energy of the signal. We used a pre-threshold period of 200 microseconds, a total time window of 1200 micro seconds and a refractory inter spike interval of 1000 micro seconds. We used the "valley seek" method for sorting and applied semi-automatic clustering using templates. These templates were constructed using manually selected waveforms that looked like action potentials. This method allowed us to reject possible artifacts and also spikes that were too small to be

reliably sorted. We applied a principal component analysis to summarize each spike waveform by a feature vector. Finally, we divided these vectors in clusters (putative neurons) using the k-means clustering feature of the software and excluded the outliers so that the clusters did not overlap.

The neural response selectivity was determined for each neuron by comparing the spike frequency for each trial across two epochs (*Baseline* and *Grasping*, described below) and 3 conditions including *Container*, *Hand* and *Mouth*. The statistical analysis consisted of a 2×3 repeated measures ANOVA (factors: epochs and conditions) followed by a Newman-Keuls post hoc test, all with a 0.05 alpha level. The two epochs had a 350 ms duration and were precisely calculated with reference to the digital events and the contact with the grasped object, and verified by an offline kinematic analysis recorded in a separate recording session (see below Movement kinematics analysis). The *Baseline* epoch corresponded to a time period in which the monkey remained still with her hand on the starting handle. This epoch lasted from 1000 to 650 ms before the monkey's release of the handle. The *Grasping* epoch was defined as starting 100 ms before the release of the handle (when the monkey's hand started to move as determined by the kinematic analysis) and ended at the object displacement onset (when the monkey had grasped the object or the piece of food and started to lift it). This event was signaled by the hand-object-table contact detection-panel. The *Grasping* epoch was defined such as to encompass the whole reaching-grasping phase without including the placing phase of the action (which was expected to necessarily be kinematically different between the conditions). The moment when the monkeys' hand touched the object (indicated by the "on" signal on the hand-object-table contact detection panel) was used to align the trials.

Neurons were classified as 1) grasping neurons, if they showed a significant main effect of Epoch; 2) as having a goal preference, if they also showed a main effect of Condition or an interaction between Epoch and Condition. Planned contrasts indicated in which condition (or conditions) the mean discharge for a grasping epoch was significantly different than the other(s).

The population analysis was carried out by constructing a spike density function for each neuron, by averaging the spike frequency for 20 ms bins across trials within each condition. The baseline activity (mean spikes per second during the *Baseline* epoch) was subtracted from the spike frequency of each bin. The data were then normalized by dividing the baseline-corrected spike frequency within each bin by the maximum absolute value found across all bins in all conditions. The result is a discharge rate ranging between -1 and 1 for each neuron. The baseline-corrected and normalized spike density function was then smoothed using a Gaussian kernel with a width of 30 ms.

The population curve was computed as the mean spike frequency in each bin over all neurons. The statistical differences between populations of neurons were calculated based on the firing rate during the *Grasping* epoch. One way repeated-measures ANOVAs used to compare the effect of condition on the population firing rate, followed by Newman-Keuls post hoc procedures.

The time at which goal-modulated neurons begin to show selectivity was calculated as follows: a baseline discharge rate was subtracted from each value (in 20 ms time bins) of the differential activity time course (preferred – non-preferred conditions). The baseline value was the mean differential activity between preferred and non-preferred conditions during the 350-ms baseline epoch plus one standard deviation. The time bin corresponding to the occurrence of a series of at least three consecutive bins with values < 0 before the peak of selectivity was taken as the onset time.

K-means clustering analysis—The standard K-means clustering algorithm (Matlab, Mathworks), with squared Euclidian distance as the metric, was applied to objectively characterize the discharge patterns of the neuronal population. Clustering was applied to the baseline-corrected, normalized, and smoothed spike density function of each neuron during the *Grasp* epoch, averaged over trials within each condition (each neuron contributed 18 time bins $\times$ 3 conditions, corresponding to the time between movement onset and object touch). The optimal number of clusters was determined using the Silhouette (Sil; Kaufman and Rousseeuw, 1990), Davies-Bouldin (DB; Davies and Bouldin, 1979), Calinski-Harabasz (CH; Calinski and Harabasz, 1974), Krzanowski-Lai (KL; Krzanowski and Lai, 1988), Hartigan (Hartigan, 1975), weighted inter-intra (Wint; Strehl, 2002), and homogeneity-separation (Chen et al., 2002) indices. Within each cluster, a one-way ANOVA was computed for each time bin of the grasping epoch to assess significant differences in discharge rate between conditions (Bonferroni corrected according to the number of time bins).

Non-metric multidimensional scaling—In order to examine the temporal trajectory of the neural population activity in each condition, we used multidimensional scaling (MDS) analyses carried out with MATLAB software (Mathworks). We used the baseline-corrected firing rate of each neuron, averaged across trials and concatenated over conditions, to construct a 267×423 dimensional matrix, X (89 time bins per trial in each of 3 conditions, 423 excitatory neurons), containing the mean firing rate of each neuron over the time course of each condition. We then computed a Euclidean distance matrix, D (267×267), where each element $D_{i,j}$ contained the Euclidean distance between rows i and j of X ($D_{i,j} = \|X_i - X_j\|^2$), and thus the distance between snapshots of population activity at different time points in 423-dimensional space. We then applied the standard non-metric MDS (NMDS) algorithm (Shepard, 1980) using D to obtain a 267×2 dimensional representation of X (we chose two dimensions based on the reduction in stress from 0.127 to 0.057 going from one to two dimensions), while maintaining the approximate relative distance between pairs of observations in the original, higher-dimensional space (Kayaert et al., 2005). We then reshaped the result to obtain one 89×2 dimensional population trajectory for each condition. NMDS makes fewer assumptions about the data compared to other techniques such as principle components analysis, and because the number of dimensions is chosen prior the analysis, there are no hidden dimensions of variation. Because the neurons from different penetrations were necessarily recorded during different trials, we estimated the variability of the population activity trajectories by using a bootstrapping technique in which we obtained the population activity by averaging each neuron's activity over 10 trials sampled randomly without replacement from the corresponding penetration. This was repeated 100 times,

resulting in 100 different 423 dimensional population activity vectors for each condition. A one-way ANOSIM (Analysis of Similarities) with Bonferroni-corrected pair-wise follow-up tests (Clarke and Gorley, 2001) implemented as part of the Fathom toolbox (Jones, 2014) was used to test for significant differences in population activity trajectories between conditions at each trial event (movement onset, handle release, object touch, object lift), using dissimilarity matrices created from pair-wise comparisons of trajectory coordinates from NMDS. ANOSIM is a nonparametric permutation test (we used 1000 permutations) which calculates an *R* test statistic indicating the degree of separation between conditions, with a value of 1 indicating complete separation and 0 indicating no separation.

Movement kinematics analysis—The kinematic parameters of reaching and grasping movements were recorded in a dedicated session. The action was filmed from a lateral view (on the monkey's left side) encompassing the hand on the starting handle and object to grasp using a 50 fps temporal resolution camera (Canon EOS-7D) with a spatial resolution of 5184 × 3456 pixels. The hand excursion in X and Y axes (sagittal and vertical axes, respectively) was measured using an open source video analysis software (Tracker). Three LEDs connected to the contact detection circuits indicated the handle release, object touch, and lifting phase of the action. They were placed in the background, within the camera field, but out of the monkey's sight. The data were first analyzed by comparing the reaching/grasping duration, as well as the velocity and movement amplitude of the wrist between the three conditions (20 trials each) using a one-way ANOVA. Data were first analyzed by comparing the reaching/grasping duration, as well as the velocity and movement amplitude of the wrist between the three conditions (20 trials each) using a one-way ANOVA. Because each trial varied in the timing of the behavioral events, the eight-dimensional trajectories (x, y positions for the wrist, CMC, index PIP, and thumb MCP) from each recorded trial were then aligned using canonical time warping (Zhou and Torre, 2009) in order to compare them between conditions. The aligned trajectories were then projected into a two-dimensional (based on the reduction in stress from 0.022 to 0.004 going from one to two dimensions) space using NMDS, with the Euclidean distance between each pair of the 60 (3 conditions, 20 trials each) temporally aligned trajectories as input. One-way ANOSIM with Bonferroni-corrected pair-wise follow-up was used to test for significant differences in kinematic trajectories between conditions at the handle release and object touch events, using dissimilarity matrices created from pair-wise comparisons of trajectory coordinates from NMDS.

RESULTS

Goal-modulated neurons

The activity of 1499 neurons was recorded from area F5 in the left hemisphere of two monkeys. A total of 437 single units (Monkey 1, 157 units; Monkey 2, 280 units) were revealed to be “hand grasping neurons” as tested with the task illustrated in Fig. 1A and statistically confirmed with the 2×3 ANOVA described in the Data analysis section above. The neurons were collected from 32 electrode penetrations and 13 different cortical depths. We found that 180 grasping neurons (41% of the recorded motor neurons) were modulated by the goal of the action and had a preference for either one or two of the conditions (Fig. 2).

We hereafter refer to these neurons as *goal-modulated*. The majority of these *goal-modulated* neurons (63%, 114/180) had a preference for only one condition. Three examples of *goal-modulated* neurons having a preference for one condition are presented in figure 3. Unit 146 had a preference for grasping followed by placing the object in a container. Unit 54 had a preference for grasping condition followed by eating the piece of food. The third neuron, Unit 174, had a preference for grasping followed by placing the object in the hand of an experimenter.

Population analysis

The population analysis for the *Goal-Modulated* neurons modulated by single goal (one condition) is shown in figure 4. As expected, the discharge in the preferred condition for these neurons is significantly stronger than in the non-preferred conditions (Fig. 4A; *Container-Goal-Modulated* neurons; $F_{(2,56)} = 45.63$; $p < 0.0001$; *Post hoc*, *Container* vs *Hand*, $p < 0.001$; *Container* vs *Mouth*, $p < 0.001$; Fig. 4B; *Hand-Goal-Modulated* neurons; $F_{(2,32)} = 31.92$; $p < 0.0001$; *Post hoc*, *Hand* vs *Container*, $p < 0.001$; *Hand* vs *Mouth*, $p < 0.001$; Fig. 4C; *Mouth-Goal-Modulated* neurons; $F_{(2,128)} = 259.27$; $p < 0.001$; *Post hoc*, *Mouth* vs *Container*, $p < 0.0001$; *Mouth* vs *Hand*, $p < 0.0001$).

The results of the population analysis of all *Goal-* and *Non-Goal-modulated* neurons are illustrated in Figure 5. Significant differences between the conditions among the *Goal-Modulated* neurons (Fig. 5A) were found (one-way repeated-measure ANOVA; $F_{(2,350)} = 10.13$; $p < 0.0001$). *Post hoc* comparisons revealed that *Goal-Modulated* neurons fired significantly more in the *Mouth* condition than in the other conditions (*Mouth* vs *Container*, $p < 0.001$; *Mouth* vs *Hand*, $p < 0.001$). Likewise, for the *Non-Goal-Modulated* neurons (Fig. 5B), significant differences between conditions at a population level were found; ($F_{(2,492)} = 12.51$; $p < 0.0001$), with *post hoc* comparisons revealing significantly more firing in the *Mouth* condition than in the other conditions (*Mouth* vs *Container*, $p < 0.0001$; *Mouth* vs *Hand*, $p < 0.0001$).

Mouth condition preference

The majority of neurons with a single preferred condition preferred the *Mouth* condition (see also Fig. 2). Likewise, a stronger discharge in the *Mouth* condition was also a characteristic of all *Goal-Modulated* neurons and was also present among the *Non-Goal-Modulated* neurons. The latter result might sound counterintuitive, but the fact that *Non-Goal-Modulated* neurons do not individually show a significant preference does not mean that their firing rate was the same for all conditions as a population.

Note that the overall preference for the *Mouth* condition is not due to oral movements. No neuron included in our analyses showed any discharge during oral movements such as mastication, and therefore we can exclude the possibility that motor preparation for mouth opening is responsible for the increase in firing activity during the grasping action in the *Mouth* condition.

Clustering analysis of neuronal activity

Since differences were found between the three conditions at a population level for both modulated and non-modulated neurons, we sought to analyze the neuronal population in more details without any of the *a priori* assumptions about neural selectivity embedded in the ANOVA-based neural classification analysis. We used cluster analysis to identify populations of neurons having different discharge patterns across conditions. The main results of this analysis are shown in figure 6A. In the central panel, each of the 423 neurons (only excitatory neurons were considered for this analysis) is plotted as a point in two dimensional space using NMDS to show their similarity in discharge pattern across conditions, with different colors denoting cluster membership (seven clusters). The optimal number of clusters was chosen according to a variety of cluster dispersion metrics (see above K-means clustering analysis). As each metric yielded a different optimal number of clusters (Sil=3, DB=7, CH=7, KL=5, Hartigan=7, Wint=3, homogeneity-separation=7), we chose the mode, seven clusters. To help visualize cluster differences, we plotted the average neuronal response of the neurons belonging to each cluster for the three conditions (Fig. 6A). The seven clusters seem to represent different aspects of the neuronal discharge during the task, i.e., preference for one or more conditions, peak response time, or a combination of both. We found three clusters reflecting the neuronal preference for the action goal with a peak of activity around the Touch event (cluster 1 for Container; cluster 2 for *Hand* and cluster 3 for *Mouth*). Two additional clusters represent the time of peak activity and a preference for *Mouth* condition (cluster 7 for peak just after Release of the handle and cluster 6 for peak around Lift Object). Cluster 5 represents neurons with a late peak of activity when the object is lifted (Lift Object) and no goal preference, while cluster 4 seems to represent late peaking activity and a preference for non-social goals (*Container* and *Mouth*).

In order to assess the similarity between neural discharge patterns within- and between-clusters, we computed a correlation-based similarity matrix (Fig. 6B) showing the mean pairwise Euclidean distance between the discharge patterns of neurons belonging to each cluster. The low distance measures in the diagonal of the matrix show that neurons in each cluster behave most similarly to neurons in the same cluster (as expected by the use of the clustering algorithm), with cluster 6 being the most homogeneous. There was also a strong similarity between cluster 6 and clusters 3, 4, and 5, all of which represented neurons with late peaks of activity. Figure 6C shows the percentage breakdown of cluster membership for each ANOVA-based classification. Note that clusters 1 to 4 contain 50% or more of the neurons that were found to be selective to *Container* (62.1% in cluster 1), *Hand* (52.9% in cluster 2), *Mouth* (52.3% in cluster 3) or *Container-Mouth* (56.3% in cluster 4), respectively. This result means that the coding of grasp goal is evident at the population level not only under *a priori* assumptions about possible goal specificity (as revealed by the ANOVAs), but also in an unbiased, unsupervised clustering of neuronal discharge profiles.

Non-metric multidimensional scaling (NMDS) assessment of population activity trajectories

The general dissimilarity of the population activity during an action with a social goal is confirmed by a non-metric multidimensional scaling (NMDS) analysis (Fig. 7). The NMDS

analysis using two dimensions resulted in a good fit to the data (stress=0.058, $R^2=0.98$, $p<0.0001$). This analysis shows that the population-level discharge pattern associated with the goal of grasping to place in the hand is qualitatively different from the one where the goal of grasping is to place in the container or eat. In this analysis, we treated the activity pattern of the entire population at each point in time (averaged over trials) as a vector in high-dimensional space, and visualized the trajectory of this vector as the trial unfolded. A bootstrapping procedure using random subsets of the trials on each iteration was used to estimate the variability of the population activity. The population activity pattern was very similar at the start of the trial in each condition (global $R=0.43$, $p=0.001$), and gradually diverged, with divergence of the *Mouth* and *Hand* conditions from the *Container* condition starting at the onset of movement (global $R=0.84$, $p=0.001$; container – hand $R=0.89$, $p=0.003$; container – mouth $R=0.99$, $p=0.003$; hand – mouth $R=0.63$, $p=0.003$) and continuing until the release of the monkey's hand from the handle following the “go” signal (global $R=0.77$, $p=0.001$; container – hand $R=0.77$, $p=0.003$; container – mouth $R=1.0$, $p=0.003$; hand – mouth $R=0.44$, $p=0.003$). Interestingly the activity pattern in the *Hand* condition was maximally different from the *Container* and *Mouth* conditions around the time of the initial touch of the object (global $R=0.74$, $p=0.001$; container – hand $R=0.98$, $p=0.003$; container – mouth $R=0.30$, $p=0.003$; hand – mouth $R=0.97$, $p=0.003$), and became more similar to the *Container* condition by the time the object was lifted (global $R=0.90$, $p=0.001$; container – hand $R=0.77$, $p=0.003$; container – mouth $R=1.0$, $p=0.003$; hand – mouth $R=0.92$, $p=0.003$). This is in spite of the fact that all conditions are kinematically similar during the grasping epoch (see Movement kinematics section below). Unsurprisingly, the population activity in the *Mouth* condition was most dissimilar to the other two conditions following the lifting of the object, which is when the kinematic differences in this action become manifest. This result shows that, at the population level, the *Hand* condition is coded differently from the *Container* and *Mouth* conditions even during the time of the greatest kinematic congruence between conditions.

Goal-modulation and discrimination latency

To better characterize the *Goal-Modulated* neurons, we analyzed the time at which they began to discriminate the goal of the action. The goal-discrimination time for each neuron is shown in figure 4 (bottom). There were no significant differences between the discrimination time latency of the three types of *Goal-Modulated* neurons (*Container*, mean=−143.40 ms, SE=18.53; *Hand*, mean=−134.20 ms, SE=29.44; *Mouth*, mean=−75.60 ms, SE=11.46; Kruskal-Wallis, $\chi^2=5.70$, $df=2$, $n.s.$).

Movement kinematics

Since the scope of the present experiment was to compare neuronal discharge between similar grasping actions that differed only by the subsequent action in which they were embedded, it was important to verify that the kinematic parameters of the reaching and grasping actions were the same across the three conditions. Results revealed no differences in movement extent (in height or in length) during the reaching phase between the three conditions. However, we found that reaching was slower in the mouth condition (*Container*: mean=787.94, SE=19.02 mm/s; *Hand*: mean=747.87, SE=15.72 mm/s, *Mouth*: mean=706.47, SE=6.88 mm/s; $F_{(2,57)}=7.59$, $p<0.005$), probably indicating a deceleration of

the reaching movement in order to grasp the cubic-shaped food morsel which has a different texture than the metallic cube used in the *Container* and *Hand* condition. This difference is, however, unlikely to have an impact on the proportion of *Goal-Modulated* neurons found in this experiment. Indeed, we found that the “grasping to eat” trials were also slower during the recording of neurons preferring the other conditions (*Container* or *Hand* condition), ruling out the possibility that the grasping velocity was a determining factor in neuronal preference. We verified if the overall neuronal firing was linked to the duration of the reaching phase in all *Goal-Modulated* neurons. We found no significant correlation between the duration of the reaching phase (from handle release to object touch) and the neuronal discharge across all the trials, for any of the conditions (Pearson correlation; *Container*: $r = -0.013$, n.s.; *Hand*: $r = -0.037$, n.s.; *Mouth*: $r = -0.046$, n.s.), thus further supporting the lack of a causal link between different kinematics parameters of the reaching-grasping and the firing rate. Additionally, the NMDS analysis (stress=0.004, $R^2=0.99$, $p < 0.0001$) demonstrated that the overall kinematic trajectories were very similar across conditions, with no separation between the trajectories at either the handle release (global $R=0.00$, $p=0.538$) or object touch events ($R=0.031$, $p=0.087$; Fig. 8).

DISCUSSION

The main finding of the present study is that, among a population of “hand grasping motor neurons” in F5 which have a preference for the final goal of an action, some neurons coded a goal that implied a social interaction between two individuals. In fact, notwithstanding the very limited interactions allowed by our experimental protocol, a number of neurons (10% of the goal-modulated neurons) had a preference for the condition *placing in the hand* of the experimenter, a condition involving a social interaction. Additionally, 76% of the neurons modulated by two conditions were modulated by the *placing in the hand* condition. In our view, these results suggest that the activity of grasping neurons in the premotor cortex is modulated by social context. These results extend previous findings (Fogassi et al., 2005; Bonini et al., 2010; Mazurek et al., 2018) showing that hand grasping neurons can differently code the same grasping act according to the goal of the action in which such act is embedded.

The activity of many motor neurons in the ventral premotor cortex is modulated by contextual cues, such as the presence of a container or the type of object to be grasped, which indicate to the monkey what the required final goal of the action is in the current experimental condition. The present findings unveiled what could be a new property of motor neurons in F5, namely a preferential modulation by a forthcoming interaction with another individual. In the present experiment, this interaction consisted of placing an object in the hand of the experimenter. The type of motor act sequence required to accomplish such a goal is not kinematically different from “placing in a container” (*Container* condition), since the kinematic trajectories were very similar across conditions. Nevertheless, the differential activity of some grasping neurons in F5 demonstrates that such a difference is indeed relevant for the monkey.

Social information is indeed crucial for non-human primates (Emery, 2000; Klein et al., 2009; Nummenmaa and Calder, 2009) as it affects motivational, affective and decision-

making processes. In a neurophysiological study, it has been shown that macaque monkeys track the behavior of other individuals during interactive tasks, and that neuronal activity in the mesial frontal cortex reflects the monitoring of both the behavior and the errors of the other individual, during self and others' actions (Yoshida et al., 2011). Despite interesting behavioral (Deaner et al., 2005; Ballesta and Duhamel, 2015) and neurophysiological investigations indicating the importance of social contexts in monkey decision making and in modulating brain activity, no studies have yet explored the impact of social cues or social interactions on the motor discharge of neurons in the monkey's premotor cortex, an important area for high-level planning of manual actions. However, socially relevant cues such as gaze direction have been shown to directly impact the discharge of grasping mirror neurons in the premotor cortex (Coudé et al., 2016). In that study, the visual discharge of some visuomotor neurons (mirror neurons) preferred a gaze direction either congruent or incongruent with the location of the grasped object. However, the monkey and the experimenter were not engaged in a direct interaction, with the monkey passively observing the experimenter as they performed grasping actions. One study has investigated the impact of social context on parietal neurons (Fujii et al., 2007). In that experiment, parietal activity was strongly tuned to the use of the arm contralateral to the recorded hemisphere when two monkeys were sitting side-by-side and could reach for and grasp food without interacting. However, when the food was put in a shared space and social conflict could emerge, the neurons developed different combinations of preferences to self and other motion (Fujii et al., 2007). These results showed that within the parietofrontal circuits involved in grasping, some neurons code space in terms of social relevance.

Interestingly, the influence of social cues on the motor system has been shown in many contexts in human studies (Sebanz et al., 2006; Sartori et al., 2011; Dolk et al., 2014; Chinellato et al., 2015). For instance, in a task where participants were required to grasp an object while observing an actor grasping with an interactive purpose, participants' grasping movements were delayed and reaching trajectories were deviated, indicating that socially relevant stimuli are acknowledged by the motor system (Chinellato et al., 2015). In another study participants were required to grasp an object and to move it towards the experimenter, or towards themselves (Scorolli et al., 2014). Kinematic analyses confirmed that reaching movements were affected by the social goal of the action and the conventional use of the object (e.g. an object typically used in functional-cooperative relation). It is worth noting that in our study the monkeys had to grasp an object while the experimenter's hand remained still, in a supine position in the reaching space of the monkey. Such an experimental design differs from the above cited human studies where subjects had to grasp an object while observing an ongoing grasping action from another individual or had to coordinate their own action with another subject. These complex dynamics might have had an impact on kinematic parameters during the grasping phases. In contrast, in our study we eliminated possible interference between the monkey grasping action and the observed experimenter movement. In addition, the long period of animal training probably contributed to reducing the variability of the kinematic parameters of the monkey's reaching-grasping movements. Nevertheless, the NMDS analysis revealed dissimilarity in neuronal population activity between the *Container* and *Hand* conditions despite the fact that they are kinematically similar. The activity pattern in the *Hand* condition was maximally different

from the *Container* and *Mouth* conditions around the time of the initial touch of the object, where kinematic parameters tend to be more similar among conditions.

Although these data show that the *Hand* differs from the *Container* condition in terms of neuronal activity, we cannot be certain that the *Hand* condition is viewed by the monkey as a socially-selective condition and not as a grasping to place the object on a different container. Even if this possibility cannot be excluded, previous studies on the functional properties of neurons in ventral premotor cortex have shown that social signals, such as gaze direction, can modulate the activity of visuomotor neurons, suggesting that social information can modulate neuronal activity in this region of the motor cortex (Coudé et al. 2016)

Most of the human kinematic studies were motivated by Gibsonian ideas that object perception directly elicits action possibilities (Gibson, 1979). Together these findings support the idea that the activation of affordances is modulated not only by physical object properties but also by the social context. Our findings complement such studies by providing key information in terms of neurophysiological mechanisms.

Mouth condition preference

The neurons preferring the *Mouth* condition have a stronger response in their preferred condition compared to the neurons preferring the *Container* or *Hand* conditions. Also, at a population level, the discharge in the *Mouth* condition is stronger among the non-modulated neurons. This could be due to the fact that the *Mouth* condition is a more natural one and corresponds to a behavior that is already present in the repertoire of the monkey. Indeed, neurons preferring “Grasping to eat” were found to represent the majority of goal-modulated neurons in previous studies (Fogassi et al., 2005; Bonini et al., 2010). Long-train intracortical microstimulation of this premotor region elicits hand-to-mouth movements similar to the actions naturally performed by the monkeys (Graziano et al., 2002, 2005). The action goal preferences of some F5 neurons shown in this study could thus constitute a possible mechanism underlying the organization of natural actions in this region. The strength of the neuronal firing in the *Mouth* condition, consisting of a natural and very motorically-familiar action, likely reflects this organization and the overlapping cortical representation of hand and mouth (Rizzolatti et al. 1988; Maranesi et al 2012).

Latency of goal discrimination

A large proportion of the goal-modulated neurons recorded in the present study are characterized by an early selectivity for the goal. Most of these neurons display differential activity before the time at which the hand makes contact with the object (78%). These neurons therefore discriminate between the forthcoming goal during the time at which the hand approaches the object and the finger aperture is “preshaped” to match the physical characteristics of the object (Pellegrino et al., 1989). Neurophysiological studies on the grip selectivity of F5 motor and visuomotor neurons have shown that grip type selectivity appears at a very early stage of grasping, and progressively increases as the grasping action progresses (Murata et al., 1997; Raos, 2005; Umiltà et al., 2007). The present data suggest that, beyond coding grip type and being influenced by the goal of the action, some F5 neurons are modulated by the social context in which the grasping action is made. Overall, it

is not surprising to find early goal-discrimination in the interactive condition because reactive, feedback-based adjustments of one's own movement likely cannot support the fine-tuned temporal contingency required by on-line interpersonal coordination. In real life interactions, cooperative and competitive agents cannot simply react to a partner's behavior but must anticipate it for a successful interaction (Knoblich and Jordan, 2003).

Possible effects of reward delay

Although the total reward quantity (liquid and solid) is the same in all conditions, the temporal proximity of the solid reward presentation to the grasping action in the *Mouth* condition is a possible factor that could have had an impact on the neuronal discharge. In the *Mouth* condition, the solid reward delivery was temporally close to the grasping action (the monkey brought the food to the mouth and ate it right after having grasped it), and the monkey received a juice reward afterward. In the *Container* and *Hand* conditions, the reward sequence was inverted. The time lapse between the two rewards were short (about 1 second) and spurious factors, such as motivation and reward expectancy can be reasonably ruled out.

Limits of the present experiments and future directions

The present experiment was a first step in investigating the influence of social interactions on PMv neuronal response. In order to avoid any visual stimulation (experimenter hand movement), the interaction between the monkey and the experimenter was kept minimal. The experimenter remained passive with the hand open, palm up, in the same location where the container was in the *Container* condition. The monkey was required to place the object in the experimenter's hand and therefore, its action was stereotypical and did not require coordination between the monkey and the experimenter. We should also take into account the fact that we have tested a limited number of conditions and this might have contributed to the reduced number of neurons selective for the social condition. For example, we did not vary the experimenter's hand configuration (e.g. an open hand but with the finger configuration required to grasp a small object from the monkey's hand) or the spatial location. It is therefore possible that other neurons could potentially show selectivity for the social condition for various combinations of hand configuration and location. Lastly, as already mentioned above, area F5 contains a considerable percentage (47%) of visuomotor neurons that are modulated by social cues (i.e. gaze; Coudé et al. 2016). The present finding further supports the idea that social cue modulation is present in area F5. The use of more elaborate interactions in future experiments could potentially reveal even stronger preferences with richer specificity patterns, which could help disentangle the role of the social cues themselves from the decision of the monkey to interact.

Acknowledgements:

This research was supported by NICHD grant (NIH P01HD064653).

REFERENCES

Ballesta S, Duhamel J-R (2015) Rudimentary empathy in macaques' social decision-making. *Proc Natl Acad Sci* 112:15516–15521 [PubMed: 26621711]

- Bonini L, Rozzi S, Ugolotti Serventi F, Simone L, Ferrari PF, Fogassi L (2010) Ventral premotor and inferior parietal cortices make distinct contribution to action organization and intention understanding. *Cereb Cortex* 20:1372–1385. [PubMed: 19805419]
- Bonini L, Ugolotti Serventi F, Simone L, Rozzi S, Ferrari PF, Fogassi L (2011) Grasping neurons of monkey parietal and premotor cortices encode action goals at distinct levels of abstraction during complex action sequences. *J Neurosci* 31:5876–5886. [PubMed: 21490229]
- Calinski T, Harabasz J (1974) A dendrite method for cluster analysis. *Commun Stat - Theory Methods* 3:1–27.
- Chen G, Jaradat SA, Banerjee N, Tanaka TS, Ko MSH, Zhang MQ (2002) Evaluation and comparison of clustering algorithms in analyzing ES cell gene expression data. *Stat Sin* 12:241–262.
- Chinellato E, Castiello U, Sartori L (2015) Motor interference in interactive contexts. *Front Psychol* 6:791. [PubMed: 26113835]
- Clarke K, Gorley R (2001) Primer version 5 PRIMER-E statistical package, Plymouth, UK
- Coudé G, Festante F, Cilia A, Loiacono V, Bimbi M, Fogassi L, Ferrari PF (2016) Mirror neurons of ventral premotor cortex are modulated by social cues provided by others ' gaze. *J Neurosci* 36:3145–3156. [PubMed: 26985026]
- Davies DL, Bouldin DW (1979) A Cluster Separation Measure. *IEEE Trans Pattern Anal Mach Intell* PAMI-1:224–227.
- Deaner RO, Khera AV, Platt ML (2005) Monkeys pay per view: Adaptive valuation of social images by rhesus macaques. *Curr Biol* 15:543–548. [PubMed: 15797023]
- Dolk T, Hommel B, Colzato LS, Schütz-Bosbach S, Prinz W, Liepelt R (2014) The joint Simon effect: a review and theoretical integration. *Front Psychol* 5:1–10. [PubMed: 24474945]
- Emery NJ (2000) The eyes have it: the neuroethology, function and evolution of social gaze. *Neurosci Biobehav Rev* 24:581–604. [PubMed: 10940436]
- Fogassi L, Ferrari PF, Gesierich B, Rozzi S, Chersi F, Rizzolatti G (2005) Parietal lobe: from action organization to intention understanding. *Science* 308:662–667. [PubMed: 15860620]
- Fujii N, Hihara S, Iriki A (2007) Dynamic social adaptation of motion-related neurons in primate parietal cortex. *PLoS One* 2:e397. [PubMed: 17460764]
- Gibson JJ (1979) The ecological approach to visual perception Boston: Houghton-Mifflin.
- Graziano MSA, Aflalo TNS, Cooke DF (2005) Arm movements evoked by electrical stimulation in the motor cortex of monkeys. *J Neurophysiol* 94:4209–4223. [PubMed: 16120657]
- Graziano MSA, Taylor CSR, Moore T, Cooke Dylan F (2002) The cortical control of movement revisited. *Neuron* 36:349–362. [PubMed: 12408840]
- Hartigan JA (1975) Clustering algorithms New York: J Wiley & Sons.
- Jones D (2014) The Fathom Toolbox for MATLAB: software for multivariate ecological and oceanographic data analysis Open source Toolbox available at: <https://www.marine.usf.edu/user/djones/matlab/matlab.html>
- Kaufman L, Rousseeuw PJ (1990) Finding groups in data: an introduction to cluster analysis Antwerp: Wiley-Interscience.
- Kayaert G, Biederman I, Vogels R (2005) Representation of regular and irregular shapes in macaque inferotemporal cortex. *Cereb Cortex* 15:1308–1321. [PubMed: 15616128]
- Klein JT, Shepherd SV, Platt ML (2009) Social attention and the brain. *Curr Biol* 19:R958–62. [PubMed: 19889376]
- Knoblich G, Jordan JS (2003) Action coordination in groups and individuals: learning anticipatory control. *J Exp Psychol Learn Mem Cogn* 29:1006–1016. [PubMed: 14516231]
- Krzanowski W, Lai Y (1988) A criterion for determining the number of groups in a data set using sum-of-squares clustering. *Biometrics* 44:23–34.
- Maranesi M, Rodà F, Bonini L, Rozzi S, Ferrari PF, Fogassi L, Coudé G (2012) Anatomic-functional organization of the ventral primary motor and premotor cortex in the macaque monkey. *Eur J Neurosci* 36:3376–3387. [PubMed: 22891812]
- Mazurek KA, Rouse AG, Schieber MH (2018) Mirror neuron populations represent sequences of behavioral epochs during both execution and observation. *J Neurosci*:3481–17.

- Murata A, Fadiga L, Fogassi L, Gallese V, Raos V, Rizzolatti G, Keysers C, Gazzola V, B PTRS, Cattaneo L, Fabbri-destro M, Rozzi S, Soekadar SR, Witkowski M, Birbaumer N, Cohen LG (1997) Object Representation in the Ventral Premotor Cortex (Area F5) of the Monkey. *J Neurophysiol* 78:2226–2230. [PubMed: 9325390]
- Nummenmaa L, Calder AJ (2009) Neural mechanisms of social attention. *Trends Cogn Sci* 13:135–143. [PubMed: 19223221]
- Pellegrino JW, Klatzky RL, McCloskey BP (1989) Timecourse of preshaping for functional responses to objects. *J Mot Behav* 21:307–316. [PubMed: 15136267]
- Raos V (2005) Functional Properties of Grasping-Related Neurons in the Ventral Premotor Area F5 of the Macaque Monkey. *J Neurophysiol* 95:709–729. [PubMed: 16251265]
- Rizzolatti G, Camarda R, Fogassi L, Gentilucci M, Luppino G, Matelli M (1988) Functional organization of inferior area 6 in the macaque monkey - II. Area F5 and the control of distal movements. *Exp Brain Res* 71:491–507. [PubMed: 3416965]
- Rizzolatti G, Cattaneo L, Fabbri-Destro M, Rozzi S (2014) Cortical mechanisms underlying the organization of goal-directed actions and mirror neuron-based action understanding. *Physiol Rev* 94:655–706. [PubMed: 24692357]
- Rizzolatti G, Fogassi L, Gallese V (2009) The mirror neuron system: A motor-based mechanism for action and intention understanding. In: *The cognitive neurosciences* (Gazzaniga M, ed), pp 625–640. Cambridge, MA: MIT Press.
- Rizzolatti G, Rizzolatti G, Fogassi L, Fogassi L, Gallese V, Gallese V (2002) Motor and cognitive functions of the ventral premotor cortex. *Curr Opin Neurobiol* 12:149–154. [PubMed: 12015230]
- Rozzi S, Ferrari PF, Bonini L, Rizzolatti G, Fogassi L (2008) Functional organization of inferior parietal lobule convexity in the macaque monkey: electrophysiological characterization of motor, sensory and mirror responses and their correlation with cytoarchitectonic areas. *Eur J Neurosci* 28:1569–1588. [PubMed: 18691325]
- Sartori L, Cavallo A, Bucchioni G, Castiello U (2011) Corticospinal excitability is specifically modulated by the social dimension of observed actions. *Exp Brain Res* 211:557–568. [PubMed: 21472443]
- Scorolli C, Miatton M, Wheaton LA, Borghi AM (2014) I give you a cup, I get a cup: A kinematic study on social intention. *Neuropsychologia* 57:196–204. [PubMed: 24680723]
- Sebanz N, Bekkering H, Knoblich G (2006) Joint action: Bodies and minds moving together. *Trends Cogn Sci* 10:70–76. [PubMed: 16406326]
- Shepard R (1980) Multidimensional Scaling, Tree-Fitting, and Clustering. *Science* (80-) 210:390–398.
- Strehl A (2002) Relationship-based clustering and cluster ensembles for high-dimensional data mining. Doctoral dissertation, University of Texas at Austin
- Umiltà M a, Escola L, Intskirveli I, Grammont F, Rochat M, Caruana F, Jezzini A, Gallese V, Rizzolatti G (2008) When pliers become fingers in the monkey motor system. *Proc Natl Acad Sci U S A* 105:2209–2213. [PubMed: 18238904]
- Umiltà MA, Brochier T, Spinks RL, Lemon RN (2007) Simultaneous recording of macaque premotor and primary motor cortex neuronal populations reveals different functional contributions to visuomotor grasp. *J Neurophysiol* 98:488–501. [PubMed: 17329624]
- Yoshida K, Saito N, Iriki A, Isoda M (2011) Representation of others' action by neurons in monkey medial frontal cortex. *Curr Biol* 21:249–253. [PubMed: 21256015]
- Zhou F, Torre F (2009) Canonical Time Warping for Alignment of Human Behavior. In: *Advances in Neural Information Processing Systems*, 22, Proceedings of the Neural Information Processing Systems Conference. (pp. 2286–2294).

SIGNIFICANCE STATEMENT

Our study suggests that the activity of grasping neurons in the premotor cortex is modulated by social context. These findings give neurophysiological support to the idea of social affordances in which the social context surrounding objects, rather than their physical properties alone, elicits action possibilities.

Figure 1.

Task and conditions. A. Task illustration. A-I) The monkey has her right hand at rest on the starting handle for at least 1000ms (baseline). A transparent barrier is in place between the monkey and the target object. A-II) The barrier is removed (“go” signal). The monkey reaches for and grasps the target object. A-III) The monkey places the object in the container (or either place in the hand of the experimenter or eat, according to the experimental condition tested). A-IV) A liquid and a solid reward is delivered if the task is successfully executed. B) Experimental conditions. B-I) *Container*: Grasping to place in a container; B-II) *Hand*: Grasping to place in the hand of an experimenter, and B-III). *Mouth*: Grasping to eat.

Of all motor neurons

- Non modulated by goal n=257 (59%)
- Modulated by goal n=180 (41%)

Of modulated neurons

- Container n=30 (17%)
- Container and Hand n=30 (17%)
- Hand n=18 (10%)
- Hand and Mouth n=20 (11%)
- Mouth n=66 (36%)
- Mouth and Container n=16 (9%)

Figure 2.

Proportion of neuronal preferences. Overall proportion of F5 motor neurons that are goal-modulated. Forty-one percent (n=180) of all of the neurons had a significant preferential modulation for one or two of the three contexts, (i.e., either when the goal was placing in a container, placing in the hand or eating).

Figure 3.

Examples of goal-modulated neurons. Unit 146 has a preference for the *Container* condition. Unit 54 has a preference for the *Mouth* condition. Unit 174 has a preference for the *Hand* condition. The gray and yellow shaded areas represent the baseline and the grasping epochs, respectively. The dark yellow ticks represent the time at which the hand released the starting handle. The hand touched the object to grasp at time 0.

Figure 4.

Goal-modulated neuron populations and discrimination latency. Neuronal response of *Goal-modulated* neurons during grasping in each condition A) *Container-modulated*, B) *Hand-modulated* and C) *Mouth-modulated*. Time zero represents the moment at which the contact between hand and object occurred during grasping. The two vertical dashed lines indicate the beginning and the end of the grasping epoch (350 ms). The thickness of the curves represents {plus minus} 1 SE. In inset, mean normalized neuronal discharge during the grasping epoch of the goal-modulated neurons (two-way ANOVA (** $p < 0.01$, *** $p < 0.001$)). Bottom, mean firing activity of all *Container*-, *Hand*- and *Mouth*-goal-modulated neurons in their preferred condition. The yellow ticks represent the time bin at which each neuron started to discriminate between preferred and non-preferred conditions. There was no significant difference in discrimination time between the conditions.

Figure 5. Population analysis of A) *Goal-modulated* and B) *Non-modulated* motor neurons during grasping in the three conditions. Same conventions as in Figure 4).

Figure 6.

Cluster analysis of the neuronal population. A) Two-dimensional NMDS plot of the seven clusters with graphs of the mean discharge of the neurons belonging to each cluster. NMDS was applied to the matrix of the Euclidian distance between each pair of neuron activity vectors. The activity vector of each neuron was defined as the spike density function averaged over trials within each condition, and concatenated across conditions. The yellow shaded areas indicate bins where the neuronal discharge was significantly different between

the three conditions. B) Mean Euclidian distance between neurons across pairs of clusters.
C) Percentage breakdown of cluster membership for each neuronal classification.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Figure 7.

A) Two dimensional representation of the motor neuron ($n = 423$) population activity trajectory as obtained by non-metric multidimensional scaling (NMDS) assessing the similarities of population activity patterns. The start of the trial is around $(-1, 0)$, the end around $(0, 0.5)$. The dark lines show the trajectory of the population activity averaged over all trials within each condition, whereas the lighter lines show the trajectories from 100 runs of NMDS using random subsets of trials. Trial event times are denoted by different symbols (circle=movement onset, diamond=handle release, square=object touch, x=object lift), with black symbols denoting the event time in the trajectories averaged over all trials, and colored symbols showing the event time during the trajectories using random trial subsets. B) Dissimilarity matrices for each trial event showing the Euclidean distance between trial event positions for each run of NMDS using random trial subsets both within and between conditions.

Figure 8.

A) First and final frames of an example trial from the *Container* condition with the single-trial and average wrist, CMC, index PIP, and thumb MCP trajectories overlaid. The initial position of the wrist was used as the origin. B) Trajectories of the wrist, CMC, index PIP, and thumb MCP in each condition. The dark lines show the average trajectory over trials within a condition, and the lighter lines show the trajectories for each trial. C) Two-dimensional representation of the eight-dimensional (x, y position of the wrist, CMC, index PIP, and thumb MCP) kinematic trajectories generated by NMDS. The start of each trial is around (-100, 0) and the end is around (125, -30). The dark lines show the average trajectory for each condition and the lighter lines show the trajectory for each trial. The handle release and object touch times are denoted by diamonds and squares, respectively, with black symbols showing the event time in the average trajectories and colored symbols showing the event time during single trial trajectories. D) Dissimilarity matrices for the hand release and object touch events showing the Euclidean distance between the NMDS-derived event positions for each trial both within and between conditions.