Published in final edited form as:

J Org Chem. 2019 June 21; 84(12): 7871–7882. doi:10.1021/acs.joc.9b00745.

Ester Formation via Symbiotic Activation Utilizing Trichloroacetimidate Electrophiles

Nivedita S. Mahajani, Rowan I. L. Meador, Tomas J. Smith, Sarah E. Canarelli, Arijit A. Adhikari, Jigisha P. Shah, Christopher M. Russo, Daniel R. Wallach, Kyle T. Howard, Alexandra M. Millimaci, and John D. Chisholm*

Department of Chemistry, Syracuse University, 1-014 Center for Science and Technology, Syracuse, NY 13244

Abstract

Trichloroacetimidates are useful reagents for the synthesis of esters under mild conditions that do not require an exogenous promoter. These conditions avoid the undesired decomposition of substrates with sensitive functional groups that are often observed with the use of strong Lewis or Brønsted acids. With heating, these reactions have been extended to benzyl esters without electron donating groups. These inexpensive and convenient methods should find application in the formation of esters in complex substrates.

Graphical Abstract

Introduction

The synthesis of esters is a vital transformation in organic chemistry, and therefore significant effort has been devoted to developing high yielding and robust protocols for ester formation. These approaches are often characterized in three archetypal pathways:¹ 1) carbonyl activation (where an alcohol is added to an activated carboxylate), 2) carboxylate alkylation (where the carboxylic acid is deprotonated to improve nucleophilicity and added to an electrophile) and 3) symbiotic activation, where reagents combine to form a reactive ion pair that leads to the ester (Figure 1). While the carbonyl activation² and carboxylate alkylation³ pathways have been deeply explored and reviewed,⁴ investigations into reagents

*Corresponding Author: jdchisho@syr.edu.

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acs.joc.xxx. Copies of ¹H NMR spectra, ¹³C NMR spectra and chiral HPLC data (PDF).

The authors declare no competing financial interest.

that proceed through symbiotic activation processes have received less attention.⁵ These processes avoid the use of strong acids and bases that often characterize the other two methods, making these transformations more amenable for use in complex polyfunctional molecules that are often prepared in natural product synthesis and pharmaceutical environments.

In order to be characterized as a symbiotic activation process both the carboxylic acid and the esterification reagent must be activated under the reaction conditions. Usually this is accomplished by the esterification reagent being basic enough to deprotonate the carboxylic acid, with the resulting cationic salt being electrophilic enough to react with the newly formed carboxylate anion. Ideally the esterification reagent would be inert to other functional groups besides the carboxylate, and only react when activated by the proton transfer. Additionally, any side-products generated from the loss of the leaving group should also be as unreactive as possible so that sensitive functionality can be readily accommodated. This minimizes the need for exogenous reagents to buffer the reaction, keeping the reaction conditions mild as the strongest acid present is the carboxylic acid starting material.

Perhaps the most well-known example of esterification by symbiotic activation is the reaction of carboxylic acids with diazoalkanes (like **1**, Figure 2).⁶ These transformations have been shown to proceed through a tight ion pair intermediate that is formed after proton transfer from the carboxylic acid⁷ and generate only unreactive nitrogen gas as a side product. Unfortunately many diazoalkanes have a reputation as being toxic and energetic, limiting their use and often forcing researchers to form these reagents in situ to avoid issues with handling.⁸ Other reagents that undergo symbiotic activation include imidazole carbamates with structures like **2**,^{1,9} alkoxy- λ^6 -sulfanenitriles like **3**,¹⁰ some amide acetals (like **4** and **5**),¹¹ 1,3,5-*O*-benzyl triazines like **6**,¹² and isoureas like **7**.¹³ Most of these reagents require heating to effect their esterifications, usually in refluxing benzene or toluene solvent, with the exception being the isoureas which provide some esters at room temperature. Many of these esterification reagents have an imidate substructure as part of the reacting functionality, closely resembling a trichloroacetimidate like **8** for example. This core facilitates the displacement as it allows for the rearrangement of the imidate imine to the corresponding acetamide carbonyl, contributing a secondary thermodynamic driving force to the esterification and often leading to higher yields under mild conditions. Additionally the basic imine of the imidate is more readily activated by weak acids than carbonyl-based alkylation agents, leading to the selection of this common structural feature.

Given the commonality of the imidate core, we speculated that trichloroacetimidates may also be effective esterification reagents under near neutral conditions. Trichloroacetimidates are known to be powerful alkylating agents for carboxylic acids and alcohols when activated with a strong Lewis or Brønsted acid catalyst.¹⁴ These acid promoters may be problematic, however, as many complex substrates do not tolerate strong acids. A number of literature examples demonstrate that ester formation with some trichloroacetimidates do not require the addition of an exogenous promoter, including descriptions of spontaneous esterification with glycosyl imidates,¹⁵ 4-methoxybenzyl trichloroacetimidate,¹⁶ and 2-phenylisopropyl trichloroacetimidate.¹⁷

Results and Discussion

Spurred by the need for mild esterification reagents from inexpensive precursors that, we recently began to further investigate these trichloroacetimidate esterifications. These studies focused on systems that do not require additional Lewis or Brønsted acid additive, and also did not require heating, to increase the chances of compatibility with complex carboxylic acids where the reagents would typically find application. Initial investigations showed that both the DPM and PMB imidates could form esters at room temperature without the need for exogenous acid.¹⁸ Promoter free imidate esterifications were shown to undergo symbiotic activation with carboxylic acids, with the reactions being sensitive to exogenous base that prevented activation of the imidate by the carboxylic acid esterification partner. These esterifications were also shown to function well with sensitive carboxylic acids which decompose under more standard esterification conditions, including β -hydroxy carboxylic acids and other polyfunctional substrates.¹⁸ The imidates are simple to prepare from inexpensive precursors, formed at room temperature from the alcohol and trichloroacetonitrile with a catalytic amount of DBU.¹⁹ In addition, the trichloroacetamide side product may be removed by washing with aqueous NaOH solution, as the pKa of this acetamide is similar to a phenol. This is a significant advantage over isourea-based esterification reagents, which usually require chromatography to remove the urea side product. Given the ready availability of trichloroacetimidates (easily synthesized from inexpensive trichloroacetonitrile and the appropriate alcohol) the scope of these transformations with respect to the imidate has now been examined. This work has determined that a surprising number of stable, unactivated trichloroacetimidates provide the corresponding esters under promoter-free conditions in excellent yield.

Initially, the esterifications were studied with a number of structurally distinct trichloroacetimidates to determine the scope with regard to trichloroacetimidate electrophile. For this study benzoic acid was utilized as a common substrate (Table 1). The most simple alkyl trichloroacetimidates did not undergo esterification at room temperature or in refluxing toluene (unreacted starting materials were recovered, entries 1–3) with the exception being the *tert*-butyl trichloroacetimidate **13** (entry 4). Oddly the esterification with **13** and benzoic acid gave a higher yield at room temperature in DCM as opposed to refluxing toluene. This may be due to the propensity of imidate **13** to undergo elimination at elevated temperatures, decomposing to isobutylene and trichloroacetamide. Indeed, this has been reported to be a limitation of this imidate in the formation of *tert*-butyl amines.²⁰ Attempts to moderate the elimination by using nitromethane as a solvent, as has been reported to facilitate substitution in other systems,^{20–21} only gave poor conversion of the imidate to the ester, with mostly starting carboxylic acid being recovered.

Allylic and propargylic trichloroacetimidates were evaluated next. While allyl imidate **14** and propargyl trichloroacetimidate **16** were unreactive (with only starting materials being observed), prenyl trichloroacetimidate **15** gave a useful 73% yield of product at room temperature (entry 6). While nearly all of the product was the prenyl ester **15a**, a trace amount of the *tert*-prenyl isomer was detected in the crude ¹H NMR. Benzylic trichloroacetimidates are of special interest, as benzyl esters are some of the most common carboxylate protecting groups utilized in organic synthesis.²² Benzyl trichloroacetimidate **17**

was unreactive at room temperature, but gave a 76% yield of product when heated in toluene. Adding electron donating groups to the benzene ring led to increased yields of the esterification, as shown with the 2,4,6-trimethylbenzyl imidate **18** and 2-methoxybenzyl trichloroacetimidate **19**. While imidate **19** did give some product at room temperature, the yield was moderate, with starting materials being recovered. Alternatively, many other electron rich benzylic trichloroacetimidates formed esters at room temperature in DCM with excellent conversion. These conditions were employed successfully for 4-methoxybenzyl trichloroacetimidate **20**, 2,4-dimethoxybenzyl trichloroacetimidate **21** and 3,4-dimethoxybenzyltrichloroacetimidate **22** (Table 1, entries 11–13). With the dimethoxybenzyl systems the esterifications were typically complete in only a few hours, demonstrating that reactivity increases observed with multiple electron donating groups were additive. The ability to form dimethoxybenzyl esters is important, as these can be removed via oxidative methods (typically with DDQ^{4a,23}) whereas other benzyl esters are inert to these conditions. The furfuryl trichloroacetimidate **24** also gave an excellent yield of the corresponding ester (entry 15) by heating in toluene, as did the phenethyl imidate **25**. The diphenylmethyl trichloroacetimidate formed the ester at room temperature, providing a virtually quantitative yield of the DPM ester. In contrast, benzylic imidates with electron-withdrawing substituents were unreactive under these promoter-free conditions, as shown by the lack of ester formation with the 4-nitrobenzyl trichloroacetimidate **23** (entry 14) which returned only unreacted starting material. From these results it's apparent that the esterifications proceed best with electron rich alkyl groups that are capable of stabilizing a positive charge or partial positive charge. The more reactive imidates can form esters at room temperature, while the less reactive systems must be heated in order for the esterification to proceed. While electronic effects were apparent, sterics also influenced reactivity as can be seen in the difference in yield when employing imidates **19** and **20**. The lower reactivity of **19** is consistent with the addition of a substituent near the reacting benzylic position slowing the esterification due to steric hindrance.

With the scope of the reaction with regard to the trichloroacetimidate determined, a brief selection of carboxylic acids were esterified with some of the functional trichloroacetimidates from Table 1 to determine the effects of the carboxylic acid structure on the esterification reaction (Table 2). Initially the effects of the steric environment near the carboxylic acid were examined. Use of diphenylacetic acid **27** provided consistently high yields of the corresponding esters. The even more sterically encumbered 2-methyl-2-phenylpropionic acid **28** provided a lower yields when the hindered *tert*-butyl imidate **13** was employed, evidently due to unfavorable steric factors when both the acid and the imidate are hindered. Additionally, a number of substrates were then evaluated with respect to functionality. The 2-bromododecanoic acid **29** was esterified in good yield by all of the imidates used. No elimination or other competing substitution reactions were observed under these reaction conditions. The ability to form esters in the presence of other protic functionality was also investigated. Mandelic acid **30** was esterified in good yield, demonstrating that the presence of an alcohol was well tolerated. The protected amino acid ($\pm$)-Boc-phenylalanine **31** was also esterified under these conditions in good yield. The propensity of alkene isomerization under these reaction conditions was also explored. Vinyl acetic acid **32** was esterified in good yield under both sets of reaction conditions without any

observed isomerization of the alkene. The exception was the *tert*-butyl ester of vinyl acetic acid **13g**, which proved to be quite volatile and therefore was difficult to obtain a high isolated yield. Cinnamic acid **33** also proved to be a good substrate, providing the corresponding esters in good yields. The *cis* alkenoic acid **34** was also investigated, as similar systems often undergoes *cis* / *trans* isomerization under esterification conditions.²⁴ No isomerization of the alkene was observed during room temperature esterifications in DCM with the tested imidates (Table 2, entries 33–36), demonstrating this a powerful method for the protection of esters without isomerization. Heterocycles like the pyridine in picolinic acid **35** were also well tolerated, with good yields being achieved with all four imidates tested.

The scope of the esterification reaction in refluxing toluene with benzyl trichloroacetimidate **17** was also explored (Table 3), as benzyl groups are common carboxylate protecting groups and given that alternative conditions are necessary for the formation of benzyl esters with imidate **17** some differences in functional group tolerance may be observed. These concerns were unfounded, however, as the same series of carboxylic acids underwent esterification in good yields. Importantly, no elimination products from the α -bromoacid **29** were observed, and no isomerization of the alkenes in carboxylates **32**, **33** or **34** was detected. The alcohol of mandelic acid **30** also remained unaffected. While some more reactive imidates (like the diphenylmethyl trichloroacetimidate **26**²⁵) have been reported to form ethers under similar conditions, imidate **17** is not reactive enough to access this reactivity manifold, and therefore alcohols are well tolerated under the reaction conditions.

The effects of these esterification reactions on a chirality center next to the carboxylate were also investigated. Chiral naproxen **27** was utilized for this section of the study, as this substrate can racemize rapidly as the chirality center is both benzylic and next to the electron-withdrawing carboxylate. Treatment of chiral naproxen with imidate **26** at room temperature in DCM gave the desired ester products in high yields (Scheme 1). Evaluation of the enantiopurity of these samples by chiral HPLC analysis showed that virtually no racemization had occurred. While these results ensured that there was no racemization in DCM at rt, it did not provide information about the possibility of racemization with less reactive benzylic imidates which required refluxing toluene to effect the esterification. To address this point, naproxen was heated in refluxing toluene with benzyl trichloroacetimidate **17** to provide a 79% yield of benzyl ester **17k**. Chiral HPLC analysis of **17k** showed that virtually no racemization had occurred with this substrate compared to a racemic sample (racemic samples of **26k** and **17k** were prepared by heating the esters with DBU in toluene).

The mechanism of the reaction was briefly probed by the use of two chiral imidate substrates in the esterification reaction. Chiral phenethyl trichloroacetimidate **25** has already been reported,²⁶ and was therefore employed in the esterification reaction. In addition, the imidate of (S)-(2-methoxyphenyl)phenylmethanol²⁷ (imidate **29**) was also prepared. During the course of the esterification with chiral **25** significant racemization was observed, with a completely racemic mixture being observed in refluxing toluene and a 71:29 mixture of enantiomers being observed in DCM at room temperature (Table 4). Similar results were observed with imidate **29**, with a nearly racemic mixture being isolated from the reaction in

refluxing toluene and a scalemic 67:33 mixture being isolated from the esterification in DCM at room temperature.

With the significant racemization that was observed with chiral imidates an SN2 type substitution mechanism was ruled out as being operative in these systems, as an SN2 mechanism proceeds with inversion. This left either a radical or a cationic mechanism as possibilities for the formation of the esters. Given that a radical substitution reaction should also proceed well with an electron poor imidate (like **23**), it seemed unlikely that a radical intermediate was involved in the transformation. The available evidence seems to implicate a cationic mechanism, as shown in Figure 3. Symbiotic activation of the imidate with the proton from carboxylic acid leads to intermediate **41**. Loss of trichloroacetamide **42** then produces the carbocation **44**, which is trapped by the carboxylate anion **43** to provide the ester product **45**. Further supporting this mechanism is that the addition of triethylamine to esterifications with imidate **20**^{18a} and **26**^{18b} effectively poisoned the reaction and halted the esterification. The more basic amine likely deprotonates the carboxylic acid, halting the activation of the imidate and disrupting the esterification. While carbocation formation is predicted to provide a racemic mixture, scalemic mixtures often result from cationic processes due to ion pairing.²⁸

While an SN1 mechanism is implicated for secondary benzylic trichloroacetimidates, less reactive trichloroacetimidates may form esters through different pathways. This is supported by the lack of Friedel-Crafts alkylation products in the results described in Table 3, even though benzyl trichloroacetimidate is known to alkylate aromatic systems like toluene when treated with strong Lewis acids which presumably proceeds through a carbocation.²⁹ Additionally, prenyl trichloroacetimidate **15** has been reported to provide a nearly 1:1 mixture of prenyl and *tert*-prenyl ethers in the presence of a strong Lewis acid (TMSOTf),³⁰ while in the esterification reaction with **15** (Table 1) the prenyl ester is highly favored and only a small amount of the *tert*-prenyl ester is observed. These results imply that some imidate esterifications with less reactive imidates may not be proceeding through a discrete carbocation. The possibility of a radical pathway was ruled out by repeating the esterification of benzoic acid with prenyl imidate **15** in the presence of 1 equiv of TEMPO, a known radical trap.³¹ This reaction provided a 50% yield of ester **15a**, so the esterification cannot be proceeding exclusively through a radical intermediate. Instead, less reactive trichloroacetimidates may esterify through an SN2 mechanism on a protonated imidate intermediate (like **41** when the R groups are small).

Conclusion

Esterification reactions of a number of structurally distinct trichloroacetimidate electrophiles have been studied. Only imidates that are precursors to stabilized carbocations function as reliable esterification reagents under promoter free conditions. Many of these systems undergo esterification at room temperature, although several others can be used by simple heating the trichloroacetimidate in refluxing toluene. No alkene isomerization or alkylation of other protic functional groups (like alcohols or amides) was observed under these esterification conditions, providing a mild method for the formation of an ester in the presence of complex functionality. The ability to form esters from imidates without the

addition of an acid or base promoter will be useful in the esterification of complex substrates that possess sensitive functionality.

Experimental

The methyl-2,2,2-trichloroacetimidate **10**, ethyl-2,2,2-trichloroacetimidate **11**, *t*-butyl-2,2,2-trichloroacetimidate **13**, allyl-2,2,2-trichloroacetimidate **14**, and benzyl-2,2,2-trichloroacetimidate **17** used in these studies were purchased from commercial sources. Cyclohexyl-2,2,2-trichloroacetimidate **12**,³² prenyl-2,2,2-trichloroacetimidate **15**,³³ propargyl-2,2,2-trichloroacetimidate **16**,³⁴ (2,4,6-trimethylphenyl)methyl-2,2,2-trichloroacetimidate **18**,³⁵ (4-methoxyphenyl)methyl-2,2,2-trichloroacetimidate **20**,^{18a} (3,4-dimethoxyphenyl)methyl-2,2,2-trichloroacetimidate **22**,³⁶ (4-nitrophenyl)methyl-2,2,2-trichloroacetimidate **23**,²¹ furfuryl-2,2,2-trichloroacetimidate **24**,³⁴ 1-phenethyl-2,2,2-trichloroacetimidate **25**,^{26a,37} diphenylmethyl-2,2,2-trichloroacetimidate **26**^{18b,38} and (2-methoxyphenyl)phenylmethyl-2,2,2-trichloroacetimidate **29**²¹ were synthesized as previously reported.

General Procedure for the synthesis of trichloroacetimidates from the corresponding alcohol:

A flame dried 25 mL round bottom flask was charged with the alcohol starting material (1 equiv) under argon. Dry DCM was then added to form a 0.5 M solution, and the flask was cooled to 0 °C. 1,8-Diazabicyclo[5.4.0]undec-7-ene (DBU, 0.2 equiv) was added to the solution, followed by trichloroacetonitrile (1.5 equiv). The reaction was monitored by TLC. After completion of the reaction, reaction mixture was concentrated and silica gel column chromatography was performed to provide the desired trichloroacetimidates.

(2-Methoxyphenyl)methyl-2,2,2-trichloroacetimidate (19).—Clear oil (1.49 g, 73%), purified by silica gel chromatography (4% ethyl acetate/1% triethylamine/ 95% hexanes). TLC R_f = 0.55 (10% ethyl acetate/ 90% hexane); ¹H NMR (300 MHz, CDCl₃) δ 8.38 (brs, 1H), 7.45 (dd, J = 7.5, 1.6 Hz, 1H), 7.32 (td, J = 8.0, 1.7 Hz, 1H), 6.98 (td, J = 7.5, 0.9 Hz, 1H), 6.91 (d, J = 8.2 Hz, 1H), 5.40 (s, 2H), 3.85 (s, 3H); ¹³C{¹H} NMR (100 MHz, CDCl₃) δ 162.8, 157.2, 129.4, 128.7, 124.0, 120.4, 110.4, 91.6, 66.6, 55.4. Anal. Calcd for C₁₀H₁₀Cl₃NO₂: C, 42.51; H, 3.57; N, 4.96. Found: C, 42.62; H, 3.74; N, 4.95.

(2,4-Dimethoxyphenyl)methyl-2,2,2-trichloroacetimidate (21).—Yellow oil (7.40 g, 99%), purified by silica gel chromatography (30% ethyl acetate/ 69% hexanes/ 1% triethylamine). ¹H NMR (400 MHz, CDCl₃) δ 8.33 (s, 1H), 7.33 (d, J = 8.4 Hz, 1H), 6.49 (t, J = 2.4 Hz, 2H), 5.30 (s, 2H), 3.81 (s, 3H), 3.81 (s, 3H); ¹³C{¹H} NMR (100 MHz, CDCl₃) δ 162.9, 161.3, 158.8, 130.6, 116.4, 104.0, 98.6, 98.4, 91.7, 66.6, 55.4; Anal. Calcd for C₁₁H₁₂Cl₃NO₃: C, 42.27; H, 3.87; N, 4.48. Found: C, 42.28; H, 3.56; N, 4.57.

General procedure for esterification by method A:

In a flame dried flask, the trichloroacetimidate (2 equiv) was dissolved in dichloromethane (0.25 M) under argon. The carboxylic acid (1 equiv) was then added and the mixture was stirred at room temperature. The reaction progress was monitored by thin layer

chromatography. After completion the reaction mixture was poured into 2N NaOH and extracted with DCM (3x). The combined organic extracts were then dried over sodium sulfate, filtered, and concentrated. The residue was purified by silica gel column chromatography to give the ester product.

General procedure for esterification by method B:

The carboxylic acid (1 equiv) and trichloroacetimidate (2 equiv) were placed in a flame dried round bottom flask under argon. Anhydrous toluene (0.25 M) was then added and the reaction was heated to reflux. The reaction progress was monitored by thin layer chromatography. After disappearance of the carboxylic acid the mixture was allowed to cool to rt, poured into 2N NaOH and extracted with DCM (3x). The combined organic extracts were then dried over sodium sulfate, filtered, and concentrated. The residue was purified by silica gel column chromatography to give the ester product.

tert-Butyl benzoate (13a).³⁹—Clear oil (0.097 g, 84%), purified by silica gel chromatography (1% ethyl acetate/99% hexane). TLC R_f = 0.53 (10% ethyl acetate/ 90% hexane); ^1H NMR (300 MHz, CDCl_3) δ 8.01–7.91 (m, 2H), 7.55–7.49 (m, 1H), 7.41 (t, J = 7.6 Hz, 2H), 1.60 (s, 9H); $^{13}\text{C}\{^1\text{H}\}$ NMR (125 MHz, CDCl_3) δ 165.8, 132.4, 132.0, 129.4, 128.2, 81.0, 28.2.

tert-Butyl 2,2-diphenylacetate (13b).⁴⁰—White solid (0.064 g, 91%), purified by silica gel chromatography (2% ethyl acetate/98% hexanes). TLC R_f = 0.68 (10% ethyl acetate/90% hexanes); ^1H NMR (400 MHz, CDCl_3) δ 7.30–7.23 (m, 10 H), 4.91 (s, 1H), 1.44 (s, 9H). $^{13}\text{C}\{^1\text{H}\}$ NMR (125 MHz, CDCl_3) δ 171.7, 139.2, 128.6, 128.5, 127.0, 81.3, 58.0, 28.0.

tert-Butyl 2-methyl-2-phenylpropanoate (13c).⁴¹—Clear oil (0.087 g, 30%), purified by silica gel chromatography (3% ethyl acetate/97% hexanes). TLC R_f = 0.71 (10% ethyl acetate/90% hexanes); ^1H NMR (300 MHz, CDCl_3) δ 7.33–7.26 (m, 5H), 1.53 (s, 6H), 1.38 (s, 9H). $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CDCl_3) δ 176.2, 145.5, 128.4, 126.6, 125.8, 80.5, 47.3, 28.0, 26.7.

tert-Butyl 2-bromododecanoate (13d).⁴²—Amorphous solid (0.091 g, 94%), purified by silica gel chromatography (100 % diethyl ether). TLC R_f = 0.54 (100% hexanes); ^1H NMR (400 MHz, CDCl_3) δ 4.08 (t, J = 7.1 Hz, 1H), 2.02–1.87 (m, 2H), 1.46 (s, 9H), 1.41–1.11 (m, 16 H), 0.86 (t, J = 5.5 Hz, 3H); $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CDCl_3) δ 169.0, 82.2, 47.9, 35.0, 31.9, 29.54, 29.47, 29.33, 29.30, 28.9, 27.8, 27.3, 22.7, 14.1.

(±)-tert-Butyl-2-hydroxy-2-phenylacetate (13e).⁴³—White solid (0.079 g, 59%), purified by silica gel chromatography (5% ethyl acetate/95% hexanes). TLC R_f = 0.65 (50% ethyl acetate/50% hexanes); ^1H NMR (400 MHz, CDCl_3) δ 7.41–7.30 (m, 5H), 5.03 (d, J = 6.0 Hz, 1H), 3.53 (d, J = 6.0 Hz, 1H), 1.40 (s, 9H); $^{13}\text{C}\{^1\text{H}\}$ NMR (125 MHz, CDCl_3) δ 172.9, 139.0, 128.4, 128.1, 126.4, 83.1, 73.0, 27.8.

(±)-(tert-Butyl-2-[(tert-butoxycarbonyl)amino]phenylpropionate (13f).⁴⁴—Clear oil (0.90 g, 90%), purified by silica gel chromatography (4% ethyl acetate/96% hexanes).

TLC R_f = 0.75 (10% ethyl acetate/90% hexanes); ^1H NMR (300 MHz, CDCl_3) δ 7.31–7.16 (m, 5H), 5.00 (brs, 1H), 4.45–4.44 (m, 1H), 3.05 (d, J = 5.8 Hz, 2H), 1.42 (s, 9H), 1.40 (s, 9H); $^{13}\text{C}\{^1\text{H}\}$ NMR (75 MHz, CDCl_3) δ 171.0, 155.1, 136.4, 129.6, 128.3, 126.8, 82.0, 79.6, 54.9, 38.6, 28.3, 27.9.

2,2-Dimethylpropyl but-3-enoate (13g).⁴⁵—Clear colorless oil (70 mg, 35% yield), purified by silica gel column chromatography (10% diethyl ether, 90% pentane). TLC R_f = 0.76 (10% ethyl acetate/ 90% hexanes); ^1H NMR (400 MHz, CDCl_3) δ 5.88–5.78 (m, 1H), 5.06 (dq, J = 14.0, 1.6 Hz, 2H), 2.93 (d, J = 6.9 Hz, 2H) 1.38 (s, 9H); $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz CDCl_3) δ 170.8, 130.9, 117.9, 80.5, 40.4, 28.0.

(E)-tert-Butyl cinnamate (13h).⁴⁶—Clear oil (0.052 g, 60%), purified by silica gel chromatography (1–4% ethyl acetate/hexanes). TLC R_f = 0.71 (10% ethyl acetate/90% hexanes); ^1H NMR (400 MHz, CDCl_3) δ 7.59 (d, J = 16.0 Hz, 1H), 7.50–7.49 (m, 2H), 7.36–7.36 (m, 3H), 6.37 (d, J = 16.0 Hz, 1H), 1.54 (s, 9H); $^{13}\text{C}\{^1\text{H}\}$ NMR (125 MHz, CDCl_3) δ 166.3, 143.5, 134.7, 129.9, 128.8, 128.0, 120.2, 80.5, 28.2.

tert-Butyl (2Z)-3-(2-methoxyphenyl)prop-2-enoate (13i).—Clear oil (0.100 g, 76%), purified by silica gel chromatography (8% ethyl acetate/92% hexanes). TLC R_f = 0.52 (20% ethyl acetate/80% hexanes); IR (film, cm^{-1}) ν_{max} 2979, 1716, 1629, 1463, 1110, 1049, 738; ^1H NMR (400 MHz, CDCl_3) δ 7.47 (d, J = 7.3 Hz, 1H), 7.31–7.27 (m, 1H), 7.07 (d, J = 12.3 Hz, 1H), 6.94–6.86 (m, 2H), 5.91 (d, J = 12.4 Hz, 1H), 3.84 (s, 3H), 1.39 (s, 9H); $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CDCl_3) δ 165.8, 157.0, 137.5, 130.6, 130.0, 124.7, 122.3, 120.0, 110.2, 80.3, 55.5, 28.0. Anal. Calcd for $\text{C}_{14}\text{H}_{18}\text{O}_3$: C, 71.77; H, 7.74. Found: C, 72.00; H, 7.36.

tert-Butyl 2-pyridinecarboxylate (13j).⁴⁷—Clear oil (0.095 g, 81%), purified by silica gel chromatography (45% ethyl acetate/55% hexanes). TLC R_f = 0.40 (50% ethyl acetate/50% hexanes); ^1H NMR (400 MHz, CDCl_3) δ 8.74 (dd, J = 4.7, 0.7 Hz, 1H), 8.19 (d, J = 7.9 Hz, 1H), 7.80 (td, J = 7.7, 1.8 Hz, 1H), 7.42 (ddd, J = 7.6, 4.8, 1.2 Hz, 1H), 1.64 (s, 9H); $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CDCl_3) δ 164.5, 150.0, 149.8, 137.0, 126.6, 125.0, 82.5, 28.3.

3-Methylbut-2-en-1-yl benzoate (5a).⁴⁸—Clear colorless oil (0.126 g, 63% yield), purified by silica gel column chromatography (10% ethyl acetate, 90% hexanes). TLC R_f = 0.60 (10% ethyl acetate/ 90% hexanes); ^1H NMR (400 MHz, CDCl_3) δ 8.08–8.06 (m, 2H), 7.58–7.53 (m, 1H), 7.44 (t, J = 7.8 Hz, 2H) 5.52–5.48 (m, 1H), 4.85 (d, J = 7.2 Hz, 2H), 1.80 (d, J = 6.3 Hz, 6H); $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz CDCl_3) δ 166.6, 139.1, 132.8, 130.5, 129.6, 128.3, 118.8, 61.9, 25.8, 18.1.

(Phenyl)methyl benzoate (17a).⁴⁹—Clear oil (0.082 g, 58%), purified by silica gel chromatography (2% ethyl acetate/98% hexane). TLC R_f = 0.57 (10% ethyl acetate/ 90% hexane); ^1H NMR (400 MHz, CDCl_3) δ 8.08 (d, J = 7.2 Hz, 2H), 7.56 (t, J = 7.4 Hz, 1H), 7.46–7.34 (m, 7H), 5.37 (s, 2H); $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CDCl_3) δ 166.4, 136.1, 133.0, 130.2, 129.7, 128.6, 128.4, 128.2, 128.2, 66.7.

(Phenyl)methyl 2,2-diphenylacetate (17b).⁵⁰—White solid (0.079 g, 88%), purified by silica gel chromatography (2% ethyl acetate/98% hexanes). TLC R_f = 0.60 (10% ethyl acetate/90% hexanes); ^1H NMR (400 MHz, CDCl_3) δ 7.29–7.25 (m, 15H), 5.17 (s, 2H), 5.07 (s, 1H); $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CDCl_3) δ 172.4, 138.7, 135.8, 128.73, 128.68, 128.24, 128.21, 127.36, 127.33, 68.0, 57.1.

(Phenyl)methyl 2-methyl-2-phenylpropanoate (17c).—Clear oil (0.157 g, 94%), purified by silica gel chromatography (4% ethyl acetate/96% hexanes). TLC R_f = 0.50 (100% hexanes); IR (film, cm^{-1}) ν_{max} 3089, 2976, 1729, 1497, 1425, 1412, 737; ^1H NMR (400 MHz, CDCl_3) δ 7.32–7.19 (m, 10H), 5.10 (s, 2H), 1.60 (s, 6H); $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CDCl_3) δ 176.5, 144.5, 136.2, 128.40, 128.36, 127.9, 127.7, 126.7, 125.7, 66.4, 46.6, 26.5. Anal. Calcd for $\text{C}_{17}\text{H}_{18}\text{O}_2$: C, 80.28; H, 7.13. Found: C, 80.58; H, 7.03.

(Phenyl)methyl 2-bromododecanoate (17d).—Clear oil (0.193 g, 80%), purified by silica gel chromatography (3% ethyl acetate/97% hexane). TLC R_f = 0.68 (10% ethyl acetate/90% hexanes); IR (film, cm^{-1}) ν_{max} 3054, 2928, 2305, 1740, 1422, 1265, 738; ^1H NMR (400 MHz, CDCl_3) δ 7.39–7.35 (m, 5H), 5.20 (s, 2H), 4.25 (t, J = 7.3 Hz, 1H), 2.06–1.95 (m, 2H), 1.35–1.21 (m, 16H), 0.88 (t, J = 6.6 Hz, 3H); $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CDCl_3) δ 169.7, 135.2, 128.8, 128.6, 128.5, 128.3, 128.2, 70.7, 67.5, 46.0, 34.9, 31.9, 29.5, 29.5, 29.3, 28.8, 27.2, 22.7, 14.1. Anal. Calcd for $\text{C}_{19}\text{H}_{29}\text{BrO}_2$: C, 61.79; H, 7.91. Found: C, 61.69; H, 8.01.

(±)-(Phenyl)methyl-2-hydroxy-2-phenylacetate (17e).^{13b}—White solid (0.209 g, 88%), purified by silica gel chromatography (10% ethyl acetate/90% hexanes). TLC R_f = 0.48 (30% ethyl acetate/70% hexanes); ^1H NMR (400 MHz, CDCl_3) δ 7.42–7.30 (m, 8H), 7.21–7.19 (m, 2H), 5.25–5.12 (m, 3H), 3.42 (d, J = 5.8 Hz, 1H); $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CDCl_3) δ 173.5, 138.2, 135.0, 128.60, 128.56, 128.5, 128.4, 128.0, 126.6, 73.0, 67.7.

Boc-phenylalanine (Phenyl)methyl ester (17f).⁵¹—Clear oil (0.251 g, 94%), purified by silica gel chromatography (10% ethyl acetate/90% hexanes). TLC R_f = 0.57 (20% ethyl acetate/80% hexanes); ^1H NMR (300 MHz, CDCl_3) δ 7.36–7.21 (m, 8H), 7.06–7.03 (m, 2H), 5.13 (q, J = 12.3 Hz, 2H), 4.97 (d, J = 7.7 Hz, 1H), 4.63 (q, J = 7.7 Hz, 1H), 3.08 (t, J = 5.2 Hz, 2H), 1.41 (s, 9H); $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CDCl_3) δ 171.8, 155.1, 135.9, 135.2, 129.4, 128.6, 128.6, 127.0, 79.9, 67.1, 54.5, 38.3, 28.3.

(Phenyl)methyl but-3-enoate (17g).⁵²—Clear oil (0.180 g, 88%), purified by silica gel chromatography (2% ethyl acetate/ 98% hexanes). TLC R_f = 0.62 (10% ethyl acetate/ 90% hexanes); ^1H NMR (400 MHz, CDCl_3) δ 7.39 (m, 5H), 6.00–5.90 (m, 1H), 5.20–5.14 (m, 4H), 3.15 (d, J = 7.0 Hz, 2H); $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CDCl_3) δ 171.4, 135.9, 130.1, 128.6, 128.3, 128.2, 118.7, 66.5, 39.1.

(Phenyl)methyl cinnamate (17h).⁵³—Viscous oil (0.120 g, 77%), purified by silica gel chromatography (2% ethyl acetate/98% hexanes). TLC R_f = 0.63 (10% ethyl acetate/90% hexanes); ^1H NMR (300 MHz, CDCl_3) δ 7.66 (d, J = 15.9 Hz, 1H), 7.47–7.44 (m, 2H), 7.36–7.27 (m, 8H), 6.42 (d, J = 16.2 Hz, 1H), 5.18 (s, 2H); $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CDCl_3) δ 166.8, 145.2, 136.1, 134.4, 130.4, 128.9, 128.6, 128.3, 128.3, 128.1, 117.9, 66.4.

Benzyl (2Z)-3-(2-methoxyphenyl)prop-2-enoate (17i).—Clear oil (0.120 g, 80%), purified by silica gel chromatography (6% ethyl acetate/94% hexanes). TLC R_f = 0.38 (20% ethyl acetate/80% hexanes); IR (film, cm^{-1}) ν_{max} 3054, 1721, 1600, 1488, 1265, 1157, 1028, 745; ^1H NMR (400 MHz, CDCl_3) δ 7.52 (d, J = 7.5 Hz, 1H), 7.32–7.18 (m, 7H), 6.89–6.85 (m, 2H) 6.03 (d, J = 12.4 Hz, 1H), 5.13 (s, 2H), 3.81 (s, 3H); $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CDCl_3) δ 166.3, 161.2, 158.9, 157.2, 138.9, 131.3, 130.8, 130.3, 124.1, 120.0, 120.0, 116.8, 110.2, 104.0, 98.5, 61.3, 55.4. Anal. Calcd for $\text{C}_{17}\text{H}_{16}\text{O}_3$: C, 76.10; H, 6.01. Found: C, 76.35; H, 5.64.

(Phenyl)methyl picolinate (17j).⁵⁴—Yellow oil (0.120 g, 57%), purified by silica gel chromatography (50% ethyl acetate/50% hexanes). TLC R_f = 0.47 (50% ethyl acetate/50% hexanes); ^1H NMR (400 MHz, CDCl_3) δ 8.77 (d, J = 4.6 Hz, 1H), 8.13 (d, J = 7.9 Hz, 1H), 7.82 (dt, J = 7.7, 1.7 Hz, 1H), 7.50–7.45 (m, 3H), 7.40–7.33 (m, 3H), 5.46 (s, 2H); $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CDCl_3) δ 165.0, 150.0, 148.0, 137.0, 135.6, 128.6, 128.6, 128.4, 126.9, 125.3, 67.5.

(S)-(Phenyl)methyl-2-(6-methoxynaphthalen-2-yl)propanoate (17k).—Yellow solid (0.110 g, 79%), purified by silica gel chromatography (2% ethyl acetate/98% hexanes). $[\alpha]_{\text{D}}^{23}$ –3.7 (c 0.35, DCM); mp = 72–73 °C; TLC R_f = 0.48 (10% ethyl acetate/90% hexanes); IR (film, cm^{-1}) ν_{max} 3055, 2982, 1733, 1634, 1265, 738; ^1H NMR (400 MHz, CDCl_3) δ 7.70–7.65 (m, 3H), 7.41 (d, J = 1.6 Hz, 1H), 7.28–7.25 (m, 5H), 7.14–7.11 (m, 2H), 5.11 (q, J = 12.4 Hz, 2H), 3.93–3.88 (m, 4H), 1.59 (d, J = 7.2 Hz, 3H); $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CDCl_3) δ 174.5, 157.6, 136.0, 135.6, 133.7, 129.3, 128.9, 128.5, 128.1, 127.9, 127.1, 126.3, 126.0, 119.0, 105.6, 66.5, 55.3, 45.5, 18.5. Anal. Calcd for $\text{C}_{17}\text{H}_{16}\text{O}_3$: C, 76.10; H, 6.01. Found: C, 76.35; H, 5.64. The er was determined by chiral HPLC (OD-H), n-hexane:*i*-PrOH = 99:1, 1 mL/min; t_1 = 10.1; t_2 = 12.2 min.

(2,4,6-Trimethylphenyl)methyl benzoate (18a).^{8f}—Clear oil (0.171 g, 82%), purified by silica gel chromatography (4–6% ethyl acetate/96–94% hexane). TLC R_f = 0.66 (20% ethyl acetate/ 80% hexane); ^1H NMR (400 MHz, CDCl_3) δ 8.03 (dd, J = 8.2, 1.0 Hz, 2H), 7.56–7.39 (m, 1H), 7.41 (t, J = 7.7 Hz, 2H), 6.92 (s, 2H), 5.42 (s, 2H), 2.42 (s, 6H), 2.30 (s, 3H); $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CDCl_3) δ 166.8, 138.5, 138.4, 132.9, 130.2, 129.7, 129.2, 129.1, 128.3, 61.6, 21.0, 19.6.

(2-Methoxyphenyl)methyl benzoate (19a).⁵⁵—Clear oil (0.080 g, 96%), purified by silica gel chromatography (1% ethyl acetate/99% pentane). TLC R_f = 0.49 (10% ethyl acetate/ 90% hexane); ^1H NMR (400 MHz, CDCl_3) δ 8.08 (d, J = 7.6 Hz, 2H), 7.52 (t, J = 7.6 Hz, 1H), 7.41 (t, J = 7.6 Hz, 3H), 7.30 (td, J = 8.0, 1.6 Hz, 1H), 6.98 (t, J = 0.4 Hz, 1H), 6.89 (d, J = 8.0 Hz, 1H), 5.42 (s, 2H), 3.83 (s, 3H); $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CDCl_3) δ 162.6, 157.6, 132.9, 130.5, 129.7, 129.53, 129.45, 128.4, 124.5, 120.5, 110.5, 62.2, 55.5.

(4-Methoxyphenyl)methyl benzoate (20a).⁵⁶—Colorless oil (0.530 g, 89%), purified by silica gel chromatography (10% ethyl acetate/ 90% hexanes). TLC R_f = 0.57 (20% ethyl acetate/ 80% hexanes); ^1H NMR (300 MHz, CDCl_3) δ 8.10–8.03 (m, 2H), 7.59–7.51 (m, 1H), 7.47–7.36 (m, 4H), 6.92 (d, J = 8.8 Hz, 2H), 5.31 (s, 2H), 3.82 (s, 3H); $^{13}\text{C}\{^1\text{H}\}$ NMR

(75 MHz, CDCl₃) δ 166.7, 159.9, 133.2, 130.5, 130.3, 129.9, 128.5, 128.4, 114.2, 66.7, 55.5.

(4-Methoxyphenyl)methyl 2,2-diphenylacetate (20b).—White solid (0.120 g, 96%), purified by silica gel chromatography (4% ethyl acetate/96% hexanes). TLC R_f = 0.50 (20% ethyl acetate/ 80% hexanes); IR (film, cm⁻¹) $\nu_{\max}$ 3054, 2986, 1734, 1613, 1516, 1265, 739, 703; ¹H NMR (400 MHz, CDCl₃) δ 7.30–7.21 (m, 12H), 6.85 (d, J = 8.7 Hz, 2H), 5.12 (s, 2H), 5.04 (s, 1H), 3.80 (s, 3H); ¹³C{¹H} NMR (100 MHz, CDCl₃) δ 172.4, 159.6, 138.6, 130.1, 128.6, 128.6, 127.8, 127.2, 113.9, 66.8, 57.1, 55.3. Anal. Calcd for C₂₂H₂₀O₃: C, 79.50; H, 6.07. Found: C, 79.34; H, 5.97.

(4-Methoxyphenyl)methyl 2-methyl-2-phenylpropanoate (20c).^{18a}—Clear oil (0.120 g, 86%), purified by silica gel chromatography (8% ethyl acetate/ 92% hexanes). TLC R_f = 0.51 (50% ethyl acetate/ 50% hexanes); ¹H NMR (400 MHz, CDCl₃) δ 7.31–7.20 (m, 5H), 7.14 (d, J = 8.6 Hz, 2H), 6.82 (d, J = 8.7 Hz, 2H), 5.03 (s, 2H), 3.77 (s, 3H), 1.57 (s, 6H); ¹³C{¹H} NMR (100 MHz, CDCl₃) 176.5, 159.4, 144.6, 129.5, 128.3, 128.3, 126.6, 125.7, 113.8, 66.3, 55.2, 46.6, 26.5.

(4-Methoxyphenyl)methyl 2-bromododecanoate (20d).—Yellow oil (0.407 g, 93%), purified by silica gel chromatography (15% ethyl acetate/ 85% hexanes). TLC R_f = 0.65 (25% ethyl acetate/ 75% hexanes); IR (film, cm⁻¹) $\nu_{\max}$ 3001, 2906, 2851, 1724, 1514, 1229; ¹H NMR (300 MHz, CDCl₃) δ 7.25 (d, J = 9.0 Hz, 2H), 6.82 (d, J = 9.0 Hz, 2H), 5.07 (s, 2H), 4.15 (t, J = 7.4 Hz, 1H), 3.73 (s, 3H), 1.98–1.91 (m, 2H), 1.30–1.10 (m, 16H), 0.82 (t, J = 6.0 Hz, 3H); ¹³C{¹H} NMR (75 MHz CDCl₃) δ 170.0, 160.8, 130.4, 127.6, 114.2, 67.6, 55.5, 46.4, 35.2, 32.1, 29.8, 29.7, 29.6, 29.1, 27.5, 22.9, 14.6. Anal Calcd for C₂₀H₃₁BrO₃: C, 60.15; H, 7.82. Found: C, 60.01; H, 7.52.

(±)-(4-Methoxyphenyl)methyl-2-hydroxy-2-phenylacetate (20e).⁵⁶—Colorless oil (0.495 g, 79%), purified by silica gel chromatography (20% ethyl acetate/ 80% hexanes). TLC R_f = 0.65 (30% ethyl acetate/ 70% hexanes); ¹H NMR (300 MHz, CDCl₃) δ 7.35–7.25 (m, 5H), 7.09 (d, J = 9.0 Hz, 2H), 6.78 (d, J = 9.0 Hz, 2H), 5.11–4.96 (m, 3H), 3.71 (s, 3H), 3.57 (d, J = 6.0 Hz, 1H); ¹³C{¹H} NMR (75 MHz CDCl₃) δ 173.8, 160.0, 138.5, 130.2, 128.8, 128.7, 127.3, 126.9, 114.2, 73.2, 67.8, 55.5.

(±)-(4-Methoxyphenyl)methyl-2-[(tert-butoxycarbonyl)amino]phenylpropionate (20f).⁵⁷—Colorless oil (0.342 g, 89%), purified by silica gel chromatography (20% ethyl acetate/ 80% hexanes). TLC R_f = 0.61 (30% ethyl acetate/ 70% hexanes); ¹H NMR (300 MHz, CDCl₃) δ 7.20–7.11 (m, 5H), 6.99–6.87 (m, 2H), 6.75 (d, J = 9.0 Hz, 2H), 5.03–4.97 (m, 3H), 4.60–4.45 (m, 1H), 3.73 (s, 3H), 3.11–2.90 (m, 2H), 1.35 (s, 9H); ¹³C{¹H} NMR (75 MHz CDCl₃) δ 173.5, 160.1, 155.5, 138.6, 132.3, 130.3, 129.5, 127.8, 127.5, 114.6, 80.2, 67.1, 55.5, 54.0, 39.1, 28.5.

(4-Methoxyphenyl)methyl but-3-enoate (20g).⁵²—Colorless oil (0.389 g, 93% yield), purified by silica gel chromatography (5% ethyl acetate/ 95% hexanes). TLC R_f = 0.50 (5% ethyl acetate/ 95% hexanes); ¹H NMR (300 MHz, CDCl₃) δ 7.23 (d, J = 9.0 Hz, 2H), 6.81 (d, J = 9.0 Hz, 2H), 5.92–5.82 (m, 1H), 5.13–5.11 (m, 1H), 5.08–5.03 (m, 1H), 5.00 (s, 2H),

3.72 (s, 3H), 3.06–3.03 (m, 2H); $^{13}\text{C}\{^1\text{H}\}$ NMR (75 MHz CDCl_3) δ 171.6, 159.9, 130.5, 130.4, 128.2, 118.8, 114.2, 66.5, 55.5, 39.4.

(4-Methoxyphenyl)methyl cinnamate (20h).⁵⁸—White solid (0.508 g, 94%), purified by silica gel chromatography (10% ethyl acetate/ 90% hexanes). TLC R_f = 0.64 (20% ethyl acetate/ 80% hexanes); mp = 61–63 °C; ^1H NMR (300 MHz, CDCl_3) δ 7.71 (d, J = 16.2 Hz, 1H), 7.55–7.48 (m, 2H), 7.41–7.33 (m, 5H), 6.92 (d, J = 8.8 Hz, 2H), 6.47 (d, J = 16.2 Hz, 1H), 5.19 (s, 2H), 3.82 (s, 3H); $^{13}\text{C}\{^1\text{H}\}$ NMR (75 MHz CDCl_3) δ 167.0, 159.8, 145.1, 134.5, 130.4, 130.3, 129.0, 128.3, 128.2, 118.1, 114.1, 66.3, 55.4.

(4-Methoxyphenyl)methyl (2Z)-3-(2-methoxyphenyl)prop-2-enoate (20i).—Clear oil (0.140 g, 84%), purified by silica gel chromatography (10% ethyl acetate/ 90% hexanes). TLC R_f = 0.42 (20% ethyl acetate/ 80% hexanes); IR (film, cm^{-1}) ν_{max} 2958, 2837, 1720, 1613, 1515, 1262, 1030, 737; ^1H NMR (400 MHz, CDCl_3) δ 7.50 (dd, J = 7.6, 1.1 Hz, 1H), 7.31–7.20 (m, 3H), 7.14 (s, 1H), 6.89–6.82 (m, 4H), 5.98 (d, J = 12.4 Hz, 1H), 5.05 (s, 2H), 3.77 (s, 6H); $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CDCl_3) δ 166.2, 159.6, 157.2, 139.4, 130.8, 130.4, 130.2, 128.1, 124.1, 120.0, 119.8, 113.9, 110.3, 65.8, 55.4, 55.3. Anal. Calcd for $\text{C}_{18}\text{H}_{18}\text{O}_4$: C, 72.47; H, 6.08. Found: C, 72.14; H, 6.11.

(4-Methoxyphenyl)methyl picolinate (20j).—Yellow oil (0.516 g, 83%), purified by silica gel chromatography (50% ethyl acetate/ 50% hexanes). TLC R_f = 0.30 (50% ethyl acetate/ 50% hexanes); IR (film, cm^{-1}) ν_{max} 3057, 3005, 2957, 2837, 1717; ^1H NMR (300 MHz, CDCl_3) δ 8.76 (d, J = 7.2 Hz, 1H), 8.12 (d, J = 8.7 Hz, 1H), 7.85–7.79 (m, 1H), 7.48–7.42 (m, 3H), 6.90 (d, J = 8.7 Hz, 2H), 5.40 (s, 2H), 3.81 (s, 3H); $^{13}\text{C}\{^1\text{H}\}$ NMR (75 MHz, CDCl_3) δ 165.2, 159.8, 150.0, 148.2, 137.0, 130.6, 127.9, 127.0, 125.3, 114.0, 67.4, 55.3. Anal. Calcd for $\text{C}_{14}\text{H}_{13}\text{NO}_3$: C, 69.12; H, 5.39; N, 5.76. Found: C, 69.52; H, 5.65; N, 5.73.

(2,4-Dimethoxyphenyl)methyl benzoate (21a).—Clear oil (0.062 g, 90%), purified by silica gel chromatography (6% ethyl acetate/ 94% hexanes). TLC R_f = 0.29 (10% ethyl acetate/ 90% hexane); IR (film, cm^{-1}) ν_{max} 3003, 2837, 1721, 1275, 712; ^1H NMR (400 MHz, CDCl_3) δ 8.05 (d, J = 7.2 Hz, 2H), 7.51 (t, J = 7.2 Hz, 1H), 7.39 (t, J = 8.0 Hz, 2H), 7.34–7.32 (m, 1H), 6.49–6.46 (m, 2H), 5.34 (s, 2H), 3.81 (s, 3H), 3.80 (s, 3H); $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CDCl_3) δ 166.7, 161.3, 159.0, 132.8, 131.2, 130.6, 129.7, 128.3, 116.9, 104.1, 98.6, 62.1, 55.5, 55.4. Anal. Calcd for $\text{C}_{16}\text{H}_{16}\text{O}_4$: C, 70.58; H, 5.92. Found: C, 70.57; H, 5.87.

(2,4-Dimethoxyphenyl)methyl 2,2-diphenylacetate (21b).—Clear oil (0.120 g, 88%), purified by silica gel chromatography (15% ethyl acetate/ 85% hexanes). TLC R_f = 0.27 (20% ethyl acetate/ 80% hexanes); IR (film, cm^{-1}) ν_{max} 1731, 1616, 1209, 1035, 701; ^1H NMR (400 MHz, CDCl_3) δ 7.30–7.23 (m, 10H), 7.15 (d, J = 8.7 Hz, 1H), 6.42–6.39 (m, 2H), 5.16 (s, 2H), 5.04 (s, 1H), 3.79 (s, 3H), 3.70 (s, 3H); $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CDCl_3) δ 172.4, 161.3, 159.0, 138.9, 131.3, 128.7, 128.5, 127.1, 116.6, 103.9, 98.5, 62.6, 57.1, 55.4, 55.3. Anal. Calcd for $\text{C}_{23}\text{H}_{22}\text{O}_4$: C, 76.22; H, 6.12. Found: C, 76.08; H, 6.10.

(2,4-Dimethoxyphenyl)methyl 2-methyl-2-phenylpropanoate (21c).—Clear oil (0.200 g, 85%), purified by silica gel chromatography (8% ethyl acetate/ 92% hexanes). TLC R_f = 0.45 (10% ethyl acetate/ 90% hexanes); IR (film, cm^{-1}) ν_{max} 3061, 2936, 1718, 1612, 1508, 1206, 1098, 1029, 731, 697; ^1H NMR (400 MHz, CDCl_3) δ 7.26–7.11 (m, 5H), 6.98 (d, J = 7.8 Hz, 1H), 6.32–6.30 (m, 2H), 5.00 (s, 2H), 3.71 (s, 3H), 3.63 (s, 3H), 1.50 (s, 6H); $^{13}\text{C}\{^1\text{H}\}$ NMR (75 MHz, CDCl_3) δ 176.6, 161.0, 158.7, 144.9, 130.5, 128.2, 126.5, 125.8, 117.1, 103.8, 98.4, 62.1, 55.4, 55.3, 46.7, 26.5. Anal. Calcd for $\text{C}_{19}\text{H}_{22}\text{O}_4$: C, 72.59; H, 7.05. Found: C, 72.36; H, 7.17.

(2,4-Dimethoxyphenyl)methyl 2-bromododecanoate (21d).—Clear oil (0.099 g, 81%), purified by silica gel chromatography (5% ethyl acetate/ 95% hexane). TLC R_f = 0.57 (20% ethyl acetate/80% hexanes); IR (film, cm^{-1}) ν_{max} 3054, 1422, 1265, 1038, 896, 739. ^1H NMR (400 MHz, CDCl_3) δ 7.24 (s, 1H), 6.48–6.46 (m, 2H), 5.18 (s, 2H), 4.22 (t, J = 7.4 Hz, 1H), 3.81 (s, 6H), 2.11–1.92 (m, 2H), 1.49–1.19 (m, 16H), 0.88 (t, J = 6.6 Hz, 3H); $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CDCl_3) δ 169.9, 161.5, 159.0, 131.4, 116.1, 104.0, 98.6, 63.1, 55.4, 55.4, 46.4, 35.0, 31.9, 29.6, 29.5, 29.3, 29.3, 28.9, 27.2, 22.7, 14.1. Anal. Calcd for $\text{C}_{21}\text{H}_{33}\text{BrO}_4$: C, 58.74; H, 7.75. Found: C, 58.57; H, 7.65.

(±)-(2,4-Dimethoxyphenyl)methyl-2-hydroxy-2-phenylacetate (21e).—White amorphous solid (0.150 g, 94%), purified by silica gel chromatography (50 % ethyl acetate/ 50% hexanes). TLC R_f = 0.37 (50% ethyl acetate/ 50% hexanes); mp = 290–292 °C; IR (film, cm^{-1}) ν_{max} 3054, 2835, 1729, 1274, 762; ^1H NMR (400 MHz, CDCl_3) δ 7.41–7.39 (m, 2H), 7.34–7.28 (m, 3H), 7.07 (d, J = 8.9 Hz, 1H), 6.39 (brs, 2H), 5.16–5.15 (m, 3H), 3.78 (s, 3H), 3.67 (s, 3H), 3.54 (brs, 1H); $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CDCl_3) δ 173.6, 161.5, 159.0, 138.5, 131.2, 128.4, 128.3, 126.6, 115.8, 103.9, 98.5, 72.9, 63.5, 55.4, 55.3. Anal. Calcd for $\text{C}_{17}\text{H}_{18}\text{O}_5$: C, 67.54; H, 6.00. Found: C, 67.72; H, 5.87.

(±)-(2,4-Dimethoxyphenyl)methyl-2-[(tert-butoxycarbonyl)amino]phenylpropionate (21f).—Clear oil (0.101 g, 80%), purified by silica gel chromatography (15 % ethyl acetate/ 85% hexanes). TLC R_f = 0.37 (40% ethyl acetate/ 60% hexanes); IR (film, cm^{-1}) ν_{max} 3054, 2985, 1709, 1421, 1261, 739; ^1H NMR (300 MHz, CD_3OD) δ 7.26–7.16 (m, 6H), 6.56 (d, J = 2.2 Hz, 1H), 6.50 (dd, J = 8.3, 2.3 Hz, 1H), 5.11 (q, J = 12.1 Hz, 2H), 4.38 (q, J = 6.0 Hz, 1H), 3.84 (s, 3H), 3.82 (s, 3H), 3.10 (dd, J = 13.9, 5.6 Hz, 1H), 2.91 (dd, J = 13.7, 8.6 Hz, 1H), 1.39 (s, 9H); $^{13}\text{C}\{^1\text{H}\}$ NMR (75 MHz, CD_3OD) δ 172.3, 161.7, 159.0, 156.4, 136.9, 131.2, 128.9, 128.0, 126.4, 116.0, 104.1, 97.9, 79.2, 62.1, 55.2, 54.6, 54.4, 37.3, 27.3. Anal. Calcd for $\text{C}_{23}\text{H}_{29}\text{NO}_6$: C, 66.49; H, 7.04; N, 3.37. Found: C, 66.20; H, 6.93; N, 3.76.

(2,4-Dimethoxyphenyl)methyl but-3-enoate (21g).—Clear oil (0.163 g, 73%), purified by silica gel chromatography (15% ethyl acetate/85% hexanes). TLC R_f = 0.40 (15% ethyl acetate/85% hexanes); IR (film, cm^{-1}) ν_{max} 2963, 2838, 2616, 1464, 1371, 1209, 739; ^1H NMR (300 MHz, CDCl_3) δ 7.23 (d, J = 8.9 Hz, 1H), 6.46 (hextet, J = 2.4 Hz, 2H), 6.00–5.87 (m, 1H), 5.18–5.11 (m, 4H), 3.81 (s, 3H), 3.80 (s, 3H), 3.11 (dt, J = 6.9, 1.4 Hz, 2H); $^{13}\text{C}\{^1\text{H}\}$ NMR (75 MHz, CDCl_3) δ 171.6, 161.3, 159.0, 131.4, 130.5, 118.4,

116.7, 104.1, 98.6, 61.9, 55.5, 55.4, 39.2. Anal. Calcd for $C_{13}H_{16}O_4$: C, 66.09; H, 6.83. Found: C, 65.91; H, 6.47.

(2,4-Dimethoxyphenyl)methyl cinnamate (21h).—Viscous oil (0.139 g, 88%), purified by silica gel chromatography (4% ethyl acetate/98% hexanes). TLC R_f = 0.38 (20% ethyl acetate/80% hexanes); IR (film, cm^{-1}) ν_{max} 1708, 1510, 1161, 737. 1H NMR (400 MHz, $CDCl_3$) δ 7.70 (d, J = 16.1 Hz, 1H), 7.51–7.49 (m, 2H), 7.37–7.35 (m, 3H), 7.30 (d, J = 8.9 Hz, 1H), 6.49–6.51–6.43 (m, 3H), 5.23 (s, 2H), 3.83 (s, 3H), 3.81 (s, 3H); $^{13}C\{^1H\}$ NMR (100 MHz, $CDCl_3$) δ 167.1, 161.3, 159.1, 144.7, 134.6, 131.5, 130.2, 128.9, 128.1, 118.4, 116.8, 104.1, 98.6, 61.8, 55.5, 55.4. Anal. Calcd for $C_{18}H_{18}O_4$: C, 72.47; H, 6.08. Found: C, 72.20; H, 5.71.

(2,4-Dimethoxyphenyl)methyl (2Z)-3-(2-methoxyphenyl)prop-2-enoate (21i).—Clear oil (0.170 g, 92%), purified by silica gel chromatography (20% ethyl acetate/80% hexanes). TLC R_f = 0.29 (20% ethyl acetate/80% hexanes); IR (film, cm^{-1}) ν_{max} 3054, 2986, 2839, 1717, 1616, 1488, 1209, 1159, 739, 705; 1H NMR (400 MHz, $CDCl_3$) δ 7.57 (d, J = 7.5 Hz, 1H), 7.31–7.26 (m, 1H), 7.18–7.15 (m, 2H), 6.90–6.85 (m, 2H), 6.44–6.41 (m, 2H), 6.01 (d, J = 12.5 Hz, 1H), 5.14 (s, 2H), 3.81 (s, 6H), 3.78 (s, 3H); $^{13}C\{^1H\}$ NMR (100 MHz, $CDCl_3$) δ 166.3, 161.2, 158.9, 157.2, 138.9, 131.3, 130.8, 130.3, 124.1, 120.00, 119.94, 116.8, 110.3, 104.0, 98.5, 61.3, 55.41, 55.40. Anal. Calcd for $C_{19}H_{20}O_5$: C, 69.50; H, 6.14. Found: C, 69.87; H, 6.47.

(2,4-Dimethoxyphenyl)methyl picolinate (21j).—Yellow oil (0.150 g, 83%), purified by silica gel chromatography (50% ethyl acetate/ 50% hexanes). TLC R_f = 0.25 (50% ethyl acetate/ 50% hexanes); IR (film, cm^{-1}) ν_{max} 2964, 1721, 1616, 1377, 927; 1H NMR (400 MHz, $CDCl_3$) δ 8.76 (dd, J = 4.7 Hz, 0.7 Hz, 1H), 8.10 (d, J = 7.8 Hz, 1H), 7.79 (td, J = 7.8, 1.7 Hz, 1H), 7.41 (ddd, J = 7.6, 4.8, 1.1 Hz 1H), 7.34 (d, J = 9.0 Hz, 1H), 6.48–6.46 (m, 2H), 5.43 (s, 2H), 3.82 (s, 3H), 3.80 (s, 3H); $^{13}C\{^1H\}$ NMR (75 MHz, $CDCl_3$) δ 165.2, 161.4, 159.1, 150.0, 148.5, 136.8, 131.6, 126.7, 125.2, 116.5, 104.1, 98.6, 62.9, 55.5, 55.4. Anal. Calcd for $C_{15}H_{15}NO_4$: C, 65.92; H, 5.53; N, 5.13. Found: C, 66.17; H, 5.54; N, 5.44.

3,4-Dimethoxybenzyl benzoate (22a).⁵⁹—Clear oil (0.17 g, 96%), purified by silica gel chromatography (5% ethyl acetate/ 95% hexanes). TLC R_f = 0.40 (20% ethyl acetate/ 80% hexane); 1H NMR (400 MHz, $CDCl_3$) δ 8.06 (d, J = 7.2 Hz, 2H), 7.52 (t, J = 7.6 Hz, 1H), 7.40 (t, J = 7.6 Hz, 2H), 7.03–6.97 (m, 2H), 6.85 (d, J = 8.0 Hz, 1H), 5.29 (s, 2H), 3.88 (s, 3H), 3.86 (s, 3H); $^{13}C\{^1H\}$ NMR (100 MHz, $CDCl_3$) δ 166.5, 149.1, 149.0, 133.0, 130.2, 129.7, 128.6, 128.4, 121.3, 111.9, 111.1, 66.8, 55.9.

(Furan-2-yl)methyl benzoate (24a).⁶⁰—Yellow oil (0.127 g, 96%), purified by silica gel chromatography (1% ethyl acetate/ 99% hexane). TLC R_f = 0.46 (10% ethyl acetate/ 90% hexane); 1H NMR (400 MHz, $CDCl_3$) δ 8.06 (d, J = 8.2 Hz, 2H), 7.57–7.53 (m, 1H), 7.45–7.41 (m, 3H), 6.49 (d, J = 3.2 Hz, 1H), 6.39–6.38 (m, 1H), 5.31 (s, 2H); $^{13}C\{^1H\}$ NMR (100 MHz, $CDCl_3$) δ 166.3, 149.6, 143.3, 133.1, 129.9, 129.8, 128.4, 110.8, 110.6, 58.5.

1-Phenethyl benzoate (25a).⁵³—Clear oil (0.140 g, 95%), purified by silica gel chromatography (2% ethyl acetate/98% hexane). TLC R_f = 0.23 (10% ethyl acetate/ 90% hexane); ^1H NMR (300 MHz, CDCl_3) δ 8.10–8.06 (m, 2H), 7.55–7.22 (m, 8H), 6.14 (q, J = 6.6 Hz, 1H), 1.66 (d, J = 6.6 Hz, 3H); $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CDCl_3) δ 165.8, 141.9, 133.0, 130.6, 129.7, 128.6, 128.4, 127.9, 126.1, 73.0, 22.5.

Diphenylmethyl benzoate (26a).—White solid (2.58 g, 99%), purified by silica gel chromatography (1% triethylamine/ 10% ethyl acetate/ 89% hexanes). mp = 88–89 °C; TLC R_f = 0.42 (10% ethyl acetate/ 90% hexanes); IR (KBr, cm^{-1}) ν_{max} 3090, 3031, 2948, 1712, 1267, 1189; ^1H NMR (300 MHz, CDCl_3) δ 8.17–8.13 (m, 2H), 7.60–7.55 (m, 1H), 7.50–7.26 (m, 12H), 7.13 (s, 1H); $^{13}\text{C}\{^1\text{H}\}$ NMR (75 MHz, CDCl_3) δ 165.8, 140.5, 133.4, 130.4, 130.0, 128.8, 128.7, 128.2, 127.4, 77.7. Anal. Calcd for $\text{C}_{20}\text{H}_{16}\text{O}_2$: C, 83.31; H, 5.59. Found: C, 83.37; H, 5.74.

Diphenylmethyl 2,2-diphenylacetate (26b).⁶¹—White solid (0.120 g, 84%), purified by silica gel chromatography (8% ethyl acetate/ 92% hexanes). TLC R_f = 0.52 (20% ethyl acetate/ 80% hexanes); ^1H NMR (400 MHz, CDCl_3) δ 7.28–7.19 (m, 20H), 6.91 (s, 1H), 5.15 (s, 1H); $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CDCl_3) δ 171.4, 140.0, 138.5, 128.8, 128.6, 128.5, 127.9, 127.3, 127.2, 57.3.

Diphenylmethyl 2-methyl-2-phenylpropanoate (26c).—Clear oil (0.120 g, 75%), purified by silica gel chromatography (6% ethyl acetate/94% hexanes). TLC R_f = 0.50 (20% ethyl acetate/ 80% hexanes); IR (film, cm^{-1}) ν_{max} 3088, 3031, 1730, 1600, 1495, 1142, 1100; ^1H NMR (400 MHz, CDCl_3) δ 7.29–7.23 (m, 11H), 7.14–7.12 (m, 4H), 6.80 (s, 1H), 1.62 (s, 6H). $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CDCl_3) δ 175.5, 144.3, 142.3, 140.3, 129.0, 128.42, 128.37, 127.7, 127.5, 127.3, 126.9, 126.7, 125.9, 80.1, 46.7, 26.3. Anal. Calcd for $\text{C}_{23}\text{H}_{22}\text{O}_2$: C, 83.60; H, 6.71. Found: C, 83.93; H, 6.99.

Diphenylmethyl 2-bromododecanoate (26d).—Clear colorless oil (0.40 g, 83%), purified by silica gel chromatography (1% triethylamine/5% ethyl acetate/ 94% hexanes). TLC R_f = 0.78 (10% ethyl acetate/90% hexanes); IR (film, cm^{-1}) ν_{max} 3064, 3032, 2924, 2854, 1741, 1495, 1454, 1257, 1144, 1080; ^1H NMR (300 MHz, CDCl_3) δ 7.38–7.26 (m, 10H), 6.89 (s, 1H), 4.31 (t, J = 7.5 Hz, 1H), 2.07–2.00 (m, 2H), 1.23 (br s, 16H), 0.88 (t, J = 6.3 Hz, 3H); $^{13}\text{C}\{^1\text{H}\}$ NMR (75 MHz CDCl_3) δ 169.0, 139.8, 139.7, 128.80, 128.79, 128.4, 127.4, 127.3, 78.5, 46.4, 35.2, 32.2, 29.8, 29.7, 29.6, 29.1, 27.4, 23.0, 14.4. Anal. Calcd for $\text{C}_{25}\text{H}_{33}\text{O}_2\text{Br}$: C, 67.41; H, 7.47. Found: C, 67.60; H, 7.66.

(±)-Diphenylmethyl-2-hydroxy-2-phenylacetate (26e).—White solid (0.58 g, 92%), purified by silica gel chromatography (1% triethylamine/ 30% ethyl acetate/ 69% hexanes). mp = 113–114 °C; TLC R_f = 0.65 (30% ethyl acetate/ 70% hexanes); IR (KBr, cm^{-1}) ν_{max} 3221, 2815, 2800, 1699, 1240; ^1H NMR (300 MHz, CDCl_3) δ 7.42–7.18 (m, 13H), 6.91–6.87 (m, 3H), 5.28 (d, J = 5.4 Hz, 1H), 3.45 (d, J = 5.7 Hz, 1H); $^{13}\text{C}\{^1\text{H}\}$ NMR (75 MHz, CDCl_3) δ 173.0, 139.5, 139.4, 138.3, 128.9, 128.83, 128.79, 128.6, 128.5, 128.1, 127.6, 127.0, 126.5, 79.0, 73.3. Anal. Calcd for $\text{C}_{21}\text{H}_{18}\text{O}_3$: C, 79.22; H, 5.70. Found: C, 79.53; H, 5.40.

(±)-Diphenylmethyl-2-[(tert-butoxycarbonyl)amino]phenylpropionate (26f).—

Clear oil (0.072 g, 54%), purified by silica gel chromatography (20 % ethyl acetate/ 80% hexanes). TLC R_f = 0.27 (20% ethyl acetate/80% hexanes); IR (film, cm^{-1}) ν_{max} 3054, 2985, 1709, 1421, 1260. ^1H NMR (300 MHz, CD_3OD) δ 7.37–7.13 (m, 15H), 6.85 (s, 1H), 4.50 (q, J = 6.1 Hz, 1H), 3.13 (dd, J = 13.8, 6.0 Hz, 1H), 2.95 (dd, J = 13.8, 8.7 Hz, 1H), 1.40 (s, 9H); $^{13}\text{C}\{^1\text{H}\}$ NMR (75 MHz, CD_3OD) δ 171.4, 156.4, 140.0, 136.8, 128.9, 128.1, 127.6, 126.9, 126.7, 126.4, 79.3, 77.8, 55.5, 37.1, 27.4. Anal. Calcd for $\text{C}_{27}\text{H}_{29}\text{NO}_4$: C, 75.15; H, 6.77; N, 3.25. Found: C, 75.16; H, 6.73; N, 3.61.

Diphenylmethyl but-3-enoate (26g).⁶²—Colorless oil (0.35 g, 76%), purified by silica gel chromatography (1% triethylamine/ 5% ethyl acetate/ 94% hexanes). TLC R_f = 0.44 (10% ethyl acetate/ 90% hexanes); IR (film, cm^{-1}) ν_{max} 3064, 3031, 2983, 2938, 1740, 1642, 1543, 1030. ^1H NMR (300 MHz, CDCl_3) δ 7.35–7.26 (m, 10H), 6.90 (s, 1H), 6.02–5.90 (m, 1H), 5.22–5.15 (m, 2H), 3.23–3.20 (m, 2H); $^{13}\text{C}\{^1\text{H}\}$ NMR (75 MHz, CDCl_3) δ 170.7, 140.5, 130.4, 128.9, 128.3, 127.4, 119.1, 77.3, 39.6.

(E)-Diphenylmethyl cinnamate (26h).⁶³—White solid (0.401 g, 93%), purified by silica gel chromatography (1% triethylamine/ 10% ethyl acetate/ 89% hexanes). mp = 74–77 °C; TLC R_f = 0.57 (10% ethyl acetate/ 90% hexanes); ^1H NMR (300 MHz, CDCl_3) δ 7.78 (d, J = 16.2 Hz, 1H), 7.57–7.54 (m, 2H), 7.44–7.26 (m, 13H), 7.05 (s, 1H), 6.58 (d, J = 15.9, 0.6 Hz, 1H); $^{13}\text{C}\{^1\text{H}\}$ NMR (75 MHz, CDCl_3) δ 166.2, 145.7, 140.5, 134.5, 130.7, 129.1, 128.8, 128.4, 128.2, 127.4, 118.2, 77.2.

Diphenylmethyl (2Z)-3-(2-methoxyphenyl)prop-2-enoate (26i).—White solid (0.099 g, 52%), purified by silica gel chromatography (4% ethyl acetate/96% hexanes); mp = 72–74 °C; TLC R_f = 0.71 (10% ethyl acetate/80% hexanes); IR (film, cm^{-1}) ν_{max} 3031, 2836, 1725, 1627, 1110; ^1H NMR (400 MHz, CDCl_3) δ 7.39 (d, J = 2.0 Hz, 1H), 7.21–7.13 (m, 12H), 6.85–6.75 (m, 3H), 6.06–6.02 (m, 1H), 3.68 (s, 3H); $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CDCl_3) δ 165.2, 157.1, 140.3, 140.0, 130.8, 130.4, 128.4, 127.8, 127.3, 124.2, 120.1, 120.0, 110.3, 76.7, 55.4. Anal. Calcd for $\text{C}_{23}\text{H}_{20}\text{O}_3$: C, 80.21; H, 5.85. Found: C, 80.38; H, 5.54.

Diphenylmethyl picolinate (26j).—White solid (0.210 g, 71%), purified by silica gel chromatography (10% acetone/ 90% hexanes). mp = 101–103 °C; TLC R_f = 0.23 (20% acetone/ 80% hexanes); IR (film, cm^{-1}) ν_{max} 3052, 1744, 1130. ^1H NMR (300 MHz, CDCl_3) δ 8.80 (dq, J = 4.8, 0.9 Hz, 1H), 8.19 (dt, J = 7.8, 1.1 Hz, 1H), 7.85 (dt, J = 7.8, 1.8 Hz, 1H), 7.50–7.44 (m, 5H), 7.39–7.27 (m, 6H), 7.22 (s, 1H); $^{13}\text{C}\{^1\text{H}\}$ NMR (75 MHz, CDCl_3) δ 164.3, 150.3, 148.3, 140.0, 137.1, 128.8, 128.2, 127.5, 127.1, 125.4, 78.2; Anal. Calcd for $\text{C}_{19}\text{H}_{15}\text{NO}_2$: C, 78.87; H, 5.23; N, 4.84. Found: C, 78.89; H, 5.36; N, 5.16.

(S)-Diphenylmethyl 2-(2-methoxynaphthalen-6-yl)propanoate ((+)-26k).—White solid (1.38 g, 89%), purified by silica gel chromatography (20% ethyl acetate/ 80% hexanes). mp = 132–134 °C; $[\alpha]_{\text{D}}^{20}$ +55.7 (c 1.1, CHCl_3); TLC R_f = 0.42 (20% ethyl acetate/ 80% hexanes); IR (KBr, cm^{-1}) ν_{max} 3059, 3000, 2942, 2359, 1723, 1604, 1162, 705; ^1H NMR (300 MHz, CDCl_3) δ 7.70–7.60 (m, 3H), 7.37 (dd, J = 8.4, 1.8 Hz, 1H), 7.29–7.04 (m, 12H), 6.84 (s, 1H), 4.01–3.93 (m, 4H), 1.59 (d, J = 6.9 Hz, 3H); $^{13}\text{C}\{^1\text{H}\}$ NMR (75 MHz, CDCl_3) δ 173.7, 157.8, 140.4, 140.3, 135.7, 133.9, 129.5, 129.1, 128.7,

128.5, 128.1, 127.9, 127.4, 127.3 126.9, 126.7, 126.3, 119.2, 105.8, 77.4, 55.5, 45.9, 18.6. Anal. Calcd for $C_{27}H_{24}O_3$: C, 81.79; H, 6.10. Found: C, 81.54; H, 6.11. The enantiomeric ratio was determined by HPLC using a chiral column (OD-H), n-hexane:i-PrOH = 99:1, 1 mL/min; compared to a racemic sample which showed two peaks t_R = 9.7 and 11.4 min.

(2-Methoxyphenyl)(phenyl)methyl [1,1'-biphenyl]-4-carboxylate (38).—White solid (73 mg, 73% yield), purified by silica gel column chromatography (10% ethyl acetate, 90% hexanes). mp = 136–137 °C; TLC R_f = 0.35 (10% ethyl acetate/ 90% hexanes); IR (film, cm^{-1}) ν_{max} 3064, 2938, 1715, 1604, 1491, 1278, 1100, 857; 1H NMR (400 MHz, $CDCl_3$) δ 8.12 (d, J = 8.2 Hz, 2H), 7.58 (d, J = 8.2 Hz, 2H), 7.54–7.52 (m, 2H), 7.44–7.15 (m, 10H), 6.91–6.80 (m, 2H), 3.75 (s, 3H); $^{13}C\{^1H\}$ NMR (100 MHz $CDCl_3$) δ 165.4, 156.5, 145.8, 140.3, 140.1, 130.3, 129.2, 129.1, 129.0, 128.9, 128.3, 128.2, 127.7, 127.3, 127.2, 127.1, 127.0, 120.7, 110.9, 72.2, 55.6. Anal Calcd for $C_{27}H_{22}O_3$: C, 82.21; H, 5.62. Found: C, 82.28; H, 5.67. The enantiomeric ratio was determined by HPLC using a chiral column (AD), n-hexane:i-PrOH = 95:5, 1 mL/min; compared to a racemic sample which showed two peaks t_R = 14.8 and 16.9 min.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This material is based upon work supported by the National Science Foundation under Grant No. CHE-1659775 (REU summer fellowship to S.E.C.). Financial support was also provided in part by the National Institute of General Medical Sciences (R15-GM116054).

References

- (1). Heller ST; Sarpong R On the reactivity of imidazole carbamates and ureas and their use as esterification and amidation reagents. *Tetrahedron* 2011, 67, 8851.
- (2). (a) Fischer E; Speier A Preparation of ethereal salts. *Ber.* 1895, 28, 3252. (b) Hassner A; Krepski LR; Alexanian V Aminopyridines as acylation catalysts for tertiary alcohols. *Tetrahedron* 1978, 34, 2069. (c) Kim S; Kim YC; Lee JI A new convenient method for the esterification of carboxylic acids. *Tetrahedron Lett.* 1983, 24, 3365. (d) Mitsunobu O The use of diethyl azodicarboxylate and triphenylphosphine in synthesis and transformation of natural products. *Synthesis* 1981, 1. (e) Swamy KCK; Kumar NNB; Balaraman E; Kumar KVPP Mitsunobu and Related Reactions: Advances and Applications. *Chem. Rev* 2009, 109, 2551. [PubMed: 19382806] (f) Diago-Meseguer J; Palomo-Coll AL; Fernandez-Lizarbe JR; Zugaza-Bilbao A A new reagent for activating carboxyl groups: preparation and reactions of N,N-bis[2-oxo-3-oxazolidinyl]phosphorodiamidic chloride. *Synthesis* 1980, 547. (g) Brechbühler H; Büchi H; Hatz E; Schreiber J; Eschenmoser A Reactions between carboxylic acids and N,N-dimethylformamide acetals—an esterification method. *Helv. Chim. Acta* 1965, 48, 1746. (h) Inanaga J; Hirata K; Saeki H; Katsuki T; Yamaguchi M A rapid esterification by mixed anhydride and its application to large-ring lactonization. *Bull. Chem. Soc. Jpn* 1979, 52, 1989. (i) Zacharie B; Connolly TP; Penney CL A Simple One-Step Conversion of Carboxylic Acids to Esters Using EEDQ. *J. Org. Chem* 1995, 60, 7072.
- (3). (a) Pfeffer PE; Silbert LS Esterification by alkylation of carboxylate salts. Influence of steric factors and other parameters on reaction rates. *J. Org. Chem* 1976, 41, 1373. (b) Jiang X; Tiwari A; Thompson M; Chen Z; Cleary TP; Lee TBK A Practical Method for N-Methylation of Indoles Using Dimethyl Carbonate. *Org. Process Res. Dev* 2001, 5, 604. (c) Chakraborti AK; Nandi AB; Grover V Chemoselective protection of carboxylic acid as methyl ester: a practical alternative to

diazomethane protocol. *J. Org. Chem* 1999, 64, 8014.(d)Raber DJ; Gariano P Jr.; Brod AO; Gariano A; Guida WC; Guida AR; Herbst MD Esterification of carboxylic acids with trialkyloxonium salts. *J. Org. Chem* 1979, 44, 1149.(e)Tummatorn J; Albinia PA; Dudley GB Synthesis of benzyl esters using 2-benzyloxy-1-methylpyridinium triflate. *J. Org. Chem* 2007, 72, 8962. [PubMed: 17935354]

- (4). (a) Howard KT; Chisholm JD Preparation and Applications of 4-Methoxybenzyl Esters in Organic Synthesis. *Org. Prep. Proced. Int* 2016, 48, 1. [PubMed: 27546912] (b)Thornton MT; Henderson LC Recent Advances in the Synthesis of Diphenylmethyl Ethers. *Org. Prep. Proced. Int* 2013, 45, 395.
- (5). Lamoureux G; Agüero C A comparison of several modern alkylating agents. *ARKIVOC* 2009, 251.
- (6). (a) Offord RE; Storey HT; Rees AR; Hayward CF; Johnson WH; Pheasey MH; Wightman DA Diazoalkanes in peptide semisynthesis. *Biochem. J* 1976, 159, 480. [PubMed: 1008812] (b)Waddell ST; Santorelli GM Mild preparation of cephalosporin allyl and p-methoxybenzyl esters using diazoalkanes. *Tetrahedron Lett.* 1996, 37, 1971.(c)Kolovos M; Froussios C O-Diphenylmethylation of alcohols and carboxylic acids using diphenylmethyl diphenyl phosphate as alkylating agent. *Tetrahedron Lett.* 1984, 25, 3909.(d)Mix KA; Raines RT Optimized Diazo Scaffold for Protein Esterification. *Org. Lett* 2015, 17, 2358. [PubMed: 25938936]
- (7). Kreevoy MM; Thomas SJ Mechanism of the reaction of diazomethane with weak acids. *J. Org. Chem* 1977, 42, 3979.
- (8). (a) Fulton JR; Aggarwal VK; de Vicente J The use of tosylhydrazone salts as a safe alternative for handling diazo compounds and their applications in organic synthesis. *Eur. J. Org. Chem* 2005, 1479.(b)Furrow ME; Myers AG A General Procedure for the Esterification of Carboxylic Acids with Diazoalkanes Generated in Situ by the Oxidation of N-tert-Butyldimethylsilylhydrazones with (Difluoriodo)benzene. *J. Am. Chem. Soc* 2004, 126, 12222. [PubMed: 15453728] (c)Morandi B; Carreira EM Iron-Catalyzed Cyclopropanation in 6 M KOH with in Situ Generation of Diazomethane. *Science* 2012, 335, 1471. [PubMed: 22442479] (d)Perusquia-Hernandez C; Lara-Issasi GR; Frontana-Urbe BA; Cuevas-Yanez E Synthesis and esterification reactions of aryl diazomethanes derived from hydrazone oxidations catalyzed by TEMPO. *Tetrahedron Lett.* 2013, 54, 3302.(e)Rulliere P; Benoit G; Allouche EMD; Charette AB Safe and Facile Access to Nonstabilized Diazoalkanes Using Continuous Flow Technology. *Angew Chem Int Ed Engl* 2018, 57, 5777. [PubMed: 29575444] (f)Squiteri RA; Shearn-Nance GP; Hein JE; Shaw JT Synthesis of Esters by in Situ Formation and Trapping of Diazoalkanes. *J. Org. Chem* 2016, 81, 5278. [PubMed: 27191835]
- (9). Heller ST; Sarpöng R Chemoselective esterification and amidation of carboxylic acids with imidazole carbamates and ureas. *Org. Lett* 2010, 12, 4572. [PubMed: 20857922]
- (10). Hao W; Fujii T; Dong T; Wakai Y; Yoshimura T Application of alkoxy- λ 6-sulfanenitriles as strong alkylating reagents. *Heteroat. Chem* 2004, 15, 193.
- (11). (a) Widmer U A convenient preparation of tert-butyl esters. *Synthesis* 1983, 135.(b)Mohacsi E A convenient preparation of methyl esters from carboxylic acids. *Synth. Commun* 1982, 12, 453. (c)Mohacsi E; Leimgruber W; Baruth H Synthesis and pharmacology of metabolically stable tert-butyl ethers of morphine and levorphanol. *J. Med. Chem* 1982, 25, 1264. [PubMed: 6292420]
- (12). Yamada K; Yoshida S; Fujita H; Kitamura M; Kunishima M O-Benzoylation of Carboxylic Acids Using 2,4,6-Tris(benzyloxy)-1,3,5-triazine (TriBOT) under Acidic or Thermal Conditions. *Eur. J. Org. Chem* 2015, 2015, 7997.
- (13). (a) Liu Y Isoureas: versatile alkylation reagents in organic chemistry. *Synlett* 2009, 1353. (b)Chighine A; Crosignani S; Arnal M-C; Bradley M; Linclau B Microwave-Assisted Ester Formation Using O-Alkylisoureas: A Convenient Method for the Synthesis of Esters with Inversion of Configuration. *J. Org. Chem* 2009, 74, 4753. [PubMed: 19462991] (c)Crosignani S; White PD; Linclau B Microwave-Accelerated O-Alkylation of Carboxylic Acids with O-Alkylisoureas. *Org. Lett* 2002, 4, 2961. [PubMed: 12182599] (d)Mathias LJ Esterification and alkylation reactions employing isoureas. *Synthesis* 1979, 561.
- (14). (a) Iversen T; Bundle DR Benzyl trichloroacetimidate, a versatile reagent for acid-catalyzed benzylation of hydroxy-groups. *J. Chem. Soc., Chem. Commun* 1981, 1240.(b)Wessel H-P; Iversen T; Bundle DR Acid-catalysed benzylation and allylation by alkyl trichloroacetimidates. *J.*

Chem. Soc. Pekin 1985, 2247.(c)Armstrong A; Brackenridge I; Jackson RFW; Kirk JM A new method for the preparation of tertiary butyl ethers and esters. *Tetrahedron Lett.* 1988, 29, 2483.
(d)Kokotos G; Chiou A Convenient synthesis of benzyl and allyl esters using benzyl and allyl 2,2,2-trichloroacetimidate. *Synthesis* 1997, 168.

- (15). (a) Schmidt RR; Michel J Simple syntheses of α - and β -O-glycosyl imidates; preparation of glycosides and disaccharides. *Angew. Chem* 1980, 92, 763.(b)Schmidt RR; Michel J O-(α -DGlucopyranosyl)trichloroacetimidate as a glucosyl donor. *J. Carbohydr. Chem* 1985, 4, 141.
- (16). (a)Shoji M; Uno T; Hayashi Y Stereoselective Total Synthesis of ent-EI-1941–2 and Epi-ent-EI-1941–2. *Org. Lett* 2004, 6, 4535. [PubMed: 15548069] (b)Shoji M; Uno T; Takeya H; Onose R; Shiina I; Osada H; Hayashi Y Enantio- and Diastereoselective Total Synthesis of EI-1941–1, –2, and –3, Inhibitors of Interleukin-1 β Converting Enzyme, and Biological Properties of Their Derivatives. *J. Org. Chem* 2005, 70, 9905. [PubMed: 16292821]
- (17). (a) Thierry J; Yue C; Potier P 2-Phenylisopropyl and t-butyl trichloroacetimidates: Useful reagents for ester preparation of N-protected amino acids under neutral conditions. *Tetrahedron Lett.* 1998, 39, 1557.(b)Respondek T; Cueny E; Kodanko JJ Cumyl Ester as the C-Terminal Protecting Group in the Enantioselective Alkylation of Glycine Benzophenone Imine. *Org. Lett* 2012, 14, 150. [PubMed: 22149572]
- (18). (a) Shah JP; Russo CM; Howard KT; Chisholm JD Spontaneous formation of PMB esters using 4-methoxybenzyl-2,2,2-trichloroacetimidate. *Tetrahedron Lett.* 2014, 55, 1740.(b)Adhikari AA; Shah JP; Howard KT; Russo CM; Wallach DR; Linaburg MR; Chisholm JD Convenient Formation of Diphenylmethyl Esters Using Diphenylmethyl Trichloroacetimidate. *Synlett* 2014, 283.
- (19). Ali, Ibrahim AI; Ashry El Sayed HE; Schmidt Richard R. Protection of Hydroxy Groups with Diphenylmethyl and 9-Fluorenyl Trichloroacetimidates – Effect on Anomeric Stereocontrol. *Eur. J. Org. Chem* 2003, 2003, 4121.
- (20). Cran JW; Vidhani DV; Krafft ME Copper-Catalyzed N-tert-Butylation of Aromatic Amines under Mild Conditions Using tert-Butyl 2,2,2-Trichloroacetimidate. *Synlett* 2014, 25, 1550.
- (21). Adhikari AA; Suzuki T; Gilbert RT; Linaburg MR; Chisholm JD Rearrangement of Benzylic Trichloroacetimidates to Benzylic Trichloroacetamides. *J. Org. Chem* 2017, 82, 3982. [PubMed: 28323428]
- (22). (a) Wuts PGM Greene's Protective Groups in Organic Synthesis; 5th ed; John Wiley & Sons: Hoboken, NJ, 2014.(b)Kocienski PJ Protecting Groups; 3rd ed; Thieme: Stuttgart, 2005.
- (23). Kononov LO; Malysheva NN; Ito Y; Ogawa T Approaches to intramolecular sialylation. 3. Synthesis of 2,4-dimethoxybenzyl ester of per-O-acetylated N-acetylneuraminic acid thioglycoside and its attempted oxidation with DDQ in the presence of nucleophiles. *Russ. Chem. Bull* 2004, 53, 254.
- (24). (a) Ichikawa A; Ono H; Furuta K; Shiotsuki T; Shinoda T Enantioselective separation of racemic juvenile hormone III by normal-phase high-performance liquid chromatography and preparation of [(2)H(3)]juvenile hormone III as an internal standard for liquid chromatography-mass spectrometry quantification. *J Chromatogr A* 2007, 1161, 252. [PubMed: 17582423] (b)Niwa H; Sakata T; Yamada K A synthesis of senecionine, a representative of hepatotoxic, macrocyclic pyrrolizidine alkaloids of retronecine type. *Bull. Chem. Soc. Jpn* 1994, 67, 1990.(c)Paterson I; de Savi C; Tudge M Synthesis of the Macrocyclic Core of Laulimalide. *Org. Lett* 2001, 3, 213. [PubMed: 11430037]
- (25). Howard KT; Duffy BC; Linaburg MR; Chisholm JD Formation of DPM ethers using O-diphenylmethyl trichloroacetimidate under thermal conditions. *Org. Biomol. Chem* 2016, 14, 1623. [PubMed: 26691695]
- (26). (a) Zhao C; Toste FD; Raymond KN; Bergman RG Nucleophilic Substitution Catalyzed by a Supramolecular Cavity Proceeds with Retention of Absolute Stereochemistry. *J. Am. Chem. Soc* 2014, 136, 14409. [PubMed: 25265509] (b)Wallach DR; Stege PC; Shah JP; Chisholm JD Bronsted acid catalyzed monoalkylation of anilines with trichloroacetimidates. *J. Org. Chem* 2015, 80, 1993. [PubMed: 25568933]
- (27). Shieh W-C; Cantrell WR Jr.; Carlson JA Asymmetric reduction of ortho-substituted benzophenones with β -chlorodiisopinocampheylborane in synthesis of enantiomerically enriched benzhydrols. *Tetrahedron Lett.* 1995, 36, 3797.

- (28). Raber DJ; Harris JM; Schleyer P. v. R. In *Ions and Ion Pairs in Organic Reactions*; Szwarc M, Ed. 1974; Vol. 2, p 247.
- (29). Zhang J; Schmidt RR O-Benzyl trichloroacetimidates having electron-withdrawing substituents in acid-catalyzed diarylmethane synthesis. *Synlett* 2006, 1729.
- (30). Skrobo B; Schlörer NE; Neudörfl J-M; Deska J Kirmse-Doyle- and Stevens-Type Rearrangements of Glutarate-Derived Oxonium Ylides. *Chem. - Eur. J* 2018, 24, 3209. [PubMed: 29231265]
- (31). (a) Beckwith ALJ; Bowry VW; Ingold KU Kinetics of nitroxide radical trapping. 1. Solvent effects. *J. Am. Chem. Soc* 1992, 114, 4983.(b)Bowry VW; Ingold KU Kinetics of nitroxide radical trapping. 2. Structural effects. *J. Am. Chem. Soc* 1992, 114, 4992.
- (32). Mou X-Q; Chen X-Y; Chen G; He G Radical-mediated intramolecular β -C(sp³)-H amidation of alkylimidates: facile synthesis of 1,2-amino alcohols. *Chem. Commun* 2018, 54, 515.
- (33). Bachi MD; Korshin EE; Hoos R; Szpilman AM; Ploypradith P; Xie S; Shapiro TA; Posner GH A Short Synthesis and Biological Evaluation of Potent and Nontoxic Antimalarial Bridged Bicyclic β -Sulfonyl-Endoperoxides. *J. Med. Chem* 2003, 46, 2516. [PubMed: 12773055]
- (34). Duffy BC; Howard KT; Chisholm JD Alkylation of thiols with trichloroacetimidates under neutral conditions. *Tetrahedron Lett.* 2015, 56, 3301.
- (35). Mahajani NS; Chisholm JD Promoter free allylation of trichloroacetimidates with allyltributylstannanes under thermal conditions to access the common 1,1'-diarylbutyl pharmacophore. *Org. Biomol. Chem* 2018, 16, 4008. [PubMed: 29766199]
- (36). Mahajani NS; Chisholm JD Synthesis of 1,1'-Diarylethanes and Related Systems by Displacement of Trichloroacetimidates with Trimethylaluminum. *J. Org. Chem* 2018, 83, 4131. [PubMed: 29478311]
- (37). Wallach DR; Chisholm JD Alkylation of Sulfonamides with Trichloroacetimidates under Thermal Conditions. *J. Org. Chem* 2016, 81, 8035. [PubMed: 27487402]
- (38). Ali IAI; El Ashry ESH; Schmidt RR Protection of Hydroxy Groups with Diphenylmethyl and 9-Fluorenyl Trichloroacetimidates - Effect on Anomeric Stereocontrol. *Eur. J. Org. Chem* 2003, 4121.
- (39). Zheng H-X; Shan X-H; Qu J-P; Kang Y-B Transition-Metal-Free Hydrogenation of Aryl Halides: From Alcohol to Aldehyde. *Org. Lett* 2017, 19, 5114. [PubMed: 28925712]
- (40). Ghorai J; Anbarasan P Rhodium Catalyzed Arylation of Diazo Compounds with Aryl Boronic Acids. *J. Org. Chem* 2015, 80, 3455. [PubMed: 25759939]
- (41). Dunsford JJ; Clark ER; Ingleson MJ Direct C(sp²)-C(sp³) Cross-Coupling of Diaryl Zinc Reagents with Benzylic, Primary, Secondary, and Tertiary Alkyl Halides. *Angew. Chem., Int. Ed* 2015, 54, 5688.
- (42). Wiener H; Gilon C An improved method for the catalytic preparation of tert-butyl esters of carboxylic and fatty acids. *J. Mol. Catal* 1986, 37, 45.
- (43). Wang P; Tao W-J; Sun X-L; Liao S; Tang Y A Highly Efficient and Enantioselective Intramolecular Cannizzaro Reaction under TOX/Cu(II) Catalysis. *J. Am. Chem. Soc* 2013, 135, 16849. [PubMed: 24161001]
- (44). Huihui KMM; Caputo JA; Melchor Z; Olivares AM; Spiewak AM; Johnson KA; DiBenedetto TA; Kim S; Ackerman LKG; Weix DJ Decarboxylative Cross-Electrophile Coupling of N-Hydroxyphthalimide Esters with Aryl Iodides. *J. Am. Chem. Soc* 2016, 138, 5016. [PubMed: 27029833]
- (45). Ryan SJ; Zhang Y; Kennan AJ Convenient access to glutamic acid side chain homologs compatible with solid phase peptide synthesis. *Org. Lett* 2005, 7, 4765. [PubMed: 16209530]
- (46). Zhang H; Huang X Ligand-Free Heck Reactions of Aryl Iodides: Significant Acceleration of the Rate through Visible Light Irradiation at Ambient Temperature. *Adv. Synth. Catal* 2016, 358, 3736.
- (47). Li X; Zou D; Zhu H; Wang Y; Li J; Wu Y; Wu Y Preparation of tert-Butyl Esters via Pd-Catalyzed tert-Butoxycarbonylation of (Hetero)aryl Boronic Acid Derivatives. *Org. Lett* 2014, 16, 1836. [PubMed: 24655126]

- (48). Moore JD; Byrne RJ; Vedantham P; Flynn DL; Hanson PR High-Load, ROMP-Generated Oligomeric Bis-acid Chlorides: Design of Soluble and Insoluble Nucleophile Scavengers. *Org. Lett* 2003, 5, 4241. [PubMed: 14601970]
- (49). Steemers L; Wijsman L; van Maarseveen JH Regio- and Stereoselective Chan-Lam-Evans Enol Esterification of Carboxylic Acids with Alkenylboroxines. *Adv. Synth. Catal* 2018, 360, 4241.
- (50). Won J-E; Kim H-K; Kim J-J; Yim H-S; Kim M-J; Kang S-B; Chung H-A; Lee S-G; Yoon Y-J Effective esterification of carboxylic acids using (6-oxo-6H-pyridazin-1-yl)phosphoric acid diethyl ester as novel coupling agents. *Tetrahedron* 2007, 63, 12720.
- (51). Ulatowski F; Jurczak J Chiral Recognition of Carboxylates by a Static Library of Thiourea Receptors with Amino Acid Arms. *J. Org. Chem* 2015, 80, 4235. [PubMed: 25867541]
- (52). Feng X; Sun A; Zhang S; Yu X; Bao M Palladium-catalyzed carboxylative coupling of benzyl chlorides with allyltributylstannane: remarkable effect of palladium nanoparticles. *Org. Lett* 2013, 15, 108. [PubMed: 23249213]
- (53). Huang H; Kang JY Mitsunobu Reaction Using Basic Amines as Pronucleophiles. *J. Org. Chem* 2017, 82, 6604. [PubMed: 28558240]
- (54). Lu B; Zhu F; Sun H-M; Shen Q Esterification of the Primary Benzylic C-H Bonds with Carboxylic Acids Catalyzed by Ionic Iron(III) Complexes Containing an Imidazolium Cation. *Org. Lett* 2017, 19, 1132. [PubMed: 28198186]
- (55). Curran SP; Connon SJ The thiolate-catalyzed intermolecular crossed Tishchenko reaction: highly chemoselective coupling of two different aromatic aldehydes. *Angew. Chem., Int. Ed* 2012, 51, 10866.
- (56). Wang MF; Golding BT; Potter GA A Convenient Preparation of p-Methoxybenzyl Esters. *Synth. Commun* 2000, 30, 4197.
- (57). Zeggaf C; Poncet J; Jouin P; Dufour MN; Castro B Isopropenyl chlorocarbonate (IPCC) in amino acid and peptide chemistry: esterification of N-protected amino acids; application to the synthesis of the depsipeptide valinomycin. *Tetrahedron* 1989, 45, 5039.
- (58). Rolfe A; Loh JK; Maity PK; Hanson PR High-load, hybrid Si-ROMP reagents. *Org. Lett* 2011, 13, 4. [PubMed: 21128690]
- (59). Ohno O; Ye M; Koyama T; Yazawa K; Mura E; Matsumoto H; Ichino T; Yamada K; Nakamura K; Ohno T; Yamaguchi K; Ishida J; Fukamizu A; Uemura D Inhibitory effects of benzyl benzoate and its derivatives on angiotensin II-induced hypertension. *Bioorg. Med. Chem* 2008, 16, 7843. [PubMed: 18672373]
- (60). Wu H; Guo W; Daniel S; Li Y; Liu C; Zeng Z Fluoride-Catalyzed Esterification of Amides. *Chem. - Eur. J* 2018, 24, 3444. [PubMed: 29369460]
- (61). Xia Y; Liu Z; Feng S; Ye F; Zhang Y; Wang J Rh(I)-Catalyzed Cross-Coupling of α -Diazoesters with Arylsiloxanes. *Org. Lett* 2015, 17, 956. [PubMed: 25671642]
- (62). Muzart J; Pale P; Pete JP; Riahi A Light-induced oxidation of η^3 -allylpalladium complexes by molecular oxygen. *Bull. Soc. Chim. Fr* 1988, 731.
- (63). Magens S; Plietker B Nucleophilic Iron Catalysis in Transesterifications: Scope and Limitations. *J. Org. Chem* 2010, 75, 3715. [PubMed: 20462205]

Figure 1.
Archetypal Esterification Pathways

Figure 2.
Alkylation Agents that Undergo Symbiotic Activation

Figure 3.
Ester formation via symbiotic activation of the trichloroacetimidate.

Scheme 1.
Esterification of Naproxen without Racemization

Table 1.

Esterifications with Different Trichloroacetimidates

$\text{R}'\text{O}-\text{C}(=\text{NH})\text{CCl}_3 + \text{Ph}-\text{C}(=\text{O})\text{OH} \xrightarrow{\text{conditions}} \text{Ph}-\text{C}(=\text{O})\text{O}-\text{R}'$				
entry	imidate	ester	yield (%) ^a	yield (%) ^b
1		MeOBz 10a	0 (48 h)	0 (18 h)
2		EtOBz 11a	0 (48 h)	0 (18 h)
3			0 (48 h)	0 (18 h)
4			84 (48 h)	17 (18 h)
5			-	0 (48 h)
6			73 ^c (16 h)	-
7			-	0 (48 h)
8			0 (48 h)	76 (18 h)
9			0 (48 h)	82 (18 h)

entry	imideate	ester	yield (%) ^a	yield (%) ^b
10			19 (48 h)	96 (19 h)
11			89 (24 h)	-
12			90 (1 h)	-
13			96 (3 h)	-
14			-	0 (48 h)
15			34 (22 h)	96 (18 h)
16			34 (48 h)	95 (23 h)
17			99 (18 h)	-

^a Isolated yield (DCM, rt).

^b Isolated yield (toluene, reflux).

^c Trace of the *tert*-prenyl isomer also detected in the ¹H NMR.

Table 2.

Reaction of Complex Carboxylic acids with Different Trichloroacetimidates

$\text{R}-\text{C}(=\text{O})\text{OH} + \text{R}'\text{O}-\text{C}(=\text{NH})\text{CCl}_3 \xrightarrow{\text{DCM, rt}} \text{R}-\text{C}(=\text{O})\text{O}-\text{R}'$				
entry	acid	imidate	time (h)	yield (%)
1	27	<i>t</i> -Bu (13)	48	81 (13b)
2	27	PMB (20)	24	96 (20b)
3	27	DMB (21)	1	88 (21b)
4	27	DPM (26)	18	76 (26b)
5	28	<i>t</i> -Bu (13)	48	60 (13c)
6	28	PMB (20)	24	93 (20d)
7	28	DMB (21)	24	81 (21d)
8	28	DPM (26)	18	83 (26d)
9	29	<i>t</i> -Bu (13)	48	94 (13d)
10	29	PMB (20)	24	93 (20d)
11	29	DMB (21)	20	81 (21d)
12	29	DPM (26)	18	83 (26d)
13	30	<i>t</i> -Bu (13)	48	59 (13e)
14	30	PMB (20)	24	79 (20e)
15	30	PMB (21)	1	94 ^a (21e)
16	30	DPM (26)	18	92 (26e)

entry	acid	imideate	time (h)	yield (%)
17		<i>t</i> -Bu (13)	48	62 (13f)
18	31	PMB (20)	24	91 (20f)
19	31	DMB (21)	24	80 (21f)
20	31	DPM (26)	3	54 ^a (26f)
21		<i>t</i> -Bu (13)	48	30 ^b (13g)
22	32	PMB (20)	24	83 (20g)
23	32	DMB (21)	0.5	73 (21g)
24	32	DPM (26)	18	76 (26g)
25		<i>t</i> -Bu (13)	48	60 (13h)
26	33	PMB (20)	24	94 (20h)
27	33	DMB (21)	24	88 (21h)
28	33	DPM (26)	18	93 (26h)
29		<i>t</i> -Bu (13)	48	76 (13i)
30	34	PMB (20)	24	84 (20i)
31	34	DMB (21)	4	92 (21i)
32	34	DPM (26)	18	72 (26i)
33		<i>t</i> -Bu (13)	48	81 (13j)
34	35	PMB (20)	24	83 (20j)
35	35	DMB (21)	24	83 (21j)
36	35	DPM (26)	18	71 (26j)

^a (only one equiv imideate was used)^b (Yield determined by ¹H NMR using a mesitylene standard as product was volatile)

Table 3.

Esterifications of Benzyl Trichloroacetimidate in Refluxing Toluene

$\text{R}-\text{C}(=\text{O})\text{OH} + \text{BnO}-\text{C}(=\text{NH})-\text{CCl}_3 \xrightarrow[\text{reflux, 18 h}]{\text{toluene}} \text{R}-\text{C}(=\text{O})\text{OBn}$		
entry	carboxylic acid	yield (%)
1	<p>27</p>	78 (17b)
2	<p>28</p>	94 (17c)
3	<p>29</p>	80 (17d)
4	<p>30</p>	88 (17e)
5	<p>31</p>	94 (17f)
6	<p>32</p>	88 (17g)
7	<p>33</p>	77 (17h)
8	<p>34</p>	80 (17i)

Table 4.

Esterification with Enantioenriched Imidates

R^1 -C₆H₄-COOH + Cl₃C-C(=O)-O-CH(R²)-Ph → R¹-C₆H₄-CO-O-CH(R²)-Ph

9 R¹ = H **25** R² = Me **25a** R¹ = H R² = Me
36 R¹ = Ph **37** R² = 2-OMePh **38** R¹ = Ph R² = 2-OMePh

entry	imide	conditions	yield (%)	e. r.
1	25	toluene, , 24 h	95	50:50 (25a)
2	25	DCM, rt, 24 h	35	71:29 (25a)
3	37	toluene, , 24 h	73	55:45 (38)
4	37	DCM, rt, 24 h	60	67:33 (38)