Published in final edited form as:

Nat Chem. 2019 May 1; 11(5): 457–462. doi:10.1038/s41557-019-0225-x.

Prebiotic Phosphorylation of 2-Thiouridine Provides Either Nucleotides or DNA Building Blocks *via* Photoreduction

Jianfeng Xu¹, Nicholas J. Green¹, Clémentine Gibard², Ramanarayanan Krishnamurthy^{2,*}, and John D. Sutherland^{1,*}

¹MRC Laboratory of Molecular Biology, Francis Crick Avenue, Cambridge Biomedical Campus, Cambridge CB2 0QH, UK

²Department of Chemistry, The Scripps Research Institute, 10550 North Torrey Pines Road, La Jolla, California 92037, USA

Abstract

Breakthroughs in the study of the origin of life have demonstrated how some of the building blocks essential to biology could have been formed under various primordial scenarios, and could therefore have contributed to the chemical evolution of life. Missing building blocks are then sometimes inferred to be products of primitive biosynthesis, which can stretch the limits of plausibility. Here, we demonstrate the synthesis of 2'-deoxy-2-thiouridine, and subsequently 2'-deoxyadenosine and 2-deoxyribose, under prebiotic conditions. 2'-Deoxy-2-thiouridine is produced by photoreduction of 2,2'-anhydro-2-thiouridine, which is in turn formed by phosphorylation of 2-thiouridine—an intermediate of prebiotic RNA synthesis. 2'-Deoxy-2-thiouridine is an effective deoxyribosylating agent and may have functioned as such in either abiotic or proto-enzyme-catalysed pathways to DNA, as demonstrated by its conversion to 2'-deoxyadenosine by reaction with adenine, and 2-deoxyribose by hydrolysis. An alternative prebiotic phosphorylation of 2-thiouridine leads to the formation of its 5'-phosphate, showing that hypotheses in which 2-thiouridine was a key component of early RNA sequences are within the bounds of synthetic credibility.

Non-canonical, sulfur-containing nucleobases are utilised by extant biology^{1,2}, and exert interesting effects on the stability, base-pairing properties and replication fidelity of nucleic acid polymers³. As such—and given the key role of sulfur in established prebiotic synthetic scenarios^{4,5}—they are potential components of a genetic system that either complemented or perhaps predated, and ultimately evolved into, enzymatically replicated canonical RNA and DNA. We recently demonstrated a viable pathway to pyrimidine ribonucleosides and

*Correspondence and requests for materials should be addressed to J.D.S. and R.K. johns@mrc-lmb.cam.ac.uk; rkrishna@scripps.edu.

Data availability

Full experimental details and data are provided in the Supplementary Information. The raw data for Supplementary Figs. 49–53 are available from the authors upon reasonable request.

Author contributions. J.D.S. and R.K. supervised the experimental research. J.X., N.J.G. and C.G. performed the experiments. All the authors co-wrote the paper and assembled the Supplementary Information.

Competing interests

The authors declare no competing interests.

Additional information

ribonucleotides, *via* thioribonucleosides (Fig. 1)⁵. Highly crystalline ribo-aminooxazoline-**1** was shown to react with cyanoacetylene **2** to give anhydronucleoside **3**, which on thiolysis furnished α -2-thioribocytidine **4**. **4** could be efficiently photoanomerised to β -2-thioribocytidine **5**, which serves as a precursor to the canonical pyrimidine ribonucleosides cytidine **6** and uridine **7**, β -2-thiouridine **8**, and oligomerisable⁶ cyclic phosphates of cytidine and uridine (**9** and **10**, respectively).

Since we had demonstrated that thioribonucleosides **5** and **8** are likely formed in primordial pyrimidine synthesis, we set out to further explore prebiotic phosphorylation conditions that might yield oligomerisable thioribonucleotides. Herein, we demonstrate that phosphorylation of 2-thiouridine **8** can either lead to an oligomerisable nucleotide or initiate a prebiotic synthesis of a deoxythionucleoside, 2'-deoxy-2-thiouridine **11**, *via* hydrogen sulfide-mediated photoreduction as the key step. We show that **11** is a productive transglycosylating reagent in an abiotic or proto-enzymatic context; for example, in the supply of 2-deoxyribose **12**, and the synthesis of a canonical 2'-deoxynucleoside, 2'-deoxyadenosine **13**. These results provide a chemical connection between RNA and DNA in a prebiotic context (that is, in the absence of biosynthesis).

Results and Discussion

5' Phosphorylation

First, we attempted the phosphorylation of 2-thiouridine **8** in hot formamide⁷, but increased the concentration of phosphate and nucleoside from 100 mM to 200 mM, compared to our previous phosphorylation of cytidine (full experimental details are included in the Supplementary Information). After heating at 100 °C for 20 h, we observed the initially formed phosphorylated product, thiouridine 5'-phosphate **14**, in 24% yield alongside the starting material **8** in 42% and some minor products (Fig. 2, Supplementary Fig. 39). In hot urea, ribonucleosides generally undergo initial (kinetic) 5'-phosphorylation⁸; over time, the 2',3'-cyclic phosphate dominates over the 5'-phosphate (thermodynamic control). The formation of phosphate monoesters in hot urea (and formamide) is reversible but slow, such that clear-cut kinetic or thermodynamic control is not shown by such systems. Szostak and coworkers^{3,9} have already shown that non-enzymatic RNA replication using the 2-methyl- and 2-amino-imidazole-activated derivatives of **14** proceeds with higher rate and fidelity than the corresponding activated uridine nucleotide, but until now a prebiotic provenance of **14** remained unestablished. These results show that the kinetic product of phosphorylation—the 5' nucleotide **14**—is indeed accessible, and would become increasingly plausible if progressively milder prebiotically plausible phosphorylation procedures are uncovered. Our synthesis of **14** shows that it could have been available for non-enzymatic incorporation into primordial RNA sequences, and provides further impetus for the investigation of the potential role of 2-thiouridine in RNA.

Thioanhydride Formation

Semi-molten urea is an alternative prebiotic phosphorylation medium to hot formamide⁸, so we next used it to attempt phosphorylation of **8** (Fig. 3a). Surprisingly, no 2',3'-cyclic nucleotide derivatives were observed. A major product had been formed, but a total of four

monophosphate ^{31}P -NMR resonances suggested that the hydroxyl groups of this product were phosphorylated to varying degrees, complexifying the NMR spectra. The crude mixture from the phosphorylation reaction was therefore treated with alkaline phosphatase, generating a single major nucleoside product. Extensive analysis of both one- and two-dimensional NMR spectra of the dephosphorylated product suggested the formation of 2',2-thioanhydronucleoside **15a**¹⁰. Previously, Hampton and Nichol synthesized a related derivative, 2',2-anhydro-uridine **16**, when uridine **7** was treated with diphenylcarbonate in hot dimethylformamide (Fig. 3b). Under those conditions, intermediate uridine 2',3'-cyclic carbonate¹¹ **17a** is a good substrate for ring opening at C2' by 2-O to give cyclonucleoside **16**. Thus, under phosphorylation conditions, we propose cyclic phosphate **18** as the intermediate in the formation of **15** ($\text{R}_1 = \text{R}_2 = \text{H}$ or PO_3H^-), with rapid ring opening by the nucleophilic sulfur atom ensuing to convert **18** to **15**. To confirm the structure of the product and support our proposed mechanism, 2',2-anhydro-2-thiouridine **15a** was unambiguously synthesized from 2-thiouridine **8** and diphenylcarbonate (Fig. 3b). The fully characterized synthetic standard was used for spiking experiments to confirm that the major product of phosphorylation of **8** (even before enzymatic dephosphorylation) was indeed **15a**. After the crude reaction products from the phosphorylation were subjected to alkaline phosphatase (Fig. 3a), addition of an NMR integration standard (sodium formate) revealed that 2',2-anhydro-2-thiouridine **15a** had been formed in 50% yield, accompanied by some glycosidic bond cleavage (2-thiouracil **19** (15%) and isocytosine **20** (7%)). To obtain the ratio of the anhydronucleoside to its phosphorylated derivatives before treatment with alkaline phosphatase, the crude mixture from the phosphorylation of **8** was subjected to analytical high performance liquid chromatography, which showed the ratio of the **15a**: **15b**: **15c** was 3.7 : 1: 1 (Supplementary Figs. 57 and 58). The same phosphorylation conditions were also applied to 2-thiocytidine **5** (Fig. 3a), which afforded 2',2-anhydro-2-thiocytidine **21**¹² (43% yield), cytidine **6** (8%), diaminopyrimidine **22** (16%), 2-thiocytosine **23** (8%) and cytosine **24** (8%). All structures were confirmed by spiking experiments, with synthetic standards made in house or purchased (Supplementary Figs. 32 and 35).

To further support a cyclic phosphate intermediate, we sought milder conditions for phosphorylation of **8**. We chose diamidophosphate (DAP), a reagent known to generate not only the corresponding 2',3'-cyclophosphates from nucleosides, but also short oligonucleotides after prolonged reaction.¹³ Indeed, reaction of 2-thionucleoside **8** with DAP and imidazole under moist-paste conditions at room temperature resulted in formation of the cyclic phosphate **18a** after one week (Fig. 4a). Monitoring the phosphorylation reaction by ^{31}P , ^1H and ^{13}C NMR, we observed the transformation of cyclophosphate **18a** to a new species that showed characteristic NMR, electrospray ionization high-resolution mass spectrometry and ultraviolet spectral signatures of anhydronucleotide **15b** (Supplementary Figs. 60–63). Furthermore, the DAP-reaction of **8** does not stop at **15b**; the 3' phosphate of **15b** reacts again with DAP leading to the corresponding 3'-amidodiphosphate of the 2',2-thioanhydronucleoside **25**. The total conversion (measured by ^1H NMR) of **8** to **15b/25** was 78% after one month at room temperature. When 4-thiouridine **26** was subjected to the same DAP phosphorylation reaction as a control, the expected cyclophosphate **27** was formed after a few days at room temperature, but no further transformation to the corresponding 2',2-anhydride was observed (Fig. 4a; Supplementary Figs. 66, 67). In this case, cyclisation

was presumably prevented due to the lower nucleophilicity of the oxygen on C2 compared with sulfur.

We also investigated the reaction of DAP (moist paste) with 2-thiocytidine **5**, which barely proceeded under the conditions employed for **8**. By lowering the pH of the reaction-paste from 7 to 4 and by raising the reaction temperature to 60 °C, we were able to detect cyclic phosphate **28**, and later were able to identify the corresponding 3'-phosphate and 3'-amidopyrophosphate of thioanhydrocytidine (**21b/29**) by ^{31}P NMR (Fig. 4b; Supplementary Figs. 64, 65). An accurate conversion was difficult to determine by ^1H -NMR, but the combined yield for **28**, **21b** and **29** was estimated at slightly greater than 10%. The greater reactivity of 2-thiouridine **8** towards DAP compared to 2-thiocytidine **5** likely stems from its greater solubility and pK_a near neutrality (8.0, N3-H)².

C2' Reduction

Our syntheses of thioanhydronucleosides **15** and **21** from 2-thionucleosides **8** and **5** are the first prebiotically plausible routes to nucleoside derivatives containing a sulfur atom connected to C2'. Such compounds are of significant interest in the origins-of-life field, because the potential conversion of a C–S to a C–H bond (a chemically facile process^{14, 15}) would result in the synthesis of 2'-deoxynucleosides – components of DNA. In modern biology, ribonucleotide reductases convert ribonucleotides to deoxyribonucleotides¹⁶. This enzymatic reduction proceeds *via* a radical mechanism with a dithiol as the stoichiometric reductant. We wondered if a mechanism involving sulfur-based reductants might have led to the formation of 2'-deoxyribonucleosides/tides on the early earth, in the absence of complex enzymes, from 2'-thio-functionalised ribonucleosides. Nagyvary and coworkers^{17–19} first experimentally investigated this idea by synthesizing 2'-thiocytidine (using an enzyme) and studying a variety of reduction procedures, including photoreduction. However, the limited published characterization data for the supposed 2'-thiocytidine substrate have since been pointed out as apparently inconsistent with the structure²⁰. Recently, Powner and coworkers²¹ revisited the idea with a proposal to modify the prebiotically plausible pyrimidine synthesis^{5, 22} by including a sulfur atom attached to C2' from the outset. This route proved unsuccessful because the sulfur-containing building blocks exhibited different reactivity to analogues from the pyrimidine synthesis. Indeed, while a route from pure ribose or arabinose to deoxyribose has recently been established²³, a plausible abiotic synthesis of 2'-deoxynucleosides themselves – whether by desulfurization or other means – has not yet been reported. 2',2'-thioanhydronucleosides **15a** and **21a** were therefore subjected to ultraviolet irradiation at 254 nm in the presence of aqueous hydrogen sulfide as a reductant (Fig. 5a). Flux measurements from the same experimental setup have been carried out within our group previously, from which we determined a flux rate of $2.5 \times 10^{16} \text{ cm}^{-2} \text{ s}^{-1}$ (19 mWcm⁻²) for the 254-nm-centred emission²⁴. Inspection of ^1H -NMR spectra showed that, under these conditions, thioanhydrocytidine **21a** gave 2-thiocytosine **23** in 29% yield after 3 h of irradiation. However, thioanhydrouridine **15a** afforded two new nucleoside derivatives as well as thiouracil **19** as major products. Promisingly, one nucleoside showed two 2'-H signals shifted upfield (2.64 ppm and 2.32 ppm) in the ^1H NMR spectrum (Supplementary Fig. 38), strongly suggesting that a deoxyribosyl moiety was present. Comparison of NMR spectra with literature values for deoxyribosyl pyrimidines suggested the formation of 2'-

deoxy-2-thiouridine **11**. We subsequently verified this assignment by synthesizing **11** *via* a literature route²⁵ and using this material to spike our samples (Supplementary Fig. 40). We speculate that the second nucleoside product is 2'- α -thio-2-thiouridine **30**, but attempts to isolate and characterize this compound have so far been thwarted by its facile decomposition to 2-thiouracil **19** as the only identifiable product. The instability of the glycosidic bonds of 2'-thioribonucleosides has previously been documented²⁶. Addition of the NMR integration standard, pentaerythritol, to the crude mixture indicated yields of 33, 25 and 28% for **11**, **19** and presumed **30**, respectively. The 5'- and 3'-phosphates of 2',2-anhydro-2-thiouridine (**15b** and **15c**, respectively) undergo the same photoreduction, in similar yields to **15a** (Supplementary Figs. 41 and 42).

A possible mechanism for this remarkable photoreduction is proposed in Figure 5b. Separate control reactions in the dark with H₂S and irradiation without H₂S gave no reduction products nor **30**, suggesting that initiation is by an electron released from hydrosulfide under ultraviolet irradiation^{4, 27, 28}, rather than homolysis of the C–S bond of **15a**. Transfer of this electron to thioanhydrouridine **15a** to form the radical anion **31**, and subsequent homolysis of the C–S bond would give intermediate **32**. The newly formed C2' radical can abstract a hydrogen atom from H₂S (pictured) or react with a hydrogen atom (not pictured), to give **11**. Alternatively, reaction with an HS[•] radical equivalent (HS[•] with loss of an electron, or H₂S₂ or a HS[•] radical, generated in previous steps) forms 2'-thionucleoside **30**. We conjecture that **21** probably does not undergo this reduction because the first step, were it to occur, would lead to a radical anion intermediate with less resonance stabilization than **31**.

Syntheses of deoxyribose and β -2'-deoxyadenosine

Since our synthesis of 2'-deoxyribonucleoside **11** takes place under the same conditions under which amino acid⁴ and sugar²⁹ precursors and RNA pyrimidines^{5, 22} are formed, we immediately questioned whether DNA building blocks such as **11**, and related (canonical) derivatives, may also have been present at the advent of life. In an attempt to form the canonical 2'-deoxyribonucleoside, deoxyuridine, we investigated the hydrolysis of **11** under the same conditions under which β -2-thioribocytidine **5** is converted to cytidine (Fig. 1)⁵. However, in the presence of phosphate buffer, **11** is cleanly hydrolysed to 2-deoxyribose **12** and thiouracil **19**. A detailed study of the hydrolysis of 2'-deoxy-2-thiouridine **11** at 60 °C revealed half-lives of 32 and 31 hours for 2'-deoxy-2-thiouridine **11** in 0.27M phosphate (pH 7.0) and acetate (pH 4.1) buffers, respectively, and 28 hours in unbuffered solution (Fig. 6a and Supplementary Figs. 43–50). The complete and clean conversion of 2'-deoxy-2-thiouridine **11** to 2-thiouracil **19** and deoxyribose **12** suggests that **11** may have been a prebiotic source of deoxyribose, both in abiotic reactions and/or for early enzymes, akin to the use of activated ribose in modern phosphoribosyltransferases^{30–32}. This reactivity of **11** presented an opportunity for a plausible synthesis of purine deoxynucleosides, *via* transglycosylation^{33–35}. Indeed, heating of **11** with excess adenine **33** in the dry state³⁶ at 100 °C for 31 h led to the formation of both α and β isomers of 2'-deoxyadenosine **13**, in 6% and 4% yield respectively, accompanied by nucleobase loss to give 2-thiouracil **19** in 72% yield (Fig 6b and Supplementary Figs. 54 and 55). Notably, replacement of **11** with 2-deoxyribose **12** led only to a complex mixture that did not contain β -2'-deoxyadenosine (Supplementary Fig. 56). Such a demonstration of a plausible abiotic pathway stemming

from the ribo-pyrimidine synthesis to DNA is significant, and further efforts are underway in our laboratory to discover more favourable transglycosylation conditions. Additionally, if promiscuous proto-enzymes with ribosylase activity existed, **11** may have been an efficient source of deoxyribose in an early biosynthesis of deoxyribonucleotides *via* transglycosylation. Thus, there are various plausible pathways by which the C2'-reduced nucleoside **11** could have facilitated the chemical progression of RNA into DNA.

Conclusions

Our investigation into the phosphorylation of prebiotic RNA pyrimidine intermediate **8** has resulted in several significant discoveries. We have shown that 5'-phosphoryl-2-thiouridine **14** can be obtained from kinetically selective phosphorylation of **8**, demonstrating its viability as a potentially important primordial non-canonical component of nucleic acids. Other phosphorylation conditions lead to the until-now elusive prebiotic functionalization of the C2' position of a nucleoside with sulfur. Moreover, we successfully reduced such a sulfur-containing derivative to a non-canonical 2'-deoxyribonucleoside, **11**, using a prebiotically plausible, hydrogen sulfide-mediated photoreduction. This photochemical reaction emulates the critical biochemical conversion of ribonucleotides to 2'-deoxyribonucleotides, but without the involvement of any complex enzymes. Finally, we have demonstrated that this deoxyribonucleoside reacts with adenine to form β -deoxyadenosine **13b**, a canonical DNA purine deoxynucleoside. Thus, 2-thioribonucleosides are key intermediates in a synthesis of both RNA and DNA building blocks, providing a long sought-after prebiotic link between DNA and its biological and chemical progenitor, RNA.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

This work was supported by the Medical Research Council (number MC_UP_A024_1009), and a grant from the Simons Foundation (number 290362 to J.D.S. and number 327124 to R.K.). We acknowledge P. B. Rimmer for discussions on the photochemical reactor setup.

References

1. Rao YS, Thimmappayya B, Cherayil JD. Thionucleotides in the transfer ribonucleic acids of *Pseudomonas aeruginosa* and *Escherichia coli*. *Biochem J.* 1972; 128:67.
2. Larsen AT, Fahrenbach AC, Sheng J, Pia J, Szostak JW. Thermodynamic insights into 2-thiouridine-enhanced RNA hybridization. *Nucleic Acids Res.* 2015; 43:7675–7687. [PubMed: 26240387]
3. Heuberger BD, Pal A, Del Frate F, Topkar VV, Szostak JW. Replacing Uridine with 2-Thiouridine Enhances the Rate and Fidelity of Nonenzymatic RNA Primer Extension. *J Am Chem Soc.* 2015; 137:2769–2775. [PubMed: 25654265]
4. Patel BH, Percivalle C, Ritson DJ, Duffy CD, Sutherland JD. Common origins of RNA, protein and lipid precursors in a cyanosulfidic protometabolism. *Nat Chem.* 2015; 7:301–307. [PubMed: 25803468]
5. Xu J, et al. A prebiotically plausible synthesis of pyrimidine beta-ribonucleosides and their phosphate derivatives involving photoanomerization. *Nat Chem.* 2017; 9:303–309. [PubMed: 28338689]

6. Verlander MS, Lohrmann R, Orgel LE. Catalysts for the self-polymerization of adenosine cyclic 2', 3'-phosphate. *J Mol Evol.* 1973; 2:303–316. [PubMed: 4377773]
7. Schoffstall AM. Prebiotic phosphorylation of nucleosides in formamide. *Orig Life.* 1976; 7:399–412. [PubMed: 1023139]
8. Lohrmann R, Orgel LE. Urea-Inorganic Phosphate Mixtures as Prebiotic Phosphorylating Agents. *Science.* 1971; 171:490–494. [PubMed: 5099649]
9. Li L, et al. Enhanced Nonenzymatic RNA Copying with 2-Aminoimidazole Activated Nucleotides. *J Am Chem Soc.* 2017; 139:1810–1813. [PubMed: 28117989]
10. Ueda T, Shibuya S. Synthesis of Sulfur-Bridged Uracil Anhydronucleosides. *Chem Pharm Bull.* 1970; 18:1076–1078.
11. Hampton A, Nichol A. Purine Ribonucleoside 2',3'-Cyclic Carbonates. Preparation and Use for the Synthesis of 5'-Monosubstituted Nucleosides. *Biochemistry.* 1966; 5:2076–2082. [PubMed: 5963450]
12. Russell AF, Prystasz M, Hamamura EK, Verheyden JP, Moffatt JG. Reactions of 2-acyloxyisobutryl halides with nucleosides. V. Reactions with cytidine and its derivatives. *J Org Chem.* 1974; 39:2182–2186. [PubMed: 4854037]
13. Gibard C, Bhowmik S, Karki M, Kim EK, Krishnamurthy R. Phosphorylation, oligomerization and self-assembly in water under potential prebiotic conditions. *Nat Chem.* 2018; 10:212–217. [PubMed: 29359747]
14. Belen'kii LI. Chemistry of organosulfur compounds : general problems. Horwood E. 1990
15. Larock, RC. Comprehensive organic transformations : a guide to functional group preparations. Third edition. John Wiley & Sons; 2018.
16. Reichard P. From RNA to DNA, why so many ribonucleotide reductases? *Science.* 1993; 260:1773–1777. [PubMed: 8511586]
17. Bradbury E, Nagyvary J. Preparation of 2'-Thio-2'-Deoxycytidine 2'-3'-Phosphorothioate. *Nucleic Acids Res.* 1976; 3:2437–2442. [PubMed: 995638]
18. Patel AD, Schrier WH, Nagyvary J. Synthesis and Properties of 2'-Deoxy-2'-Thiocytidine. *J Org Chem.* 1980; 45:4830–4834.
19. Wolman, Y, editor. Origin of life : proceedings of the Third ISSOL meeting and the Sixth ICOL meeting, Jerusalem, June 22-27, 1980. D. Reidel Pub. Co; 1981.
20. Divakar KJ, Mottoh A, Reese CB, Sanghvi YS. Approaches to the synthesis of 2'-thio analogues of pyrimidine ribosides. *J Chem Soc, Perkin Trans.* 1990; 1:969–974.
21. Powner MW, Zheng SL, Szostak JW. Multicomponent Assembly of Proposed DNA Precursors in Water. *J Am Chem Soc.* 2012; 134:13889–13895. [PubMed: 22839703]
22. Powner MW, Gerland B, Sutherland JD. Synthesis of activated pyrimidine ribonucleotides in prebiotically plausible conditions. *Nature.* 2009; 459:239–242. [PubMed: 19444213]
23. Ritson DJ, Sutherland JD. Conversion of Biosynthetic Precursors of RNA to Those of DNA by Photoredox Chemistry. *J Mol Evol.* 2014; 78:245–250. [PubMed: 24736972]
24. Rimmer PB, et al. The origin of RNA precursors on exoplanets. *Science Advances.* 2018; 4doi: 10.1126/sciadv.aar3302
25. Kuimelis RG, Hope H, Nambiar KP. A Stereoselective Synthesis of α - and β -2'-Deoxy-2-thiouridine. *Nucleosides and Nucleotides.* 1993; 12:737–755.
26. Johnson R, Reese CB, Peizhuo Z. Lability of Glycosidic Linkages of 2'-Thio-Ribonucleosides. *Tetrahedron.* 1995; 51:5093–5098.
27. Ritson DJ, Sutherland JD. Synthesis of Aldehydic Ribonucleotide and Amino Acid Precursors by Photoredox Chemistry. *Angew Chem Int Ed.* 2013; 52:5845–5847.
28. Sauer MC, Crowell RA, Shkrob IA. Electron Photodetachment from Aqueous Anions. 1. Quantum Yields for Generation of Hydrated Electron by 193 and 248 nm Laser Photoexcitation of Miscellaneous Inorganic Anions. *The Journal of Physical Chemistry A.* 2004; 108:5490–5502.
29. Ritson D, Sutherland JD. Prebiotic synthesis of simple sugars by photoredox systems chemistry. *Nat Chem.* 2004; 4:895–899.
30. Unrau PJ, Bartel DP. RNA-catalysed nucleotide synthesis. *Nature.* 1998; 395:260–263. [PubMed: 9751052]

31. Chapple KE, Bartel DP, Unrau PJ. Combinatorial minimization and secondary structure determination of a nucleotide synthase ribozyme. *RNA*. 2003; 9:1208–1220. [PubMed: 13130135]
32. Lau MW, Cadieux KE, Unrau PJ. Isolation of fast purine nucleotide synthase ribozymes. *J Am Chem Soc*. 2004; 126:15686–15693. [PubMed: 15571390]
33. Miyaki M, Saito A, Shimizu B. N→N Alkyl and Glycosyl Migrations of Purines and Pyrimidines. IV. trans-Glycosylation from Pyrimidines to Purines. (A Novel Synthetic Method of Purine Nucleosides and Nucleotides). *Chem Pharm Bull*. 1970; 18:2459–2468.
34. Vorbrüggen, H, Ruh-Pohlentz, C. Handbook of nucleoside synthesis. Wiley; 2001.
35. Nam I, Nam HG, Zare RN. Abiotic synthesis of purine and pyrimidine ribonucleosides in aqueous microdroplets. *Proc Natl Acad Sci USA*. 2018; 115:36–40. [PubMed: 29255025]
36. Fuller WD, Orgel LE, Sanchez RA. Studies in Prebiotic Synthesis .7. Solid-State Synthesis of Purine Nucleosides. *J Mol Evol*. 1972; 1:249–257. [PubMed: 4681226]

Fig. 1. Summary of previous and present work.

Previous work demonstrated the synthesis of the canonical pyrimidine RNA nucleotides **9** and **10** alongside other RNA building blocks, such as 2-thiouridine **8**. This work (boxed) demonstrates a link between **8** and a range of DNA building blocks, *via* prebiotic phosphorylation and photoreduction. hv, UV irradiation; P_{i1} , $\text{NH}_4\text{H}_2\text{PO}_4$ /urea.

Fig. 2. 5' phosphorylation of 2-thiouridine **8 in concentrated formamide solution.**

Prebiotically plausible conversion of **8** into 2-thiouridine-5'-phosphate **14** in hot formamide potentially enables incorporation of **8** into primordial RNA sequences, by previously developed prebiotic activation of nucleoside phosphates and oligomerisation chemistry^{3,9}.

Fig. 3. Prebiotic and synthetic routes to (thio)anhydronucleosides.

a, Synthesis of the thioanhydronucleosides of uridine **15** and cytidine **21** under prebiotic phosphorylation conditions. 2-Thiouridine **8** and 2-thiocytidine **5** both undergo phosphorylation to form cyclophosphate intermediate **18** shown for the uridine variant. These intermediates undergo rearrangement to give structures of type **15** and **21** respectively. **15d** was not directly observed, but is inferred from the mechanism. In addition to rearrangement, the nucleosides undergo some cleavage of the glycosidic C–N bond to yield free nucleobases **19** and **23**, as well as their hydrolysis/ammonolysis products **20**, **22** and **24**, respectively. **b**, Hampton and Nichol's synthesis of 2',2-anhydrouridine **16** and our use of the same conditions to synthesize the sulfur analogue **15a** as a standard. **17a** and **17b** are the proposed intermediates leading to the formation of **16** and **15a**, respectively. P_2 , $\text{Na}_2\text{HPO}_4/\text{NH}_4\text{Cl}/\text{NH}_4\text{HCO}_3$.

Fig. 4. Phosphorylation of thionucleosides under mild conditions with DAP.

a, Phosphorylation of 2-thiouridine **8** with DAP leads to the formation of thioanhydrouridine-3'-phosphate **15b** and corresponding 3'-amidodiphosphate **25** *via* the intermediate cyclophosphate **18a**. The same phosphorylation with 4-thiouridine **26** results in the cyclophosphate **27**, which can undergo no further transformation. **b**, Phosphorylation of 2-thiocytidine **5** with DAP under slightly more forcing conditions affords thioanhydrocytidine-3'-phosphate **21b** and corresponding 3'-amidodiphosphate **29**, again *via* an intermediate, cyclophosphate **28**.

Fig. 5. Photoinduced reduction of thioanhydouridine with hydrosulfide (HS^-) and its proposed mechanism.

a. Photoreduction of **15a** gives the major product 2'-deoxy-2-thiouridine **11**, as well as 2-thiouracil **19**, which is possibly formed *via* facile nucleobase loss from proposed intermediate **30** (see ref. ²⁶). Nucleotides **15b** and **15c**—minor products of the phosphorylation of **8** in Fig. 3—undergo photoreduction under the same conditions, in similar yields (Supplementary Figs. 41 and 42). **b.** Our proposed mechanism is initiated by photodetachment of an electron from aqueous hydrosulfide to generate a solvated electron.

Reduction of **15a** by this electron generates radical anion **31**, which undergoes C–S bond homolysis to generate stabilised radical anion **32**. **32** can react with a hydrogen atom donor or combine with a HS• radical to give major isolated product **11** or proposed intermediate **30**, respectively, which we observed in crude spectra but were unable to isolate.

Fig. 6. Hydrolysis and transglycosylation reactions of 2'-deoxy-2-thiouridine 11.

a, Hydrolysis of **11** at different pH values gives deoxyribose **12** and the corresponding nucleobase, 2-thiouracil **19**. **b**, Transglycosylation in the dry state with adenine **33** gives a mixture of the α - and β - stereoisomers of the canonical deoxynucleoside, deoxyadenosine **13**, as well as nucleobase loss to yield **19**.