after stopping treatment) into account? Will there be an attempt to reach a consensus on how to limit coadministration of these antimicrobials or, like so much of CF management, will individual clinicians choose their own (diverse) paths? When new therapies come on line, there are always “early adopters” who embrace innovations and more conservative clinicians who are inclined to want to see additional data before they change their practice. Will these behaviors hold true for these data suggesting a clinically impactful aminoglycoside–macrolide antagonism? Will there be early adopters who will begin to avoid the combination immediately and others who will hold out for additional data?

These results have broader implications that warrant consideration. First, as the CF community comes to grips with the realization that *in vitro* antimicrobial susceptibility testing of patient bacterial isolates is not predictive of a clinical response to antimicrobial treatment (7), we can’t lose sight of an unassailable truth of antimicrobial treatment in CF: killing bacteria is an important (if not the sole) mechanism by which antimicrobials provide benefit in CF. In this situation, the same type of *in vitro* testing that performs poorly in predicting individual clinical responses provides a possible mechanism by which macrolides may be inhibiting clinically relevant aminoglycoside activity (4). These results suggest that decreased

sensitivity of infecting bacteria to antimicrobials matters clinically—an observation that is distinct from deficiencies in antimicrobial susceptibility testing as a predictor of clinical response in the individual. Second, it has never been truer that we are unlikely to recognize interactions we don’t look for: how many other such interactions are occurring in CF management today? Our best hope for learning the answer is to continue to conduct thoughtful retrospective analyses of existing single-center, registry, and clinical trial data, regardless of the level of evidence such studies provide. ■

Author disclosures are available with the text of this article at www.atsjournals.org.

References

- Somayaji R, Russell R, Cogen JD, Goss CH, Nick SE, Saavedra MT, *et al*. Oral azithromycin use and the recovery of lung function from pulmonary exacerbations treated with intravenous tobramycin or colistimethate in adults with cystic fibrosis. *Ann Am Thorac Soc* 2019;16:853–860.
- Klingel M, Stanojevic S, Tullis E, Ratjen F, Waters V. The effect of oral azithromycin on the response to pulmonary exacerbations treated with intravenous tobramycin in children with cystic fibrosis. *Ann Am Thorac Soc* 2019;16:861–867.
- Nick JA, Moskowitz SM, Chmiel JF, Forssén AV, Kim SH, Saavedra MT, *et al*. Azithromycin may antagonize inhaled tobramycin when targeting *Pseudomonas aeruginosa* in cystic fibrosis. *Ann Am Thorac Soc* 2014;11:342–350.
- Nichols DP, Happoldt CL, Bratcher PE, Caceres SM, Chmiel JF, Malcolm KC, *et al*. Impact of azithromycin on the clinical and antimicrobial effectiveness of tobramycin in the treatment of cystic fibrosis. *J Cyst Fibros* 2017;16:358–366.
- Zhao J, Schloss PD, Kalikin LM, Carmody LA, Foster BK, Petrosino JF, *et al*. Decade-long bacterial community dynamics in cystic fibrosis airways. *Proc Natl Acad Sci USA* 2012;109:5809–5814.
- Hoffman LR, Déziel E, D’Argenio DA, Lépine F, Emerson J, McNamara S, *et al*. Selection for *Staphylococcus aureus* small-colony variants due to growth in the presence of *Pseudomonas aeruginosa*. *Proc Natl Acad Sci USA* 2006;103:19890–19895.
- Somayaji R, Parkins MD, Shah A, Martiniano SL, Tunney MM, Kahle JS, *et al*. Antimicrobial Resistance in Cystic Fibrosis International Working Group. Antimicrobial susceptibility testing (AST) and associated clinical outcomes in individuals with cystic fibrosis: a systematic review. *J Cyst Fibros* 2019;18:233–240.

Copyright © 2019 by the American Thoracic Society

Preschool Wheezing Phenotypes Exhibit Heterogeneity in Disease Expression and Prognosis

Leonard B. Bacharier, M.D.¹, and Theresa W. Guilbert, M.D., M.S.^{2,3}

¹Robert C. Strunk Endowed Chair for Lung and Respiratory Research, Department of Pediatrics, Washington University School of Medicine in St. Louis, St. Louis, Missouri; ²Department of Pediatrics, University of Cincinnati, Cincinnati, Ohio; and ³Pulmonary Division, Cincinnati Children’s Hospital Medical Center, Cincinnati, Ohio

Wheezing during the first years of life is common; fortunately, only a minority of children with early-life wheezing will progress to having recurrent or persistent wheezing as well as asthma (1, 2). Children with preschool wheezing exhibit multiple

facets of disease heterogeneity, including multiple trajectories of disease expression and prognosis. Several phenotypes of early-life wheezing have been identified, and these phenotypes have distinct but often overlapping characteristics. Children may switch between phenotypes or evolve into another (1, 3). It is important to gain a better understanding of these phenotypes, as each phenotype may result from a different causal pathway (i.e., endotype) or have a specific pattern of response to specific

therapies, and to identify children who may benefit from specific disease-prevention strategies.

In this issue of *AnnalsATS*, Oksel and colleagues (pp. 868–876) conducted a substantial analysis of pooled data from five population-based cohorts of children who were followed from birth to adolescence (4). By pooling the data and following the subjects into adolescence, the authors were able to overcome the potential limitations of individual cohort analyses, which may differ

Supported by grants from the National Institutes of Health (L.B.B. and T.W.G.).

DOI: 10.1513/AnnalsATS.201903-234ED

in the temporal trajectories of disease, distribution of wheezing phenotypes, and timing and frequency of visits.

The strength of this study is that the authors performed several latent class analyses to describe the phenotypes using robust methodologies with high-quality and complete datasets. First, they performed an analysis of children with complete data from five visits at infancy, early childhood, preschool/early school age, middle childhood, and adolescence, involving 7,719 subjects. This analysis confirmed the phenotypes first described by Martinez and colleagues in the TCRS (Tucson Children's Respiratory Study) (1) (early-onset–middle-childhood remitting wheeze), which were later described in ALSPAC (Avon Longitudinal Study of Parents and Children). The confirmation of these two ALSPAC phenotypes is not unexpected, as ALSPAC represented the largest of the five cohorts included in this analysis. The school-age-onset remitting wheeze phenotype identified in ALSPAC was not confirmed, which may represent an artifact of pooling data across the cohorts into five clinical visits rather than the more frequent visits used in ALSPAC, and overall the remitting phenotypes were not as stable as the TCRS phenotypes.

The analysis was then repeated in a larger cohort that included children with two or more visits ($n = 15,941$), with similar results. Moreover, the authors conducted an analysis using four rather than five classes, and another using four cohorts while excluding the largest ALSPAC cohort, again with similar results. They also included the probability of each subject belonging to each phenotype in the analyses. This is a novel approach, and was possible because of the large size of the pooled cohort. These

results highlight the fact that significant within-class heterogeneity exists at the individual level, particularly in the late-onset wheeze group and early-onset preschool remitting wheeze groups. These children may not have fit easily into one of the five categories, and approximately 12% of the cohort did not fit neatly into one of these five categories. Not unsurprisingly, the persistent-wheeze group had the most consistent and stable class membership.

Compared with the never/infrequent-wheezing group, lower lung function was seen in all phenotypes, including episodic early-life wheezing, suggesting that early-life wheezing may not carry an entirely benign prognosis. This was also demonstrated in the TCRS cohort. If this low function is progressive over time, it may put these individuals at risk for chronic obstructive pulmonary disease as well as premature death. The persistent-wheezing and late-onset-wheezing groups exhibited the highest prevalence of current asthma in adolescence (adjusted odds ratio [adj OR], 23.02 and 22.25, respectively) and use of asthma medications (adj OR, 23.41 and 17.75, respectively).

Persistent wheeze, mid-childhood remitting wheeze, and late-onset wheeze were all significantly associated with allergic sensitization measured at one time point. Both the timing of allergic sensitization and the pattern of wheezing may be important, as demonstrated in a recent analysis by Bacharier and colleagues of a high-risk, inner-city cohort of 442 children who were followed between birth and 7 years of age, in which phenotypes were defined with the use of latent class analysis models (5). Allergic sensitization measured over several time points, frequency of wheezing episodes, and early-life environmental exposures were found to differentiate five early-onset wheezing phenotypes.

Key factors that identified the persistent-wheeze and late-onset groups were low lung function, a family history of asthma, maternal smoke exposure during pregnancy, and early allergic sensitization. It is notable that only smoke exposure is potentially modifiable. The early-onset–middle-childhood remitting and persistent wheeze groups were also associated with a history of eczema (adj OR, 1.42 and 2.47, respectively).

The study has a number of limitations, as acknowledged by the authors. All of the cohorts were based in the United Kingdom, and presumably the majority of the subjects were white individuals, so these findings may not hold true in more diverse populations; this requires additional study. These phenotypes are latent and thus not easily observable, and there is some controversy as to whether they should even be referred to as phenotypes. Thus, it is not possible to accurately identify the phenotype of an individual child in clinical practice. Several asthma prediction models have been studied which often include risk factors that were associated with the persistent-wheeze class in this analysis (sensitization, parental history of asthma, severity and frequency of wheezing, and maternal smoke exposure during pregnancy) (6–12). The late-onset group is another at-risk group for asthma management, but these individuals may present at an age too late for prevention.

Overall, the authors should be congratulated for their extensive analysis of five pooled cohorts, which has helped to better define the complexities underlying preschool-wheezing phenotypes. Clinically effective strategies are still needed to prospectively identify young children who will have the persistent-wheezing phenotype, which is associated with the highest prevalence of adolescent asthma, and these children would likely receive the most benefit from asthma therapy and prevention strategies. Two prevention trials in children at high risk for having a persistent-wheezing phenotype and subsequent asthma based on a personal or family history of atopic disease and/or recurrent wheezing are currently under way. One is a primary study testing the prevention of wheezing/lower-respiratory-tract illness using bacterial lyophilisates in high-risk infants and young children (ClinicalTrials.gov Identifier: NCT02148796). The other is a secondary prevention trial testing the hypothesis that anti-IgE therapy might decrease the risk for subsequent persistent asthma in aeroallergen-sensitized children with a history of recurrent wheezing (ClinicalTrials.gov Identifier: NCT02570984). ■

Author disclosures are available with the text of this article at www.atsjournals.org.

References

- Martinez FD, Wright AL, Taussig LM, Holberg CJ, Halonen M, Morgan WJ; The Group Health Medical Associates. Asthma and wheezing in the first six years of life. *N Engl J Med* 1995;332:133–138.
- Ly NP, Gold DR, Weiss ST, Celedón JC. Recurrent wheeze in early childhood and asthma among children at risk for atopy. *Pediatrics* 2006;117:e1132–e1138.
- Bacharier LB. The recurrently wheezing preschool child—benign or asthma in the making? *Ann Allergy Asthma Immunol* 2015;115:463–470.
- Oksel C, Granell R, Haider S, Fontanella S, Simpson A, Turner S, et al.; STELAR and Breathing Together investigators. Distinguishing wheezing phenotypes from infancy to adolescence: a pooled analysis of five birth cohorts. *Ann Am Thorac Soc* 2019;16:868–876.
- Bacharier LB, Beigelman A, Calatroni A, Jackson DJ, Gergen PJ, O'Connor GT, et al. Longitudinal phenotypes of respiratory health in a high-risk urban birth cohort. *Am J Respir Crit Care Med* 2019;199:71–82.
- Castro-Rodríguez JA, Holberg CJ, Wright AL, Martinez FD. A clinical index to define risk of asthma in young children with recurrent wheezing. *Am J Respir Crit Care Med* 2000;162:1403–1406.
- Chang TS, Lemanske RF Jr, Guilbert TW, Gern JE, Coen MH, Evans MD, et al. Evaluation of the modified asthma predictive index in high-risk preschool children. *J Allergy Clin Immunol Pract* 2013;1:152–156.
- Kurukulaaratchy RJ, Matthews S, Holgate ST, Arshad SH. Predicting persistent disease among children who wheeze during early life. *Eur Respir J* 2003;22:767–771.
- Hafkamp-de Groen E, Lingsma HF, Caudri D, Levie D, Wijga A, Koppelman GH, et al. Predicting asthma in preschool children with asthma-like symptoms: validating and updating the PIAMA risk score. *J Allergy Clin Immunol* 2013;132:1303–1310.
- van der Mark LB, van Wonderen KE, Mohrs J, van Aalderen WM, ter Riet G, Bindels PJ. Predicting asthma in preschool children at high risk presenting in primary care: development of a clinical asthma prediction score. *Prim Care Respir J* 2014;23:52–59.
- Biagini Myers JM, Schaubberger E, He H, Martin LJ, Kroner J, Hill GM, et al. A pediatric asthma risk score to better predict asthma development in young children. *J Allergy Clin Immunol* 2018;pii: S0091-6749(18)31577-X.
- Smit HA, Pinart M, Antó JM, Keil T, Bousquet J, Carlsen KH, et al. Childhood asthma prediction models: a systematic review. *Lancet Respir Med* 2015;3:973–984.

Copyright © 2019 by the American Thoracic Society

Communicating to Collaborate: Overlooked Requirements for Implementation Success

Erika L. Crable, M.P.H.^{1,2}, Mari-Lynn Drainoni, Ph.D., M.Ed.^{1,2,4,5}, and Allan J. Walkey, M.D., M.Sc.^{1,2,3*}

¹Evans Center for Implementation and Improvement Sciences, Department of Medicine, Boston University School of Medicine, Boston, Massachusetts; ²Department of Health Law, Policy and Management, Boston University School Public Health, Boston, Massachusetts; ³The Pulmonary Center, Department of Medicine, Boston University School of Medicine, Boston, Massachusetts; ⁴Section of Infectious Diseases, Department of Medicine, Boston University School of Medicine, Boston, Massachusetts; and ⁵Center for Healthcare Organization and Implementation Research, Edith Nourse Rogers Memorial VA Hospital, Bedford, Massachusetts

ORCID ID: 0000-0002-2605-5339 (E.L.C.).

Large gaps exist in the use of evidence-based practices in intensive care units (ICUs). For example, the ABCDEF bundle is used in fewer than 8% of all ICU days (1), and less than two-thirds of patients with acute respiratory distress syndrome receive low tidal volume ventilation (2). Better strategies are needed to increase knowledge transfer and evidence uptake in the ICU.

Understanding the barriers and facilitators to practice change is an essential step in the evaluation of evidence uptake and implementation (3). In this issue of the *AnnalsATS*, Johnson and colleagues (pp. 877–885) use qualitative methods to evaluate barriers and

facilitators in the process of evidence transfer from academic ICUs to partner community hospitals, offering new insights into implementation within understudied rural community hospital ICUs (4). Their study clearly shows the importance of effective communication and collaborative team environments as preconditions for implementation success.

In Johnson and colleagues' study, each community hospital partnered with an external team (an "innovations team") from an affiliated academic hospital to facilitate implementation of a quality improvement program aimed at enhancing uptake of evidence-based practices (4). Community hospitals identified "champions," local nurses and physicians, to promote staff buy-in and coordinate improvement efforts with the innovations team. Johnson and colleagues conducted thematic analysis of semistructured interviews with the community hospital champions and the external innovations

team that revealed that both inner ICU team dynamics (e.g., work culture, leadership styles, and staff readiness for change), as well as outer setting factors (e.g., interactions with the external innovations team, policy and professional guidelines), influenced implementation (4). Nine of the eleven themes that emerged from interviews with the participants in improvement initiatives emphasized effective communication and collaborative work cultures as intertwined elements required for implementation success. For example, two-way

*A.J.W. is Associate Editor of *AnnalsATS*. His participation complies with American Thoracic Society requirements for recusal from review and decisions for authored works.

DOI: 10.1513/AnnalsATS.201903-269ED