Published in final edited form as:

Psychiatr Serv. 2019 July 01; 70(7): 578–585. doi:10.1176/appi.ps.201800455.

Mental Health Status, Need, and Unmet Need for Mental Health Services among U.S. Pacific Islanders

Andrew M. Subica, Ph.D.^a, Nia Aitaoto, Ph.D.^b, Bruce G. Link, Ph.D.^c, Ann Marie Yamada, Ph.D.^d, Benjamin F. Henwood, Ph.D.^d, and Greer Sullivan, M.D., MSPH.^e

^aUniversity of California, Riverside Center for Healthy Communities, School of Medicine

^bUniversity of Utah, College of Health

^cUniversity of California, Riverside School of Public Policy

^dUniversity of Southern California, USC Suzanne Dworak-Peck School of Social Work

^eBorrego Health

Abstract

Objective: This study examined the mental health status, need for services, and unmet need of community-dwelling Native Hawaiian and other Pacific Islander (NHPI) adults.

Method: Survey data were collected from 223 NHPI adults of Samoan or Marshallese heritage. Surveys were translated into Samoan and Marshallese using back translation and cultural expert feedback. Participants' severity of depression, anxiety, and alcohol use, plus their perceived need for, and avoidance/delay of, mental health services were obtained. Logistic regressions calculated adjusted odds ratios for past-year: (1) perceived need for services, and (2) avoidance/delay of needed services, controlling for depression, anxiety, and alcohol use.

Results: Participants' screened prevalence of major depression, generalized anxiety disorder, and alcohol use disorder was 21%, 12%, and 22% respectively. Over the past year, 35% and 26% of all participants reported needing services and avoiding/delaying needed services, respectively. Urban Samoan and rural Marshallese participants did not significantly vary in mean depression, anxiety, or alcohol use despite demonstrating significant demographic differences. Female gender and greater familiarity/contact with persons with mental illness significantly predicted greater odds for needing services, and avoiding/delaying services.

Conclusion: Community-dwelling NHPI participants reported heavy burden of depression, anxiety, and alcohol use, and high perceived need for services yet low levels of help-seeking. The large unmet need found in the study sample suggests a gap may exist between service need and engagement in U.S. NHPI communities that could be targeted using culturally tailored approaches that promote engagement in care.

Introduction

Native Hawaiians and other Pacific Islanders (NHPI) comprise one of only seven federally designated U.S. racial/ethnic groups and are the third fastest growing in size (increasing 40.1% from 2000 to 2010).¹ Although over 1.4 million NHPIs reside in the U.S.¹ and over 10 million in the Pacific Islands,² little is known about NHPIs' health status (e.g., heart disease, cancer) and even less about their mental health—especially among NHPI adults—except they appear to be at high risk for problems such as depression, substance use, and suicide.^{3–9}

Despite this risk, extant data suggest that NHPIs are underrepresented in mental health services.^{3,10–14} For example, national admissions data indicate NHPIs and Asian Americans collectively seek substance use treatment at half the rate of other racial groups (5.3% vs. 10.4% for non-NHPIs and non-Asian Americans) and account for only 1% of admissions¹⁵ despite composing 7% of the U.S. population.¹ In Los Angeles County, which has a large, growing NHPI population of over 65,000,¹ few NHPIs have been reported to enter public treatment (0.09% in 2011, 0.07% in 2015),¹³ and only 4 NHPIs enrolled in a NHPI-specific County integrated care program over a two-year period.¹⁴ Due to NHPI adults' suspected resistance to mental health services and research, we know little about NHPI communities' mental health burden or their need for services.

NHPIs are a diverse racial population consisting of over 15 subgroups originating from the Pacific Islands (e.g., Hawaii, Samoa, Guam). These subgroups share cultural elements that emphasize collectivism, closeness to the natural world, reverence for the past via legends, chants, and storytelling shared over generations, and hierarchical social order led by chiefs or elders. Mirroring indigenous American Indians and Alaskan Natives, NHPI populations have been radically altered by U.S. colonialism. For instance, the U.S. overthrow of the Hawaiian Monarchy caused profound Native Hawaiian cultural loss and continued social, economic, and political marginalization in Hawaii,¹⁶ while the U.S. military's extensive nuclear testing in the Marshall Islands—equaling 7,200 Hiroshima-sized bombs¹⁷—contaminated generations of Marshallese people, lands, and food sources in radioactive fallout.^{17,18} Accordingly, NHPIs suffer from legacies of adverse colonization and historical traumatization that may contribute to heightened mental health risk.

With support from the National Institute of Mental Health, we conducted one of the first studies to quantify the level of mental health status and need among community NHPI adults; aiming to identify the key contributors of NHPIs' perceived need for, and avoidance/delay of, mental health services to better understand how to engage them in care. We surveyed urban-dwelling Samoans in Los Angeles County as they represent the largest NHPI community in the continental U.S. and rural Marshallese in Northwest Arkansas as they represent the fastest growing community. While all NHPI subgroups share cultural similarities, Samoans and Marshallese are also among the most divergent NHPI subgroups across numerous characteristics that could affect their mental health including immigration patterns/status, colonization experiences, and U.S. context (e.g., urban vs. rural, West vs. South region); rendering them ideal for comparison.

Conceptual Model of NHPI Mental Health Help-Seeking

Because underserved racial/ethnic minority populations are believed to seek help along different pathways than non-Hispanic whites due to their differing cultural factors/context,¹⁹ we drew from existing racial/ethnic minority help-seeking models by Cauce and colleagues¹⁹ and Eiraldi and colleagues²⁰ to create an NHPI-specific conceptual model of mental health help-seeking. Guiding our study, our conceptual model outlined several factors we predicted from existing models and help-seeking literature with non-NHPI populations, coupled with NHPI experts' feedback, may predict NHPIs' perceived need for, and avoidance/delay of services.

Our model posits that an NHPI individual's perceived need for services reflects his/her recognition that he/she has a mental health problem that requires treatment; a recognition labeled "problem recognition" in the help-seeking literature.^{19–21} In our model, NHPI problem recognition (characterized by our perceived need for services variable) may be influenced by a variety of factors including demographics (e.g., age, gender, formal school education),²¹ cultural factors (e.g., nativity and acculturation to the Pacific Islands or U.S.),^{19,22,23} presence and severity of mental health symptomology, and familiarity with NHPIs with mental health problems.^{19,23,24} Similarly, our model posits these same demographic, cultural, mental health, and familiarity variables will also influence an individual's decision to seek or avoid/delay help; a component of help-seeking distinct from problem recognition. It is important to note that this NHPI help-seeking model focuses on individual and not service system factors (e.g., lack of culturally responsive services), which are likely to interact with our individual-level factors to affect NHPI help-seeking.

This exploratory study's purpose is to characterize community NHPIs' mental health status and need for services. Drawing from existing literature indicating depression and anxiety are the most prevalent mental health problems and alcohol use disorder is the most common substance use problem affecting the U.S. general population,^{25,26} combined with the sparse NHPI mental health literature indicating depression, anxiety, and alcohol use may be problematic in NHPI communities,^{3,5,9} we anticipate screened prevalence of depression, anxiety, and alcohol use will be high in our study sample. But because NHPIs are underrepresented in treatment,^{13–15} we expect: (a) perceived need for mental health services will be low; and (b) many participants will report avoiding or delaying needed services.

Procedures

Procedures were approved by the Institutional Review Board at [redacted]. Data were collected from October 2017-January 2018 from community-dwelling Samoan and Marshallese adults age 18 years or older in Los Angeles County and Northwest Arkansas, respectively. To establish our sampling frame, because our populations were hard-to-reach and therefore probability sampling was not feasible, we systematically applied a non-probabilistic, respondent-driven sampling approach successfully used in prior mental health services research with hard-to-reach minority populations^{27,28} to collect our data. Specifically, to obtain as representative a community sample as possible, we trained community members in research who then purposively recruited participants of varying ages and genders—guided by a preset recruitment table—from diverse settings (e.g., churches of

different denominations, women's groups, cultural organizations for younger NHPIs). Participants were recruited in-person or through referral and completed informed consent and surveys in their preferred language (English, Samoan, Marshallese) via self-report or interview, receiving \$15 gift cards for their time.

Measures

For translation, measures were reviewed by NHPI experts for cultural content validity, translated from English to Samoan and Marshallese by experienced translators, and back-translated to English by additional translators blinded to the original measures. Original and back-translated measures were compared for discrepancies, which were resolved through consensus discussion between investigators, experts, and translators.

Past-year perceived need for mental health services and avoidance/delay of services were determined using two well-established items with strong reliability²⁶ from the Agency for Healthcare Research and Quality's national Medical Expenditure Panel Survey:²⁹ (a) "*In the past 12 months, was there a time when you wanted to talk with or seek help about stress, depression, or problems with emotions?*" and (b) "*Did you delay or not get the care you thought you needed?*"

Demographic and cultural factors of age, gender, formal education, nativity (Pacific Islands vs. U.S.), and acculturation were measured. U.S. nativity was defined as being raised in the U.S. since age four based on feedback from NHPI cultural experts. NHPI and U.S. acculturation levels were measured using the PIACCULT scale,³⁰ a 22-item scale—adapted from the General Ethnicity Questionnaire³¹—shown to associate with many NHPI social and health outcomes.^{32,33} Internal consistency was good for the 11-item NHPI-specific scale ($\alpha=.81$) and 11-item U.S.-specific scale ($\alpha=.83$).

Depression symptomology was evaluated via PHQ-9,³⁴ a 9-item tool with strong reliability and validity with diverse cultural groups³⁵ that assesses the past two-week presence of cardinal depression symptoms. Anxiety symptomology was captured via the GAD-7,³⁶ a well-validated 7-item measure of general anxiety symptoms experienced in the past two weeks. The PHQ-9 and GAD-7 are cumulatively scored with a 10-point screening diagnostic threshold for major depressive disorder and generalized anxiety disorder, respectively.^{37–39} Alcohol use was screened with the AUDIT-C,⁴⁰ a widely-used 3-item measure for identifying alcohol-related problems in adults. Scoring is cumulative with diagnostic cutoffs for alcohol use disorder of 4+ for men and 3+ for women. Internal consistencies for the PHQ-9, GAD-7, and AUDIT-C were strong with α 's of .86, .88, and .89, respectively.

Familiarity/contact with persons with mental illness was measured via the Level of Contact Report,^{41,42} which presents 12 situations of varying degrees of intimacy with people with severe mental illness (e.g., "*I have a relative with mental illness*"). Participants were rank scored (1–12) by the most intimate situation endorsed^{41,42} ranging from lowest ("*I have never observed a person that I was aware had a severe mental illness*"=1) to highest familiarity/contact ("*I have a severe mental illness*"=12).

Statistical Analyses

Analyses were performed in SPSS v.24.⁴³ Non-normal variables were logarithmically transformed. Descriptive statistics calculated sample frequencies, means, standard deviations, and 95% confidence intervals. Chi-square and independent *t* tests examined Samoan vs. Marshallese differences. Two logistic regression models were run. Dependent variables were past-year need for mental health services and avoidance/delay of needed services. Predictor variables for both models included: (a) age, gender, formal education, nativity, and acculturation, (b) severity of depression, anxiety, and alcohol use, and (c) level of familiarity/contact with persons with mental illness.

Results

Participant characteristics are presented in Table 1. Of 223 NHPI participants ($M_{\text{age}} = 40.9 \pm 16.0$ years), 57% were women and 55% were Samoan. Mean NHPI acculturation exceeded U.S. acculturation ($p < 0.05$). Mean familiarity/contact with persons with mental illness was 5.8 ± 3.7 —consistent with working with someone with severe mental illness.

Mean depression and anxiety severity were high and 21% and 12% of participants exceeded the 10-point diagnostic thresholds for major depression and generalized anxiety disorder, respectively—8% of participants exceeded both diagnostic thresholds, suggesting comorbid major depression and generalized anxiety disorder. For alcohol use, 22% of participants (29% of men, 18% of women) met diagnostic thresholds for alcohol use disorder; comorbidity with major depression and generalized anxiety disorder were 4% and 2%, respectively. No gender differences emerged for mean severity of depression and anxiety but men had significantly higher mean alcohol use than women ($M = 2.27 \pm 2.93$ vs. 1.18 ± 2.18 ; $t(217) = -0.27$, $p < 0.01$).

Over the past year, 35% and 26% of participants reported needing services and avoiding/delaying services, respectively. Fifty-nine percent of participants screening positive for depression, 54% for generalized anxiety disorder, and 35% for alcohol use disorder reported needing services.

Comparing Marshallese and Samoan subgroups, Marshallese participants were younger with significantly ($p < 0.05$) lower levels of: formal education (e.g., 26% had below high school formal education vs. 4% of Samoans); U.S. nativity (19% vs. 59%); and U.S. acculturation. No significant differences were found for mean depression, anxiety, or alcohol use but Marshallese participants had higher prevalence of needing past-year services ($p < 0.05$) and avoiding/delaying services ($p < 0.05$).

Logistic Regressions

To inform future NHPI service engagement approaches, logistic regressions explored which demographic, cultural, mental health, and other key variables accounted for participants' high past-year perceived need for, and avoidance/delay of, services.

Table 2 presents adjusted odds ratios (AOR) for past-year perceived need for services. For predicting past-year perceived need: (a) women had 4.4 times greater AOR than men

($p<0.01$), (b) completing high school and entering college both had 0.2 times lower AORs vs. not completing high school ($p<0.05$), (c) one-point increase in depression had 1.2 times greater AOR ($p<0.05$), and (d) one-point increase in mental illness familiarity/contact had 1.2 times greater AOR ($p<0.01$).

Table 3 presents adjusted odds ratios (AOR) for past-year avoidance/delay of services. For predicting past-year avoidance/delay of services: (a) women had 2.9 times greater AOR than men ($p<0.05$), (b) completing high school and completing college had 0.1 lower AORs vs. not completing high school ($p<0.05$), and (c) one-point increase in mental illness familiarity/contact had 1.3 times greater AOR ($p<0.01$).

Discussion

As hypothesized, although the presence of depression, anxiety, and alcohol use was high in our study, mental health help-seeking was relatively low. One of every four participants reported avoiding/delaying needed services in the past year; signaling a high degree of community unmet mental health need.

Unsurprisingly, more formal education associated with lower likelihood of avoiding/delaying needed services, suggesting formal education may be a useful tool for increasing help-seeking among NHPs. In contrast, female gender and greater familiarity/contact with persons with mental illness associated with greater likelihood for avoiding/delaying services. Accordingly, individuals demonstrating these characteristics may be at especially high risk for having unmet mental health need.

In particular, NHP women had far greater likelihood of avoiding/delaying services than men despite perceiving a greater need for services. Because many cultures (including NHPs) are more accepting of women than men expressing emotional distress and using emotion-focused coping,^{19,44} NHP women may be recognizing the need to seek help for their mental health problems (i.e., problem recognition) more readily than NHP men, but relying on the greater menu of informal coping options afforded NHP women (e.g., mobilizing social support, women's groups) in lieu of formal care.

Similarly, while increased contact with persons with mental illness associated with greater perceived need for services, it also associated with greater avoidance/delay of services. We suspect this seemingly contradictory finding may help to explain the push-and-pull pattern noted among our participants, who reported high perceived need but low help-seeking, by suggesting that gaining greater mental health awareness through exposure to persons with mental illness may be a double-edged sword for NHPs. That is, boosting mental health awareness may make NHP individuals more aware that their mental health problems require formal help,^{19,22,23} while also making more salient the risk of encountering stigma and discrimination^{45,46} associated with seeking help; thereby suppressing help-seeking. Given that increasing mental health awareness and contact is an important element in many anti-stigma interventions, further research is needed to clarify the role of mental health stigma and awareness on NHP help-seeking.

With regard to NHPs' mental health burden, while keeping in mind that we did not assess a probability sample, a large number of participants screened positive for major depression, generalized anxiety disorder, or alcohol use disorder. Compared to the U.S. adult general population, participants screened positive at three times the national major depression rate,⁴⁷ over two times the national generalized anxiety disorder rate,³⁷ and four times the national alcohol use disorder rate.^{48,49} Accordingly, our community NHP populations appear more comparable to clinical primary care populations³⁶ than to the general population.

Yet, not all disorders elicited help-seeking equally as relatively fewer NHPs who screened positive for alcohol use disorders perceived a need for services compared to NHPs experiencing major depression or generalized anxiety disorder. While this may in part be due to our perceived need item not specifically referencing alcohol use treatment, some of this discrepancy likely stems from NHP cultural norms that are generally accepting of risky drinking;^{50,51} possibly leading to lower perceived need by NHPs for what is considered problem alcohol use in U.S. populations. Unfortunately, because alcohol use disorders have many "downstream" effects including heightened risk for fatal and non-fatal accidents, domestic violence, and suicide,^{52–54} the implications of participants' high screened alcohol use disorder rates but low perceived need for services could have broad social and health consequences for U.S. NHPs.

Between our Samoan and Marshallese subgroups, no significant differences emerged in their levels of depression, anxiety, and alcohol use despite differences in many of the factors we anticipated might influence their mental health such as age, formal education, nativity, and acculturation. Findings of similar depression, anxiety, and alcohol use problems in these highly divergent NHP subgroups living in very different parts of the U.S support exploring the mental health needs of other NHP subgroups (e.g., Native Hawaiians, Chuukese) for commonalities that may inform the development of generalizable interventions for NHPs.

From our findings, we offer two thoughts for addressing NHPs' unmet need. First, the high screened rate of alcohol use disorders among NHP participants coupled with most participants with alcohol use disorders not believing they needed services underscores the importance of tailoring alcohol prevention interventions to NHPs so fewer services are needed. This may include interventions that combat NHP myths about alcohol use (e.g., education on normal vs. binge drinking)^{55,56} or train NHPs to resist alcohol offers from relatives and friends.^{57,58} Second, participants' observed resistance to seeking needed services suggests conventional engagement approaches may not work with community NHPs. Because the presence of many systems-level barriers (e.g., absence of interpreters) including a lack of NHP-tailored outreach and engagement likely creates an unwelcoming atmosphere for NHPs who wish to seek services—thereby contributing to limited NHP help-seeking—we recommend public-sector settings develop and implement culturally grounded engagement approaches for NHPs.^{59–61} This may be accomplished by first establishing relationships with NHP leaders/organizations to access community NHPs, then training lay NHPs as community health workers/peer navigators^{62–64} to deliver mental health education and anti-stigma interventions in community settings.

Limitations

As one of the first community studies of NHPI mental health, several limitations exist. First, using respondent-driven sampling to research two hard-to-reach populations may have introduced bias, potentially reducing generalizability of our findings to the overall Samoan, Marshallese, and NHPI populations. Second, we did not screen for all mental health problems that may be affecting NHPIs including substance use disorders and suicide. Third, we did not test all possible model relationships, including the potential relationship between perceived need for services and avoidance/delay of services. Finally, although we carefully prepared our measures for these specific populations, measurement error may exist in symptom under or overreporting. Yet, despite these limitations, we believe our data is compelling since so little is known about NHPI mental health, and their rates of problems in our data appear so high that even with some estimate errors, NHPIs' prevalence of mental health problems are likely to remain elevated relative to the general population.

Conclusions

Our findings reveal considerable mental health burden and high unmet need among individuals from two large NHPI communities. While developing culturally tailored engagement approaches may decrease this unmet need, as our exploratory study is among the first to rigorously quantify this burden within NHPI communities, more research is needed to identify best practices for engaging these underserved communities in care. This includes investigating other community-dwelling NHPI populations and exploring the major service barriers (e.g., stigma) that engagement interventions must address to reduce NHPI communities' heavy unmet need.

Disclosures and Acknowledgements:

The authors of this study declare no conflicts of interest. We would like to thank the staff from our community partners, the Office of Samoan Affairs and the Arkansas Coalition of the Marshallese, for their hard work and contributions to this project.

This project was supported by funding from the National Institutes of Health/National Institute of Mental Health (R21 MH110814). The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health or National Institute of Mental Health.

References

1. U.S. Census Bureau (2010). American Factfinder. Available from: <http://factfinder.census.gov/>.
2. PRISM. Pacific Island populations. Downloaded June 11, 2018 at: <https://prism.spc.int/regional-data-and-tools/population-statistics/169-pacific-island-populations>.
3. Asian Americans Advancing Justice, Orange County, & Asian Pacific Islander Community Alliance. (2014). A Community of Contrasts : Asian Americans, Native Hawaiians and Pacific Islanders in Orange County. Los Angeles, CA.
4. Crabbe KM (1998). Etiology of depression among Native Hawai'ians. *Pacific Health Dialog*, 5(2), 341–345.
5. Subica AM, & Wu LT (2018). Substance use and suicide in Pacific Islander, American Indian, and multiracial youth. *American Journal of Preventive Medicine*, 54(6), 795–805. [PubMed: 29656915]
6. Yuen NY, Nahulu LB, Hishinuma ES, Miyamoto RH (2000). Cultural identification and attempted suicide in Native Hawaiian adolescents. *Journal of the Academy of Child Adolescent Psychiatry*, 39, 360–367.

7. Bourke T (2001). Suicide in Samoa. *Pacific Health Dialog*, 8, 213–219. [PubMed: 12017827]
8. De Leo D, Milner A, Xiangdon W (2009). Suicidal behavior in the Western Pacific region: Characteristics and trends. *Suicide and Life-Threatening Behavior*, 39, 72–81. [PubMed: 19298152]
9. Ta VM, Chen TC (2008). Substance abuse among Native Hawaiian women in the United States: A review of current literature and recommendations for future research. *Journal of Psychoactive Drugs*, 40, 411–422.
10. Park VT, Kaholokula JK, Chao PJ, Antonio M (2018). Depression and help-seeking among Native Hawaiian women. *The Journal of Behavioral Health Services & Research*, 45, 454–468. [PubMed: 29280087]
11. U.S. Department of Health and Human Services. (2001). *Mental Health: Culture, Race, and Ethnicity (a Supplement to Mental Health: A Report of the Surgeon General)*. Rockville, MD: U.S. Department of Health and Human Services, Substance Abuse and Mental Health Services Administration, Center for Mental Health Services.
12. Centers for Disease Control and Prevention. (2011). *National Diabetes Fact Sheet, 2011*. Washington, D.C.: U.S. Department of Health and Human Services, Prevention CDC.
13. Los Angeles County Department of Public Health, Substance Abuse Prevention and Control. (2016). Los Angeles County Participant Reporting System Los Angeles, CA: Los Angeles County Department of Public Health Available from: <http://publichealth.lacounty.gov/sapc/bulletins/bulletin.html>
14. Los Angeles County Department of Mental Health, Health Services Research Center, Harder and Company, & University of Southern California School of Social Work. (2014). *County of Los Angeles MHSA Innovations Program Annual Report*. Los Angeles, CA: Los Angeles County Department of Mental Health.
15. Substance Abuse and Mental Health Services Administration, Office of Applied Studies. (2010). *Treatment Episode Data Set (TEDS) Report: Gender Differences among Asian and Pacific Islander Treatment Admissions Aged 18–25*. Rockville, MD: Substance Abuse and Mental Health Services Administration
16. Kaholokula JK (2007). *Colonialism, acculturation and depression among K naka Maoli of Hawai'i Penina Uliuli: Confronting challenges in mental health for Pacific Peoples*. Honolulu, HI: University of Hawaii Press.
17. Yamada S (2004). Cancer, Reproductive Abnormalities, and Diabetes in Micronesia: The Effect of Nuclear Testing. *Pacific Health Dialog*, 11, 216–221. [PubMed: 16281703]
18. McElfish PA, Hallgren E, & Yamada S (2015). Effect of US health policies on health care access for Marshallese Migrants. *American Journal of Public Health*, 105, 637–643. [PubMed: 25713965]
19. Cauce AM, Domenech-Rodríguez M, Paradise M, Cochran BN, Shea JM, Srebnik D, & Baydar N (2002). Cultural and contextual influences in mental health help seeking: A focus on ethnic minority youth. *Journal of Consulting and Clinical Psychology*, 70(1), 44. [PubMed: 11860055]
20. Eiraldi RB, Mazzuca LB, Clarke AT, & Power TJ (2006). Service utilization among ethnic minority children with ADHD: A model of help-seeking behavior. *Administration and Policy in Mental Health and Mental Health Services Research*, 33(5), 607–622. [PubMed: 16862391]
21. Kuhl J, Jarkon-Horlick L, & Morrissey RF (1997). Measuring barriers to help-seeking behavior in adolescents. *Journal of Youth and Adolescence*, 26(6), 637–650.
22. Alegria M, Robles R, Freeman DH, Vera M, Jimenez A, Rios C, & Rios R (1991). Patterns of mental health utilization among Puerto Rican poor. *American Journal of Public Health*, 81, 875–879. [PubMed: 2053664]
23. Pescosolido BA (1992). Beyond rational choice: The social dynamics of how people seek help. *American Journal of Sociology*, 97, 1096–1138.
24. Srebnik D, Cauce AM, & Baydar N (1996). Help-seeking pathways for children and adolescents. *Journal of Emotional and Behavioral Disorders*, 4, 210–220.
25. Kessler RC, Chiu WT, Demler O, & Walters EE (2005). Prevalence, severity, and comorbidity of 12-month DSM-IV disorders in the National Comorbidity Survey Replication. *Archives of General Psychiatry*, 62, 617–627. [PubMed: 15939839]

26. Kessler RC, McGonagle KA, Zhao S, Nelson CB, Hughes M, Eshleman S, ... & Kendler KS (1994). Lifetime and 12-month prevalence of DSM-III-R psychiatric disorders in the United States: results from the National Comorbidity Survey. *Archives of General Psychiatry*, 51, 8–19 [PubMed: 8279933]
27. Alvarez RA, Vasquez E, Mayorga CC, Feaster DJ, & Mitrani VB (2006). Increasing minority research participation through community organization outreach. *Western Journal of Nursing Research*, 28, 541–560. [PubMed: 16829637]
28. Breland-Noble AM, Board APAA. (2012). Community and treatment engagement of depressed African American youth: The Aakoma Floa pilot. *Journal of Clinical Psychology in Medical Settings*, 19, 41–48. [PubMed: 22354616]
29. Agency For Healthcare Research. Medical Expenditure Panel Survey Survey Questionnaires 2011 Available from: http://www.meps.ahrq.gov/mepsweb/survey_comp/survey.jsp#Questionnaires.
30. Borrows J, Williams M, Schluter P, Paterson J, & Langitoto Helu S (2011). Pacific Islands Families Study: The association of infant health risk indicators and acculturation of Pacific Island mothers living in New Zealand. *Journal of Cross-Cultural Psychology*, 42(5), 699–724.
31. Tsai JL, Ying Y, & Lee PA (2000). The meaning of “being Chinese” and “being American”: Variation among Chinese American young adults. *Journal of Cross-Cultural Psychology*, 31(3), 302–322.
32. Abbott M, & Williams M (2006). Postnatal depressive symptoms among Pacific mothers in Auckland: Prevalence and risk factors. *Australian and New Zealand Journal of Psychiatry*, 40(3), 230–238. [PubMed: 16476150]
33. Low P, Paterson J, Woulde S, Carter S, Williams M, & Percival T (2005). Factors affecting antenatal care attendance by mothers of Pacific infants living in New Zealand. *New Zealand Medical Journal*, 118, 1489.
34. Kroenke K, Spitzer RL, & Williams JB (2001). The PHQ-9: validity of a brief depression severity measure. *Journal of general internal medicine*, 16(9), 606–613. [PubMed: 11556941]
35. Huang FY, Chung H, Kroenke K, Delucchi KL, & Spitzer RL (2006). Using the patient health questionnaire-9 to measure depression among racially and ethnically diverse primary care patients. *Journal of general internal medicine*, 21(6), 547–552. [PubMed: 16808734]
36. Spitzer RL, Kroenke K, Williams JB, & Löwe B (2006). A brief measure for assessing generalized anxiety disorder: the generalized anxiety disorder-7. *Archives of internal medicine*, 166(10), 1092–1097. [PubMed: 16717171]
37. Löwe B, Decker O, Müller S, Brähler E, Schellberg D, Herzog W, & Herzberg PY (2008). Validation and standardization of the Generalized Anxiety Disorder Screener (generalized anxiety disorder-7) in the general population. *Medical care*, 266–274. [PubMed: 18388841]
38. Kroenke K, & Spitzer RL (2002). The PHQ-9: a new depression diagnostic and severity measure. *Psychiatric annals*, 32(9), 509–515.
39. Kroenke K, Spitzer RL, Williams JB, Monahan PO, & Löwe B (2007). Anxiety disorders in primary care: prevalence, impairment, comorbidity, and detection. *Annals of internal medicine*, 146(5), 317–325. [PubMed: 17339617]
40. Bush K, Kivlahan DR, McDonell MB, Fihn SD, Bradley KA. The Audit Alcohol Consumption Questions (Audit-C): An Effective Brief Screening Test for Problem Drinking. Ambulatory Care Quality Improvement Project (Acquip). Alcohol Use Disorders Identification Test. *Arch Intern Med*. 1998;158(16):1789–1795. [PubMed: 9738608]
41. Corrigan PW, Baks A, Green A, et al.: Prejudice, social distance, and familiarity with severe mental illness. *Schizophrenia Bulletin*, in press.
42. Holmes EP, Corrigan PW, Williams P, et al. (1999). Changing attitudes about schizophrenia. *Schizophrenia Bulletin*, 25, 447–456. [PubMed: 10478780]
43. Corp IBM. (2016). IBM SPSS version 24. Armonk, NY: IBM Corp.
44. Copeland EP, & Hess RS (1995). Differences in young adolescents' coping strategies based on gender and ethnicity. *The Journal of Early Adolescence*, 15(2), 203–219.
45. Lee S, Chiu MYL, Tsang A, Chiu H, & Kleinman A (2006). Stigmatizing experience and structural discrimination associated with the treatment of schizophrenia in Hong Kong. *Social Science & Medicine*, 62, 1685–1696. [PubMed: 16174547]

46. Yang HS, Kleinman A, Link BG, Phelan JC, Lee S, 7 Good B. (2007). Culture and stigma: Adding moral experience to stigma theory. *Social Science & Medicine*, 64, 1524–1535. [PubMed: 17188411]
47. Substance Abuse and Mental Health Services Administration. (2017). Key substance use and mental health indicators in the United States: Results from the 2016 National Survey on Drug Use and Health (HHS Publication No. SMA 17–5044, NSDUH Series H-52). Rockville, MD: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration Retrieved from <https://www.samhsa.gov/data/>
48. Substance Abuse and Mental Health Services Administration (SAMHSA). 2015 National Survey on Drug Use and Health (NSDUH). Table 5.6A—Substance Use Disorder in Past Year among Persons Aged 18 or Older, by Demographic Characteristics: Numbers in Thousands, 2014 and 2015. Available at: <https://www.samhsa.gov/data/sites/default/files/NSDUH-DetTabs-2015/NSDUHDetTabs-2015/NSDUH-DetTabs-2015.htm#tab5-6a>. Accessed 1/18/17.
49. Substance Abuse and Mental Health Services Administration (SAMHSA). 2015 National Survey on Drug Use and Health (NSDUH). Table 5.6B—Substance Use Disorder in Past Year among Persons Aged 18 or Older, by Demographic Characteristics: Percentages, 2014 and 2015. Available at: <https://www.samhsa.gov/data/sites/default/files/NSDUH-DetTabs-2015/NSDUHDetTabs-2015/NSDUH-DetTabs-2015.htm#tab5-6b>. Accessed 1/18/17.
50. Huakau J, Asiasiga L, Ford M, Pledger M, Casswell S, Suaalii-Sauni T, Lima I. (2005). New Zealand Pacific Peoples' drinking style: Too much or nothing at all? *New Zealand Medical Journal*, 118, 1216.
51. Ministry of Health. (2009). Alcohol use in New Zealand: Key results of the 2007/08 New Zealand Alcohol and Drug Use Survey. Wellington, NZ: Ministry of Health.
52. Klostermann KC, & Fals-Stewart W (2006). Intimate partner violence and alcohol use: Exploring the role of drinking in partner violence and its implications for intervention. *Aggression and Violent Behavior*, 11(6), 587–597.
53. Rehm J, Mathers C, Popova S, Thavorncharoensap M, Teerawattananon Y, & Patra J (2009). Global burden of disease and injury and economic cost attributable to alcohol use and alcohol-use disorders. *The Lancet*, 373(9682), 2223–2233.
54. Wilcox HC, Conner KR, & Caine ED (2004). Association of alcohol and drug use disorders and completed suicide: an empirical review of cohort studies. *Drug and Alcohol Dependence*, 76, S11–S19. [PubMed: 15555812]
55. Santisteban DA, Coatsworth JD, Perez-Vidal A, Kurtines WM, Schwartz SJ, LaPerriere A, & Szapocznik J (2003). Efficacy of brief strategic family therapy in modifying Hispanic adolescent behavior problems and substance use. *Journal of Family Psychology*, 17(1), 121. [PubMed: 12666468]
56. Coatsworth JD, Pantin H, & Szapocznik J (2002). Familias Unidas: A family-centered ecodevelopmental intervention to reduce risk for problem behavior among Hispanic adolescents. *Clinical Child and Family Psychology Review*, 5(2), 113–132. [PubMed: 12093012]
57. Helm S, Okamoto SK, Maddock J, Hayes D, Lowery T, & Rajan R (2013). Insights in public health: Developing the Ho'ouana Pono substance use prevention curriculum: Collaborating with Hawaiian Youth and communities. *Hawaiian Journal of Medicine and Public Health*, 72, 66–69.
58. Okamoto SK, Kulis S, Helm S, Lauricella M, & Valdez JK (2016) An evaluation of the Ho'ouana Pono curriculum: A pilot study of culturally grounded substance abuse prevention for rural Hawaiian Youth. *Journal of Health Care for the Poor and Underserved*, 27, 815–833. [PubMed: 27180710]
59. Goebert D, Alvarez A, Andrade NN, Balberde-Kamalii J, Carlton BS, Chock S, ... Sugimoto-Matsuda J. (2018). Hope, help, and healing: Culturally embedded approaches to suicide prevention, intervention and postvention services with native Hawaiian youth. *Psychological Services*, 15, 332–339. [PubMed: 30080092]
60. Okamoto SK, Kulis S, Marsiglia FF, Steiker LKH, & Dustman P (2014). A continuum of approaches toward developing culturally focused prevention interventions: From adaptation to grounding. *The Journal of Primary Prevention*, 35, 103–112 [PubMed: 24322970]

61. Okamoto SK, Helm S, Pel S, McClain LL, Hill AP, Hayashida JK (2014). Developing empirically based, culturally grounded drug prevention interventions for indigenous youth populations. *The Journal of Behavioral Health Services & Research*, 41, 8–19. [PubMed: 23188485]
62. Kelly E, Fulginiti A, Pahwa R, Tallen L, Duan L, & Brekke JS (2014). A pilot test of a peer navigator intervention for improving the health of individuals with serious mental illness. *Community Mental Health Journal*, 50(4), 435–446. [PubMed: 23744292]
63. Witmer A, Seifer SD, Finocchio L, Leslie J, & O'Neil EH (1995). Community health workers: integral members of the health care work force. *American Journal of Public Health*, 85, 1055–1058. [PubMed: 7625495]
64. Rosenthal EL, Brownstein JN, Rush CH, Hirsch GR, Willaert AM, Scott JR, ... & Fox DJ (2010). Community health workers: part of the solution. *Health Affairs*, 29(7), 1338–1342. [PubMed: 20606185]

Highlights:

- Little is known about mental health in Pacific Islander communities.
- Pacific Islander participants reported high levels of depression, anxiety, and alcohol use.
- 36% of participants needed services yet 26% delayed the needed services.

Table 1:

Characteristics and descriptive statistics for NHPI participant sample

Variables	Total Sample N = 223			Samoan n = 123			Marshallse n = 100			χ^2 or t between Samoans and Marshallse	df	N	p value
	%	n	SD	%	n	SD	%	n	SD				
Gender (Female)	57	127		62	76		52	51					
Formal education													
< High school	14	30		4	5		26	25					
High school	44	95		40	49		48	46					
Some college	32	69		39	47		23	22					
College	11	24		17	21		3	3					
Nativity (U.S.)	41	82		59*	65		18*	17		$\chi^2 = 34.27$	1	202	$p < .05$
Needed Care in Past Year	35	78		29*	36		43*	42		$\chi^2 = 4.85$	1	221	$p < .05$
Avoided/Delayed Care	26	56		20*	24		33*	32		$\chi^2 = 4.88$	1	218	$p < .05$
	Mean	SD		Mean	SD		Mean	SD					
Age	40.87	16.01		44.66*	16.68		36.30*	13.93		$t = 3.95$	214		$p < .01$
Acculturation													
Pacific Islander	46.25	5.89		46.55	5.55		45.84	6.34					
United States	40.30	7.19		42.82*	5.63		37.02*	7.69		$t = 6.40$	214		$p < .01$
Depression (PHQ-9)	5.87	5.10		5.20	4.93		6.72	5.21					
General Anxiety (GAD-7)	4.72	4.38		4.16*	3.92		5.44*	4.84		$t = -2.16$	216		$p < .01$
Alcohol (AUDIT-C)	1.64	2.58		1.81	2.53		1.43	2.64					
Level of Contact	5.83	3.68		5.27	3.69		6.53	3.57					

* Significant difference at .05 level between Samoan and Marshallse participants

PHQ-9 = Patient Health Questionnaire – 9; GAD-7 = Generalized Anxiety Disorder – 7; AUDIT-C = Alcohol Use Disorders Identification Test- Concise

Table 2:

Logistic regressions for perceived need for mental health services in past year

Variables	AOR	95% CI
DV = Perceived Need for Mental Health Services		
Demographic		
Age	0.66	0.43–1.01
Gender (reference: Male)		
Female	4.35 **	1.73–10.93 **
Formal education (reference: < High school)		
High school	0.22 *	0.06–0.78 *
Some college	0.17 *	0.04–0.76 *
College	0.32	0.06–1.59
Cultural		
Nativity (reference: U.S.)		
Pacific Islands	0.53	0.18–1.58
Acculturation		
Pacific Islander	1.03	0.95–1.11
United States	0.99	0.95–1.11
Symptoms		
Depression (PHQ-9)	1.18 **	1.05–1.32 **
General Anxiety (GAD-7)	0.92	0.80–1.04
Alcohol (AUDIT-C)	1.07	0.90–1.28
Familiarity with Mental Illness		
Level of Contact with Mental Illness	1.23 **	1.09–1.39 **

* $p < .05$;** $p < .01$

AOR = Adjusted Odds Ratio; CI = Confidence Interval; DV = Dependent Variable; PHQ-9 = Patient Health Questionnaire – 9; GAD-7 = Generalized Anxiety Disorder – 7; AUDIT-C = Alcohol Use Disorders Identification Test – Concise

Table 3:

Logistic regression for avoiding/delaying needed mental health services in past year

Variables	AOR	95% CI
DV = Avoided/Delayed Needed Services		
Demographic		
Age	0.86	0.54–1.39
Gender (reference: Male)		
Female	2.89 *	1.03–8.06 *
Formal education (reference: < High school)		
High school	0.14 **	0.04–0.55 **
Some college	0.22	0.43–1.07
College	0.14 *	0.02–1.01 *
Cultural		
Nativity (reference: U.S.)		
Pacific Islands	3.65	0.96–13.84
Acculturation		
Pacific Islander	1.05	0.96–1.15
United States	1.02	0.94–1.10
Symptoms		
Depression (PHQ-9)	1.11	0.99–1.25
General Anxiety (GAD-7)	1.02	0.89–1.17
Alcohol (AUDIT-C)	1.17	0.95–1.43
Familiarity with Mental Illness		
Level of Contact with Mental Illness	1.30 **	1.12–1.50 **

*
 $p < .05$;**
 $p < .01$

AOR = Adjusted Odds Ratio; CI = Confidence Interval; DV = Dependent Variable; PHQ-9 = Patient Health Questionnaire – 9; GAD-7 = Generalized Anxiety Disorder – 7; AUDIT-C = Alcohol Use Disorders Identification Test – Concise