in situ Smad3 expression in the lungs of patients with idiopathic PAH and control subjects, PAs were laser microdissected, followed by qPCR. Smad3 mRNA (trend, statistically not significant) and miR-130a/301b ($P < 0.01$) expression was higher in the PAs and plexiform lesions of patients with idiopathic PAH compared with control subjects (Figure 1E). Thus, our human data support the notion that boosted signaling of TGF- β 1 and its canonical downstream effector Smad3 in human PAs is crucial for PAH development and severity.

We designed this focused study to address the controversy regarding TGF- β 1-mediated Smad3 signaling in PAH. We show that chronic TGF- β 1 signaling *in vivo* leads to spontaneous PAH in mice; is associated with canonical Smad3 activation (Smad3 phosphorylation) but not Smad3 downregulation; and drives PAH by means of proliferation, inflammation, and metabolism in the lung. Our results are supported by several other important publications (2, 6–8, 10). In contrast to the study by Zabini and colleagues (9), we cannot confirm a loss of Smad3 with chronic TGF- β 1 signaling in our PAH rodent models (TG-TGF- β 1 mouse and SuHx rat). Zabini and colleagues focused only on total Smad3 expression and did not investigate Smad3 phosphorylation, which is the hallmark of canonical TGF- β 1 activation. It is certainly possible that in PAH, downregulation of Smad3 mRNA and/or protein expression occurs in pulmonary arterial endothelial cells or other non-SMCs, such as pericytes and fibroblasts; however, experimental evidence is currently lacking, and resolving this issue would require additional studies.

A better understanding of the functional antagonism of TGF- β 1 and BMP2 (2) (and other BMPs) and their downstream pathways is crucial for our understanding of PAH. Emerging discoveries in this field will guide the future development of more efficient therapies for PAH, which currently is incurable. Taken together, our results indicate that chronic TGF- β 1 signaling in PAH induces sustained canonical Smad3 signaling in PASMCs, and that such signals correlate with the hemodynamic and morphological PAH phenotype in rodents. ■

Author disclosures are available with the text of this letter at www.atsjournals.org.

Acknowledgment: The authors thank Jonas Geldner for technical assistance.

Laurent Calvier, Ph.D.
Philippe Chouvarine, Ph.D.
Ekaterina Legchenko, Ph.D.
Department of Pediatric Cardiology and Critical Care
Hannover Medical School
Hannover, Germany

Gabor Kokeny, M.D., Ph.D.
Miklos M. Mozes, M.D., Ph.D.
Department of Pathophysiology
Semmelweis University
Budapest, Hungary

Georg Hansmann, M.D., Ph.D.*
Department of Pediatric Cardiology and Critical Care
Hannover Medical School
Hannover, Germany

ORCID ID: 0000-0003-0709-3935 (G.H.).

*Corresponding author (e-mail: georg.hansmann@gmail.com).

References

- Alarcón C, Zaromytidou AI, Xi Q, Gao S, Yu J, Fujisawa S, *et al*. Nuclear CDKs drive Smad transcriptional activation and turnover in BMP and TGF- β pathways. *Cell* 2009;139:757–769.
- Calvier L, Chouvarine P, Legchenko E, Hoffmann N, Geldner J, Borchert P, *et al*. PPAR γ links BMP2 and TGF β 1 pathways in vascular smooth muscle cells, regulating cell proliferation and glucose metabolism. *Cell Metab* 2017;25:1118–1134.e7.
- Akhurst RJ, Hata A. Targeting the TGF β signalling pathway in disease. *Nat Rev Drug Discov* 2012;11:790–811.
- Morrell NW, Bloch DB, ten Dijke P, Goumans M-JTH, Hata A, Smith J, *et al*. Targeting BMP signalling in cardiovascular disease and anaemia. *Nat Rev Cardiol* 2016;13:106–120.
- Calvier L, Miana M, Reboul P, Cachofeiro V, Martinez-Martinez E, de Boer RA, *et al*. Galectin-3 mediates aldosterone-induced vascular fibrosis. *Arterioscler Thromb Vasc Biol* 2013;33:67–75.
- Zakrzewicz A, Kouri FM, Nejman B, Kwapiszewska G, Hecker M, Sandu R, *et al*. The transforming growth factor- β /Smad2,3 signalling axis is impaired in experimental pulmonary hypertension. *Eur Respir J* 2007;29:1094–1104.
- Zhang N, Dong M, Luo Y, Zhao F, Li Y. Danshensu prevents hypoxic pulmonary hypertension in rats by inhibiting the proliferation of pulmonary artery smooth muscle cells via TGF- β -smad3-associated pathway. *Eur J Pharmacol* 2018;820:1–7.
- Graham BB, Chabon J, Gebreab L, Poole J, Debella E, Davis L, *et al*. Transforming growth factor- β signaling promotes pulmonary hypertension caused by *Schistosoma mansoni*. *Circulation* 2013;128:1354–1364.
- Zabini D, Granton E, Hu Y, Miranda MZ, Weichert U, Breuils Bonnet S, *et al*. Loss of SMAD3 promotes vascular remodeling in pulmonary arterial hypertension via MRTF disinhibition. *Am J Respir Crit Care Med* 2018;197:244–260.
- Legchenko E, Chouvarine P, Borchert P, Fernandez-Gonzalez A, Snay E, Meier M, *et al*. PPAR γ agonist pioglitazone reverses pulmonary hypertension and prevents right heart failure via fatty acid oxidation. *Sci Transl Med* 2018;10:pil:ea00303.

Copyright © 2019 by the American Thoracic Society

IL-6Ra in Smooth Muscle Cells Protects against *Schistosoma*- and Hypoxia-induced Pulmonary Hypertension

To the Editor:

The role of IL-6 signaling in pulmonary hypertension (PH) is generally believed to be deleterious. Patients with idiopathic and chronic obstructive pulmonary disease-associated PH have

Supported by American Heart Association grant 17POST33670045 (R.K.), the American Thoracic Society Foundation/Pulmonary Hypertension Association, and National Institutes of Health grants P01HL014985 (R.M.T. and B.B.G.), R03HL133306 (B.B.G.), and R01HL135872 (B.B.G.). Schistosome-infected mice were provided by the National Institute of Allergy and Infectious Diseases Schistosomiasis Resource Center at the Biomedical Research Institute (Rockville, MD) through National Institute of Allergy and Infectious Diseases/National Institutes of Health contract HHSN272201000005I for distribution through BEI Resources.

This letter has a data supplement, which is accessible from this issue's table of contents at www.atsjournals.org.

increased serum IL-6 (1–3). In rodents, overexpression of IL-6 induces spontaneous PH, which worsens with additional hypoxia (4). Ablation of IL-6 or the IL-6 receptor protects against hypoxia-induced PH in the mouse (5, 6), and pharmacological blockade is beneficial in rat models of PH (6). However, IL-6 may also be protective in the context of inflammation-triggered diseases, including schistosomiasis-associated PH and systemic vascular diseases (7–9). There is a currently ongoing trial in subjects with PH of tocilizumab, which blocks the IL-6 receptor (10). We sought to determine the role of IL-6 signaling in pulmonary smooth muscle cells in schistosomiasis- and hypoxia-induced PH using compartment-specific deletion of a critical receptor component, IL-6Ra (or CD126).

Methods

We purchased *Il6ra*^{flox/flox} and *Myh11*-CreER^{T2} mice (The Jackson Laboratory), and crossed the lines to generate *Il6ra*^{flox/flox}; *Myh11*-CreER^{T2} mice. All animal experiments were approved by the University of Colorado Institutional Animal Care and Use Committee. For further details, see the data supplement.

Results

We treated *Il6ra*^{flox/flox}; *Myh11*-CreER^{T2} mice with tamoxifen and quantified the deletion efficiency by qPCR of smooth muscle cells obtained from dissected and digested pulmonary vessels grown on smooth muscle-selective media (>90% of the cells were α -SMA⁺). We found that on average, 59% of the *Il6ra* DNA was deleted. We challenged tamoxifen-treated *Il6ra*^{flox/flox}; *Myh11*-CreER^{T2} mice with either *Schistosoma* or hypoxia, and found that the PH severity as measured by right ventricle systolic pressure (RVSP) was worse with either stimuli as compared with either stimuli after corn oil administration alone (Figure 1A). We corroborated the finding of more severe PH in tamoxifen-treated mice by observing increased pulmonary vascular media thickness as assessed by immunostaining for either α -smooth muscle actin (α -SMA) (Figure 1B) or smooth muscle myosin heavy chain (Figure E1 in the data supplement) in *Schistosoma*-challenged mice that had received tamoxifen.

We recently observed that thrombospondin-1 (TSP-1) activation of transforming growth factor β is critical for *Schistosoma*- and hypoxia-induced PH (11). TSP-1 can be regulated by IL-6 (12). We determined that mice with the IL-6Ra deletion had an increase in TSP-1 expression (Figure E2), potentially contributing to the increase in PH severity observed.

We noted that corn oil-treated *Il6ra*^{flox/flox}; *Myh11*-CreER^{T2} mice challenged with *Schistosoma* did not develop increased RVSP (Figure 1A), as previously documented in wild-type mice (7, 11, 13). We observed that *Myh11*-CreER^{T2} mice treated with corn oil developed increased RVSP upon exposure to either *Schistosoma* or hypoxia (Figure 1C). However, otherwise untreated *Il6ra*^{flox/flox} mice developed increased RVSP only upon exposure to hypoxia, and not to *Schistosoma* (Figure 1D), which is suggestive of a genetic abnormality in this line that suppresses type 2 immune response-driven PH, but remained dominant in the *Il6ra*^{flox/flox}; *Myh11*-CreER^{T2} phenotype. We quantified the concentration of IL-13 in *Schistosoma*-exposed *Il6ra*^{flox/flox}; *Myh11*-CreER^{T2} mice (Figure E3) and observed that it was less than one-third of that previously found in *Schistosoma*-exposed wild-type mice (13). An alternative potential confounder is that corn oil alone could induce PH, but we found no

change in RVSP in *Myh11*-CreER^{T2} mice treated with corn oil (Figure E4). Another potential explanation is that tamoxifen could induce PH, but we found no change in RVSP in *Il6ra*^{flox/flox} mice treated with tamoxifen (Figure E5).

Discussion

IL-6 can be released by multiple cell types, including endothelial cells, smooth muscle cells, fibroblasts, and leukocytes. IL-6 has two downstream signaling pathways that signal via hexameric formation with the ligand and two receptor proteins, IL-6Ra and gp130 (14). gp130 is ubiquitously expressed and membrane bound. However, IL-6Ra can be either membrane bound or alternatively spliced and released extracellularly. In the classical signaling pathway (*cis*), IL-6Ra is membrane bound, and in the *trans* signaling pathway, IL-6Ra is entirely extracellular. Both signaling pathways induce expression of downstream targets via the Janus tyrosine kinase–signal transducer and activator of transcription 3 pathway. Recently, it was reported that both *cis* and *trans* IL-6 signaling is increased in pulmonary vascular smooth muscle cells in patients and animals with PH (6).

Here, we observed that *Il6ra* deletion in smooth muscle cells resulted in increased severity of both *Schistosoma*- and hypoxia-induced PH. This phenotype is most likely mediated by IL-6Ra *cis* signaling in the vascular smooth muscle compartment, as persistent IL-6Ra expression by other cells likely compensates for IL-6Ra *trans* signaling, although inadequate access to the smooth muscle cells could also suppress *trans* signaling. *Myh11* expression is specific for smooth muscle cells, which aside from vascular smooth muscle cells include airway, intestine, and bladder smooth muscle cells. *Il6ra* deletion in these cells specifically avoids altered IL-6Ra expression by other cells that express it, including osteoclasts, T cells, fibroblasts, and macrophages. We did not observe a change in the phenotype of airway smooth muscle cells by histology (Figure E6), although altered IL-6 signaling in these cells could affect the nearby vascular phenotype.

These results are in line with our prior observation that *Il6*^{−/−} mice had increased PH severity upon exposure to *Schistosoma* (7). However, the observed deleterious effect of IL-6Ra deletion in hypoxia-induced PH contradicts the results that were recently published by Tamura and colleagues, who reported that *Il6ra*^{flox/flox}; Sm22a-Cre mice were significantly protected from hypoxia-induced PH (6). There are several potential reasons for the difference in results between these two genotypes, including possible developmental abnormalities with germline deletion; different cell compartment expression (for example, Sm22a, encoded by *Tagln*, is also expressed in fibroblasts [15]); different efficiencies of deletion (59% in our model; the deletion efficiency was not reported by Tamura and colleagues) driving a nonlinear dose effect, with harm at moderate deletion and protection at potentially greater deletion; different environments (our experiments were performed at Denver elevation); and sex differences (we used male mice; the sex of the mice was not reported by Tamura and colleagues).

In summary, we observed that inducible deletion of *Il6ra* in the smooth muscle cells of mice resulted in increased severity of hypoxia- and *Schistosoma*-induced PH. There may be specific contexts in which IL-6 signaling is deleterious or protective in PH. ■

Figure 1. Effect of *Il6ra* deletion in smooth muscle cells on *Schistosoma*-induced pulmonary hypertension (Schisto-PH) and hypoxia-induced PH (hypoxia-PH). (A) Right ventricle systolic pressure (RVSP) in control, *Schistosoma*-challenged and hypoxia-challenged *Il6ra^{fllox/fllox};Myh11-CreER^{T2}* mice ($n=5-10$ /group; two-way ANOVA: $P<0.0001$ for both PH stimulus and tamoxifen treatment effects; *post hoc* Tukey tests shown only for tamoxifen treatment effect). (B) Representative images of pulmonary arteries stained for α -smooth muscle actin (α -SMA) (green: α -SMA; blue: DAPI; scale bars: 50 μ m) and quantitative fractional media thickness in control and *Schistosoma*-challenged *Il6ra^{fllox/fllox};Myh11-CreER^{T2}* mice ($n=6-10$ /group; two-way ANOVA: $P<0.0001$ for PH stimulus effect and $P<0.05$ for tamoxifen treatment effect; *post hoc* Tukey tests shown only for tamoxifen treatment effect [all *post hoc* testing results are shown in Figure E7]; 61 mean [range 44–91] number of arteries analyzed per group). (C) RVSP in control, *Schistosoma*-challenged and hypoxia-challenged *Myh11-CreER^{T2}* mice ($n=5-7$ /group; one-way ANOVA: $P<0.0005$; *post hoc* Tukey test shown). (D) RVSP in control, *Schistosoma*-challenged and hypoxia-challenged *Il6ra^{fllox/fllox}* mice ($n=4-6$ /group; one-way ANOVA: $P<0.0001$; *post hoc* Tukey test shown). Mean $\pm$ SEM is plotted (* $P<0.05$ and **** $P<0.001$).

Author disclosures are available with the text of this letter at www.atsjournals.org.

Claudia Mickael, Ph.D.
 Rahul Kumar, Ph.D.
 Daniel Hernandez-Saavedra, Ph.D.
 Biruk Kassa, B.S.
 Linda Sanders, B.S.
 Dan Koyanagi, B.S.
 Sue Gu, M.D.
 Michael H. Lee, M.D.
 Rubin M. Tuder, M.D.
 Brian B. Graham, M.D.*
 University of Colorado Anschutz Medical Campus
 Aurora, Colorado

ORCID ID: 0000-0001-7541-2585 (B.B.G.).

*Corresponding author (e-mail: brian.graham@ucdenver.edu).

References

1. Chaouat A, Savale L, Chouaid C, Tu L, Sztrymf B, Canuet M, *et al*. Role for interleukin-6 in COPD-related pulmonary hypertension. *Chest* 2009;136:678–687.
2. Humbert M, Monti G, Brenot F, Sitbon O, Portier A, Grangeot-Keros L, *et al*. Increased interleukin-1 and interleukin-6 serum concentrations in severe primary pulmonary hypertension. *Am J Respir Crit Care Med* 1995;151:1628–1631.
3. Jasiewicz M, Knapp M, Waszkiewicz E, Ptaszynska-Kopczynska K, Szpakowicz A, Sobkowicz B, *et al*. Enhanced IL-6 trans-signaling in pulmonary arterial hypertension and its potential role in disease-related systemic damage. *Cytokine* 2015;76:187–192.
4. Steiner MK, Syrkina OL, Kolliputi N, Mark EJ, Hales CA, Waxman AB. Interleukin-6 overexpression induces pulmonary hypertension. *Circ Res* 2009;104:236–244.
5. Savale L, Tu L, Rideau D, Izziki M, Maitre B, Adnot S, *et al*. Impact of interleukin-6 on hypoxia-induced pulmonary hypertension and lung inflammation in mice. *Respir Res* 2009;10:6.
6. Tamura Y, Phan C, Tu L, Le Hiress M, Thuillet R, Jutant EM, *et al*. Ectopic upregulation of membrane-bound IL6R drives vascular remodeling in pulmonary arterial hypertension. *J Clin Invest* 2018;128:1956–1970.
7. Graham BB, Chabon J, Kumar R, Kolosionek E, Gebreab L, Debella E, *et al*. Protective role of IL-6 in vascular remodeling in Schistosoma pulmonary hypertension. *Am J Respir Cell Mol Biol* 2013;49:951–959.
8. Madan M, Bishayi B, Hoge M, Amar S. Atheroprotective role of interleukin-6 in diet- and/or pathogen-associated atherosclerosis using an ApoE heterozygote murine model. *Atherosclerosis* 2008;197:504–514.
9. Fisman EZ, Tenenbaum A. The ubiquitous interleukin-6: a time for reappraisal. *Cardiovasc Diabetol* 2010;9:62.
10. Hernández-Sánchez J, Harlow L, Church C, Gaine S, Knightbridge E, Bunclark K, *et al*. Clinical trial protocol for TRANSFORM-UK: a therapeutic open-label study of tocilizumab in the treatment of pulmonary arterial hypertension. *Pulm Circ* 2018;8:2045893217735820.
11. Kumar R, Mickael C, Kassa B, Gebreab L, Robinson JC, Koyanagi DE, *et al*. TGF- β activation by bone marrow-derived thrombospondin-1 causes Schistosoma- and hypoxia-induced pulmonary hypertension. *Nat Commun* 2017;8:15494.
12. Kim SA, Kang JH, Cho I, Bae SW, Hong KJ. Cell-type specific regulation of thrombospondin-1 expression and its promoter activity by regulatory agents. *Exp Mol Med* 2001;33:117–123.
13. Kumar R, Mickael C, Chabon J, Gebreab L, Rutebemberwa A, Garcia AR, *et al*. The causal role of IL-4 and IL-13 in Schistosoma mansoni pulmonary hypertension. *Am J Respir Crit Care Med* 2015;192:998–1008.
14. Lacroix M, Rousseau F, Guilhot F, Malinge P, Magistrelli G, Herren S, *et al*. Novel insights into interleukin 6 (IL-6) Cis- and trans-signaling pathways by differentially manipulating the assembly of the IL-6 signaling complex. *J Biol Chem* 2015;290:26943–26953.
15. Lawson D, Harrison M, Shapland C. Fibroblast transgelin and smooth muscle SM22alpha are the same protein, the expression of which is down-regulated in many cell lines. *Cell Motil Cytoskeleton* 1997;38:250–257.

Copyright © 2019 by the American Thoracic Society