DOTA-MGS5, a New Cholecystokinin-2 Receptor-Targeting Peptide Analog with an Optimized Targeting Profile for Theranostic Use

Maximilian Klingler¹, Dominik Summer¹, Christine Rangger¹, Roland Haubner¹, Julie Foster², Jane Sosabowski², Clemens Decristoforo¹, Irene Virgolini¹, and Elisabeth von Guggenberg¹

¹Department of Nuclear Medicine, Medical University of Innsbruck, Innsbruck, Austria; and ²Centre for Molecular Oncology, Barts Cancer Institute, Queen Mary University of London, London, United Kingdom

Molecular imaging and targeted radiotherapy with radiolabeled cholecystokinin-2 receptor (CCK2R) targeting peptide probes holds high promise to improve the clinical management of patients with metastatic medullary thyroid carcinoma and other CCK2R-expressing malignancies. Low stability and suboptimal targeting of currently available radiolabeled peptide analogs has prompted us to seek new stabilization strategies. In this study, we present a new minigastrin analog with site-specific C-terminal modifications showing a highly optimized targeting profile. **Methods:** DOTA-D-Glu-Ala-Tyr-Gly-Trp-(N-Me)Nle-Asp-1-Nal-NH₂ (DOTA-MGS5) radiolabeled with ¹¹¹In, ⁶⁸Ga, and ¹⁷⁷Lu was evaluated in extensive in vitro stability studies. For ¹⁷⁷Lu-DOTA-MGS5, additional metabolic studies were performed on BALB/c mice. Receptor affinity and cell uptake were studied in A431 human epidermoid carcinoma cells transfected with human CCK2R (A431-CCK2R), as well as the same cell line transfected with the empty vector (A431-mock). A431-CCK2R/A431-mock xenografted athymic BALB/c nude mice were used for biodistribution studies and small-animal SPECT/CT. **Results:** DOTA-MGS5 radiolabeled with ¹¹¹In and ¹⁷⁷Lu showed a highly increased stability against enzymatic degradation in different media up to 24 h of incubation. Similar results were observed for ⁶⁸Ga-DOTA-MGS5 incubated up to 4 h. In the blood of mice injected with ¹⁷⁷Lu-DOTA-MGS5, at least 70% intact radiopeptide was detected up to 1 h after injection. The unlabeled peptide and the complexes with the natural isotopes showed retained receptor affinity, and the radiopeptides showed unexpectedly high cell uptake in A431-CCK2R cells (>60% at 4 h). Regardless of the radiometal used for labeling, impressively high uptake in A431-CCK2R xenografts was found (~20% injected activity/g 1–4 h after injection), whereas the uptake in A431-mock xenografts was negligible. Low background activity and favorable tumor-to-kidney ratios (4–6) allowed for high image contrast in small-animal SPECT/CT. **Conclusion:** The excellent targeting properties of DOTA-MGS5 support future clinical studies evaluating the diagnostic and therapeutic potential in patients with progressive or metastatic medullary thyroid carcinoma, as well as other advanced-stage CCK2R-expressing malignancies.

Key Words: cholecystokinin-2 receptor; minigastrin; molecular imaging; radiometals; targeted radiotherapy

J Nucl Med 2019; 60:1010–1016

DOI: 10.2967/jnumed.118.221283

Cholecystokinin-2 receptors (CCK2R) are expressed at high incidence in medullary thyroid carcinomas (MTC, > 90%), small cell lung cancers (>50%), astrocytomas (>60%), insulinomas, stromal ovarian cancers, gastrointestinal stromal tumors, and more than 20% of gastroenteropancreatic tumors (1,2). CCK2R targeting, therefore, offers a highly personalized diagnostic and therapeutic approach for different cancers when safe and effective systemic therapies are lacking. This is especially the case for patients with unresectable, locally advanced, or metastatic MTC. Because of substantial side effects, targeted treatment with the multikinase inhibitors vandetanib and cabozantinib is restricted to patients with progressive and symptomatic disease (3). In patients with metastasized MTC, a higher tumor detection rate (>90%) was found for CCK2R imaging in comparison to somatostatin receptor imaging (40%) (4). Also, for about 20% of the patients with carcinoids and other neuroendocrine tumors, an additive value of CCK2R imaging was reported (5). Initial targeted radiotherapy in MTC patients showed encouraging therapeutic effects; nephrotoxicity was, however, found to be a major concern (6).

CCK2R-targeting peptide probes are based on the natural ligands cholecystokinin and gastrin. Gastrin analogs generally show higher receptor-specific uptake, but the kidney uptake is also high. Renal accumulation was successfully reduced by depletion of the penta-Glu sequence in minigastrin but was connected with reduced physiologic half-life and poor diagnostic efficacy (7). A high variety of synthetic modifications in the N-terminal part of the peptide sequence was investigated to increase enzymatic stability, reduce kidney retention, and improve tumor targeting (8,9). In the C-terminal sequence Trp-Met-Asp-Phe-NH₂, known to be essential for receptor binding (10), only replacement of oxidation-sensitive Met allowed maintenance of receptor affinity (8,11).

For 2 novel radiopeptides, clinical trials have recently been initiated. In 4 MTC patients, ¹¹¹In-labeled DOTA-(D-Glu)₆-Ala-Tyr-Gly-Trp-Met-Asp-Phe-NH₂ (CP04) visualized all known tumor lesions (12). Also, ¹⁷⁷Lu-PP-F11N, derived from CP04 by replacement of Met with Nle, showed high CCK2R-specific tumor uptake in

Received Oct. 15, 2018; revision accepted Nov. 13, 2018.

For correspondence or reprints contact: Elisabeth von Guggenberg, Department of Nuclear Medicine, Medical University of Innsbruck, Anichstrasse 35, 6020 Innsbruck, Austria.

E-mail: elisabeth.von-guggenberg@i-med.ac.at

Published online Dec. 7, 2018.

COPYRIGHT © 2019 by the Society of Nuclear Medicine and Molecular Imaging.

a patient with progressive, metastatic MTC (13). Despite physiologic uptake observed in stomach, colon, kidneys, and urinary bladder, the dosimetry studies indicate the feasibility of targeted radiotherapy.

A main issue remaining unsolved is the limited enzymatic stability of these linear peptide analogs *in vivo*. Rapid cleavage in the C-terminal receptor-specific sequence occurs, and no improvement in tumor uptake can be achieved by coadministration of protease inhibitors (13). We have recently described a new stabilization strategy based on site-specific amino acid modifications with unnatural aromatic amino acids and N-methylated amino acids preventing the metabolism of the C-terminal receptor-specific sequence. When applied to the truncated minigastrin analog DOTA-MG11 missing the penta-Glu motif (7), these modifications led to compounds with high CCK2R affinity, improved bioavailability, limited kidney retention, and highly increased tumor uptake (14).

In the present study, we have further refined our approach and present the new minigastrin analog DOTA-D-Glu-Ala-Tyr-Gly-Trp-(N-Me)Nle-Asp-1-Nal-NH₂ (DOTA-MGS5) with an optimized targeting profile. Specific modifications were applied in the C-terminal receptor-specific sequence of the peptide, namely replacement of Met with Nle and of Phe with 1-naphthyl-alanine (1-Nal), as well as N-methylation of the peptide bond between Trp and Nle. A comprehensive preclinical evaluation of the DOTA-conjugated peptide analog radiolabeled with ¹¹¹In, ⁶⁸Ga, and ¹⁷⁷Lu was performed. These studies included extensive *in vitro* and *in vivo* stability studies, receptor affinity and cell uptake studies, and tumor targeting and biodistribution studies in an animal tumor model, highlighting the specific features of this novel CCK2R-targeting peptide analog.

MATERIALS AND METHODS

Materials

All commercially obtained chemicals were of analytic grade and used without further purification. ¹¹¹InCl₃ was purchased from Mal-inkrodt Medical. Non-carrier-added ¹⁷⁷LuCl₃ produced from highly enriched ¹⁷⁶Yb was purchased from Isotope Technologies Garching. ⁶⁸GaCl₃ was obtained from a commercial ⁶⁸Ge/⁶⁸Ga generator with nominal activity of 1,850 MBq (Eckert and Ziegler Eurotope) eluted with 0.1N HCl solution (Rotem). DOTA-MG11 (DOTA-D-Glu-Ala-Tyr-Gly-Trp-Met-Asp-Phe-NH₂) and CP04 used for comparative studies were purchased from piCHEM. The A431 human epidermoid carcinoma cell line stably transfected with human CCK2R (A431-CCK2R), as well as the same cell line transfected with the empty vector (A431-mock), was kindly provided by Dr. Luigi Aloj and cultured as described elsewhere (15).

Peptide Synthesis

DOTA-MGS5 (DOTA-D-Glu-Ala-Tyr-Gly-Trp-(N-Me)Nle-Asp-1-Nal-NH₂) was synthesized and analyzed following a previously described protocol (14). The chemical structure is displayed in Figure 1.

Radiolabeling and Characterization *In Vitro*

Standard protocols were used for labeling with the different radiometals (9,14,16). Radiochemical purity of the radiopeptides was analyzed using a high-performance liquid chromatography system described elsewhere (9) together with a water/acetonitrile/0.1% trifluoroacetic acid gradient with increasing concentrations of acetonitrile (¹¹¹In and ¹⁷⁷Lu: 0–3 min, 10%; 3–18 min, 10%–55%; 18–20 min, 55%–80%; 20–21 min, 80%–10%; 20–25 min, 10%; ⁶⁸Ga: 0–2 min, 20%; 2–12 min, 20%–70%; 12–12.1 min, 70%–20%; 12.1–15 min, 20%).

The stability of ¹¹¹In-, ⁶⁸Ga- and ¹⁷⁷Lu-labeled DOTA-MGS5 was investigated after incubation in phosphate-buffered saline (PBS; *n* = 1)

FIGURE 1. Chemical structure of DOTA-MGS5.

and fresh human serum (1–2 nmol of peptide/mL, *n* = 2). A PBS sample of the radiolabeled peptides (~0.05 nmol/mL) was used to determine the distribution coefficient (log *D*; *n* = 8). A serum sample (~0.5 nmol/mL) was used to assess the protein binding by size-exclusion chromatography (*n* = 2). For comparison, the same assays were performed with ¹¹¹In-CP04. The enzymatic degradation of radiolabeled DOTA-MGS5 (*n* = 2 for ¹¹¹In; *n* = 1 for ⁶⁸Ga and ¹⁷⁷Lu) was further assessed in rat liver and kidney homogenates (~0.5 nmol of peptide/mL). Previously published protocols were followed for this *in vitro* characterization (14,17).

Cell Uptake and Receptor Binding Studies

The receptor affinity of DOTA-MGS5 (*n* = 3), as well as CP04, DOTA-MG11, and pentagastrin (*n* = 1), was evaluated in competition assays against ¹²⁵I-Tyr¹²-gastrin-I on A431-CCK2R cells. Binding assays, including also receptor binding studies with DOTA-MGS5 complexed with ^{nat}In, ^{nat}Lu and ^{nat}Ga (*n* = 1), were performed as described elsewhere (16). Half maximal inhibitory concentrations (IC₅₀s) were calculated following nonlinear regression using Origin software (Microcal Origin, version 6.1).

Internalization and externalization experiments were performed on A431-CCK2R and A431-mock cells following previously published protocols (14,18). For internalization experiments, the cells were incubated with the radiopeptides (final peptide concentration of 0.4 nM) for up to 4 h, and the radioactivity of the lysed cells was determined in relation to the total radioactivity added (% internalized radioactivity). For externalization experiments, after 2 h of incubation the medium was replaced with 3 mL of fresh medium (unblocked) or medium containing 1 mM pentagastrin (blocked) and the efflux of radioactivity evaluated in relation to the cell uptake (% externalized radioactivity). Nonspecific binding was assessed in A431-mock cells. In an additional experiment, the cell uptake in dependence on the peptide concentration within the range of 0.04–160 nM was investigated. Furthermore, comparative internalization studies at 2 h of incubation were performed for DOTA-MGS5 and CP04 radiolabeled with different radiometals, including blocking of the receptor-specific uptake with 1 μM human minigastrin.

Animal Studies Evaluating the *In Vivo* Stability and Tumor Uptake

Metabolic and biodistribution studies were performed in accordance with the ethical standards of the institution and approved by the Austrian Ministry of Science (BMFWF-66.011/0075-WF/V/3b/2016).

The metabolic stability of ¹⁷⁷Lu-DOTA-MGS5 (10, 30, and 60 min after injection; *n* = 1) and ¹⁷⁷Lu-CP04 (30 min after injection; *n* = 1) was evaluated in 5- to 7-wk-old female BALB/c mice (Charles River). Mice were injected intravenously through a lateral tail vein with 20–45 MBq of the ¹⁷⁷Lu-labeled peptide (≤2 nmol total peptide), and the percentage of intact radiopeptide in blood was analyzed as described elsewhere (14,18).

Biodistribution studies were performed on female athymic BALB/c nude mice (Charles River) injected subcutaneously with A431-CCK2R

TABLE 1
In Vitro Characteristics of Radiolabeled DOTA-Peptides

Radiolabeled peptide	log D	Protein binding* (%)	Stability in PBS* (% intact radiopeptide)	Stability in serum* (% intact radiopeptide)
⁶⁸ Ga-DOTA-MGS5	-2.37 ± 0.09	23.0 ± 0.3	95.5	96.6 ± 0.8
¹¹¹ In-DOTA-MGS5	-2.03 ± 0.12	44.3 ± 0.3	97.7	97.2 ± 0.6
¹⁷⁷ Lu-DOTA-MGS5	-2.18 ± 0.11	39.0 ± 1.6	95.5	96.5 ± 0.1
¹¹¹ In-CP04	-4.70 ± 0.35	33.0 ± 0.3	96.0	88.2 ± 0.1

*¹¹¹In and ¹⁷⁷Lu: 24 h of incubation; ⁶⁸Ga: 4 h of incubation.

and A431-mock cells (2×10^6 in 200 μ L) at an age of 6–8 wk. After 7–11 d, when tumors reached a volume of about 0.3 mL (1 mouse did not develop the A431-mock xenograft), the mice were injected intravenously via a lateral tail vein with about 0.02 nmol of radiolabeled peptide (~ 0.05 MBq for ¹¹¹In, ~ 0.8 MBq for ⁶⁸Ga, ~ 0.4 MBq for ¹⁷⁷Lu) in groups of 4. After sacrifice at 1 or 4 h after injection, tumors and other tissues were removed and weighed, and the radioactivity was measured in the γ -counter (Perkin Elmer Life Sciences and Analytic Sciences) together with a standard. Results were expressed as percentage of injected activity per gram of tissue (%IA/g), and tumor-to-organ activity ratios were calculated for selected tissues. Statistical analysis (independent-samples *t* test, *P* < 0.05) was performed using Origin software.

The imaging properties of ¹¹¹In-DOTA-MGS5 were evaluated by small-animal SPECT/CT. Three xenografted athymic BALB/c nude mice (Charles River) were injected with 10 MBq (0.3 nmol) of the radiolabeled peptide. Whole-body SPECT/CT images were obtained at 1 and 4 h after injection under inhaled anesthesia (1 L/min 1.5% isoflurane in O₂) in the prone position on a heated bed using a small-animal SPECT/CT 4-head camera (Bioscan). SPECT images were acquired in helical scanning mode with 24 projections using 1.4-mm-pinhole collimators and a voxel size of $100 \times 100 \mu\text{m}$, and CT images were acquired with a 45-kVp x-ray source (180 projections over 6 min) (14). Region of interest analysis of the coregistered SPECT and CT images was performed using VivoQuant image analysis software (inviCRO LLC). After being scanned, the animals were sacrificed to determine the tracer uptake in tumors and other tissues.

RESULTS

Peptide Synthesis and Radiolabeling

DOTA-MGS5 was synthesized with moderate yield and high purity of at least 95%. Radiolabeling with the different radiometals

generally resulted in radiochemical purity of at least 95% at a molar activity of no more than 35 GBq/ μmol for ¹¹¹In, no more than 40 GBq/ μmol for ¹⁷⁷Lu, and no more than 60 GBq/ μmol for ⁶⁸Ga.

Characterization of the Radiopeptides In Vitro

Table 1 summarizes the in vitro characteristics of radiolabeled DOTA-MGS5 in comparison with ¹¹¹In-CP04. For ¹¹¹In-DOTA-MGS5 and ¹⁷⁷Lu-DOTA-MGS5, the stability in PBS and human serum was analyzed for up to 24 h after incubation. Because of the short half-life of ⁶⁸Ga, ⁶⁸Ga-DOTA-MGS5 was incubated only for up to 4 h. The radiolabeled complexes showed a high stability in aqueous solution. Also, in human serum more than 95% intact radiopeptide was found. Comparative studies with ¹¹¹In-CP04 showed a high stability in PBS, whereas a lower stability (<90% intact radiopeptide) was found in serum. ⁶⁸Ga-DOTA-MGS5, ¹⁷⁷Lu-DOTA-MGS5, and ¹¹¹In-DOTA-MGS5 showed a higher log D value than ¹¹¹In-CP04, confirming a somewhat lower hydrophilicity. Protein binding was, however, comparable to ¹¹¹In-CP04. A much higher degree of degradation, with less than 30% intact radiopeptide, after 2 h was found in rat liver homogenates than in human serum. ⁶⁸Ga-DOTA-MGS5 showed a faster degradation over time than the ¹¹¹In- and ¹⁷⁷Lu-labeled radiopeptides. In rat kidney homogenates, all radiopeptides were almost completely degraded (Fig. 2).

Receptor Binding and Internalization Studies

A high CCK2R affinity with IC₅₀s of less than 1 nM was observed for DOTA-MGS5 and for the complexes with the natural isotopes ^{nat}In, ^{nat}Lu, and ^{nat}Ga. A similar binding affinity was also found for pentagastrin, DOTA-MG11, and CP04 (Table 2).

With ¹¹¹In-DOTA-MGS5, remarkably high cell uptake values of more than 60% were reached in A431-CCK2R cells after 4 h of incubation (Fig. 3A). Efflux studies showed only a minor release of radioactivity (<10%) under unblocked assay conditions, whereas much higher values of more than 40% were reached when the reuptake was blocked by addition of pentagastrin (Fig. 3B). Stable uptake values of more than 50% were observed within a concentration range of 0.04–1.6 nM. At a concentration of 4 nM, the uptake was slightly reduced to approximately 40% and further dropped to 5%–10% at 40 nM and 160 nM (Fig. 3C). The nonspecific uptake in A431-mock cells was always below 1.5%. DOTA-MGS5 radiolabeled with ¹¹¹In, ⁶⁸Ga, and ¹⁷⁷Lu showed a considerably increased cell uptake of 50%–60% in A431-CCK2R cells

FIGURE 2. Intact radiopeptide in rat homogenates of liver (A) and kidney (B).

TABLE 2
IC50 as Calculated from Competition Assays Against
¹²⁵I-Tyr¹²-Gastrin-I

Peptide analog	IC50 (nM)
DOTA-MGS5	0.4 ± 0.2
^{nat} In-DOTA-MGS5	0.2 ± 0.04
^{nat} Ga-DOTA-MGS5	0.6 ± 0.2
^{nat} Lu-DOTA-MGS5	0.6 ± 0.1
CP04	0.4 ± 0.1
DOTA-MG11	0.9 ± 0.3
Pentagastrin	1.0 ± 0.2

at 2 h after incubation (Fig. 4A) when compared with radiolabeled CP04 (15%–30%; Fig. 4B). Blocking studies with human mini-gastrin showed uptake values below 1.5%, confirming the receptor specificity (Fig. 4C).

Metabolic Studies and Biodistribution Studies In Vivo

In BALB/c mice injected with ¹⁷⁷Lu-DOTA-MGS5, 86% intact radiolabeled peptide was present in blood at 10 min after injection and still 77% and 70% intact radiolabeled peptide could be detected at 30 and 60 min after injection, respectively. In comparative studies performed with ¹⁷⁷Lu-CP04 at 30 min after injection, only 5% intact radiolabeled peptide was observed in blood. Representative radiochromatograms at 30 min after injection are displayed in Figure 5.

In the biodistribution studies, DOTA-MGS5 radiolabeled with different radiometals was rapidly cleared from the body, mainly through the kidneys, and a low nonspecific uptake generally occurred in most tissues (Fig. 6; Supplemental Table 1; supplemental materials are available at <http://jnm.snmjournals.org>). In blood, lung, heart, spleen, and liver, ⁶⁸Ga-DOTA-MGS5 showed a slightly higher uptake than ¹¹¹In-DOTA-MGS5 at 1 h after injection ($P > 0.05$). In CCK2R-expressing stomach and pancreas, the uptake of ⁶⁸Ga-DOTA-MGS5 was lower but was significantly reduced only for stomach ($P < 0.01$). The uptake of ¹¹¹In-DOTA-MGS5 significantly decreased over time in blood, lung, heart, spleen, liver, and kidney ($P < 0.03$). When ¹¹¹In-DOTA-MGS5 and ¹⁷⁷Lu-DOTA-MGS5 were compared at 4 h after injection, only in CCK2R-expressing pancreas ($P < 0.02$) was a significant difference found. The uptake in kidneys remained at less than 6 %IA/g

for all radiolabeled peptides and time points studied. Impressive tumor uptake and tumor retention were observed for ¹¹¹In-DOTA-MGS5 in A431-CCK2R xenografts, with values of 19.53 ± 5.42 %IA/g at 1 h after injection and 23.49 ± 1.25 %IA/g at 4 h. A similar tumor uptake was also observed with ⁶⁸Ga-DOTA-MGS5 (23.25 ± 4.70 %IA/g, 1 h after injection) and ¹⁷⁷Lu-DOTA-MGS5 (24.45 ± 3.11 %IA/g, 4 h after injection), whereas only minimal uptake occurred in A431-mock xenografts (<1 %IA/g). When comparing the uptake of ¹¹¹In-DOTA-MGS5 at 0.02 nmol (biodistribution, 4 h after injection) and 0.3 nmol (imaging, ~5 h after injection), that in A431-CCK2R xenografts was reduced by 40% ($P < 0.02$) at the higher peptide amount. Also, the uptake in CCK2R-expressing pancreas ($P < 0.01$) and stomach ($P < 0.02$), as well as intestine ($P < 0.01$) and kidneys ($P < 0.01$), was significantly lower. The low activity levels in blood and normal tissue combined with strong uptake and prolonged retention in the tumor over time resulted in high tumor-to-organ activity ratios, especially at 4 h after injection, with mean values of 256–572 for blood, 6.11–6.53 for kidneys, and 3.08–4.74 for stomach (Supplemental Table 2).

Small-animal SPECT/CT imaging with ¹¹¹In-DOTA-MGS5 ultimately confirmed the favorable biodistribution profile (Fig. 7). On the maximum-intensity projections, a low background activity and high image contrast were observed. At 1 h after injection, the most prominent structure was the A431-CCK2R tumor. Excretion of radioactivity via the kidneys and bladder could also be seen. No uptake was observed in A431-mock tumors, which were visible only on CT. At 4 h after injection, the bladder was almost emptied and the kidneys could no longer be seen, whereas the tumor uptake remained comparatively high (0.41 and 0.36 MBq at 1 and 4 h, respectively). From the region-of-interest analysis performed on 2 mice, a tumor-to-kidney ratio of more than 2 and more than 4 was calculated for 1 and 4 h after injection, respectively.

DISCUSSION

Despite extensive radiopharmaceutical development, radiolabeled CCK2R-targeting peptide analogs have not yet been introduced into routine clinical practice. In the present study, we describe the novel minigastrin analog DOTA-MGS5 with site-specific stabilizations in the C-terminal receptor-specific sequence. It has recently been reported that the radiolabeled minigastrin analogs currently investigated in clinical trials are rapidly metabolized at this site by different proteases (13). The preclinical characterization of DOTA-MGS5 radiolabeled with ¹¹¹In, ⁶⁸Ga, and ¹⁷⁷Lu shows that

FIGURE 3. Cell uptake of ¹¹¹In-DOTA-MGS5. (A) Internalized activity in A431-CCK2R and A431-mock cells. (B) Efflux of radioactivity from A431-CCK2R cells. (C) Internalization in dependence on peptide concentration.

FIGURE 4. Cell internalization of DOTA-MGS5 (A) and CP04 (B), as well as blocking studies for DOTA-MGS5 (C).

this new peptide analog fulfils important criteria for diagnostic and therapeutic use in nuclear medicine: excellent target-to-nontarget ratios, specific binding to tumor tissue, and high metabolic stability (19). To our knowledge, this is the first report on a radiolabeled CCK2R-targeting peptide probe with such an intense tumor uptake along with favorable tumor-to-kidney ratios.

Because most of the CCK2R-targeting radiopeptides previously developed have been studied in the A431-CCK2R/A431-mock mouse tumor model (9,14,20), this model was used for the pre-clinical evaluation of DOTA-MGS5. An impressively high and persistent tumor uptake (>20 %IA/g) was observed in A431-CCK2R xenografts together with considerably improved tumor-to-kidney ratios (>6). The uptake was clearly receptor-specific, as shown by the limited uptake in A431-mock xenografts (<1 %IA/g), and was confirmed in blocking studies on A431-CCK2R cells in vitro. The extraordinary targeting properties also become clear from the small-animal SPECT/CT imaging performed with ^{111}In -DOTA-MGS5 at a 10-fold higher peptide amount, even though partial saturation effects occurred in A431-CCK2R tumors and in CCK2R-expressing tissues.

DOTA-MGS5 also seems highly promising for targeted radiotherapy, as the tumor uptake of ^{177}Lu -DOTA-MGS5 is more than tripled in comparison with the radiopeptides currently used in clinical trials. ^{177}Lu -CP04 and ^{177}Lu -PP-F11N show a tumor uptake of 6.70 ± 0.56 and 6.94 ± 0.82 %IA/g in A431-mock xenografts when injected at a comparable peptide dose of 0.01 nmol. In xenografts based on human MTC cells (MZ-CRC-1), in line with a

higher number of binding sites, a higher uptake (10–20 %IA/g) was observed for both radiopeptides. Still, tumor-to-kidney ratios with values of 3–4 were clearly inferior (13). In comparative cell uptake studies using CP04 and DOTA-MGS5 radiolabeled with different radiometals, we could confirm the highly increased cell internalization of DOTA-MGS5.

When analyzing our data, different parameters explain the optimized targeting profile of radiolabeled DOTA-MGS5.

Even though the C-terminal sequence Trp-X-Asp-Phe-NH₂ (with X being Met or a Met-like substituent) (7) was postulated to be essential for receptor binding, DOTA-MGS5, with methylation of the peptide bond between Trp and Nle and the voluminous amino acid 1-Nal at the C terminus, showed high binding affinity. For CCK4 analogs, the exchange of Met with (N-Me)Nle resulted in increased CCK2R affinity (21), whereas the additional substitution of Phe with 1-Nal abolished this positive effect (22). As reported previously also for another minigastrin analog (23), we saw no metal-related changes on receptor affinity and cell uptake. We found evidence in the literature that the C terminus of gastrin assumes a well-defined β -turn that interacts with a hydrophobic pocket of the CCK2R (24). Especially, the aromaticity of Phe was found

FIGURE 5. Radiometabolites as analyzed by radio-high-performance liquid chromatography from blood of BALB/c mice 30 min after injection with ^{177}Lu -DOTA-MGS5 (A) and ^{177}Lu -DOTA-CP04 (B). Dashed line shows radiochromatogram of radiolabeled peptide before injection.

FIGURE 6. Biodistribution of DOTA-MGS5 radiolabeled with different radiometals in A431-CCK2R/A431-mock xenograft-bearing BALB/c nude mice. Data are mean $\pm$ SD ($n = 4$). p.i. = after injection.

FIGURE 7. Small-animal SPECT/CT images at 1 and 4 h after injection of BALB/c nude mice xenografted with A431-CCK2R (left) and A431-mock cells (right) injected with ^{111}In -DOTA-MGS5. SPECT is in color scale ranging from 0.0015 to 0.011 kBq/voxel; CT is in gray scale.

to increase polar receptor–ligand interactions (25). The amino acid substitutions applied in DOTA-MGS5 increase the lipophilicity of the radiopeptide, possibly promoting the receptor interaction.

We could show that the specific substitutions in the C-terminal sequence of DOTA-MGS5 successfully prevent the metabolic degradation in vivo. In the blood of BALB/c mice injected with ^{177}Lu -DOTA-MGS5, 70% intact radiopeptide was still found at 1 h after injection. Such a stabilization could not be achieved with N-terminal modifications (13,26). This improved stability against enzymatic degradation in vivo was not an unexpected finding. Methylation of the peptide bond between Trp and Nle and introduction of voluminous amino acids at the C terminus were already investigated for CCK4 analogs, achieving compounds that are protected from degradation in rat brain homogenates (23). Higher bioavailability is likely to increment the amount of intact radiopeptide reaching the target site. In the efflux studies from A431-CCK2R cells, we found a much higher efflux of radioactivity under blocked assay conditions, suggesting that the intact radiopeptide is excreted from the cells and that only under unblocked assay conditions can reuptake occur.

Interestingly, DOTA-MGS5, because of its lower hydrophilicity, revealed a 3-fold increased protein binding in vitro compared with ^{111}In -DOTA-MG11, with values of 41.0% versus 10.1% at 4 h (14). Also, ^{111}In -CP04 showed an increased protein binding of 33.0%. The combination of increased enzymatic stability and increased protein binding enhancing the initial systemic circulation seems to be responsible for the reinforced specific CCK2R interaction at the tumor site. This hypothesis is supported by our recent findings with another ^{111}In -labeled minigastrin analog showing C-terminal modifications. ^{111}In -DOTA-MGS4 substituted with (N-Me)Nle and (N-Me)Phe (instead of 1-Nal) showed enzymatic stability

similar to ^{111}In -DOTA-MGS5 but a much lower protein binding. The blood levels of ^{111}In -DOTA-MGS5 at 1 h after injection (1.19 ± 0.29 %IA/g) were more than 5 times higher than those of ^{111}In -DOTA-MGS4 (0.2 ± 0.07 %IA/g). Consequently, ^{111}In -DOTA-MGS5 showed a 2-fold higher tumor uptake in A431-CCK2R xenografts than did ^{111}In -DOTA-MGS4 (14). At the later time point of 4 h after injection, the blood values of ^{111}In -DOTA-MGS5 and ^{177}Lu -DOTA-MGS5 declined to low levels of about 0.1 %IA/g, comparable to ^{111}In -DOTA-MGS4 (0.08 ± 0.09 %IA/g), indicating that no prolonged systemic circulation occurred. Also, the 3-fold improvement in tumor uptake of ^{111}In -CP04 when compared with ^{111}In -DOTA-MG11 could be attributed to its higher protein binding rather than stabilization against metabolic degradation.

The impressive targeting properties become clear from the small-animal SPECT/CT imaging performed with ^{111}In -DOTA-MGS5 and indicate that with this new CCK2R-targeting peptide analog a higher absorbed dose may be reached in the tumor also in targeted radiotherapy with ^{177}Lu -DOTA-MGS5. The absorbed dose of the kidneys could potentially be reduced by the use of kidney protective agents such as Gelofusine (B. Braun) (27). Also, the higher uptake observed in the stomach when compared with ^{177}Lu -CP04 and ^{177}Lu -PP-F11N (13) seems not to be limiting for a therapeutic application, as it slowly decreases with time and in dependence on the injected peptide amount. In radiation therapy, doses of less than 45 Gy are generally accepted for the stomach without unacceptable complications (28). Besides improved molecular imaging, DOTA-MGS5 seems therefore especially promising for targeted radiotherapy in patients with advanced MTC and other CCK2R-expressing malignancies.

CONCLUSION

The high and persistent tumor uptake and favorable tumor to-background activity ratios, including kidneys, give high promise that new radiopharmaceuticals based on DOTA-MGS5 will be powerful peptide probes in the localization and treatment of patients with CCK2R-expressing tumors. DOTA-MGS5 radiolabeled with different radiometals— ^{111}In for SPECT, ^{68}Ga for PET, or β -emitting radionuclides such as ^{177}Lu —seems therefore most suited for future clinical studies evaluating the diagnostic and therapeutic potential in patients with progressive or metastatic MTC, as well as other advanced-stage CCK2R-expressing malignancies.

DISCLOSURE

This study was financially supported by the Austrian Science Fund (FWF) (project P 27844). Elisabeth von Guggenberg and Maximilian Klingler are inventors of the novel intellectual property presented in this article; the patent application submitted by the Medical University of Innsbruck is currently under evaluation. No other potential conflict of interest relevant to this article was reported.

REFERENCES

1. Reubi JC, Schaer JC, Waser B. Cholecystokinin(CCK)-A and CCK-B/gastrin receptors in human tumors. *Cancer Res.* 1997;57:1377–1386.
2. Sanchez C, Escriveau C, Clerc P, et al. Characterization of a novel five-transmembrane domain cholecystokinin-2 receptor splice variant identified in human tumors. *Mol Cell Endocrinol.* 2012;349:170–179.
3. Brose MS, Bible KC, Chow LQM, et al. Management of treatment-related toxicities in advanced medullary thyroid cancer. *Cancer Treat Rev.* 2018;66:64–73.
4. Gotthardt M, Béhé MP, Beuter D, et al. Improved tumour detection by gastrin receptor scintigraphy in patients with metastasised medullary thyroid carcinoma. *Eur J Nucl Med Mol Imaging.* 2006;33:1273–1279.

5. Gotthardt M, Béhé MP, Grass J, et al. Added value of gastrin receptor scintigraphy in comparison to somatostatin receptor scintigraphy in patients with carcinoids and other neuroendocrine tumours. *Endocr Relat Cancer*. 2006;13:1203–1211.
6. Béhé M, Behr TM. Cholecystokinin-B (CCK-B)/gastrin receptor targeting peptides for staging and therapy of medullary thyroid cancer and other CCK-B receptor expressing malignancies. *Biopolymers*. 2002;66:399–418.
7. Fröberg AC, De Jong M, Nock BA, et al. Comparison of three radiolabelled peptide analogues for CCK-2 receptor scintigraphy in medullary thyroid carcinoma. *Eur J Nucl Med Mol Imaging*. 2009;36:1265–1272.
8. Laverman P, Joosten L, Eek A, et al. Comparative biodistribution of 12 ¹¹¹In-labelled gastrin/CCK2 receptor-targeting peptides. *Eur J Nucl Med Mol Imaging*. 2011;38:1410–1416.
9. Summer D, Rangger C, Klingler M, et al. Exploiting the concept of multivalency with ⁶⁸Ga- and ⁸⁹Zr-labelled fusarinine C-minigastrin bioconjugates for targeting CCK2R expression. *Contrast Media Mol Imaging*. 2018;2018:3171794.
10. Roosenburg S, Laverman P, van Delft FL, Boerman OC. Radiolabeled CCK/gastrin peptides for imaging and therapy of CCK2 receptor-expressing tumors. *Amino Acids*. 2011;41:1049–1058.
11. Grob NM, Behe M, von Guggenberg E, Schibli R, Mindt TL. Methoxinine: an alternative stable amino acid substitute for oxidation-sensitive methionine in radiolabelled peptide conjugates. *J Pept Sci*. 2017;23:38–44.
12. Konijnenberg M, Erba PA, Mikolajczak R, et al. First biosafety, biodistribution and dosimetry study of the gastrin analogue ¹¹¹In-CP04 in medullary thyroid cancer: phase I clinical trial, GRANT-T-MTC. *Eur J Nucl Med Mol Imaging* [abstract]. 2017;44(suppl 2):S258.
13. Sauter AW, Mansi R, Hassiepen U, et al. Targeting of the cholecystokinin-2 receptor with the minigastrin analog ¹⁷⁷Lu-DOTA-PP-F11N: does the use of protease inhibitors further improve in vivo distribution? *J Nucl Med*. July 12, 2018 [Epub ahead of print].
14. Klingler M, Decristoforo C, Rangger C, et al. Site-specific stabilization of minigastrin analogs against enzymatic degradation for enhanced cholecystokinin-2 receptor targeting. *Theranostics*. 2018;8:2896–2908.
15. Aloj L, Caracò C, Panico M, et al. In vitro and in vivo evaluation of ¹¹¹In-DTPAGlu-G-CCK8 for cholecystokinin-B receptor imaging. *J Nucl Med*. 2004;45:485–494.
16. Rangger C, Klingler M, Balogh L, et al. ¹⁷⁷Lu labeled cyclic minigastrin analogues with therapeutic activity in CCK2R expressing tumors: preclinical evaluation of a kit formulation. *Mol Pharm*. 2017;14:3045–3058.
17. von Guggenberg E, Behe M, Behr TM, Saurer M, Seppi T, Decristoforo C. ^{99m}Tc-labeling and in vitro and in vivo evaluation of HYNIC- and (N_α-His)acetic acid-modified [D-Glu¹]-minigastrin. *Bioconjug Chem*. 2004;15:864–871.
18. von Guggenberg E, Sallegger W, Helbok A, et al. Cyclic minigastrin analogues for gastrin receptor scintigraphy with technetium-99m: preclinical evaluation. *J Med Chem*. 2009;52:4786–4793.
19. Fani M, Maecke HR, Okarvi SM. Radiolabeled peptides: valuable tools for the detection and treatment of cancer. *Theranostics*. 2012;2:481–501.
20. von Guggenberg E, Rangger C, Sosabowski J, et al. Preclinical evaluation of radiolabeled DOTA-derivatized cyclic minigastrin analogs for targeting cholecystokinin receptor expressing malignancies. *Mol Imaging Biol*. 2012;14:366–375.
21. Bellier B, Million ME, DaNascimento S, et al. Replacement of glycine with dicarbonyl and related moieties in analogues of the C-terminal pentapeptide of cholecystokinin: CCK₂ agonists displaying a novel binding mode. *J Med Chem*. 2000;43:3614–3623.
22. Corringier PJ, Weng JH, Ducos B, et al. CCK-B agonist or antagonist activities of structurally hindered and peptidase-resistant Boc-CCK4 derivatives. *J Med Chem*. 1993;36:166–172.
23. Lipiński PFJ, Garnuszek P, Maurin M, et al. Structural studies on radiopharmaceutical DOTA-minigastrin analogue (CP04) complexes and their interaction with CCK2 receptor. *EJNMMI Res*. 2018;8:33.
24. Giragossian C, Schaschke N, Moroder L, Mierke DF. Conformational and molecular modeling studies of beta-cyclodextrin-heptagastrin and the third extracellular loop of the cholecystokinin 2 receptor. *Biochemistry*. 2004;43:2724–2731.
25. Ahmed SI, Wibowo F, Gembitsky DS, Bozsó Z, Murphy RF, Lovas S. Importance of the C-terminal phenylalanine of gastrin for binding to the human CCK2 receptor. *J Pept Res*. 2001;58:332–337.
26. Ocak M, Helbok A, Rangger C, et al. Comparison of biological stability and metabolism of CCK2 receptor targeting peptides, a collaborative project under COST BM0607. *Eur J Nucl Med Mol Imaging*. 2011;38:1426–1435.
27. Gotthardt M, van Eerd-Vismale J, Oyen WJG, et al. Indication for different mechanisms of kidney uptake of radiolabeled peptides. *J Nucl Med*. 2007;48:596–601.
28. Marks LB, Yorke ED, Jackson A, et al. Use of normal tissue complication probability models in the clinic. *Int J Radiat Oncol Biol Phys*. 2010;76(suppl):S10–S19.