Olufunmilayo O. Adebayo¹

Meinig School of Biomedical Engineering,
Cornell University,
Ithaca, NY 14853

Derek T. Holyoak¹

Meinig School of Biomedical Engineering,
Cornell University,
Ithaca, NY 14853

Marjolein C. H. van der Meulen²

Meinig School of Biomedical Engineering,
Cornell University,
113 Weill Hall,
Ithaca, NY 14853;

Sibley School of Mechanical and
Aerospace Engineering,
Cornell University,
Ithaca, NY 14853;

Research Division,
Hospital for Special Surgery,
New York, NY 10021
e-mail: mcv3@cornell.edu

Mechanobiological Mechanisms of Load-Induced Osteoarthritis in the Mouse Knee

Osteoarthritis (OA) is a degenerative joint disease that affects millions of people worldwide, yet its disease mechanism is not clearly understood. Animal models have been established to study disease progression by initiating OA through modified joint mechanics or altered biological activity within the joint. However, animal models often do not have the capability to directly relate the mechanical environment to joint damage. This review focuses on a novel in vivo approach based on controlled, cyclic tibial compression to induce OA in the mouse knee. First, we discuss the development of the load-induced OA model, its different loading configurations, and other techniques used by research laboratories around the world. Next, we review the lessons learned regarding the mechanobiological mechanisms of load-induced OA and relate these findings to the current understanding of the disease. Then, we discuss the role of specific genetic and cellular pathways involved in load-induced OA progression and the contribution of altered tissue properties to the joint response to mechanical loading. Finally, we propose using this approach to test the therapeutic efficacy of novel treatment strategies for OA. Ultimately, elucidating the mechanobiological mechanisms of load-induced OA will aid in developing targeted treatments for this disabling disease. [DOI: 10.1115/1.4043970]

Introduction

Osteoarthritis (OA) is a degenerative joint disease that affects approximately 27 million individuals in the U.S. and is predicted to affect 25% of the adult population by 2030 [1,2]. With highest prevalence in the knee [3], OA is characterized by cartilage degradation, subchondral bone changes, and osteophyte formation [4]. Clinically, the disease presents as radiographic narrowing of the joint space, debilitating pain and stiffness, and loss of joint function [5]. Based on clinical evidence, a wide range of risk factors have been identified, including obesity [6,7], aging [8], excessive mechanical loading during occupational activities [9,10], prior joint injuries [11,12], and genetic abnormalities [11]. Further, clinical data suggest that OA progression results in four different and overlapping phenotypes: inflammatory OA, cartilage-driven OA, bone-driven OA, and traumatic/acute OA [13], thus confounding the clinical assessment of disease progression. Although mechanical forces play a major role in OA initiation [11,14,15], the exact disease etiology is unknown, thus limiting effective preventative measures and clinical treatment options prior to surgical intervention, an end-stage treatment.

To better understand disease mechanism, preclinical animal models have been developed to recapitulate OA initiation and progression. These approaches can be classified under invasive and noninvasive categories (Fig. 1). Invasive models primarily use chemically induced cartilage degeneration or surgical injuries. Noninvasive models involve repetitive joint loading, load-induced impact injury, or spontaneously occurring OA [16–18].

Preclinical Animal Models of Osteoarthritis

Preclinical animal models have been used extensively to recapitulate and study the progression of osteoarthritis [16–20]. These models have included mice, rats, rabbits, dogs, sheep, and horses. These models, particularly those using small animals, enable rapid, large-scale studies of disease progression, with lower costs

and a greater ability to elucidate disease mechanisms compared to clinical studies. Furthermore, genetic manipulation of mouse models allows study of the effect of particular genetic pathways on OA progression. Although similarities in cartilage composition between articular joints of humans and animals allow for understanding of clinical OA, generalization to the human case may be limited due to differences in joint size, gait, and time course of disease progression [21,22]. However, preclinical models, using both invasive and noninvasive methods, provide appropriate platforms to begin to elucidate the mechanobiology of OA initiation and progression.

Invasive Preclinical Models of Osteoarthritis. Within the category of invasive models, chemically induced cartilage degeneration approaches involve intra-articular injection of collagenase [23–25], transforming growth factor- β [26], monosodium iodoacetate [27–31], or papain [32–34]. These models have provided critical knowledge about key molecular and cellular pathways during disease initiation. However, these models do not fully recapitulate human OA progression, as the outcome is often rapid and more severe, limiting their value for understanding clinical disease pathology [16,17].

Surgical models of OA better mimic clinical injuries and subsequent OA progression. Models, such as the anterior cruciate ligament (ACL) transection [35–38], meniscectomy [36,37], and the destabilization of the medial meniscus [39,40], use surgical methods to disrupt or transect stabilizing ligaments in the knee, leading to altered joint mechanics. With surgical trauma, OA develops over time, enabling investigation of the entire time course of post-traumatic OA progression. These surgical injuries have been implemented in a variety of animal models [16], and have significantly contributed to the current knowledge of post-traumatic OA progression. However, the difficulty in differentiating the surgical inflammatory and healing responses from the progression of OA is a major limitation of surgical models that confounds the understanding of disease etiology.

Noninvasive Preclinical Models of Osteoarthritis. Noninvasive models of OA have been developed to elucidate the mechanisms of disease initiation and progression without the

¹O. O. Adebayo and D. T. Holyoak have contributed equally to this paper.

²Corresponding author.

Manuscript received December 21, 2018; final manuscript received June 4, 2019; published online June 18, 2019. Editor: Beth A. Winkelstein.

Fig. 1 Overview of invasive and noninvasive preclinical OA models

confounding effects of surgery [41]. These models include genetic/spontaneous OA development [42,43], noninvasive injury models [44–47], and the recently developed cyclic load-induced model [19,20]. Genetic/spontaneous models involve the use of animals, such as the guinea pig [43], or specific strains of mice (e.g., C57Bl/6 and BALB/c) [48] that spontaneously develop joint damage, or mice that develop the disease throughout their lifespan as a result of genetic manipulation, through under- or overexpression of specific genes. Advantages of these approaches include the lack of external intervention and ability to explore specific genetic pathways to target with potential therapies. A limitation of these approaches is that spontaneous OA can take months or even years to develop. Genetically altered models can develop damage rapidly; however, such genetic manipulation may not reflect the human case [49,50].

Noninvasive injury models use high-magnitude compressive impact loads to induce an intra-articular fracture of the tibial plateau [44] or a rupture of the ACL [46], which both lead to OA initiation and progression. These approaches recapitulate the clinical joint injury scenario without invasive measures, a key advantage. On the other hand, the exact mechanical environment of the joint within these models is unknown. Thus, the relationship between the altered joint mechanics and OA initiation and progression is difficult to understand.

The cyclic load-induced model is a unique platform to specifically elucidate the relationship between mechanical forces and pathology initiation and progression. Primarily used in mice, applying controlled, cyclic compression to the tibia develops OA in the knee joint [19,20]. In this review, we focus on this load-induced model in the mouse, highlighting its development, characterization, and application to study the mechanical perturbation of cellular pathways and the mechanobiological mechanism of load-induced OA.

Development of the Load-Induced Osteoarthritis Model

Cyclic Tibial Compression: From Bone Adaptation to Cartilage Damage. Although *in vivo* cyclic compression of the tibia is a relatively novel technique to recapitulate OA-like pathology in mice [19,20], previously this model has been used

extensively to study the skeletal response to loading [51,52]. Cyclic tibial compression enables simultaneous and specific examination of cancellous and cortical bone responses to mechanical forces in osteoporosis and fracture healing models. These advantages also render the model appropriate for examining cartilage changes with loading, allowing for noninvasive, controlled compression of the articular cartilage in the knee without ligament rupture. In addition, temporal changes in both bone (Figs. 2(a)–2(c)) and cartilage (Figs. 2(d)–2(f)) can be examined using microcomputed tomography and histology, providing the opportunity to investigate the crosstalk between joint tissues in response to repetitive mechanical loading.

Adapted from the ulnar loading model [53], cyclic tibial compression applies repetitive axial loading at the ankle and knee joints to reliably induce anabolic responses in the tibial cancellous metaphysis and cortex [51,54,55]. Specifically, a loading system or device is used to apply cyclic compression to the lower limb while the mouse is anesthetized [56]. Custom-built devices include an electromagnetic actuator, load cell, and feedback control. The actuator applies cyclic loads according to voltage output from an amplifier, and the load cell provides force feedback control (Fig. 3). The system delivers controlled compressive forces to the tibia without dislocation of the knee or ankle joints.

Initially, load-induced OA models were developed and described in two laboratories, with slightly different device configurations and loading regimens [19,20]. The first configuration allows for normal flexion of the knee (normal flexion configuration, Fig. 3(b)) [20], and the second configuration constrains the knee in deep flexion (deep flexion configuration, Fig. 3(b)) [19]. These two configurations involve different loading regimens and experimental parameters, providing a unique opportunity to compare the effects of age, mouse strain, load magnitude, and loading frequency and duration on load-induced OA initiation and progression.

The initial study using the deep flexion configuration [19] used 8-week old male CBA mice, with a 9.0 N peak load applied for 40 cycles. Loading was conducted 3 days/week for 1 day, and 2 and 5 weeks (Fig. 3(c)). Initial experiments using the normal flexion configuration studied OA development and progression in 10- and 26-week old male C57Bl/6 mice, with a 9.0 N peak load, applied for 1200 cycles. Loading was conducted 5 days/week for 1, 2, and

Fig. 2 Analysis of bone and cartilage following load-induced OA. Microcomputed tomography analyses include: (a) subchondral bone thickness, (b) epiphyseal cancellous morphology, and (c) metaphyseal cancellous morphology. Histological analyses include (d) cartilage morphology, (e) local subchondral bone thickness, and (f) osteophyte size and maturity.

Fig. 3 (a) Schematic of limb in compressive loading device. The fixture positioning the foot is attached to an electromagnetic actuator that applies cyclic compression to the tibia according to voltage output from a LabVIEW-controlled amplifier. The fixture holding the knee connects to a load cell that measures the applied compressive force and sends feedback to the LabVIEW program to adjust the output voltage accordingly and control the actuator. (b) Schematic of a mouse limb in the deep flexion (blue) and normal flexion (red) configurations. The loading device applies compression according to a specified loading waveform. (c) Applied cyclic tibial loading waveforms for the deep flexion (blue) and normal flexion (red) configurations. Figure adapted from Poulet et al. [19] and Ko et al. [20].

6 weeks (Fig. 3(c)) [20]. In the normal configuration, a 9.0 N peak compressive load engenders $1200 \mu\epsilon$ at the midshaft of the tibia, measured in vivo by strain gauges [54]. This strain value is within the range of physiological strains engendered during locomotion in mice and most vertebrates [53,57,58].

Load-Induced Osteoarthritis Model Outcomes. Noninvasive mechanical loading recapitulated OA-like pathology under a variety of experimental conditions. Loading produced lesions and

erosion of the cartilage that progressed with duration and magnitude of loading (Fig. 4(a)). Osteophyte formation occurred with both the normal and deep loading configurations (Fig. 4(a)). Furthermore, 9.0 N loading induced posterior subchondral bone plate thickening in both configurations. Thus, cyclic tibial compression recapitulated key features of human OA pathology regardless of loading approach.

Specifically, in the normal flexion configuration, loading increased histological cartilage degradation scores 1.8-fold, with the most significant changes occurring after 6 weeks of loading

Fig. 4 (a) Cartilage matrix and osteophyte changes in the tibia after cyclic tibial compression. The left tibiae of adult mice were loaded for 1, 2, and 6 weeks. The nonloaded contralateral limb at 6 weeks load duration served as control. (b) Scoring of Safranin O-fast green staining of the articular cartilage on the medial tibial plateau indicated damage to the cartilage matrix occurred following loading and was exacerbated with longer durations. Figure adapted from Ko et al. [20].

(Fig. 4(b)). In the deep flexion configuration, loading increased mean histological scores from 0.11 to 3.64, reflecting cartilage damage with lesions extending to the intermediate articular cartilage zone [19].

Bone adaptation was present with both configurations and differed at early and later time points. In normal flexion cancellous bone volume fraction decreased in the epiphysis by 14% and was unchanged in the metaphysis with 6 weeks of loading [20]. In contrast, both epiphyseal and metaphyseal cancellous bone volume changed in the deep flexion configuration [59]. After 5 weeks of repetitive loading in deep flexion, epiphyseal trabecular bone volume fraction increased 31% in the lateral compartments of both loaded and nonloaded contralateral limbs.

Cyclic tibial compression not only mimicked the hallmarks of OA, but also led to other important changes in the tissues surrounding the joint. In the deep flexion configuration, loading also induced meniscal chondroplasia, synovial lining thickening, increased hypertrophy in the anterior cruciate ligament insertion points [19], and gait changes [59]. Interestingly, gait patterns were changed only in the nonloaded contralateral limbs. Specifically, stance time, stride length, and paw area were increased in contralateral limbs after loading and unchanged in loaded limbs. These outcomes were not reported for normal flexion. Changes in gait have been reported in clinical studies with OA progression, though none have specifically examined the contralateral limb [60–62]. In addition, pain has not been reported in any load-induced studies and could be another critical measure and mechanism [63,64].

Although the trends in OA progression are similar in both the normal and deep flexion configurations, the morphological changes in joint tissues are strikingly different. Specifically, normal flexion primarily induces articular cartilage erosion and osteophyte formation at the medial tibial plateau. In deep flexion, articular cartilage erosion and osteophyte formation mainly occur in the lateral femoral condyle. The differences likely reflect the substantial influence of mechanical loading regimens and joint kinematics on the biological response of the knee joint. In addition to differences in the degree of knee flexion, loading waveforms, and experimental durations, the knee joint kinematics may influence disease initiation and progression in both configurations.

Knee Kinematics and Joint Stresses. The mechanics of the normal and deep flexion configurations of the load-induced OA model have been examined using rigid body kinematics and computational modeling, respectively. For normal flexion, the

mechanics of the loading model were characterized using quasi-static X-ray imaging, with roentgen stereophotogrammetric analysis [65]. Under controlled compressive loading, the tibia reproducibly translated primarily proximally and anteriorly, with minimal medial–lateral translation and minimal rotation about all axes. Furthermore, the femoral condyles did not translate relative to the tibia, but rather increased knee flexion by approximately 7 deg as the load magnitude increased to 9 N. Thus, in the normal flexion configuration, controlled tibial loading produced compressive and shear forces to the proximal tibial articulating surface through translation along the proximal and anterior directions, and flexion of the knee (Fig. 5). These findings correlated with the thinning of the cartilage surface in the posterior tibia observed within this model. Furthermore, lateral motion was minimal during loading, confirming that stabilizing ligaments remained intact.

In the deep flexion configuration, contact stresses in the femoral cartilage during compressive loading were characterized using a finite element model and quasi-static X-ray imaging measurements [66]. The contact area was reproducible with deep flexion loading, allowing for reliable predictive models of cartilage damage. Furthermore, whereas the magnitudes of the contact pressures were similar in the tibia and femur (14.5 MPa peak stress), the contact area was smaller in the femur. Such differences in contact area suggest a stress concentration during cyclic loading and may explain the initiation of spontaneous cartilage lesions on the femoral condyles after loading with the deep flexion configuration. Further, the contact pressures measured are within the range

Fig. 5 The load-induced OA model demonstrates controlled and consistent kinematics and contact mechanics. In the normal flexion configuration, the tibia translates in the proximal and anterior directions and knee flexion increases with applied loading.

Load-Induced Osteoarthritis

Fig. 6 Schematic of cellular and molecular changes in load-induced OA progression. Cyclic tibial compression induces a cascade of destructive events including active aggrecanases and collagenases that break down key components of the cartilage matrix, resulting in increased production of COMP. Subchondral bone changes and synovium cell proliferation also occur in the load-induced OA model. Proteins shown are likely involved in OA initiation and progression but have not been confirmed in the load-induced OA model including active MMPs, IL-1, TNF α , and TGF- β (red). Adapted from Goldring and Marcu [71].

of the failure strength of cartilage in other cyclic compression studies (3–50 MPa) [67–69], thus explaining the resulting cartilage damage. Performing similar studies in the normal loading configuration would enable correlation of areas of high stresses with areas of cartilage damage. Understanding contact mechanics will enable more robust comparisons of the normal and deep flexion configurations and may explain the different biological responses with each loading approach.

Understanding Disease Pathology With Load-Induced Osteoarthritis

Although differences between the normal and deep flexion configurations are evident, both approaches enable exploration of genetic and molecular pathways, and potential treatments for load-induced OA. Cyclic tibial compression has been applied to different mouse genetic backgrounds/strains to better understand the role of cartilage and bone tissue properties in disease progression [17,66,70]. Future studies could explore changes in the progression of load-induced OA in mice with altered cellular and genetic pathways.

Elucidating Mechanobiological Mechanisms of Load-Induced Osteoarthritis. The load-induced OA model allows us to directly relate areas of acute cartilage damage to specific cellular signaling in response to loading (Fig. 6). To this end, cellular responses to mechanical loading in the model have been characterized [19,47,72]. Specifically, a single bout of loading has been used to elucidate the cellular and tissue mechanism by which OA initiates in both configurations. In normal flexion, a single bout of 1200 cycles of compressive loading did not induce spontaneous cartilage lesions, but rather initiated cell-mediated processes that led to OA without additional injurious loading [72]. Immediately after loading, cartilage appeared normal in both control and loaded limbs. However, after 1 and 2 weeks of normal activity, cartilage damage, cartilage thinning, and pre-osteophyte formation were present at the medial aspect of the tibia plateau. In addition, loading induced thinning of the subchondral bone plate in

loaded limbs 1 and 2 weeks after the loading bout. Based on immunohistochemical staining, chondrocyte apoptosis was not different between control and loaded joints, suggesting loading had no acute effects on chondrocyte viability. Furthermore, synovial hyperplasia was not evident after the single loading bout. However, osteoclast activity in the underlying subchondral bone increased after loading, explaining the decreased subchondral bone plate thickness. Ultimately, because hyperplasia and chondrocyte apoptosis did not change with loading, the tissue response to loading was most likely mediated by cells resident in the joint.

In contrast to the normal flexion configuration, a single bout of loading in deep flexion initiated spontaneous lesions and clefts in the articular cartilage surface [19]. Left untreated, the lesions remained but did not progress further; however, when loading continued after the single bout, the cartilage lesions grew, subsequently leading to more severe OA-like pathology.

Using the deep flexion configuration, OA pathology was investigated in 8-week old C57Bl/6 mice after single loading bouts of 60 cycles with magnitudes of 3 N, 6 N, and 9 N [47]. In contrast to prior studies, the 9 N load level ruptured the ACL, demonstrating the result of a slight variation in joint angle or loading protocol in the mechanical response of the joint tissues. With 3 N and 6 N loads, articular cartilage lesions occurred primarily on the lateral condyle of the femur. Lesion size increased with load magnitude but did not increase with loading duration from 5 to 14 days after the loading bout.

In addition to examining the morphological changes that occurred in the cartilage in response to loading, a few key cellular factors have been related to cartilage degradation [47]. Chondrocyte apoptosis increased with loading and was colocalized with the cartilage lesion. In addition, synovitis scores increased with load magnitude. Within the cartilage, loading had little to no effect on type II collagen amount or location, but significantly changed the amount and distribution of aggrecan. In addition, the distribution of cartilage oligomeric matrix protein (COMP) extended to the middle zone of the cartilage in loaded limbs, but was only present in the superficial zone in control limbs. Chondrocytes in deeper zones and synovial cells produced COMP in response to mechanical stress, suggesting that COMP plays an important role in load-induced cartilage degradation.

Altered Tissue Material Properties. The role of subchondral bone tissue material properties in OA has been of interest in OA research for many years [73,74]. In general, increased bone mass and stiffness have been assumed to increase stresses in cartilage with loading and contribute to the development of OA. These questions can be addressed directly in these loading-based mouse models through manipulation of tissue properties with genetics or drug treatment.

The role of altered subchondral bone properties in load-induced OA was investigated using the normal flexion configuration with two mouse strains: the C57Bl/6 strain with low bone mass and the FVB strain with high bone mass [70]. The tibiae of FVB mice had a thicker subchondral bone plate and higher epiphyseal cancellous bone mass and tissue mineral density compared to C57Bl/6 mice. In response to loading, FVB mice developed less severe and less progressive cartilage damage and osteophytes (Fig. 7). Lower bone mass attenuated load-induced cartilage damage. However, FVB mice also had thicker articular cartilage compared to C57Bl/6 mice. Thicker cartilage has been associated with lower contact stresses during tibial compression and therefore may have contributed to the lower load-induced cartilage damage [66]. Other mouse-strain specific differences could also contribute to this result. Nonetheless, future work needs to investigate correlations between subchondral bone properties and the severity of load-induced OA pathology.

In addition to bone, cartilage tissue material properties also may play a role in OA development and progression. The *cho/+* mouse is haploinsufficient for *Col11a1*, has abnormal collagen II

Fig. 7 Cartilage matrix changes in the tibia after cyclic tibial compression in mice with (a) altered subchondral bone [70] and (b) altered cartilage tissue material properties [75]. In both studies, increased cartilage thickness resulted in decreased load-induced OA severity, regardless of tissue material properties. (a) The FVB strain has higher bone mass compared to the C57Bl/6 (B6) strain. The FVB strain developed less severe load-induced cartilage damage compared to B6 knees. (b) The cho/+ mouse developed spontaneous OA at 6 months of age, indicated by decreased proteoglycan content compared to wildtype (WT) littermates [75]. However, cho/+ developed less severe load-induced OA after 6 weeks of 9 N loading compared to WT. Scale bars = 50 μ m. Figure adapted from Adebayo et al. [70] and Holyoak et al. [75].

fibrils, and has less stiff tissue material properties in the cartilage matrix [76]. The role of these altered cartilage tissue material properties in load-induced OA was investigated using the normal flexion loading configuration with cho/+ mice and littermate controls [75]. Compared to littermate controls, cho/+ mice developed less severe load-induced cartilage damage (Fig. 7). Cho/+ mice also had thicker cartilage and thinner, less dense subchondral cortical bone. The differences in cartilage thickness and subchondral bone may have contributed to the reduced severity of load-induced cartilage damage. Based on the results from studies with altered bone [70] and cartilage [75] material properties, increased cartilage thickness may play a protective role in load-induced OA progression. Further research is required to determine the specific tissue properties that most strongly influence load-induced OA progression.

Bone and cartilage properties could be altered using a variety of other approaches. For example, mice deficient in collagen IX spontaneously developed OA-like features in their cartilage matrix [77]. Understanding the effects of collagen IX and XI and other protein deficiencies in the cartilage matrix on the severity of load-induced OA may elucidate the mechanical function of each protein in the cartilage matrix. Osteoporosis treatments alter bone mass and will alter the stresses in the joint with loading [78]. Furthermore, these treatments are FDA-approved and could be used to modulate cartilage properties clinically, as described below. Overall, the role of altered bone and cartilage tissue properties in load-induced OA initiation and progression can be determined from these studies, allowing potential targeted treatment options to be identified. Finally, bone and cartilage properties are focused on here, and the role of other joint tissues such as the meniscus or synovium should be explored.

Altered Cellular and Genetic Pathways. Understanding the role of genetic pathways in load-induced OA pathology may aid in developing therapies. The Str/ort mouse is a genetically unique strain that spontaneously develops knee OA lesions at skeletal maturity [66]. When cyclic tibial compression was applied using the deep flexion configuration, Str/ort mice did not have enhanced vulnerability to mechanical damage and did not develop load-induced OA. These results were similar to cho/+ mice loaded in normal flexion, which produced less severe load-induced OA despite having abnormal collagen II fibrils. Understanding the Str/ort pathophysiology may help to differentiate spontaneous and load-induced OA mechanisms. Similar approaches with other mouse strains and altered genetics may help to determine the role of genetics in load-induced OA progression.

Key signaling pathways, such as the Wnt, bone morphogenetic protein, transforming growth factor- β , estrogen receptor- α , matrix metalloproteinase, and vascular endothelial growth factor pathways, could be examined to elucidate their role in load-induced OA progression. Specifically, overexpression, knocking down or out, or altering candidate genes in the mouse would allow for in-depth characterization of OA pathophysiology with in vivo loading. These studies would provide valuable knowledge about OA initiation at the cellular and molecular level and potentially reveal critical signaling pathways for treatment options. For example, loading of transgenic mice could be used to investigate molecules involved in the inhibition of the Wnt signaling pathway. Although the Wnt pathway plays a key role in joint development and maintenance [79], elevated levels of Wnt antagonists have been associated with less severe OA [80–82]. Inhibiting or attenuating Wnt signaling could test whether inhibitors of this pathway are viable for preventing or treating load-induced OA.

The role of inflammation and obesity in exacerbating load-induced OA progression should also be explored. With a growing obese population accounting for a large portion of the rise in the prevalence of OA [83], the relationship between obesity, systemic inflammation, and OA development and progression needs to be established. Naturally occurring obesity in mice can lead to systemic inflammation, either by genetic manipulation (*ob/ob* mouse) [84] or modification of diet [85]. Tibial loading at moderate load magnitudes led to more severe load-induced cartilage damage in mice with severe obesity due to high-fat diet compared to mice on a normal diet. However, load-induced OA severity was not exacerbated with moderate obesity due to metabolic syndrome resulting from an altered gut microbiome [86]. Future experiments need to examine not only the interactions of loading, obesity, and systemic inflammation in inducing OA, but also the underlying mechanisms. Ultimately, an understanding is needed not only of clinically relevant aspects of disease pathology, but also of potential preventative measures and treatment options.

Testing Novel Therapeutics With the Load-Induced Osteoarthritis Model

The load-induced OA model provides an *in vivo* platform to test the therapeutic efficacy of traditional and novel treatment strategies for OA [70,87]. For example, viscosupplementation of hyaluronic acid (HA) is a common clinical treatment to delay total joint replacement [88]. Local intra-articular delivery of a HA derivative had no effect on load-induced cartilage proteoglycan loss, chondrocyte apoptosis, or synovitis after tibial compression in deep flexion at 6 N [87]. However, the HA derivative had subtle protective effects on load-induced changes in cancellous bone. Alendronate (ALN) is a bisphosphonate that inhibits bone remodeling and is commonly used for osteoporosis treatment, but the inhibition of bone remodeling also may be effective for OA treatment [74,89]. Systemic delivery of ALN had no effect on load-induced cartilage degradation or bone changes after tibial compression in normal flexion at 9 N in C57BL/6 mice. However, ALN treatment attenuated the progression of OA in FVB mice, a strain with higher bone mass. The results of both the HA and ALN studies were counter to the hypotheses and suggest that other therapeutic strategies may be necessary to slow the progression of load-induced OA.

Finally, novel biomaterials-based and therapeutic approaches can also be tested with load-induced OA. Recent strategies to improve drug delivery include microparticles, nanoparticles, and hydrogels [90]. Micro and nanoparticles can be engineered to target localized delivery to the joint tissues, resulting in longer retention and higher efficacy of drugs [17,91,92]. Injectable hydrogels provide a unique platform with tunable mechanical properties and degradation profiles [93,94]. In addition, the most recent advances in OA-targeted therapeutics, including green fluorescent proteins [95] and biomimetic proteoglycans [96], can be tested in isolation or in combination with drug delivery systems. In summary, the combination of therapeutics and biomaterials-based intra-articular drug delivery systems need to be examined in load-induced OA.

Limitations of the Load-Induced Osteoarthritis Model

Preclinical models are valuable tools to study disease pathology and emerging treatments. By definition all models have limitations that are outweighed by their strengths and/or limit the specific application. The load-induced OA model recapitulates the progression of OA-like pathology in humans based on the degradation of cartilage, formation of osteophytes, changes in subchondral bone, and other tissue/molecular results discussed in this review. However, one must consider differences in human and mouse cartilage physiology and structure such as degree of cellularity and zonal structure. The specific loading conditions of applied to the mouse tibia also cannot be directly translated to the human knee. For example, 1200 cycles of daily physiological loading will not result in human OA. In the mouse, a OA-like pathology results from a combination of factors in the experiments, including the direction of the applied load, the angle of the mouse knee, and the applied load waveform.

Although cyclic tibial compression has been extensively used to study the detrimental effects of loading, dynamic loading can also be beneficial to joint health. Dynamic low-level knee joint loading in the mediolateral direction attenuated post-traumatic OA in mice [97]. In contrast, unloading the joint resulted in cartilage thinning [98], but unloading knee braces have been associated with a delay for the need of surgery in humans [99]. Similarly, static loading did not result in significant cartilage degradation in mouse due to the viscoelastic nature of cartilage [100]. In addition, dynamic loading is used in cartilage tissue engineering to encourage fluid/nutrient flow and to regenerate cartilage tissue [101]. Clearly, a wide spectrum of loading conditions exists, and each loading condition results in a unique physiological response.

To date, the study of load-induced OA progression in mice has been limited to the knee. Other joints, including the hip, ankle, temporomandibular joint, and spinal facets may respond differently to cyclic compression. The normal kinematics of these other joints with loading differs considerably from the knee. Therefore, the results discussed in this review cannot be generalized to other joints.

Summary and Conclusions

Osteoarthritis pathology has been studied in several invasive and noninvasive *in vivo* models including surgical injury, chemical injection, spontaneous OA, impact injury, and load-induced OA models. Models need to be considered in terms of their advantages; load-induced OA models are valuable tools to explore the relationship between subfailure mechanical forces and OA progression, without confounding and invasive surgery [31]. The application of tibial compression in the load-induced OA model provides controlled, consistent kinematics to the joint that lead to cellular responses and subsequent progression of OA-like pathology (Fig. 8). Two load-induced OA configurations have been established that primarily differ in the joint angle at the knee when loads are applied. Subtle differences in the outcome of loading at both the tissue and cellular levels between these

Fig. 8 Summary of load-induced OA model. Cyclic compressive loading of the mouse tibia produces consistent kinematics and contact stresses at the knee joint, initiating robust cellular and tissue-level responses that result in OA-like pathology in the tissues of the knee joint.

configurations will be critical to relate the joint kinematics, tissue stresses, and biological mechanisms associated with load-induced joint pathology. Multiple studies have begun to characterize the mechanobiology of load-induced OA progression. Future work will elucidate specific mechanobiological mechanisms and continue to explore the cellular and molecular pathways of OA initiation and progression. Considering the joint as a system to understand the role of altered tissue properties and genetic pathways in disease progression will enable potential treatment or prevention options for OA.

Acknowledgment

Support for the work summarized in this review was provided by NIH R21-AR064034, NSF GRFP, Sloan Foundation, U.S. Department of Education GAANN Fellowship (P200A150273), Robert and Helen Appel Fellowship, and the Clark and Kirby Foundations.

Funding Data

- Sloan Foundation (Funder ID: 10.13039/100000879).
- Foundation for the National Institutes of Health (Grant No. NIH R21-AR064034; Funder ID: 10.13039/100000009).
- National Science Foundation (Funder ID: 10.13039/100000001).
- U.S. Department of Education GAANN Fellowship (Grant No. P200A150273; Funder ID: 10.13039/100000138).

References

- [1] Lawrence, R. C., Felson, D. T., Helmick, C. G., Arnold, L. M., Choi, H., Deyo, R. A., Gabriel, S., Hirsch, R., Hochberg, M. C., Hunder, G. G., Jordan, J. M., Katz, J. N., Kremers, H. M., and Wolfe, F., 2008, "Estimates of the Prevalence of Arthritis and Other Rheumatic Conditions in the United States—Part II," *Arthritis Rheumatol.*, **58**(1), pp. 26–35.
- [2] Hootman, J. M., and Helmick, C. G., 2006, "Projections of US Prevalence of Arthritis and Associated Activity Limitations," *Arthritis Rheumatol.*, **54**(1), pp. 226–229.
- [3] Felson, D. T., 1988, "Epidemiology of Hip and Knee Osteoarthritis," *Epidemiol. Rev.*, **10**, pp. 1–28.
- [4] Wieland, H. A., Michaelis, M., Kirschbaum, B. J., and Rudolph, K. A., 2005, "Osteoarthritis—An Untreatable Disease?," *Nat. Rev. Drug Discov.*, **4**(4), pp. 331–344.
- [5] Hunter, D. J., and Felson, D. T., 2006, "Osteoarthritis," *BMJ*, **332**(7542), pp. 639–642.
- [6] Felson, D. T., Anderson, J. J., Naimark, A., Walker, A. M., and Meenan, R. F., 1988, "Obesity and Knee Osteoarthritis," *Ann. Intern. Med.*, **109**(1), p. 18.
- [7] Messier, S. P., Loeser, R. F., Miller, G. D., Morgan, T. M., Rejeski, W. J., Sevcik, M. A., Ettinger, W. H., Pahor, M., and Williamson, J. D., 2004, "Exercise and Dietary Weight Loss in Overweight and Obese Older Adults With Knee Osteoarthritis: The Arthritis, Diet, and Activity Promotion Trial," *Arthritis Rheumatol.*, **50**(5), pp. 1501–1510.
- [8] Buckwalter, J. A., Saltzman, C., and Brown, T., 2004, "The Impact of Osteoarthritis: Implications for Research," *Clin. Orthop. Relat. Res.*, **427**, pp. S6–S15.
- [9] Kaila-Kangas, L., Arokoski, J., Impivaara, O., Viikari-Juntura, E., Leino-Arjas, P., Luukkainen, R., and Heliövaara, M., 2011, "Associations of Hip Osteoarthritis With History of Recurrent Exposure to Manual Handling of Loads Over 20 kg and Work Participation: A Population-Based Study of Men and Women," *Occup. Environ. Med.*, **68**(10), pp. 734–738.
- [10] Cameron, K. L., Hsiao, M. S., Owens, B. D., Burks, R., and Svoboda, S. J., 2011, "Incidence of Physician-Diagnosed Osteoarthritis Among Active Duty United States Military Service Members," *Arthritis Rheumatol.*, **63**(10), pp. 2974–2982.
- [11] Felson, D. T., Lawrence, R. C., Dieppe, P. A., Hirsch, R., Helmick, C. G., Jordan, J. M., Kingdon, R. S., Lane, N. E., Nevitt, M. C., Zhang, Y., Sowers, M., McAlindon, T., Spector, T. D., Poole, A. R., Yanovski, S. Z., Ateshian, G., Sharma, L., Buckwalter, J. A., Brandt, K. D., and Fries, J. F., 2000, "Osteoarthritis: New Insights—Part I: The Disease and Its Risk Factors," *Ann. Intern. Med.*, **133**(8), p. 635.
- [12] Lohmander, L. S., Englund, P. M., Dahl, L. L., and Roos, E. M., 2007, "The Long-Term Consequence of Anterior Cruciate Ligament and Meniscus Injuries: Osteoarthritis," *Am. J. Sports Med.*, **35**(10), pp. 1756–1769.
- [13] Mobasheri, A., Rayman, M. P., Gualillo, O., Sellam, J., Van Der Kraan, P., and Fearon, U., 2017, "The Role of Metabolism in the Pathogenesis of Osteoarthritis," *Nat. Rev. Rheumatol.*, **13**(5), pp. 302–311.
- [14] Arokoski, J., Kiviranta, I., Jurvelin, J., Tamm, M., and Helminen, H. J., 1993, "Long-Distance Running Causes Site-Dependent Decrease of Cartilage

- Glycosaminoglycan Content in the Knee Joints of Beagle Dogs," *Arthritis Rheumatol.*, **36**(10), pp. 1451–1459.
- [15] Lapveteläinen, T., Nevalainen, T., Parkkinen, J. J., Arokoski, J., Kiraly, K., Hyttinen, M., Halonen, P., and Helminen, H. J., 1995, "Lifelong Moderate Running Training Increases the Incidence and Severity of Osteoarthritis in the Knee Joint of C57BL Mice," *Anat. Rec.*, **242**(2), pp. 159–165.
- [16] Christiansen, B. A., Guilak, F., Lockwood, K. A., Olson, S. A., Pitsillides, A. A., Sandell, L. J., Silva, M. J., van der Meulen, M. C. H., and Haudenschild, D. R., 2015, "Non-Invasive Mouse Models of Post-Traumatic Osteoarthritis," *Osteoarthritis Cartilage*, **23**(10), pp. 1627–1638.
- [17] Holyoak, D. T., Tian, Y. F., van der Meulen, M. C. H., and Singh, A., 2016, "Osteoarthritis: Pathology, Mouse Models, and Nanoparticle Injectable Systems for Targeted Treatment," *Ann. Biomed. Eng.*, **44**(6), pp. 2062–2075.
- [18] Bende, A. M., 2001, "Animal Models of Osteoarthritis," *J. Musculoskeletal Neuronal Interact.*, **1**(4), pp. 363–376.
- [19] Poulet, B., Hamilton, R. W., Shefelbine, S., and Pitsillides, A. A., 2011, "Characterizing a Novel and Adjustable Noninvasive Murine Joint Loading Model," *Arthritis Rheumatol.*, **63**(1), pp. 137–147.
- [20] Ko, F. C., Dragomir, C., Plumb, D. A., Goldring, S. R., Wright, T. M., Goldring, M. B., and van der Meulen, M. C. H., 2013, "In Vivo Cyclic Compression Causes Cartilage Degeneration and Subchondral Bone Changes in Mouse Tibiae," *Arthritis Rheumatol.*, **65**(6), pp. 1569–1578.
- [21] Bajpayee, A. G., Scheu, M., Grodzinsky, A. J., and Porter, R. M., 2015, "A Rabbit Model Demonstrates the Influence of Cartilage Thickness on Intra-Articular Drug Delivery and Retention Within Cartilage," *J. Orthop. Res.*, **33**(5), pp. 660–667.
- [22] Gregory, M. H., Capito, N., Kuroki, K., Stoker, A. M., Cook, J. L., and Sherman, S. L., 2012, "A Review of Translational Animal Models for Knee Osteoarthritis," *Arthritis*, **2012**, p. 764621.
- [23] van der Kraan, P. M., Vitters, E. L., van de Putte, L. B., and van den Berg, W. B., 1989, "Development of Osteoarthritic Lesions in Mice by 'Metabolic' and 'Mechanical' Alterations in the Knee Joints," *Am. J. Pathol.*, **135**(6), pp. 1001–1014.
- [24] Kikuchi, T., Sakuta, T., and Yamaguchi, T., 1998, "Intra-Articular Injection of Collagenase Induces Experimental Osteoarthritis in Mature Rabbits," *Osteoarthritis Cartilage*, **6**(3), pp. 177–186.
- [25] van der Kraan, P. M., Vitters, E. L., van Beuningen, H. M., van de Putte, L. B., and van den Berg, W. B., 1990, "Degenerative Knee Joint Lesions in Mice After a Single Intra-Articular Collagenase Injection: A New Model of Osteoarthritis," *J. Exp. Pathol. (Oxford)*, **71**(1), pp. 19–31.
- [26] van Beuningen, H., Glansbeek, H., van der Kraan, P., and van den Berg, W., 2000, "Osteoarthritis-Like Changes in the Murine Knee Joint Resulting From Intra-Articular Transforming Growth Factor- β Injections," *Osteoarthritis Cartilage*, **8**(1), pp. 25–33.
- [27] Clements, K. M., Ball, A. D., Jones, H. B., Brinckmann, S., Read, S. J., and Murray, F., 2009, "Cellular and Histopathological Changes in the Infrapatellar Fat Pad in the Monoiodoacetate Model of Osteoarthritis Pain," *Osteoarthritis Cartilage*, **17**(6), pp. 805–812.
- [28] Guzman, R. E., Evans, M. G., Bove, S., Morenko, B., and Kilgore, K., 2003, "Mono-Iodoacetate-Induced Histologic Changes in Subchondral Bone and Articular Cartilage of Rat Femorotibial Joints: An Animal Model of Osteoarthritis," *Toxicol. Pathol.*, **31**(6), pp. 619–624.
- [29] Bove, S. E., Calcaterra, S. L., Brooker, R. M., Huber, C. M., Guzman, R. E., Juneau, P. L., Schrier, D. J., and Kilgore, K. S., 2003, "Weight Bearing as a Measure of Disease Progression and Efficacy of Anti-Inflammatory Compounds in a Model of Monosodium Iodoacetate-Induced Osteoarthritis," *Osteoarthritis Cartilage*, **11**(11), pp. 821–830.
- [30] Combe, R., Bramwell, S., and Field, M. J., 2004, "The Monosodium Iodoacetate Model of Osteoarthritis: A Model of Chronic Nociceptive Pain in Rats?," *Neurosci. Lett.*, **370**(2–3), pp. 236–240.
- [31] Little, C. B., and Zaki, S., 2012, "What Constitutes an 'Animal Model of Osteoarthritis'—The Need for Consensus?," *Osteoarthritis Cartilage*, **20**(4), pp. 261–267.
- [32] Inoue, S., and Glimcher, M. J., 1982, "The Reaction of Cartilage and Osteophyte Formation After the Intraarticular Injection of Papain," *Nihon Seikeigeka Gakkai Zasshi*, **56**(5), pp. 415–430.
- [33] Murat, M., Karadam, B., Ozkal, S., Karatosun, V., and Gidener, S., 2007, "Quantification of Papain-Induced Rat Osteoarthritis in Relation to Time With the Mankin Score," *Acta Orthop. Traumatol. Turc.*, **41**(3), pp. 233–237.
- [34] Bentley, G., 1971, "Papain-Induced Degenerative Arthritis of the Hip in Rabbits," *J. Bone Jt. Surg. Br.*, **53**(2), pp. 324–337.
- [35] Pond, M. J., and Nuki, G., 1973, "Experimentally-Induced Osteoarthritis in the Dog," *Ann. Rheumatol. Dis.*, **32**(4), pp. 387–388.
- [36] Kamekura, S., Hoshi, K., Shimoaka, T., Chung, U., Chikuda, H., Yamada, T., Uchida, M., Ogata, N., Seichi, A., Nakamura, K., and Kawaguchi, H., 2005, "Osteoarthritis Development in Novel Experimental Mouse Models Induced by Knee Joint Instability," *Osteoarthritis Cartilage*, **13**(7), pp. 632–641.
- [37] Hayami, T., Pickarski, M., Zhuo, Y., Wesolowski, G. A., Rodan, G. A., and Duong, L. T., 2006, "Characterization of Articular Cartilage and Subchondral Bone Changes in the Rat Anterior Cruciate Ligament Transection and Meniscectomized Models of Osteoarthritis," *Bone*, **38**(2), pp. 234–243.
- [38] McLernain, D. D., Appleton, C. T. G., Litchfield, R. B., Pitelka, V., Henry, J. L., Bernier, S. M., Beier, F., and Holdsworth, D. W., 2008, "Study of Subchondral Bone Adaptations in a Rodent Surgical Model of OA Using In Vivo Micro-Computed Tomography," *Osteoarthritis Cartilage*, **16**(4), pp. 458–469.

- [39] Glasson, S. S., Blanchet, T. J., and Morris, E. A., 2007, "The Surgical Destabilization of the Medial Meniscus (DMM) Model of Osteoarthritis in the 129/SvEv Mouse," *Osteoarthritis Cartilage*, **15**(9), pp. 1061–1069.
- [40] Culley, K. L., Dragomir, C. L., and Chang, J., 2015, *Mouse Models of Osteoarthritis: Surgical Model of Posttraumatic Osteoarthritis Induced by Destabilization of the Medial Meniscus*, Humana Press, New York, pp. 143–173.
- [41] Poulet, B., 2016, "Non-Invasive Loading Model of Murine Osteoarthritis," *Curr. Rheumatol. Rep.*, **18**(7), p. 40.
- [42] McCoy, A. M., 2015, "Animal Models of Osteoarthritis: Comparisons and Key Considerations," *Vet. Pathol.*, **52**(5), pp. 803–818.
- [43] Jimenez, P. A., Glasson, S. S., Trubetskoy, O. V., and Haimes, H. B., 1997, "Spontaneous Osteoarthritis in Dunkin Hartley Guinea Pigs: Histologic, Radiologic, and Biochemical Changes," *Lab Anim. Sci.*, **47**(6), pp. 598–601.
- [44] Furman, B. D., Strand, J., Hembree, W. C., Ward, B. D., Guilak, F., and Olson, S. A., 2007, "Joint Degeneration Following Closed Intraarticular Fracture in the Mouse Knee: A Model of Posttraumatic Arthritis," *J. Orthop. Res.*, **25**(5), pp. 578–592.
- [45] Lewis, J. S., Hembree, W. C., Furman, B. D., Tippets, L., Cattel, D., Huebner, J. L., Little, D., DeFrate, L. E., Kraus, V. B., Guilak, F., and Olson, S. A., 2011, "Acute Joint Pathology and Synovial Inflammation is Associated With Increased Intra-Articular Fracture Severity in the Mouse Knee," *Osteoarthritis Cartilage*, **19**(7), pp. 864–873.
- [46] Christiansen, B. A., Anderson, M. J., Lee, C. A., Williams, J. C., Yik, J. H. N., and Haudenschild, D. R., 2012, "Musculoskeletal Changes Following Non-Invasive Knee Injury Using a Novel Mouse Model of Post-Traumatic Osteoarthritis," *Osteoarthritis Cartilage*, **20**(7), pp. 773–782.
- [47] Wu, P., Holguin, N., Silva, M. J., Fu, M., Liao, W., and Sandell, L. J., 2014, "Early Response of Mouse Joint Tissue to Noninvasive Knee Injury Suggests Treatment Targets," *Arthritis Rheumatol.*, **66**(5), pp. 1256–1265.
- [48] Stoop, R., van der Kraan, P. M., Buma, P., Hollander, A. P., Billingham, R. C., Poole, A. R., and van den Berg, W. B., 1999, "Type II Collagen Degradation in Spontaneous Osteoarthritis in C57BL/6 and BALB/c Mice," *Arthritis Rheumatol.*, **42**(11), pp. 2381–2389.
- [49] Esapa, C. T., Hough, T. A., Testori, S., Head, R. A., Crane, E. A., Chan, C. P., Evans, H., Bassett, J. D., Tylzanowski, P., McNally, E. G., Carr, A. J., Boyde, A., Howell, P. G., Clark, A., Williams, G. R., Brown, M. A., Croucher, P. I., Nesbit, M. A., Brown, S. D., Cox, R. D., Cheeseman, M. T., and Thakker, R. V., 2012, "A Mouse Model for Spondyloepiphyseal Dysplasia Congenita With Secondary Osteoarthritis Due to a Col2a1 Mutation," *J. Bone Miner. Res.*, **27**(2), pp. 413–428.
- [50] Vasheghani, F., Zhang, Y., Li, Y.-H., Blati, M., Fahmi, H., Lussier, B., Roughley, P., Lagares, D., Endisha, H., Saffar, B., Lajeunesse, D., Marshall, W. K., Rampersaud, Y. R., Mahomed, N. N., Gandhi, R., Pelletier, J.-P., Martel-Pelletier, J., and Kapoor, M., 2015, "PPAR γ Deficiency Results in Severe, Accelerated Osteoarthritis Associated With Aberrant mTOR Signaling in the Articular Cartilage," *Ann. Rheumatol. Dis.*, **74**(3), pp. 569–578.
- [51] Fritton, J. C., Myers, E. R., Wright, T. M., and Van Der Meulen, M., 2005, "Loading Induces Site-Specific Increases in Mineral Content Assessed by Microcomputed Tomography of the Mouse Tibia," *Bone*, **36**(6), pp. 1030–1038.
- [52] De Souza, R. L., Matsuura, M., Eckstein, F., Rawlinson, S. C. F., Lanyon, L. E., and Pitsillides, A. A., 2005, "Non-Invasive Axial Loading of Mouse Tibiae Increases Cortical Bone Formation and Modifies Trabecular Organization: A New Model to Study Cortical and Cancellous Compartments in a Single Loaded Element," *Bone*, **37**(6), pp. 810–818.
- [53] Lee, K., Maxwell, A., and Lanyon, L., 2002, "Validation of a Technique for Studying Functional Adaptation of the Mouse Ulna in Response to Mechanical Loading," *Bone*, **31**(3), pp. 407–412.
- [54] Lynch, M. E., Main, R. P., Xu, Q., Schmicker, T. L., Schaffler, M. B., Wright, T. M., and van der Meulen, M. C. H., 2011, "Tibial Compression is Anabolic in the Adult Mouse Skeleton Despite Reduced Responsiveness With Aging," *Bone*, **49**(3), pp. 439–446.
- [55] Sugiyama, T., Saxon, L. K., Zaman, G., Moustafa, A., Sinters, A., Price, J. S., and Lanyon, L. E., 2008, "Mechanical Loading Enhances the Anabolic Effects of Intermittent Parathyroid Hormone (1–34) on Trabecular and Cortical Bone in Mice," *Bone*, **43**(2), pp. 238–248.
- [56] Melville, K. M., Robling, A. G., and van der Meulen, M., 2015, "In Vivo Axial Loading of the Mouse Tibia," *Methods Mol. Biol.*, **1226**, pp. 99–115.
- [57] Mosley, J. R., March, B. M., Lynch, J., and Lanyon, L. E., 1997, "Strain Magnitude Related Changes in Whole Bone Architecture in Growing Rats," *Bone*, **20**(3), pp. 191–198.
- [58] Rubin, C. T., and Lanyon, L. E., 1985, "Regulation of Bone Mass by Mechanical Strain Magnitude," *Calcif. Tissue Int.*, **37**(4), pp. 411–417.
- [59] Poulet, B., de Souza, R., Kent, A. V., Saxon, L., Barker, O., Wilson, A., Chang, Y.-M., Cake, M., and Pitsillides, A. A., 2015, "Intermittent Applied Mechanical Loading Induces Subchondral Bone Thickening That May Be Intensified Locally by Contiguous Articular Cartilage Lesions," *Osteoarthritis Cartilage*, **23**(6), pp. 940–948.
- [60] Astephen Wilson, J. L., Deluzio, K. J., Dunbar, M. J., Caldwell, G. E., and Hubley-Kozey, C. L., 2011, "The Association Between Knee Joint Biomechanics and Neuromuscular Control and Moderate Knee Osteoarthritis Radiographic and Pain Severity," *Osteoarthritis Cartilage*, **19**(2), pp. 186–193.
- [61] Chang, A., Hurwitz, D., Dunlop, D., Song, J., Cahue, S., Hayes, K., and Sharma, L., 2007, "The Relationship Between Toe-Out Angle During Gait and Progression of Medial Tibiofemoral Osteoarthritis," *Ann. Rheumatol. Dis.*, **66**(10), pp. 1271–1275.
- [62] Mündermann, A., Dyrby, C. O., and Andriacchi, T. P., 2005, "Secondary Gait Changes in Patients With Medial Compartment Knee Osteoarthritis: Increased Load at the Ankle, Knee, and Hip During Walking," *Arthritis Rheumatol.*, **52**(9), pp. 2835–2844.
- [63] Miller, R. E., and Malfait, A.-M., 2017, "Osteoarthritis Pain: What are we Learning From Animal Models?," *Best Pract. Res. Clin. Rheumatol.*, **31**(5), pp. 676–687.
- [64] Malfait, A. M., 2017, "Why We Should Study Pain in Animal Models of Rheumatic Diseases," *Clin. Exp. Rheumatol.*, **35**(5), pp. S37–S39.
- [65] Adebayo, O. O., Ko, F. C., Goldring, S. R., Goldring, M. B., Wright, T. M., and van der Meulen, M., 2017, "Kinematics of Meniscal- and ACL-Transected Mouse Knees During Controlled Tibial Compressive Loading Captured Using Roentgen Stereophotogrammetry," *J. Orthop. Res.*, **35**(2), pp. 353–360.
- [66] Poulet, B., Westerhof, T. A. T., Hamilton, R. W., Shelfelbine, S. J., and Pitsillides, A. A., 2013, "Spontaneous Osteoarthritis in Str/Ort Mice is Unlikely Due to Greater Vulnerability to Mechanical Trauma," *Osteoarthritis Cartilage*, **21**(5), pp. 756–763.
- [67] Sadeghi, H., Espino, D. M., and Shepherd, D. E., 2015, "Variation in Viscoelastic Properties of Bovine Articular Cartilage Below, Up to and Above Healthy Gait-Relevant Loading Frequencies," *Proc. Inst. Mech. Eng. Part H*, **229**(2), pp. 115–123.
- [68] Sadeghi, H., Espino, D. M., and Shepherd, D. E. T., 2017, "Fatigue Strength of Bovine Articular Cartilage-on-Bone Under Three-Point Bending: The Effect of Loading Frequency," *BMC Musculoskeletal Disord.*, **18**(1), p. 142.
- [69] Kerin, A. J., Wisnom, M. R., and Adams, M. A., 1998, "The Compressive Strength of Articular Cartilage," *Proc. Inst. Mech. Eng. Part H*, **212**(4), pp. 273–280.
- [70] Adebayo, O. O., Ko, F. C., Wan, P. T., Goldring, S. R., Goldring, M. B., Wright, T. M., and van der Meulen, M. C. H., 2017, "Role of Subchondral Bone Properties and Changes in Development of Load-Induced Osteoarthritis in Mice," *Osteoarthritis Cartilage*, **25**(12), pp. 2108–2118.
- [71] Goldring, M. B., and Marcu, K. B., 2009, "Cartilage Homeostasis in Health and Rheumatic Diseases," *Arthritis Res. Ther.*, **11**(3), p. 224.
- [72] Ko, F. C., Dragomir, C. L., Plumb, D. A., Hsia, A. W., Adebayo, O. O., Goldring, S. R., Wright, T. M., Goldring, M. B., and van der Meulen, M. C. H., 2016, "Progressive Cell-Mediated Changes in Articular Cartilage and Bone in Mice Are Initiated by a Single Session of Controlled Cyclic Compressive Loading," *J. Orthop. Res.*, **34**(11), pp. 1941–1949.
- [73] Radin, E. L., and Rose, R. M., 1986, "Role of Subchondral Bone in the Initiation and Progression of Cartilage Damage," *Clin. Orthop. Relat. Res.*, **213**, pp. 34–40.
- [74] Burr, D. B., and Utreja, A., 2018, "Editorial: Wnt Signaling Related to Subchondral Bone Density and Cartilage Degradation in Osteoarthritis," *Arthritis Rheumatol.*, **70**(2), pp. 157–161.
- [75] Holyoak, D. T., Otero, M., Armar, N. A., Ziemian, S. N., Otto, A., Cullinane, D., Wright, T. M., Goldring, S. R., Goldring, M. B., and van der Meulen, M. C. H., 2018, "Collagen XI Mutation Lowers Susceptibility to Load-Induced Cartilage Damage in Mice," *J. Orthop. Res.*, **36**(2), pp. 711–720.
- [76] Xu, L., Flahiff, C. M., Waldman, B. A., Wu, D., Olsen, B. R., Setton, L. A., and Li, Y., 2003, "Osteoarthritis-Like Changes and Decreased Mechanical Function of Articular Cartilage in the Joints of Mice With the Chondrodysplasia Gene (Cho)," *Arthritis Rheumatol.*, **48**(9), pp. 2509–2518.
- [77] Hu, K., Xu, L., Cao, L., Flahiff, C. M., Brussiau, J., Ho, K., Setton, L. A., Youn, I., Guilak, F., Olsen, B. R., and Li, Y., 2006, "Pathogenesis of Osteoarthritis-Like Changes in the Joints of Mice Deficient in Type IX Collagen," *Arthritis Rheumatol.*, **54**(9), pp. 2891–2900.
- [78] Buckland-Wright, J. C., Messent, E. A., Bingham, C. O., Ward, R. J., and Tonkin, C., 2006, "A 2 yr Longitudinal Radiographic Study Examining the Effect of a Bisphosphonate (Risedronate) Upon Subchondral Bone Loss in Osteoarthritic Knee Patients," *Rheumatology*, **46**(2), pp. 257–264.
- [79] Guo, X., Day, T. F., Jiang, X., Garrett, B. L., Topol, L., and Yang, Y., 2004, "Wnt/Beta-Catenin Signaling is Sufficient and Necessary for Synovial Joint Formation," *Genes Dev.*, **18**(19), pp. 2404–2417.
- [80] Lane, N. E., Nevitt, M. C., Lui, L.-Y., de Leon, P., and Corr, M., 2007, "Wnt Signaling Antagonists are Potential Prognostic Biomarkers for the Progression of Radiographic Hip Osteoarthritis in Elderly Caucasian Women," *Arthritis Rheumatol.*, **56**(10), pp. 3319–3325.
- [81] Hopwood, B., Tsykin, A., Findlay, D. M., and Fazzalari, N. L., 2007, "Microarray Gene Expression Profiling of Osteoarthritic Bone Suggests Altered Bone Remodelling, WNT and Transforming Growth Factor-Beta/Bone Morphogenic Protein Signalling," *Arthritis Res. Ther.*, **9**(5), p. R100.
- [82] Power, J., Poole, K. E. S., van Bezooijen, R., Doube, M., Caballero-Alías, A. M., Lowik, C., Papapoulos, S., Reeve, J., and Loveridge, N., 2010, "Sclerostin and the Regulation of Bone Formation: Effects in Hip Osteoarthritis and Femoral Neck Fracture," *J. Bone Miner. Res.*, **25**(8), pp. 1867–1876.
- [83] Kulkarni, K., Karssiens, T., Kumar, V., and Pandit, H., 2016, "Obesity and Osteoarthritis," *Maturitas*, **89**, pp. 22–28.
- [84] Enser, M., 1972, "Clearing-Factor Lipase in Obese Hyperglycaemic Mice (ob-ob)," *Biochem. J.*, **129**(2), pp. 447–453.
- [85] Surwit, R. S., Kuhn, C. M., Cochrane, C., McCubbin, J. A., and Feinglos, M. N., 1988, "Diet-Induced Type II Diabetes in C57BL/6J Mice," *Diabetes*, **37**(9), pp. 1163–1167.
- [86] Guss, J. D., Ziemian, S. N., Luna, M., Sandoval, T. N., Holyoak, D. T., Guisado, G. G., Roubert, S., Callahan, R. L., Brito, I. L., van der Meulen, M. C. H., Goldring, S. R., and Hernandez, C. J., 2019, "The Effects of Metabolic Syndrome, Obesity, and the Gut Microbiome on Load-Induced Osteoarthritis," *Osteoarthritis Cartilage*, **27**(1), pp. 129–139.

- [87] Duan, X., Sandell, L. J., Chinzei, N., Holguin, N., Silva, M. J., Schiavinato, A., and Rai, M. F., 2017, "Therapeutic Efficacy of Intra-Articular Hyaluronan Derivative and Platelet-Rich Plasma in Mice Following Axial Tibial Loading," *PLoS One*, **12**(4), p. e0175682.
- [88] Altman, R., Lim, S., Steen, R. G., and Dasa, V., 2015, "Hyaluronic Acid Injections Are Associated With Delay of Total Knee Replacement Surgery in Patients With Knee Osteoarthritis: Evidence From a Large U.S. Health Claims Database," *PLoS One*, **10**(12), p. e0145776.
- [89] Hayami, T., Pickarski, M., Wesolowski, G. A., McLane, J., Bone, A., Destefano, J., Rodan, G. A., and Duong, L. T., 2004, "The Role of Subchondral Bone Remodeling in Osteoarthritis: Reduction of Cartilage Degeneration and Prevention of Osteophyte Formation by Alendronate in the Rat Anterior Cruciate Ligament Transection Model," *Arthritis Rheumatol.*, **50**(4), pp. 1193–1206.
- [90] Kang, M. L., and Im, I., 2014, "Drug Delivery Systems for Intra-Articular Treatment of Osteoarthritis," *Expert Opin. Drug Deliv.*, **11**(2), pp. 269–282.
- [91] Whitmire, R. E., Scott Wilson, D., Singh, A., Levenston, M. E., Murthy, N., and García, A. J., 2012, "Self-Assembling Nanoparticles for Intra-Articular Delivery of Anti-Inflammatory Proteins," *Biomaterials*, **33**(30), pp. 7665–7675.
- [92] Singh, A., Agarwal, R., Diaz-Ruiz, C. A., Willett, N. J., Wang, P., Lee, L. A., Wang, Q., Guldberg, R. E., and García, A. J., 2014, "Nanoengineered Particles for Enhanced Intra-Articular Retention and Delivery of Proteins," *Adv. Healthcare Mater.*, **3**(10), pp. 1562–1567.
- [93] He, Z., Wang, B., Hu, C., and Zhao, J., 2017, "An Overview of Hydrogel-Based Intra-Articular Drug Delivery for the Treatment of Osteoarthritis," *Colloids Surf. B Biointerfaces*, **154**, pp. 33–39.
- [94] Seliktar, D., 2012, "Designing Cell-Compatible Hydrogels for Biomedical Applications," *Science*, **336**(6085), pp. 1124–1128.
- [95] Krishnan, Y., Rees, H. A., Rossitto, C. P., Kim, S.-E., Hung, H.-H. K., Frank, E. H., Olsen, B. D., Liu, D. R., Hammond, P. T., and Grodzinsky, A. J., 2018, "Green Fluorescent Proteins Engineered for Cartilage-Targeted Drug Delivery: Insights for Transport Into Highly Charged Avascular Tissues," *Biomaterials*, **183**, pp. 218–233.
- [96] Phillips, E. R., Prudnikova, K., Bui, T., Taylor, A. J., Galindo, D. A., Huneke, R. B., Hou, J. S., Mulcahey, M. K., and Marcolongo, M. S., 2019, "Biomimetic Proteoglycans Can Molecularly Engineer Early Osteoarthritic Cartilage In Vivo," *J. Orthop. Res.*, **37**(2), pp. 403–411.
- [97] Hamamura, K., Zhang, P., Zhao, L., Shim, J. W., Chen, A., Dodge, T. R., Wan, Q., Shih, H., Na, S., Lin, C. C., Sun, H. B., and Yokota, H., 2013, "Knee Loading Reduces MMP13 Activity in the Mouse Cartilage," *BMC Musculoskeletal Disord.*, **14**(1), p. 312.
- [98] Nomura, M., Sakitani, N., Iwasawa, H., Kohara, Y., Takano, S., Wakimoto, Y., Kuroki, H., and Moriyama, H., 2017, "Thinning of Articular Cartilage After Joint Unloading or Immobilization. An Experimental Investigation of the Pathogenesis in Mice," *Osteoarthritis Cartilage*, **25**(5), pp. 727–736.
- [99] Mistry, D. A., Chandratreya, A., and Lee, P. Y. F., 2018, "An Update on Unloading Knee Braces in the Treatment of Unicompartmental Knee Osteoarthritis From the Last 10 Years: A Literature Review," *Surg. J.*, **4**(3), pp. 110–118.
- [100] Abusara, Z., Von Kossel, M., and Herzog, W., 2016, "In Vivo Dynamic Deformation of Articular Cartilage in Intact Joints Loaded by Controlled Muscular Contractions," *PLoS One*, **11**(1), p. e0147547.
- [101] Correia, C., Pereira, A. L., Duarte, A. R. C., Frias, A. M., Pedro, A. J., Oliveira, J. T., Sousa, R. A., and Reis, R. L., 2012, "Dynamic Culturing of Cartilage Tissue: The Significance of Hydrostatic Pressure," *Tissue Eng.*, **18**(19–20), pp. 1979–1991.