Published in final edited form as:

Ann Surg Oncol. 2019 February ; 26(2): 329–335. doi:10.1245/s10434-018-6866-y.

Multifocal/Multicentric Ipsilateral Invasive Breast Carcinomas with Similar Histology: Is Multigene Testing of All Individual Foci Necessary?

Anne Grabenstetter¹, Edi Brogi¹, Joanne F Chou², Monica Morrow³, Maura Dickler⁴, Larry Norton⁵, and Hannah Y Wen¹

¹Department of Pathology, Memorial Sloan Kettering Cancer Center, New York, NY 10065, USA

²Department of Epidemiology & Biostatistics, Memorial Sloan Kettering Cancer Center, New York, NY 10065, USA

³Department of Surgery, Memorial Sloan Kettering Cancer Center, New York, NY 10065, USA

⁴Lilly Oncology, Eli Lilly and Company, New York, NY 10016, USA

⁵Department of Medicine, Memorial Sloan Kettering Cancer Center, New York, NY 10065, USA

Abstract

Background.—Multiple synchronous ipsilateral invasive breast carcinomas (BCs) with similar histology usually have concordant receptor status. It is unknown whether individual foci with similar histology also share molecular and biological similarities or are heterogeneous. This study examines the concordance of the 21-gene recurrence score (RS) in multiple synchronous morphologically similar ipsilateral BCs.

Methods.—We identified patients with multiple ipsilateral BCs and available RS treated at our institution from 1/2014 to 6/2018. BCs were divided into 3 groups based on RS: 1) RS in same risk category, 2) RS in different risk categories but within 2-unit difference (e.g. RS 17 and RS 19), and 3) RS in different risk categories and a change of >2 units. BCs in groups 1 and 2 were considered concordant (no significant clinical impact) and BCs in group 3 were discordant (variation affects management).

Results.—A total of 53 patients met the study criteria. RS was concordant in 46 (87%) cases. Seven (13%) cases were discordant (group 3). Of these, 3 (43%, 3/7) had biopsy cavity changes (BXC) adjacent to the BC with the highest RS. In two cases the focus with higher RS had a lower percentage of PR-positive tumor cells. In two cases extensive DCIS was associated with BC focus with the lower RS.

Conclusions.—Morphologically similar multifocal ipsilateral BCs have concordant RS in 87% (46/53) of cases. Our results suggest that in cases of morphologically similar multifocal BC testing of a single focus provides accurate prognostic and predictive information.

Corresponding author: Anne Grabenstetter, MD, Department of Pathology, Memorial Sloan Kettering Cancer Center, 1275 York Avenue, New York, NY 10065, Phone: 212-639-4826, grabensa@mskcc.org.

Disclosure/Conflict of Interest

Authors Dr. Monica Morrow and Dr. Maura Dickler have consulted for Genomic Health.

Keywords

Oncotype DX; recurrence score; multifocal/multicentric; breast cancer; 21-gene recurrence score assay; progesterone receptor; biopsy cavity; TAILORx

With the advancement of preoperative imaging, multiple ipsilateral foci of invasive breast carcinoma (BC), defined as multifocal or multicentric carcinoma, are identified more frequently^{1,2}, and the incidence of multifocal/multicentric BC ranges from 6 to 77%³. Multifocal/multicentric carcinoma can be due to intramammary spread of a single primary BC or to multiple synchronous independent primary BCs^{4–8}. There is limited data on whether multiple synchronous tumors of similar histology have significant biological and molecular differences. The 21-gene recurrence score assay (Oncotype DX™, Genomic Health, Redwood City, CA) is a commercially available prognostic and predictive assay which measures the expression levels of 21 genes (16 cancer-related and 5 reference genes) by quantitative reverse transcriptase polymerase chain reaction. The assay generates a numeric recurrence score (RS), on a scale from 0–100, to predict the 10-year risk of distant recurrence and magnitude of chemotherapy benefit in patients with early stage, estrogen receptor-positive (ER+), human epidermal growth factor receptor 2-negative (HER2-) breast cancer. The RS stratifies patients into three risk categories: low (RS <18), intermediate (RS 18–30), and high (RS ≥31)⁹. Adjuvant chemotherapy is added to endocrine therapy for patients with high RS, while the benefit is estimated to be too low to outweigh the side effects in those with low RS¹⁰. The prospective randomized clinical trial TAILORx (Trial Assigning Individualized Options for Treatment) was designed to determine the benefits of adjuvant chemotherapy in patients with BC of intermediate RS¹¹. In the TAILORx study, the risk categories are defined as: low, RS <11, intermediate, RS 11–25, and high, RS >25. These categories differ slightly from the originally defined RS categories and are based on the threshold for recommending adjuvant chemotherapy which is when the risk of local and distant recurrence is 5–10%, the level of risk associated with RS 11¹². The TAILORx trial concluded that there was no chemotherapy benefit for RS <25 in women over the age of 50 and no benefit for RS ≥15 in women 50 years or younger¹³.

The National Comprehensive Cancer Network and American Society of Clinical Oncology recommend using RS to guide adjuvant systemic therapy in patients with ER+, HER2-, lymph node negative invasive BC that are ≤0.5 cm^{14,15}. The importance of genomic assays for risk stratification has been acknowledged in the 8th edition of the American Joint Commission on Cancer (AJCC) staging guidelines¹⁶ which incorporates RS into BC staging. It remains unclear whether the RS of morphologically similar multifocal/multicentric BCs differ significantly to warrant testing of multiple individual foci or if testing of a single representative tumor would provide adequate and accurate information. This study examines the concordance of the RS in multiple synchronous ipsilateral BC with similar histology.

Methods

At our center, all lymph node negative ER+, HER2- invasive BC measuring ≤0.5 cm in patients suitable for chemotherapy are routinely sent for the 21-gene RS assay. The tissue block(s) for testing are selected by the pathologist at the time of initial histopathologic

review. In cases with multiple synchronous foci, a sample of each individual focus is sent for testing if requested by the treating physician.

Through a retrospective search of our pathology database, we identified patients with multiple synchronous ipsilateral BCs who were treated at our institution between January 2014 and June 2018 and had available RS results. Multifocal tumors were defined as infiltrating carcinomas in the same breast, macroscopically distinct and measurable using available clinical and pathologic techniques¹⁷. Clinicopathologic features were obtained from pathology reports, including patient age at diagnosis, histologic subtype, size, grade, and ER and progesterone receptor (PR) expression level by immunohistochemistry. We also reviewed the original 21-gene assay reports and recorded the RS results. Cases reported as morphologically similar were included; if no comment regarding morphologic similarity was noted, hematoxylin and eosin stained (H&E) slides of the tumor blocks submitted for analysis were reviewed and assessed for morphologic similarity.

Based on the RS of each tumor focus, BCs were divided into three groups: 1) BCs with RS in same risk category, 2) BCs with RS in different risk categories but within a 2-unit difference (e.g. RS 17 and RS 19), and 3) BCs with RS in different risk categories and a difference greater than 2 units. The selection of a 2 units cut off was based on the analytic validation studies of the Oncotype DX assay showing a standard deviation of 2 RS units on a 100 unit scale^{9,18}. BCs in groups 1 and 2 were considered concordant because there was no significant clinical impact due to the variation in RS result. BCs in group 3 were considered discordant because the RS variation was significant enough to impact therapeutic decisions. The H&E slides for all group 3 cases were reviewed by two pathologists (AG, HYW). The institutional review board approved the study.

Protocol for assessment of the 21-gene recurrence score in cases of multifocal invasive carcinoma

According to verbal information obtained from a representative of the company that performs the 21-gene assay¹⁹, when multiple tumor foci are received the reviewing pathologist determines which focus has the most worrisome histologic features, based on size and morphology, and selects that sample for testing first. In those carcinomas with similar morphology, the largest focus is tested first. If the RS is less than 25, the next sample is reflexively tested. When the assay result is greater than 25 the submitting clinician is contacted to advise whether testing of additional foci is needed.

Results

Patient selection and tumor characteristics

We identified a total of 108 patients with multiple ipsilateral BC and available RS results. Fifty-three patients were excluded due to morphologic dissimilarity of the BCs, and two had tumors insufficient for testing. Fifty-three patients met the inclusion criteria. The patient median age was 52 years (range 32–73). The majority of patients (81%, 43/53) underwent mastectomy while 19% (10/53) had breast conservation with wide local excision. BC types include ductal (n=36), lobular (n=14) and mixed ductal and lobular (n=3). The median tumor

size was 1.2 cm (range 0.5–4.6 cm). Most BCs were Nottingham Grade 2/3 (79%, 42/53). In eight patients (15%) the BCs were Grade 1/3, and in three patients (6%) the BCs were Grade 3/3. Most (83%, 44/53) patients had two tumor foci tested, eight (15%) had three foci, and one patient (2%) had four separate invasive foci submitted for analysis.

Recurrence score

A total of 116 BC samples from 53 patients were tested for RS. The median RS of the carcinomas was 14 (range 0–42). Most tumors (70%, 81/116) had a low RS (≤ 17). Thirty-three (28%) had intermediate RS (18–30) and two (2%) had a high RS (≥ 31).

Overall, the RS was concordant in 87% (46/53) of cases, including forty-four (83%) in group 1, and two (4%) in group 2. The median difference in RS in group 1 was 2 (range 0–11). One patient in this group had two foci of BC with RS scores of 0 and 11. The median difference in group 2 was 1.5 (range 1–2). Seven (13%) cases were discordant (group 3, Table 1) and had a median RS difference of 9 (range 4–22).

Pathology review of RS discordant cases

In three (43%) of the 7 cases with BC foci having discordant RS, we found that the tissue sections of the BC that yielded higher RS contained biopsy cavity changes (BXC) (Figure 1). In all three cases the BC focus without BXC yielded a low RS and the BC focus with BXC yielded intermediate RS. In four group 3 cases no BXC were identified. In two cases, both invasive ductal carcinoma (IDC), grade 2/3, immunohistochemical detection of PR expression showed varying results in each BC focus. One case showed nuclear positivity in 70% of cells in the focus with RS 13 and 40% in the focus with RS 23 (Figure 2). A second case with foci having RS 1 and RS 23 showed nuclear positivity in 80% and 5% of cells, respectively. The remaining two discordant cases were IDC with associated intermediate nuclear grade ductal carcinoma in situ (DCIS). In both cases the foci with lower RS had extensive DCIS admixed with the invasive carcinoma and no DCIS near the focus with higher RS.

TAILORx criteria

Using TAILORx cutoffs, most BCs (70%, 81/116) had an intermediate RS (11–25), with 25% (29/116) having a low RS (≤ 11) and 5% (6/116) a high RS (≥ 26). RS was concordant in 81% (43/53) of cases, all in group 1. The median difference in RS in group 1 was 2 (range 0–11). RS was discordant in 19% (10/53) of cases (Table 2). The median difference in RS in group 3 was 6.5 (range 5–22). Of the ten group 3 cases, 60% (6/10) were in women older than 50 years and all had RS ≤ 25 and 40% (4/10) were in women younger than 50 and three had RS ≥ 26 . One case in a 39 years old woman had two BC with high risk RS 26 and 27 and one BC with intermediate RS 22. On morphologic review, all three BCs were cytologically similar (MBR nuclear grade 3) and had similar mitotic activity (MBR mitotic count 2). The focus with RS 22 showed more tubule formation (MBR histologic grade 2) than the foci with RS 26 and 27 (MBR histologic grade 3).

Discussion

According to the College of American Pathologists recommendations, when there are multiple synchronous ipsilateral invasive carcinomas of the same histology, the largest invasive carcinoma is selected for classification and receptor assessment²⁰. However, when multigene prognostic assays are utilized to guide decisions on adjuvant systemic therapy, there is no consensus on whether testing of all individual foci is necessary. At our institution the treating physician decides whether one or multiple foci of BC are submitted for the 21-gene assay. Our data show that there is good correlation of RS with 87% concordance in patients with multifocal morphologically similar BC.

Seven (13%) cases in our cohort showed discordant results; all had low RS in one BC focus and intermediate RS in the other. Upon re-review of the H&E slides, biopsy cavity changes (BXC) were found adjacent to or admixed with the focus associated with higher RS in 3 of the 7 discordant cases. Studies from Genomic Health²¹ and others^{22–24} have shown that BXC have a significant effect on the expression of individual genes in the 21-gene assay which can lead to the false elevation of the RS in low to intermediate grade tumors and a decrease in the RS in poorly differentiated tumors. Furthermore, Genomic Health showed microdissected BXC had an average 13.4 ± 2.1 higher RS compared with the microdissected invasive cancers²¹. However, it should be noted that even with macro- or micro-dissection there are intervening elements, such as inflammatory cells and reactive stromal fibroblasts, which cannot be adequately separated from tumor tissue and can potentially alter the RS. Acs et al. reported increased RS in low grade invasive cancers associated with cellular stroma and/or inflammatory cells having an elevated proliferation index²⁵. These findings emphasize the importance of avoiding or at least minimizing BXC and other mitotically active stromal components in the tissue samples selected for multigene assays.

Together with the proliferation gene group the ER-related gene group, which includes PR, is one of the most influential of the multiple gene groups assessed by the Oncotype DX assay²⁶. PR expression is regulated by estrogen and is used to predict a functional ER pathway and therefore the likelihood of response to endocrine therapies²⁷. Several studies have shown that PR negative tumors have more aggressive profiles, benefit less from tamoxifen therapy, and have worse overall prognosis compared with PR positive tumors^{27–29}. Tang et al. found a similar correlation between PR expression and RS, reporting a significantly higher RS in PR-negative than in PR-positive BCs³⁰. Further validating these findings, an inverse relationship between PR expression level and RS independent of tumor grade has been identified^{22,28,31}. Accordingly, two of the discordant cases in our study had different PR expression in each BC focus tested, all of which were IDC, grade 2/3. Despite having similar histology, each focus with a low risk RS had a greater percentage of nuclear staining for PR, while the focus with intermediate risk RS showed less nuclear PR positivity. These differences in PR expression could be due to tumor heterogeneity or different clonality among tumor foci.

Two discordant cases showed a more predominant in situ component admixed with the BC focus having the lower RS. To our knowledge, there is no data examining how DCIS may

affect the RS assay result. Whenever possible, it is advised to avoid tissue blocks containing extensive DCIS. Further investigation of the impact of DCIS on the RS assay is warranted.

Previous studies have evaluated the RS in multifocal breast cancers by evaluating bilateral tumors³² or a mix of bilateral and ipsilateral BCs³³ which may or may not have been histologically different. Karsten et al. examined synchronous bilateral tumors and found a 33% rate of discordance, ultimately recommending testing of all foci³². This is consistent with our practice of staging synchronous bilateral primary carcinomas separately, according to the AJCC staging guidelines. Toole et al found that 50% of bilateral tumors and 22% of ipsilateral tumors had sufficiently different RS to warrant changes in therapeutic recommendations. This included bilateral BCs of different histologic subtype (75%) and ipsilateral BCs with different histologies (22%). However, they found a statistically significant difference in tumor histology and grade when comparing those patients who had discordant results versus those results that did not change management³³.

The TAILORx trial randomized patients with RS 11–25 to treatment with a combination of chemotherapy and endocrine therapy versus endocrine therapy alone. Analysis of the study results found that adjuvant chemotherapy is not beneficial in patients with RS <25 and age over 50 years. However, women under age 50 with BC having a mid-range RS score (RS 16–25) may still derive some benefit from chemotherapy¹³. Based on recent data from the TAILORx trial, 98% (52/53) of our cases were concordant in terms of adjuvant treatment recommendations. In the one discordant case all three BC foci were in a woman under 50 and had RS >15 (RS 22, 26, and 27). The clinical management of this patient would be unchanged under the current guidelines.

Our study looked only at patients with ipsilateral morphologically similar tumor foci and is the largest series reported so far. Our data show a good correlation of RS between multiple tumors with 87% concordance. In the seven discordant cases, three had associated BXC, known to artifactually increase the RS. In two cases, the change in RS was due to varying expression of PR. In two cases, the presence of DCIS may have affected the RS assay. Our findings 1) emphasize the importance of avoiding the biopsy site when selecting the tumor tissue blocks for testing and 2) suggest that testing of a single focus from a multifocal BC is sufficient to provide accurate prognostic information and to direct treatment, provided the foci are morphologically similar and have similar ER and PR expression.

Acknowledgment:

We thank Ms. Donna Thompson for her invaluable assistance with data collection.

References

1. Houssami N, Ciatto S, Macaskill P, et al. Accuracy and surgical impact of magnetic resonance imaging in breast cancer staging: systematic review and meta-analysis in detection of multifocal and multicentric cancer. *J Clin Oncol.* 2008;26(19):3248–3258. [PubMed: 18474876]
2. Wilkinson LS, Given-Wilson R, Hall T, Potts H, Sharma AK, Smith E. Increasing the diagnosis of multifocal primary breast cancer by the use of bilateral whole-breast ultrasound. *Clin Radiol.* 2005;60(5):573–578. [PubMed: 15851045]

3. Salgado R, Aftimos P, Sotiriou C, Desmedt C. Evolving paradigms in multifocal breast cancer. *Semin Cancer Biol.* 2015;31:111–118. [PubMed: 25066860]
4. Andea AA, Wallis T, Newman LA, Bouwman D, Dey J, Visscher DW. Pathologic analysis of tumor size and lymph node status in multifocal/multicentric breast carcinoma. *Cancer.* 2002;94(5):1383–1390. [PubMed: 11920492]
5. Buggi F, Folli S, Curcio A, et al. Multicentric/multifocal breast cancer with a single histotype: is the biological characterization of all individual foci justified? *Ann Oncol.* 2012;23(8):2042–2046. [PubMed: 22219015]
6. Choi Y, Kim EJ, Seol H, et al. The hormone receptor, human epidermal growth factor receptor 2, and molecular subtype status of individual tumor foci in multifocal/multicentric invasive ductal carcinoma of breast. *Hum Pathol.* 2012;43(1):48–55. [PubMed: 21733550]
7. Dawson PJ, Baekey PA, Clark RA. Mechanisms of multifocal breast cancer: an immunocytochemical study. *Hum Pathol.* 1995;26(9):965–969. [PubMed: 7672796]
8. Middleton LP, Vlastos G, Mirza NQ, Eva S, Sahin AA. Multicentric mammary carcinoma: evidence of monoclonal proliferation. *Cancer.* 2002;94(7):1910–1916. [PubMed: 11932891]
9. Paik S, Shak S, Tang G, et al. A multigene assay to predict recurrence of tamoxifen-treated, node-negative breast cancer. *N Engl J Med.* 2004;351(27):2817–2826. [PubMed: 15591335]
10. Paik S, Tang G, Shak S, et al. Gene expression and benefit of chemotherapy in women with node-negative, estrogen receptor-positive breast cancer. *J Clin Oncol.* 2006;24(23):3726–3734. [PubMed: 16720680]
11. Sparano JA, Gray RJ, Makower DF, et al. Prospective Validation of a 21-Gene Expression Assay in Breast Cancer. *N Engl J Med.* 2015;373(21):2005–2014. [PubMed: 26412349]
12. Sparano JA. TAILORx: trial assigning individualized options for treatment (Rx). *Clin Breast Cancer.* 2006;7(4):347–350. [PubMed: 17092406]
13. Sparano JA, Gray RJ, Makower DF, et al. Adjuvant Chemotherapy Guided by a 21-Gene Expression Assay in Breast Cancer. *N Engl J Med.* 2018;379(2):111–121. [PubMed: 29860917]
14. Gradishar WJ, Anderson BO, Balassanian R, et al. NCCN Guidelines Insights: Breast Cancer, Version 1.2017. *J Natl Compr Canc Netw.* 2017;15(4):433–451. [PubMed: 28404755]
15. Harris LN, Ismaila N, McShane LM, et al. Use of Biomarkers to Guide Decisions on Adjuvant Systemic Therapy for Women With Early-Stage Invasive Breast Cancer: American Society of Clinical Oncology Clinical Practice Guideline. *J Clin Oncol.* 2016;34(10):1134–1150. [PubMed: 26858339]
16. Giuliano AE, Connolly JL, Edge SB, et al. Breast Cancer-Major changes in the American Joint Committee on Cancer eighth edition cancer staging manual. *CA Cancer J Clin.* 2017;67(4):290–303. [PubMed: 28294295]
17. Amin M, Edge S, Green F, et al., ed AJCC Cancer Staging Manual Eighth Edition. Chicago, Illinois 2017 Amin M, ed.
18. Cronin M, Sangli C, Liu ML, et al. Analytical validation of the Oncotype DX genomic diagnostic test for recurrence prognosis and therapeutic response prediction in node-negative, estrogen receptor-positive breast cancer. *Clin Chem.* 2007;53(6):1084–1091. [PubMed: 17463177]
19. Oncotype IQ (2018). Oncotype DX breast recurrence score information for healthcare professionals. Available: www.oncotypeiq.com/en-US.
20. Lester SC, Bose S, Chen YY, et al. Protocol for the examination of specimens from patients with invasive carcinoma of the breast. *Arch Pathol Lab Med.* 2009;133(10):1515–1538. [PubMed: 19792042]
21. Baehner FQC, Pomeroy C, Cherbavaz D, Shak S. Biopsy Cavities in Breast Cancer Specimens: Impact on Quantitative RT-PCR Gene Expression Profiles and Recurrence Risk Assessment (abstract). *Mod Pathol.* 2009;22:28A–29A.
22. Allison KH, Kandalaft PL, Sitlani CM, Dintzis SM, Gown AM. Routine pathologic parameters can predict Oncotype DX recurrence scores in subsets of ER positive patients: who does not always need testing? *Breast Cancer Res Treat.* 2012;131(2):413–424. [PubMed: 21369717]
23. Hanna MG, Bleiweiss IJ, Nayak A, Jaffer S. Correlation of Oncotype DX Recurrence Score with Histomorphology and Immunohistochemistry in over 500 Patients. *Int J Breast Cancer.* 2017;2017:1257078. [PubMed: 28168058]

24. Hou Y, Zynger DL, Li X, Li Z. Comparison of Oncotype DX With Modified Magee Equation Recurrence Scores in Low-Grade Invasive Carcinoma of Breast. *Am J Clin Pathol*. 2017;148(2): 167–172. [PubMed: 28898988]
25. Acs G, Esposito NN, Kiluk J, Loftus L, Laronga C. A mitotically active, cellular tumor stroma and/or inflammatory cells associated with tumor cells may contribute to intermediate or high Oncotype DX Recurrence Scores in low-grade invasive breast carcinomas. *Mod Pathol*. 2012;25(4):556–566. [PubMed: 22173289]
26. Sparano JA, Paik S. Development of the 21-gene assay and its application in clinical practice and clinical trials. *J Clin Oncol*. 2008;26(5):721–728. [PubMed: 18258979]
27. Ahmed SS, Thike AA, Zhang K, Lim JC, Tan PH. Clinicopathological characteristics of oestrogen receptor negative, progesterone receptor positive breast cancers: re-evaluating subsets within this group. *J Clin Pathol*. 2017;70(4):320–326. [PubMed: 27612503]
28. Chaudhary LN, Jawa Z, Szabo A, Visotcky A, Chitambar CR. Relevance of progesterone receptor immunohistochemical staining to Oncotype DX recurrence score. *Hematol Oncol Stem Cell Ther*. 2016;9(2):48–54. [PubMed: 26808222]
29. Rakha EA, El-Sayed ME, Green AR, et al. Biologic and clinical characteristics of breast cancer with single hormone receptor positive phenotype. *J Clin Oncol*. 2007;25(30):4772–4778. [PubMed: 17876012]
30. Tang P, Wang J, Hicks DG, et al. A lower Allred score for progesterone receptor is strongly associated with a higher recurrence score of 21-gene assay in breast cancer. *Cancer Invest*. 2010;28(9):978–982. [PubMed: 20690804]
31. Clark BZ, Dabbs DJ, Cooper KL, Bhargava R. Impact of progesterone receptor semiquantitative immunohistochemical result on Oncotype DX recurrence score: a quality assurance study of 1074 cases. *Appl Immunohistochem Mol Morphol*. 2013;21(4):287–291. [PubMed: 23060300]
32. Karsten M, Stempel M, Radosa J, Patil S, King TA. Oncotype DX in Bilateral Synchronous Primary Invasive Breast Cancer. *Ann Surg Oncol*. 2016;23(2):471–476. [PubMed: 26340863]
33. Toole MJ, Kidwell KM, Van Poznak C. Oncotype dx results in multiple primary breast cancers. *Breast Cancer (Auckl)*. 2014;8:1–6. [PubMed: 24453493]

Synopsis

We examined the concordance of the Oncotype DX recurrence score (RS) in multiple synchronous ipsilateral breast carcinomas with similar histology. Our results show concordant RS in 87% (46/53) of cases suggesting testing of a single focus provides accurate prognostic information.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Figure 1:
Example of Group 3 case with BXC associated with higher RS. a: Well differentiated ductal carcinoma, 0.6 cm, with RS 11 (H&E, 200X); b: Well differentiated ductal carcinoma with similar morphology but adjacent BXC (outlined), 0.8 cm, with RS 20 (H&E, 20X); inset: higher power showing similar morphology (H&E, 200X)

Figure 2:

Example of Group 3 case with different PR expression. a: Ductal carcinoma with micropapillary features, 1.0 cm, with RS 13 (H&E, 100X); b: PR IHC with 70% nuclear positivity (40X); c: Ductal carcinoma with micropapillary features, 0.8 cm, with RS 23 (H&E, 100X); d: PR IHC with 40% nuclear positivity (40X)

Table 1:

Summary of Group 3 cases – Oncotype DX criteria

Case	Tumor type	MBR grade	Tumor size (cm)	Recurrence Score	Risk Classification
1	Lobular with pleomorphic features	2	1.8	15	Low
			3.0	19 [*]	Intermediate
2	Ductal, NOS	1	0.6	11	Low
			0.8	20 [*]	Intermediate
3	Mammary with mixed ductal and lobular features	2	1.0	12	Low
			1.2	19 [*]	Intermediate
			0.9	21 [*]	Intermediate
4	Ductal with micropapillary features	2	1.0	13 [†]	Low
			0.8	23	Intermediate
5	Ductal, NOS	2	1.4	1 [†]	Low
			0.6	23	Intermediate
6	Ductal, NOS	1	0.8	14	Low
			0.6	19	Intermediate
7	Ductal, NOS	2	1.5	14	Low
			1.1	19	Intermediate
			1.9	11	Low

Note: MBR – Modified Bloom Richardson

^{*} Focus with adjacent BXC[†] Focus with higher PR positivity

Table 2:

Summary of Group 3 cases - TAILORx criteria

Case	Tumor type	MBR Grade	Patient age (yrs)	Tumor size (cm)	Recurrence Score
1	Ductal, NOS	1	43	1.1	5
				1.0	12
2	Ductal, NOS	3	39	2.4	26
				1.8	27
				1.9	22
3	Ductal, NOS	2	69	2.3	11
				0.6	6
				0.5	5
4	Classic lobular	2	44	1.7	12
				1.4	7
5	Ductal with focal micropapillary features	2	56	1.1	15
				0.6	7
6	Classic lobular	2	52	2.0	14
				1.0	8
7	Classic lobular	2	53	3.4	11
				1.1	0
8	Ductal, NOS	2	43	1.1	7
				1.0	15
9	Ductal, NOS	2	53	1.4	1
				0.6	23
10	Mammary with mixed ductal and lobular features	2	60	4.6	10
				1.0	13