Published in final edited form as:

Pregnancy Hypertens. 2019 January ; 15: 7–11. doi:10.1016/j.preghy.2018.10.007.

Placental CD4⁺ T Cells Isolated from Preeclamptic Women Cause Preeclampsia-Like Symptoms in Pregnant Nude-Athymic Rats

Ashlyn C. Harmon¹, Tarek Ibrahim¹, Denise C. Cornelius^{1,2}, Lorena M. Amaral¹, Mark W. Cunningham Jr.¹, Kedra Wallace³, and Babbette LaMarca^{1,3}

¹University of Mississippi Medical Center, Department of Pharmacology and Toxicology, Jackson, MS.

²University of Mississippi Medical Center, Department of Emergency Medicine, Jackson, MS.

³University of Mississippi Medical Center, Department of Obstetrics and Gynecology, Jackson, MS.

Abstract

Preeclampsia (PE), new onset hypertension during pregnancy, is associated with a proinflammatory profile compared to normal pregnancy (NP). We hypothesize that CD4⁺ T cells from PE patient placentas cause PE symptoms during pregnancy compared to those from NP women. CD4⁺ T cells were isolated from placentas of PE and NP women using anti-CD4 magnetic separation, cultured in TexMACS medium at 37°C in 5% CO₂, and injected intraperitoneally into nude-athymic rats on day 12 of gestation. On day 18, carotid catheters were implanted and on day 19, mean arterial pressure (MAP) was measured and blood and tissues were collected. MAP was 125±2 mmHg in rats with NP CD4⁺T cells but increased to 140±4 mmHg in rats with PE CD4⁺T cells. Significant differences in circulating cytokines tumor necrosis factor- α (TNF- α), interleukin-17 (IL-17) and soluble fms-like tyrosine kinase-1 (sFlt-1) were found with PE vs NP CD4⁺ T cells (TNF- α - PE=167.4 pg/mL, NP=79.4 pg/mL; IL-17-PE= 7.054 pg/mL, NP=3.185 pg/mL; sFlt-1-PE=90.7 pg/mL, NP=58.2 pg/mL). In addition, renal cortical endothelin-1 (ET-1) mRNA expression increased 4.5 fold in rats with PE CD4⁺ T cells versus those receiving to NP CD4⁺ T cells. These data indicate an important role for placental PE CD4⁺ T cells to cause many characteristics of PE during pregnancy.

Corresponding Author: Babbette LaMarca, Ph.D., Professor, Thesis Director, Maternal Fetal Medicine Fellowship Program, Department of Pharmacology & Physiology & Ob/Gyn, Center for Excellence in Cardiovascular-Renal Research, University of Mississippi Medical Center, 2500 N. State St., Jackson MS 39216, bblamarca@umc.edu.

Ashlyn C. Harmon: Carried out the experiments and drafted original manuscript. Revised and edited the final version.

Tarek Ibrahim: Collected data necessary to the project, thus completing and finalizing the work presented.

Denise C. Cornelius: Performed experiments and analyzed data, revised manuscript and approved final manuscript.

Lorena M. Amaral: Performed experiments and data analysis.

Mark W. Cunningham Jr.: Performed experiments and data analysis.

Kedra Wallace: Edited and revised manuscript.

Babbette LaMarca: Conceived the hypothesis, designed the experiments, and supervised the execution and completion of the project.

Conflict of Interest/ Disclosure Statement

There are no disclosures or conflicts of interest.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Keywords

Preeclampsia; Pregnancy; T Cells; Inflammation; Renal

Introduction

Preeclampsia (PE) affects 5-10% of pregnant women each year worldwide. In the United States alone, the disease accounts for 18% of maternal deaths each year and is the number one reason doctors deliver babies prematurely(4). It is diagnosed by hypertension after week 20 of pregnancy and may be accompanied by proteinuria and edema. Women with PE present with characteristics of chronic inflammation with inflammatory CD4⁺ T cells and inflammatory cytokines, TNF- α and IL-17 are increased during the disease, while regulatory T cells and anti-inflammatory cytokines are suppressed. The underlying cause of PE is not known, however, evidence supporting the role of the placenta in PE has been presented numerous times(16,18), and it has been shown that delivery of the placenta from mothers with PE reduces the symptoms of the disease(7). Therefore, the objective of this study was to compare the effect of placental PE patient CD4⁺ T cells to that of placental NP CD4⁺ T cells to cause hypertension and other factors that are associated with the pathology of this disease.

Zenclussen et al. (2004) demonstrated with adoptive transfer of splenocytes isolated from normal pregnant mice, cultured to promote differentiation of Th-1 like cells that secreted Th-1 specific cytokines provoked symptoms of PE(15). In addition to these findings, data from our laboratory indicate a role for T cell-mediated pathophysiology during PE ((13,14,16, 17, 19)). CD4⁺T cells from the reduced uterine perfusion pressure (RUPP) rat model of PE were adoptively transferred into normal pregnant Sprague-Dawley rats. RUPP CD4⁺ T cells leads to many of the characteristic pathophysiology seen during PE, such as a significant increase in blood pressure, while there was no significant increase in blood pressure seen in the normal pregnant recipients of normal pregnant CD4⁺ T cells(16). Adoptive transfer of RUPP CD4⁺T cells into normal pregnant rats also led to an increase in circulating inflammatory cytokines, anti-angiogenic factor sFlt-1 and the vasoconstrictor ET-1, which has been previously shown to play a key role in the development of hypertension during PE(13,14,16). Although the importance of inflammatory T cells in the pathogenesis of PE has been established by adoptive transfer studies using animal models, no studies examining the importance of PE patient placental T cells in causing characteristics of PE have ever been performed.

Materials and Methods

Human Participant Selection

Twelve (6 normal pregnant; 6 PE) pregnant women undergoing cesarean-section at the Wiser Hospital for Women and Infants at the University of Mississippi Medical Center (UMMC) in Jackson, MS, were enrolled in the current study. PE was defined according to the 2002 ACOG recommendations of a systolic blood pressure ≥ 140 mmHg and a diastolic pressure ≥ 90 mmHg in previously normotensive women with the presence of proteinuria defined as

excretion of 0.3g in a 24hr specimen(1). All PE patients were treated with labetalol, hydralazine and/or Procardia to decrease their blood pressure and were prophylactically infused with magnesium sulfate upon hospital admission. Women with multiple gestations, chronic hypertension, pre-existing immune disorder (i.e. systemic lupus erythematosus), current tobacco/nicotine/alcohol usage or illegal substance abuse were excluded from the study. Prior to delivery all women signed an UMMC Institutional Review Board (2010-0301) approved consent and their placentas were collected and processed within 30 minutes of delivery.

Isolation, Expansion and Injection of Placental CD4⁺ T cells

On day 1 of isolation, NP and PE placental explants (~4-5g) were collected from the maternal side of the placenta, rinsed in PBS+5% penicillin/streptomycin (Pen/Strep) to remove surface blood, followed by tissue digestion in 4% collagenase I, 0.25U DNaseI and 5% Pen/Strep in RPMI1640 buffer for an hour at 37°C while stirring. The digest was strained with 100µm filters and centrifuged at 1200 RPM for 10 minutes. The cell pellet was reconstituted with Hanks Balance Salt Solution (HBSS) and layered carefully over 3 mL of Histopaque solution (Sigma-Aldrich, St. Louis, MO) and centrifuged for 30 minutes at 1200 RPM with no brake at room temperature. The lymphocyte layer was visualized and removed from the Histopaque via aspiration and placed in a clean tube before addition of 10 mL of RPMI+5%Fetal calf serum (FCS)+ 5%Pen/Strep. Lymphocytes were centrifuged at 3200RPM with brake at room temperature for 5 minutes to remove any residual Histopaque. The cell pellet was resuspended in degassed MACS running buffer (Miltenyi Biotec, San Diego, CA), followed by mixing with CD4 microbeads at 4°C for 30min. The lymphocyte mixture was centrifuged at 3200 RPM for 10 minutes and the cell pellet was resuspended in MACS running buffer and filtered with a 70µm filter. The filtered lymphocyte mixture was passed through MS columns (Miltenyi Biotec) on a MACS magnetic stand which allowed the CD4-labelled lymphocyte microbeads to adhere to the MS column and all non-labeled cells passed through the column. CD4⁺ T cells were mechanically released from the columns with the manufacture provided elutant (Miltenyi Biotec) and the resulting CD4⁺- labeled T cells were then incubated in TexMACs media (Miltenyi Biotec) + 1.022ng/mL of IL-2 overnight. IL-2 is needed to maintain viability of T cells. Day 2 of isolation, cells were collected and centrifuged at 1600 RPM, counted using a cell counter from Beckman-Coulter, then stored frozen in liquid nitrogen until needed for injection.

Adoptive transfer of T cells into pregnant rats

Timed-pregnant nude-athymic rats from Harlan (Indianapolis, IN) were used in this study. Nude-athymic rats were chosen because they lack a thymus, meaning they do not produce T cells and will not reject the human T cells injected into them. However, this does render them immunocompromised and thus, animals were continuously housed in a temperature-controlled barrier suite (23 °C) with a 12:12-hour light/dark cycle. All experimental procedures performed in this study were in accordance with the National Institutes of Health guidelines for use and care of animals. All protocols were approved by the Institutional Animal Care and Use Committee (IACUC) at the University of Mississippi Medical Center.

The day prior to injecting nude-athymic rats with isolated human placental CD4⁺ T cells, vials of frozen cells were removed from storage in liquid nitrogen and thawed quickly using ambient temperature water. Cells were washed in HBSS and incubated overnight for further expansion. On the day of injection, CD4⁺ T cells were removed from incubation, counted, then suspended in sterile saline at a concentration of 1×10^6 cells per 500 μ L. In order to be consistent with our other publications performing the adoptive transfer technique (12,13,16,17), on gestational day 12, normal pregnant, nude-athymic rats were anesthetized using isoflurane and quickly injected intraperitoneally with 1×10^6 placental CD4⁺ T cells isolated from either one NP or one PE patient. A group of normal pregnant (NP) nude athymic rats was used as a blood pressure control in which no CD4⁺ T cells were injected.

Measurement of Blood Pressure in Conscious, Nude-Athymic Rats

Under isoflurane anesthesia, on day 18 of gestation, carotid arterial catheters were inserted for blood pressure measurements. The catheters inserted are V3 tubing (SCI) which is tunneled to the back of the neck and exteriorized. On day 19 of gestation, arterial blood pressure was analyzed after placing the rats in individual restraining cages. Arterial pressure was monitored with a pressure transducer (Cobe III Transducer CDX Sema) and recorded continuously for one hour after a 30 min. stabilization period. The nude-athymic rats were placed under isoflurane anesthesia prior to blood and tissue collection. Kidneys and placentas were harvested and weighed prior to being stored for future analysis. Pups were removed and weights were recorded. Serum and plasma were isolated via centrifugation.

Determination Endothelin Expression in the Renal Cortex

Real time PCR (qRT-PCR) was utilized to determine tissue preproendothelin-1 levels. Kidneys were isolated, weighed and quickly frozen in liquid nitrogen. The tissues were then stored at -80°C . Total RNA was extracted from the tissues using the RNeasy Protect Mini Kit (Qiagen) and the isolation procedure was performed according to the manufacturer provided instructions. cDNA was synthesized from 1 μ g of RNA using the iScript cDNA Synthesis Kit (BioRad). qRT-PCR was performed using iQ SYBR Green Supermix (BioRad) and the CFX96 Touch Real-Time PCR Detection System (BioRad). The following primer sequences provided by Life technologies were used for Preproendothelin (PPET) as previously described: forward 1, ctaggctaacgcatccttg, and reverse 1, tctttgtctgcttgcc. Levels of mRNA were calculated using the mathematical formula $2^{-\text{ct}}$ ($2^{\text{avg. ct gene of interest} - \text{avg ct beta actin}}$) recommended by Applied Biosystems (Applied Biosystems User Bulletin, No. 2, 1997).

Measurement of Circulating sFlt-1, TNF alpha and IL-17

Circulating levels of the anti-angiogenic factor sFlt-1, TNF alpha and IL-17 were measured using a commercially available enzyme-linked immunosorbent assay (ELISA) kit (Quantikine, R&D Systems, Minneapolis, MN). Appropriate amounts of plasma from each sample were analyzed according to the manufacturer's instructions. The MDD for sFlt-1 assay is 15.2 pg/mL and intra-assay/inter-assay precision is 7.3% CV of 4.8% CV, respectively. The MDD for TNF alpha and IL-17 is 5pg/mL and intra-assay precision is 3.1% CV and 3.3% CV, respectively, inter-assay precision is 9.3% CV and 4.9% CV, respectively.

Statistical Analysis

All of the data are expressed as mean $\pm$ SEM. Comparisons of control with experimental groups were analyzed by the student *t*-test and analysis of variance (ANOVA) with Tukey's post hoc analysis. A value of $P < 0.05$ was considered statistically significant.

Results

Participant's Data

The participants' clinical characteristics are presented in Table 1. There were no statistically significant differences in maternal age between women with PE and NP women ($p=0.78$), maternal race ($p>0.99$) or in maternal BMI at the time of delivery ($p=0.29$; Table 1). PE women did delivery significantly earlier compared to NP women ($p=0.04$) which corresponded to a significantly lower fetal birthweight among infants born to PE women compared to infants born to NP women ($p=0.003$; Table 1). Women with PE had significantly increased systolic ($p=0.003$) and diastolic ($p=0.04$) blood pressures compared to NP women (Table 1).

MAP was Increased in Recipients of PE T Cells

MAP was measured in pregnant, nude-athymic rat recipients of NP placental CD4⁺ T cells and PE placental CD4⁺ T cells. MAP was significantly increased to 140 ± 4 mmHg in recipients of PE T cells ($n=6$) vs. normal pregnant (NP) nudes ($n=3$) (140 ± 4 vs. 123 ± 5 mmHg, $p<0.05$) (Figure 1). MAP was 125 ± 2 mmHg in recipients of NP T cells ($n=6$). No change in MAP was noted between NP nude-athymic rats and nude-athymic rats given NP CD4⁺ T cells ($n=6$) (123 ± 5 vs 125 ± 2 mmHg, ns).

TNF- α and IL-17 are Increased in Recipients of PE T cells

TNF- α production increased to 167.4 ± 30.3 pg/mL in pregnant, nude-athymic recipients of placental PE CD4⁺ T cells ($n=6$) versus 79.4 ± 44.6 pg/mL in nude-athymic recipients of placental NP CD4⁺ T cells ($n=6$) ($p<0.05$) (Figure 2A). IL-17 significantly increased to 7.054 ± 1.446 pg/mL in recipients of PE CD4⁺ T cells versus 3.185 ± 0.6623 pg/mL recipients of NP T Cells ($p<0.05$) (Figure 2B).

Circulating sFlt-1 and Renal Cortical PPET-1 Increased in Recipients of PE T cells

Circulating sFlt-1 production was 71.5 ± 14.7 pg/mL in recipients of NP CD4⁺T cells ($n=6$) and was significantly increased to 90.7 ± 6.5 pg/mL in recipients of PE CD4⁺ T cells ($n=6$) (Figure 3A). PPET-1 expression was significantly increased in the renal cortex of PE CD4⁺ T cells recipients by a greater than 4 fold change compared to nude-athymic rats injected with NP T cells ($p<0.05$) (Figure 3B).

Discussion

As poor placentation leading to placental ischemia is considered to be one of the primary pathophysiological events leading to the development of PE, it is important to try to identify some of the molecular mechanisms associated with placental ischemia (6,15). The idea that the imbalance in Tregulatory and Th17 cells can contribute to disease pathology has only

recently started to be explored in animal models, however several clinical studies have reported imbalances in these subsets (3,11). Based on these studies along with those conducted in our lab we set out to determine if CD4⁺ T cells could lead to the changes blood pressure, inflammation and renal function that accompanies PE.

We demonstrate in this study for the first time that PE CD4⁺ T cells isolated from the placentas of PE patients increase circulating inflammatory cytokines TNF- α and IL-17 and sFlt-1 when injected into normal pregnant nude athymic rats compared NP recipients of NP CD4⁺ T cells. We have previously shown TNF- α to cause hypertension associated with oxidative stress, endothelin, and the antiangiogenic factor sFlt-1, while IL-17 stimulates ROS and sFlt-1 in association with hypertension during pregnancy. (4, 12,13,16,17). We have also shown that blocking TNF- α or IL-17 lowers blood pressure and these vasoactive pathways in the RUPP rat model of PE (2, 8). Importantly, when we infuse either TNF- α or sFlt-1 into normal pregnant rats, ET-1 is stimulated and is instrumental in causing the resultant hypertension, which is ameliorated with ETA blockade(4).

The current study is the first study to show that PE placental CD4⁺T cells are important to stimulate an elevation in blood pressure during pregnancy by activating ET-1 and sFlt-1 pathways. Moreover, the increase in blood pressure with PE placental T cells injected into normal pregnant, nude-athymic rats, was significantly greater than that observed with NP placental CD4⁺ T cells. We recognize there were no changes in pup weight, which may be due to the late gestation injection, however injection of CD4⁺T cells on gestation day 12 of pregnancy is consistent with our previous adoptive transfer studies. Nevertheless, it is a good model in which to study placental PE CD4⁺ T cells as mediators of hypertension and the PE phenotype during pregnancy.

A number of underlying mechanisms may be utilized by CD4⁺ T cells to induce these vasoactive factors via direct or indirect actions. We have previously shown that activated T cells form the RUPP rat model of PE secrete sFlt-1, IL-17 and TNF- α , which could be mechanism where by these cytokines are elevated in the rats receiving PE T cells compared to those receiving NP T cells(13). In addition to secreting sFlt-1, CD4⁺ T cells may induce the vasoactive factors sFlt-1 and PPET-1 through indirect mechanisms. The CD4⁺ T cells from RUPP rats secrete TNF α and directly causes increased vascular and placental levels of TNF α . Increased TNF- α activates endothelial cells, decreases nitric oxide synthase mRNA, and increases the production ET-1 by stimulating expression of PPET-1 mRNA (7-10,18). The increased population of T cells also stimulate B cells to produce agonistic angiotensin II type 1 receptor autoantibodies (AT1-AA). AT1-AAs induce signaling by the angiotensin II type 1 receptor (AT1-receptor) leading to intracellular cascades and promoter activations in the nucleus to cause upregulation of vasoactive factors including PPET-1 and sFlt1(5). The CD4⁺ T cells can directly or indirectly stimulate pathways to induce the PE phenotype observed in this adoptive transfer model.

This study establishes a role for human preeclamptic CD4⁺ T cells in mediating PE-like symptoms and adds further evidence to existing studies examining the role of inflammatory T cells and their contribution to the disease. This study not only established a role for the altered CD4⁺ T cell population in human preeclamptic patients, but it also strongly suggests

that the increase in blood pressure and the pathophysiology of preeclampsia is further mediated by the production of sFlt-1 and ET-1. With the increase in IL-17 in the recipient rats, future studies will further examine a role of TH17 cells to mediate some of these effects.

Acknowledgments

Funding Sources

This work was supported by NIH grants HL78147 and HL51971 and HLHD067541 and P20GM121334 awarded to BL. DCC is supported by NIH grant HL130456 and the American Heart Association 16SDG27520000.

References

1. ACOG. Diagnosis and management of preeclampsia and eclampsia. Practice Bulletin 33: 2002.
2. Cornelius DC, Hogg JP, Scott J, Wallace K, Herse F, Moseley J, Wallukat G, Dechend R, and LaMarca B. Administration of interleukin-17 soluble receptor C suppresses TH17 cells, oxidative stress, and hypertension in response to placental ischemia during pregnancy. *Hypertension* 62: 1068–1073, 2013. [PubMed: 24060899]
3. Darmochwal-Kolarz D, Kludka-Sternik M, Tabarkiewicz J, Kolarz B, Rolinski J, Leszczynska-Gorzela B, Oleszczuk J. The predominance of Th17 lymphocytes and decreased number and function of Treg cells in preeclampsia. *J Rep Immun* 2012: 93(2):75–81.
4. Dhillon P, Wallace K, Herse F, Scott J, Wallukat G, Heath J, Mosely J, Martin JN Jr., Dechend R, and LaMarca B. IL-17-mediated oxidative stress is an important stimulator of AT1-AA and hypertension during pregnancy. *American journal of physiology Regulatory, integrative and comparative physiology* 303: R353–358, 2012.
5. Herse F, and LaMarca B. Angiotensin II type 1 receptor autoantibody (AT1-AA)-mediated pregnancy hypertension. *American journal of reproductive immunology* 69: 413–418, 2013. [PubMed: 23279165]
6. Karumanchi SA, Granger JP. Preeclampsia and pregnancy-related hypertensive disorders. *Hypertension* 2016;67(2):238–242. [PubMed: 26693822]
7. LaMarca B, Cornelius D, and Wallace K. Elucidating immune mechanisms causing hypertension during pregnancy. *Physiology (Bethesda)* 28: 225–233, 2013. [PubMed: 23817797]
8. LaMarca B, Speed J, Fournier L, Babcock SA, Berry H, Cockrell K, and Granger JP. Hypertension in response to chronic reductions in uterine perfusion in pregnant rats: effect of tumor necrosis factor-alpha blockade. *Hypertension* 52: 1161–1167, 2008. [PubMed: 18981324]
9. LaMarca B, Wallace K, and Granger J. Role of angiotensin II type I receptor agonistic autoantibodies (AT1-AA) in preeclampsia. *Current opinion in pharmacology* 11: 175–179, 2011. [PubMed: 21317038]
10. LaMarca BB, Cockrell K, Sullivan E, Bennett W, and Granger JP. Role of endothelin in mediating tumor necrosis factor-induced hypertension in pregnant rats. *Hypertension* 46: 82–86, 2005. [PubMed: 15928030]
11. Jianjun Z, Yali H, Zhiqun W, Mingming Z, Xia Z. Imbalance of T-cell transcription factors contributes to the Th1 Type Immunity predominant in pre-eclampsia. *Am J Rep Imm.* 2010;63(1): 38–45.
12. Novotny S, Wallace K, Herse F, Moseley J, Darby M, Heath J, Gill J, Wallukat G, Martin JN, Dechend R, and LaMarca B. CD4 T Cells Play a Critical Role in Mediating Hypertension in Response to Placental Ischemia. *Journal of hypertension : open access* 2: 2013.
13. Novotny SR, Wallace K, Heath J, Moseley J, Dhillon P, Weimer A, Wallukat G, Herse F, Wenzel K, Martin JN Jr., Dechend R, and LaMarca B. Activating autoantibodies to the angiotensin II type I receptor play an important role in mediating hypertension in response to adoptive transfer of CD4+ T lymphocytes from placental ischemic rats. *American journal of physiology Regulatory, integrative and comparative physiology* 302: R1197–1201, 2012.

14. Roberts JM, and Gammill HS. Preeclampsia: recent insights. *Hypertension* 46: 1243–1249, 2005. [PubMed: 16230510]
15. Sava RI, March KL, Pepine CJ. Hypertension in pregnancy: Taking cues from pathophysiology for clinical practice. *Clin Cardiol.* 2018;41(2):220–227. [PubMed: 29485737]
16. Wallace K, Novotny S, Heath J, Moseley J, Martin J, Owens M, and Lamarca B. Hypertension in response to CD4+ T cells from reduced uterine perfusion pregnant rats is associated with activation of the endothelin-1 system. *American journal of physiology Regulatory, integrative and comparative physiology* 303: R144–149, 2012.
17. Wallace K, Richards S, Dhillon P, Weimer A, Edholm ES, Bengten E, Wilson M, Martin JN Jr., and LaMarca B. CD4+ T-helper cells stimulated in response to placental ischemia mediate hypertension during pregnancy. *Hypertension* 57: 949–955, 2011. [PubMed: 21464392]
18. Yoshizumi M, Perrella MA, Burnett JC Jr., and Lee ME. Tumor necrosis factor downregulates an endothelial nitric oxide synthase mRNA by shortening its half-life. *Circ Res* 73: 205–209, 1993. [PubMed: 7685252]
19. Zenclussen AC, Fest S, Joachim R, Klapp BF, and Arck PC. Introducing a mouse model for pre-eclampsia: adoptive transfer of activated Th1 cells leads to pre-eclampsia-like symptoms exclusively in pregnant mice. *European journal of immunology* 34: 377–387, 2004. [PubMed: 14768042]

Highlights

The placenta has long been thought to be the culprit of much of the pathology of preeclampsia. This study highlights the role of CD4⁺ T cells from the placenta of preeclamptic patients to cause many facets of the disease such as elevated cytokines vasoactive pathways and blood pressure during pregnancy. Importantly, adoptive transfer of placental CD4⁺ T cells from normal pregnant mothers had no effect to increase blood pressure during pregnancy. This study indicates the importance of placental inflammation to cause many facets of preeclampsia and thus highlights new areas of cytokine therapy that may be of interest in treating this disease, such as TNF alpha and IL-17.

Figure 1. Circulating inflammatory cytokines TNF- α and IL-17 are increased in response to adoptive transfer of PE CD4⁺ T cells compared to recipients of normal pregnant CD4⁺ T cells. **A)** Plasma TNF- α production had a tendency to increase in pregnant, nude-athymic recipients of placental PE CD4⁺ T cells vs. nude-athymic recipients of NP CD4⁺ T cells (not significant). **B)** IL-17 was increased nude-athymic recipients of placental PE T cells vs nude-athymic recipients of NP T cells. Statistical changes was determined using student *t*-test, **p*<0.05 vs NP.

Figure 2. Renal and circulating vasoconstrictors are increased in Nude-Athymic Recipients of Preeclamptic CD4⁺ T cells.

A) Plasma sFlt-1 increased in nude-athymic recipients of placental PE CD4⁺ T cells vs. nude-athymic recipients of NP CD4⁺ T cells. **B)** PPET-1 expression was significantly increased by 4 fold in the renal cortex of nude-athymic recipients of placental PE CD4⁺ T cells vs. nude-athymic recipients of NP CD4⁺ T cells. Statistical changes was determined using student *t-test*, **p*<0.05 vs NP.

Figure 3. Blood Pressure is increased in Nude-Athymic Recipients of Preeclamptic CD4⁺ T cells compared to NP CD4⁺ T cells.

MAP was significantly increased in recipients of PE CD4⁺ T cells vs. recipients of NP T cells. No change in MAP between NP nude-athymic rats and nude-athymic rats given NP CD4⁺ T cells. Statistical changes was determined using one-way ANOVA with Turkey's post hoc analysis, *p<0.05 vs NP.

Table 1.

Clinical characteristics of study participants. Data is expressed as mean $\pm$ standard error mean.

Variables	Pre-eclampsia	Normal Pregnant	P-value
Maternal Age	25.0 $\pm$ 1.9yrs	25.8 $\pm$ 2.1yrs	0.78
% African American	83%	100%	>0.99
Gestational age at delivery	34.7 $\pm$ 1.6wks	38.8 $\pm$ 0.7wks	0.04
Fetal Birthweight	1679 $\pm$ 312.3g	3259 $\pm$ 243.3g	0.003
% Multiparous	100%	67%	0.46
Body mass index	30.65 $\pm$ 3.5kg/m ²	37.1 $\pm$ 4.7 kg/m ²	0.29
Systolic blood pressure	156.5 $\pm$ 5.8mmHg	125.3 $\pm$ 5.6mmHg	0.003
Diastolic blood pressure	94.5 $\pm$ 6.1mmHg	73.8 $\pm$ 6.1mmHg	0.04