United Kingdom CSF Disorders Day 2018

Jessica K. Walker ^a, Alexandra J. Sinclair ^b, and Susan P. Mollan ^c

^aCollege of Medical and Dental Sciences, Medical School, University of Birmingham, Birmingham, UK; ^bMetabolic Neurology, Institute of Metabolism and Systems Research, University of Birmingham, Birmingham, UK; ^cBirmingham Neuro-ophthalmology Unit, University Hospital Birmingham, Birmingham, UK

ABSTRACT

The multi-specialty approach to managing patients with cerebrospinal fluid (CSF) disorders requires neurology, neurosurgery, neuroradiology and neuro-ophthalmology. The annual United Kingdom CSF disorders meeting presented the first guidelines for idiopathic intracranial hypertension focusing on protecting the vision, managing the underlying disease and reducing headache morbidity. The most recent scientific concepts in CSF disorders were presented, as was the applied understanding of shunt and valve technology.

ARTICLE HISTORY

Received 1 December 2018
Accepted 16 December 2018

KEYWORDS

Cerebrospinal fluid; IIH; CSF diversion; headache

The United Kingdom Cerebrospinal Fluid (CSF) symposium is an annual multi-disciplinary event aimed at all members of the multi-disciplinary team. Attendees included specialists in neurosurgery, neurology, neuroradiology and neuro-ophthalmology. The symposium was organised by Miss Susan Mollan, Consultant Neuro-ophthalmologist at University Hospitals Birmingham and Dr Alexandra J. Sinclair, National Institute for Health Research (NIHR) Clinician Scientist, Institute of Metabolism and Systems Research at University of Birmingham with help from Miss Jessica Walker, Medical Student at the University of Birmingham (Figure 1).

The 2018 symposium aims were:

- To facilitate discussion within the multi-specialty team regarding the more challenging and controversial issues in managing CSF disorders.
- To provide a synopsis of the latest scientific and clinical trial research.
- To launch the Idiopathic Intracranial Hypertension Consensus Guidelines.¹

Speed science section

In the speed science section, the latest work of basic scientists and clinicians was presented. Miss Jessica Katanga, a medical student supervised by Dr Arani

Nitkunan at St George's and Croydon Hospitals, was the winner of this section for presenting an audit of the management of idiopathic intracranial hypertension (IIH) in the context of the new consensus guidelines.¹ In the consecutive retrospective audit they showed that implementation of neuroimaging and lumbar puncture was high but improvement was needed in routine measurement of body mass index (BMI) and documentation of teratogenicity and side effects of medications prescribed for IIH or headache. Venography performed within 24 hours was achieved in just over half of patients; this was echoed in the audience discussion as the availability of venography within 24 hours around the UK is challenging. The neuroradiology experts explained that this standard was not based on current practice but what should be a “gold standard” of care for papilloedema patients, as the detection of a venous sinus thrombosis would change management. Additionally, it would be more cost effective to undergo the MRI and MRV at the same time rather than separately.

Dr Nicola Adderley from the Institute of Applied Health Research, University of Birmingham, presented a case control study investigating the risk of cardiovascular disease in women with IIH. Her results showed that there was twice the risk of composite cardiovascular events independent of BMI, suggesting

Figure 1. Miss Susan Mollan (left) and Dr Alexandra Sinclair (third from left) with the chair and trustee of IIH UK.

cardiovascular risk in IIH may not exclusively be driven by obesity. A PhD student at the Open University, Ester Pascual-Baixaul, presented the effect of a high fat diet on CSF secretion in an animal model. There was a positive correlation between a high percentage of body fat and CSF secretion and between CSF secretion and testosterone levels.

Dr Anna Anthonypillai, a foundation doctor at St Thomas' Hospital, London audited the cases of pseudotumour cerebri (PTC) syndrome secondary to anaemia. Of 98 patients seen at one unit with PTC, only 10 had anaemia (Hb < 11 g/dl) and 4 had anaemia without precipitous weight gain or other causal factors for IIH. Treating these four patients with iron supplementation led to a reduction in papilloedema.

The final speed science presentation was by Linda D'Antona from University College London who examined the role of spontaneous venous pulsation as a non-invasive measure of intracranial pressure using the infrared video feature on Spectralis optical coherence tomography.

Latest in trials for CSF disorders

Dr Keira Markey from the University of Birmingham presented the results of the IIH:DT, a phase II randomised control trial (RCT) assessing the safety, tolerability and efficacy of an 11 β -hydroxysteroid dehydrogenase type 1 inhibitor, AZD4017, in IIH. This is the first trial of a novel therapy for IIH.

AZD4017 was safe and tolerated over 12 weeks. In vivo efficacy was demonstrated in urinary, hepatic and cerebrospinal fluid biomarkers. Possible clinical benefits were noted, but a further phase III clinical trial will be required to demonstrate efficacy.

Mr Conor Mallucci, Consultant Neurosurgeon from Alder Hey Hospital, Liverpool gave an overview of the British antibiotic and silver impregnated catheters for ventriculoperitoneal shunts (BASICS) trial. This multi-centre RCT funded by the NIHR HTA programme randomised 1,605 patients between standard ventriculoperitoneal shunts (VPS), antibiotic impregnated VPS and silver impregnated VPS with a primary outcome of time to VPS infection following insertion of the first de novo VPS. The results of the BASICS trial will have clinical impact for all CSF disorders requiring a VPS.

IIH consensus guidelines

There was a series of three presentations regarding the guidelines.¹ Miss Susan Mollan presented the process of engagement of the key stakeholders including patients, the patient charity IIH UK, the Society of British Neurological Surgeons, the Association of British Neurologists, the Royal College of Ophthalmologists and key international opinion leaders in the field. The aim of the guidance was to equip healthcare professionals to correctly diagnose and optimally manage those with IIH

Figure 2. Keynote speaker Christophe Miethke meets representatives from IIH UK. IIH UK generously supports the UK CSF Annual Meeting.

Figure 3. Dr Abd Tahrani delivers audience captivating talk on obesity, barriers and solutions.

within the context of the published literature. Where evidence did not exist the guidelines were developed by Delphi rounds of UK specialists in neurology, neurosurgery and ophthalmology. The importance of primarily protecting vision was highlighted and flow diagrams of the consensus guidance were presented.²

Mr Ahmed Toma, Consultant Neurosurgeon, The National Hospital for Neurology and Neurosurgery, London presented the neurosurgical approach to IIH. He educated on how shunts had evolved within the management for IIH. Looking at the recent evidence, Mr Toma posed that shunts were not the solution, but a temporising measure, due to surgical complications and revision rates. He underlined that CSF diversion in IIH was to protect vision within the context of acute visual deterioration, and should only be used exceptionally for headache. The resulting discussion included comments from colleagues, both in the UK and Denmark, about the utility of optic nerve sheath fenestration as an alternative.

Focusing on the management of headache in IIH was Dr Brendan Davies, Consultant Neurologist, University Hospital North Midlands, Stoke-on-Trent. Managing the headache morbidity in the IIH patient group has been shown to improve quality of life. Headaches in IIH typically have a migrainous phenotype and this is helpful for recommending acute and preventative therapies. He highlighted that medication-overuse headache was important to deal with.

Keynote lecture

Mr Christoph Miethke from Miethke GmbH & Co. KG, Potsdam, Germany, gave a detailed lecture which described the physics and mechanisms behind shunts and valve technology (Figure 2). Demonstrating his points with videos, he explained the pressure in a straw and evolved this example to allow the audience to understand pressures in open and closed systems and then described some of the more intricate details underlying valve technology in an accessible way for the audience.

From the multi-disciplinary team

Mr George Tsermoulas, Consultant Neurosurgeon from University Hospitals Birmingham presented on the clinical utility of CSF infusion studies for Normal Pressure Hydrocephalus (NPH) and possible shunt malfunction. He emphasised the importance of diagnosing NPH as a reversible cause of disability.

Dr Abd Tahrani, NIHR Clinician Scientist from the University of Birmingham, presented a thought-provoking lecture explaining that obesity should be classed as a complex relapsing disease, which is typically poorly managed and leads to multi-system complications (Figure 3). He explained the hormone and energy changes which makes maintaining weight loss difficult. He presented the current trends in the use of bariatric surgery and medical therapies needed to prevent regain of weight. One key message was regarding society's attitude to obesity and he shared a powerful radio-interview exemplifying the discrimination faced by those with obesity and appealed to all health care professionals in the room to recognise obesity stigma and ensure it did not interfere with clinical practice.

ORCID

Jessica K. Walker <http://orcid.org/0000-0003-2586-7172>
 Alexandra J. Sinclair <http://orcid.org/0000-0003-2777-5132>
 Susan P. Mollan <http://orcid.org/0000-0002-6314-4437>

References

1. Mollan SP, Davies B, Silver NC, Shaw S, Mallucci CL, Wakerley BR, Krishnan A, Chavda SV, Ramalingam S, Edwards J, Hemmings K, Williamson M, Burdon MA, Hassan-Smith G, Digre K, Liu GT, Jensen RH, Sinclair AJ. Idiopathic intracranial hypertension: consensus guidelines on management. *J Neurol Neurosurg Psychiatry*. 2018;89(10):1088–1100. doi:10.1136/jnnp-2017-317440.
2. Mollan SP, Hornby C, Mitchell J, Sinclair AJ. Evaluation and management of adult idiopathic intracranial hypertension. *Pract Neurol*. 2018;18(6):485–488. doi:10.1136/practneurol-2018-002009.