Published in final edited form as:

Cancer Causes Control. 2019 March ; 30(3): 281–290. doi:10.1007/s10552-019-1135-3.

Healthy behavioral choices and cancer screening in persons living with HIV/AIDS are different by sex and years since HIV diagnosis

Akemi T. Wijayabahu¹, Zhi Zhou¹, Robert L. Cook¹, Babette Brumback², Nicole Ennis³, and Lusine Yaghjian¹

¹Department of Epidemiology, College of Public Health and Health Professions and College of Medicine, University of Florida, Gainesville, United States of America

²Department of Biostatistics, College of Public Health and Health Professions and College of Medicine, University of Florida, Gainesville, United States of America

³Department of Clinical and Health Psychology, College of Public Health and Health Professions and College of Medicine, University of Florida, Gainesville, United States of America

Abstract

Purpose: The prevalence of non-AIDS related malignancies is on the rise among people aging with HIV population, but the evidence on healthy behaviors including cancer screening practices in this population subgroup is extremely limited. Therefore, we investigated the prevalence of healthy behaviors and sex-specific cancer screening among persons living with HIV, by sex and time since HIV diagnosis.

Methods: We included 517 persons living with HIV from the Florida Cohort. Data were obtained from the baseline and follow-up questionnaires, electronic medical records, and Enhanced HIV/AIDS Reporting System. The prevalence of self-reported, age-appropriate cancer screening (anal, colorectal, prostate, breast and cervical) and healthy behaviors (sustaining healthy body weight, refraining from smoking and alcohol and engaging in physical activity) was compared by sex and years since HIV diagnosis (13 vs. >13 years).

Results: In the analyses by sex, females were more likely to be obese than males (56.5% vs. 22.2%, $p < 0.0001$). Distribution of healthy behaviors did not differ by time since diagnosis among

Corresponding Author: Lusine Yaghjian, MD, MPH, PhD, Assistant Professor, Department of Epidemiology, College of Public Health and Health Professions, College of Medicine, University of Florida, 2004 Mowry Road, PO Box 100321, Gainesville, Florida 32610-0231, Phone: 352-294-5940, lyaghjian@ufl.edu.

Compliance with Ethical Standards

Disclosure of potential conflicts of interest

The authors declare that they have no conflict of interest.

Research involving Human Participants and/or Animals

All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards.”

Informed consent

This study was approved by the University of Florida, Florida International University, and the Florida Department of Health Institutional Review Boards (IRBs). All participants provided written informed consent.

ORCID 0000-0002-1626-5340

males and females. In the analysis of age-appropriate screening among males, 64.8% never had an anal Pap-smear, 36.2% never had a colonoscopy, and 38.9% never had prostate cancer screening. In the analysis of age-appropriate screening among females, 50.0% never had an anal Pap-smear, 46.5% never had a colonoscopy, 7.9% never had a cervical Pap-smear and 12.7% never had a mammogram. The difference in anal Pap-smear by sex was statistically significant ($p<0.0001$). Among males, the age-adjusted prevalence of never having a colonoscopy was higher in those who had HIV for ≥ 13 years (50.8% vs. 30.6%, $p=0.03$).

Conclusion: Prevalence of selected healthy behaviors and cancer screening differed by sex and/or years since HIV diagnosis suggesting a need for tailored cancer prevention efforts among persons living with HIV via long-term sex-specific interventions.

Keywords

cancer screening; healthy behaviors; HIV/AIDS

Introduction

In the last two decades, advancements in antiretroviral therapy and the implementation of population-based interventions resulted in improved survival among persons living with HIV/AIDS in the US [1–3]. HIV continues to be a major public health and economic burden in the US, accounting for more than 1.2 million persons live with the disease and nearly half of that population being aged 50 or older [4]. Older adults living with HIV/AIDS are at an increased risk for developing AIDS-defining (Kaposi sarcoma, non-Hodgkin lymphoma, and cervical cancer) and non-AIDS-defining malignancies (anal cancer, liver cancer, lung cancer, Hodgkin lymphoma) as well as other co-morbidities including metabolic syndrome and cardiovascular diseases [5, 6]. The evidence on the risk of colorectal, breast and prostate cancers among individuals with HIV as compared to the general population remains very limited and inconsistent [6–10], but, individuals with HIV appear to have poorer outcomes after colon, breast and prostate cancers as compared to the general population [11–13]. Thus, this large proportion of older individuals presents challenges for chronic disease prevention and early cancer detection.

Comparison of the cancer rates among persons living with HIV (HIV/AIDS Cancer Match [HACM] Study) and those for the general US population from Surveillance, Epidemiology, and End Results program, in 2010 showed that AIDS-defining cancers represented one third of all incident cancers among persons living with HIV; two thirds were represented by non-AIDS-defining cancers [14]. There was a 50% excess in total cancer incidence (both AIDS defining and non-AIDS defining malignancies) in people living with HIV/AIDS when compared to the general US population [14].

Several healthy lifestyle factors have been previously suggested as general guidelines for cancer prevention, including smoking cessation, compliance with recommended alcohol consumption limits (≤ 7 drinks per week for females and ≤ 14 drinks per week for males), sustaining normal Body Mass Index (BMI), engaging in adequate physical activity and adherence to recommended cancer screening [15, 16]. In addition, screening for potential

co-infections such as Hepatitis-C virus (HCV) and long term adherence to recommended anti-retroviral treatment are recommended for people living with HIV/AIDS [5, 17].

The prevalence of healthy behaviors for cancer prevention among HIV/AIDS populations is poorly described. Previous studies in the general population and among cancer survivors suggest that these behaviors and adherence to cancer screening practices may differ by gender, race, education, and, among cancer survivors, also by the time since diagnosis [18–22]. The evidence on these differences among persons living with HIV remains extremely limited, but a few previous studies suggest that some of these behaviors and screening practices may differ by sex and time since HIV diagnosis in HIV/AIDS populations [23–25]. To address these knowledge gaps, we examined the prevalence of selected healthy behaviors including never smoking or smoking cessation, adhering to recommended alcohol consumption limits, maintaining a healthy body weight, engaging in physical activity and compliance with age-appropriate cancer screening practices in a cohort of persons living with HIV in Florida. We next compared distributions of these factors by sex and time since HIV diagnosis. We focus on differences by sex and time since HIV diagnosis as in our study population the distribution of other factors that may influence these behaviors and screening practices would not allow us to have sufficient number of individuals in certain strata thus preventing us from obtaining meaningful results.

Methods

Study population

This study was conducted using the data collected by the Florida Cohort between 2014 and 2017. Details of this cohort has been previously described in detail [26]. Briefly, the Florida Cohort recruited individuals aged 18 or older who are living with HIV/AIDS to assess demographic, behavioral and social determinants associated with the disease outcomes and access to care. The study uses a convenience sample recruitment strategy in a collaborative network of county health departments and community clinics including sites at Lake City, Gainesville, Tampa, Orlando, Sanford, Ft. Lauderdale, and Miami in the State of Florida.

At the time of recruitment, all participants completed a baseline questionnaire on demographics, selected lifestyle factors, and medication use (available from <http://sharc-research.org/projects/#Cohort>). The follow-up questionnaire was administered on average 8 months after the recruitment (inter quartile range 6–11 months) to collect information on detailed demographics, socioeconomic characteristics, basic health parameters, access to health care, risk behaviors related to HIV/AIDS (including substance use and sexual behaviors), and cancer screening. Additionally, for each participant in the Florida Cohort, medical history (such as HIV viral load, CD4 T-cell count, height and weight) was collected through electronic medical record abstraction and linkage to the Enhanced HIV/AIDS Reporting System (eHARS) database which is a real time medical information collection application provided by the Centers for Disease Control and Prevention [27].

From 902 participants in the Florida Cohort, 57% (N=517) completed the follow-up questionnaire and were included in the current analysis. Characteristics of individuals in this analysis and all cohort participants were similar. This study was approved by the University

of Florida, Florida International University, and the Florida Department of Health Institutional Review Boards (IRBs). All participants provided written informed consent.

Healthy behaviors and cancer screening

We included the following lifestyle and behavioral factors in the analysis: body mass index (BMI), smoking status, alcohol use, ability to do physical activity, and age-appropriate screening for female breast (mammogram), cervical (cervical Pap-smear), colorectal (colonoscopy), prostate and anal (anal Pap-smear) cancers.

Height and weight from the medical records were used to calculate BMI (kg/m^2). BMI was categorized as underweight ($<18.5 \text{ kg}/\text{m}^2$), normal weight ($18.5\text{--}24.9 \text{ kg}/\text{m}^2$), overweight ($25.0\text{--}29.9 \text{ kg}/\text{m}^2$) and obese ($\geq 30.0 \text{ kg}/\text{m}^2$) [28]. Alcohol use within past six months and physical activity within the past week were self-reported on the follow-up questionnaire. Tobacco smoking status was self-reported on the baseline and follow-up questionnaire. Because the baseline questionnaire collected information on current and ever smoking but the follow-up questionnaire only asked about current smoking status, information from both questionnaires were combined to categorize the smoking status as never, former and current smoking. Alcohol drinking was defined according to the Dietary Guidelines for Americans for 2015–2020 as follows: no alcohol consumption, current use below the recommended limit (≤ 7 drinks per week for females or ≤ 14 drinks per week for males) and current use above the recommended limit [29]. Physical activity in the Florida Cohort was assessed using the frequency of engaging in physical activity that was long enough to work up a sweat. Finally, we calculated the duration since HIV diagnosis using the date of HIV diagnosis provided by eHARS.

In the follow-up questionnaire, participants were asked to indicate the frequency of screening for each of the cancers as none, within past 3 years, and more than 3 years ago. Based on American Cancer Society (ACS) screening recommendations for the general public, sex-specific screening practices were described only for individuals who reached the recommended age for starting the screening for each screening modality (age 50 years for colon and prostate cancer, age 40 years for breast cancer and age 21 years for cervical cancer) [30, 31]. However, there are no national guidelines for routine anal cancer screening in the general population due to its very low prevalence [32, 33].

Statistical analysis

The study population was first described in terms of important socio-demographic characteristics (age, sex, race, education, health insurance, sexual orientation reported at baseline, marital status reported at baseline) and HIV/AIDS-related measures (CD4 cell count categories based on clinical significance of HIV continuum: <199 cells/ml, 200 to 349 cells/ml and more than 350 cells/ml [34], use of HIV anti-retroviral drugs and Hepatitis-C virus test status at baseline). The crude and age-adjusted prevalence of selected healthy behaviors and cancer screening was then described overall as well as by sex (male or female) and years since HIV diagnosis (dichotomized based on the median in the study population, ≤ 13 vs. >13 years). The prevalence across the strata was compared using Chi square test or Fisher's exact test where appropriate (for crude rates) or Mantel–Haenszel test

(for adjusted rates). All analyses were conducted using SAS version 9.4 (SAS Institute, Inc., Cary, NC, USA).

Results

In our study population, majority of the individuals were older than 50 years (54.5%) and males (61.1 %). Overall, 36.3% of the participants were obese, 9.0% were non-compliant to recommended alcohol intake limits, 50.7% were current smokers and 12.4% reported never engaging in physical activity that was long enough to work up a sweat. Characteristics of the 517 study participants, overall and by sex, are presented in Table 1.

The prevalence of healthy behaviors is presented in Table 2. In the analysis by sex, prevalence of obesity among females was significantly higher than that among males (56.5% vs. 22.2%, $p<0.0001$). The prevalence of alcohol consumption above the recommended limit was similar in both males and females (8.3% vs. 8.5%), but males were more likely to consume alcohol below recommended limit (61.0 vs. 43.2, $p<0.001$). No differences were noted for the prevalence of current smoking and never engaging in physical activity by sex. These prevalence patterns remained similar in age-adjusted analysis (Table 3). In the analysis by the years since HIV diagnosis, the prevalence of overweight or obese BMI among males was significantly higher in those who had HIV diagnosis for more than 13 years (overweight: 41.7% vs. 28.2%; obese: 25.0% vs. 20.1%, $p=0.02$) (Table 2). After adjustment for age, these differences in BMI were no longer significant (Table 3). We found no statistically significant differences in the prevalence of alcohol consumption, smoking, and physical activity by the time since HIV diagnosis among males and females (Table 2). These findings remained unchanged after age-adjustment (Table 3).

Prevalence of sex-specific, age-appropriate cancer screening practices among individuals at the screening recommended age is presented in Table 4. Among males, 64.8% never had an anal Pap-smear, 36.2% never had a colonoscopy, and 38.9% never had prostate cancer screening. Among females, 50.0% never had an anal Pap-smear (p for difference by sex <0.0001), 46.5% never had a colonoscopy (p for difference by sex $=0.28$), 7.9% never had a cervical Pap-smear, and 12.7% never had a mammogram. These results were similar after adjustment for age (Table 5). In the analysis by the years since HIV diagnosis, males who had HIV for ≤ 13 years were more likely to never have had colonoscopy as compared to those with time since diagnosis >13 years (49.1% vs. 28.3%, $p=0.02$). These differences in colonoscopy were still significant after age-adjustment ($p=0.03$) (Table 5). None of the other screening procedures showed statistically significant differences by the time since HIV diagnosis among males and females (Table 4) and these findings remained similar after adjustment for age (Table 5).

Discussion

This study describes disparities in healthy behaviors and cancer screening practices by sex and years since HIV diagnosis among persons living with HIV in Florida. Our findings suggest that the prevalence of obesity and selected cancer screening practices differ by sex

and years since HIV diagnosis suggesting a need for tailored cancer prevention efforts via long-term sex-specific interventions among persons living with HIV.

Health compromising behaviors such as not maintaining a normal body weight, physical inactivity, smoking, heavy drinking, and not adhering to recommended cancer screening practices can increase the risk of chronic diseases including cancer [15, 35, 36]. Previous studies have also shown that individuals with health compromising behaviors such as smoking and heavy drinking were further more likely to underutilize recommended cancer screening [37–39]. However, the evidence on healthy behaviors and screening practices among persons living with HIV remains extremely limited. A study by Mdodo et al. has shown that people with HIV have a greater prevalence of health compromising behaviors as compared to the general population [40], which may explain the increased prevalence of chronic diseases and non-AIDS defining cancers in persons living with HIV [6, 14, 33, 41, 42]. Therefore, tailored preventive approaches are needed in this population subgroup to alter unhealthy behaviors and improve long-term health outcomes [43]. Our study adds to the very limited evidence on the prevalence of these risk factors and adherence to cancer screening among persons living with HIV thus providing further insights into potential behavioral disparities between persons living with HIV and general population.

Our study population had similar prevalence of obesity and overweight BMI as compared to the general population (65.5% vs. 62.2%, respectively)[44]. However, in our study we found that the prevalence of obesity and overweight BMI was higher among females as compared to males in contrast to the findings in the general population where males appear to have higher prevalence (69.0% vs. 55.4%) [44]. Additionally, among males in the study population, the prevalence of obesity and overweight BMI was higher in those who had HIV diagnosis for >13 years. Núñez-Rocha et al. reported time since HIV diagnosis as a positive and independent predictor of obesity and overweight BMI among persons living with HIV [23]. Although further studies are needed to confirm our findings, the results suggest that the prevalence of obesity and overweight BMI may differ by sex and time since HIV diagnosis.

Persons living with HIV appeared to have a higher prevalence of non-compliance to the recommended limits of alcohol intake when compared to the general population (8.2% vs. 5.4%, respectively) [44]. Our findings on differences in alcohol consumption by sex were similar to those from the general population (non-compliance to drinking limits: 6.1% for males vs. 4.8% for females) [44]. The prevalence of current smoking was similar among males and females in our study population, while in the general population prevalence of current smoking was found to be slightly higher among males as compared to females (22.4% vs. 18.0%, respectively) [44].

We found a higher prevalence of never having physical activity among females as compared to males. Although we cannot directly compare physical activity status of the general population to the study population due to the different format of the physical activity-related items of our questionnaire and existing HIV related disabilities, prevalence of inactivity by sex in our study population (14.5% for females vs. 10.8% for males) had a similar trend as that in the general population (36.1% for females vs. 31.4% for males) [44].

In our study, we used cancer screening guidelines of the American Cancer Society (ACS) to investigate differences in cancer screening practices [16, 36]. Applicability of ACS screening guidelines to persons living with HIV has been confirmed in previous studies [33, 45]. However, there are no national recommendations for routine anal cancer screening in the general population due to its very low prevalence [32, 33]. Previous studies also do not report on anal cancer screening in the general population [41]. According to the U.S. Department of Health and Human Services, some experts recommend regular digital rectal exams and anal cytology for men and women who live with HIV[46]. The New York State AIDS Institute recommends baseline and annual screening with anal cytology for men living with HIV who have sex with men, women with a history of cervical or vulvar dysplasia, and persons with a history of anal condylomas, and the annual digital ano-rectal examination (DARE) for all persons with HIV [27] due to the strong association between HIV and the risk of anal cancer [33, 47]. A recent review by Goedert et al recommends anal pap cytology testing or DARE only for HIV infected men who have sex with men [32]. Finally, the U.S. Department of Veterans Affairs recommends all men and women living with HIV to be screened at baseline and annually using anal Pap-smear or digital rectal examination [48].

We found that males appeared to be more likely to adhere to colorectal cancer screening recommendations as compared to females, but these differences were not statistically significant. These patterns were consistent with the previously reported results for the general population (32.2% for males and 29.8% for females) [49, 50]. In addition, among male participants at the screening recommended age for colorectal cancer those with 13 years since HIV diagnosis were more likely to report never having a colonoscopy as compared to those with >13 years since HIV diagnosis.

In our study, females were more likely to undergo anal cancer screening as compared to males. Overall, the evidence on anal cancer screening in persons living with HIV by sex is extremely limited. However, due to the absence of established national anal cancer screening guidelines, inconsistencies in expert recommendations, and absence of estimates from the general population, the prevalence rates of anal cancer screening in our study must be interpreted with caution. Although Wells et al. reported that women are less likely to be screened for anal cancer compared to men (OR=0.2; $p=0.007$, adjusted for age and heterosexual/men who have sex with men status), we cannot directly compare these findings to our results because of the differences in statistical approaches [41]. Further, no data are available on agreement between self-reported and confirmed history of anal Pap-smear and thus potential misclassification cannot be assessed. Finally, it is also possible that providers implementing cervical Pap-smear simultaneously obtain a sample for anal Pap-smear in this high-risk population of women, but it is impossible to know for sure if this is the reason for the observed patterns.

Prevalence of ever having prostate cancer screening among our male participants was greater than the prevalence of prostate specific antigen (PSA) testing among men in the general population (60.4% vs. 44.1%, respectively) [36]. These findings are consistent with the previously reported prevalence of prostate cancer screening among persons living with HIV [51].

In our study, the prevalence of ever having a mammogram among women was greater than that in the general population (87.6% vs. 64.0%, respectively) [52]. Similarly, the prevalence of cervical Pap-smear screening in the study population appeared to be greater than that in the general population (92.7% vs. 70.2%, respectively) [52]. From this study, we cannot determine the reasons for higher utilization of cervical Pap-smear screening and the adherence to the recommended screening intervals among women living with HIV as compared to the general population. Our findings may be in part explained by the larger number of encounters with primary care physicians in women diagnosed with HIV as compared to the general population [53, 54].

Our study adds to the very limited evidence on the prevalence of healthy behaviors and cancer screening practices in persons living with HIV using an established cohort with comprehensive data collection and prospective follow-up. Our findings further strengthen the understanding of sex-related disparities in healthy behaviors and cancer screening practices, and add to the very limited evidence on these differences by years since HIV diagnosis. All HIV diagnoses in Florida Cohort have been confirmed with medical records. Similarly, BMI data were also available from the medical records. However, our study has a few limitations. The format of the questions on cancer screening in Florida Cohort (available from <http://sharc-research.org/projects/#Cohort>) differed from the questions used to assess cancer screening in the general population thus limiting to some degree the direct comparison between our findings and those from the general population. Second, the potential for misclassification of self-reported screening practices and healthy behaviors cannot be ruled out. However, previous studies suggest good agreement between self-reported vs. medical record-confirmed screening practices that differs by cancer type (overall agreement: 83% for colorectal cancer screening; 61% for prostate cancer screening; 62% for mammography; and 61% for cervical Pap-smear) and some healthy behaviors (95% for smoking status and 86% for alcohol intake) [55, 56]. The correlation coefficient for self-reported and objectively measured physical activity was reported to be moderate (spearman correlation coefficient=0.56) [57].

In conclusion, we examined the prevalence of healthy behaviors and cancer screening among persons living with HIV. Although future studies are needed to further confirm our findings by sex and years since HIV diagnosis, the results suggest that specific subgroups of persons living with HIV may benefit from long-term, tailored public health interventions aimed at improved lifestyle and cancer screening in this population segment.

Acknowledgements

This work was supported by NIAAA grant U24 AA022002, and by the State of Florida via the University of Miami Miller School of Medicine Institute for AIDS and Emerging Infectious Diseases. The Enhanced HIV/AIDS Reporting System data were provided from the HIV data center, Infectious Disease Surveillance Division, Florida Department of Health.

References

1. Chirch LM, Hasham M, and Kuchel GA (2014). HIV and aging: a clinical journey from Koch's postulate to the chronic disease model and the contribution of geriatric syndromes. *Curr Opin HIV AIDS*. 9:405–11. [PubMed: 24824883]

2. Deeks SG, Lewin SR, and Havlir DV (2013). The end of AIDS: HIV infection as a chronic disease. *Lancet*. 382:1525–33. [PubMed: 24152939]
3. Risher KA, Kapoor S, Daramola AM, Paz-Bailey G, Skarbinski J, Doyle K, and Shah M (2017). Challenges in the Evaluation of Interventions to Improve Engagement Along the HIV Care Continuum in the United States: A Systematic Review. *AIDS and Behavior*:1–23.
4. Center for Disease Control and Prevention. HIV Among People Aged 50 and Over. HIV/AIDS age 2017; Available from: <https://www.cdc.gov/hiv/group/age/olderamericans/> Access date: April 10, 2018.
5. Parka LS H-RR, Silverberg MJ, Crothers KA, Dubrow R. (2016). Prevalence of non-HIV cancer risk factors in persons living with HIV/AIDS. *AIDS*. 30:273–91. [PubMed: 26691548]
6. Shiels MS and Engels EA (2017). Evolving epidemiology of HIV-associated malignancies. *Curr Opin HIV AIDS*. 12:6–11. [PubMed: 27749369]
7. Coghill AE, Engels EA, Schymura MJ, Mahale P, and Shiels MS (2018). Risk of Breast, Prostate, and Colorectal Cancer Diagnoses Among HIV-Infected Individuals in the United States. *Journal of the National Cancer Institute*. 110:959–966. [PubMed: 29529223]
8. Patel P, Hanson DL, Sullivan PS, and et al. (2008). Incidence of types of cancer among hiv-infected persons compared with the general population in the united states, 1992–2003. *Annals of Internal Medicine*. 148:728–736. [PubMed: 18490686]
9. Shiels MS, Islam JY, Rosenberg PS, Hall HI, Jacobson E, and Engels EA (2018). Projected Cancer Incidence Rates and Burden of Incident Cancer Cases in HIV-Infected Adults in the United States Through 2030. *Annals of Internal Medicine*. 168:866–873. [PubMed: 29801099]
10. Presti C, Haslinger M, Wehner P (2017) Breast Cancer in HIV-positive Patients: A Multi-Institutional Retrospective Review. *J Prev Med Healthc* 1(2): 1009
11. Silberstein J, Downs T, Lakin C, and Kane CJ (2008). HIV and prostate cancer: a systematic review of the literature. *Prostate Cancer And Prostatic Diseases*. 12:6. [PubMed: 18711409]
12. Gomez A, Montero AJ, and Hurley J (2015). Clinical outcomes in breast cancer patients with HIV/AIDS: a retrospective study. *Breast Cancer Research and Treatment*. 149:781–788. [PubMed: 25663516]
13. Sigel C, Cavalcanti MS, Daniel T, Vakiani E, Shia J, and Sigel K (2016). Clinicopathologic Features of Colorectal Carcinoma in HIV-Positive Patients. *Cancer epidemiology, biomarkers & prevention : a publication of the American Association for Cancer Research, cosponsored by the American Society of Preventive Oncology*. 25:1098–1104.
14. Robbins HA, Pfeiffer RM, Shiels MS, Li J, Hall HI, and Engels EA (2015). Excess cancers among HIV-infected people in the United States. *J Natl Cancer Inst*. 107.
15. Colditz GA, Wolin KY, and Gehlert S (2012). Applying What We Know to Accelerate Cancer Prevention. *Sci Transl Med*. 4:127rv4.
16. Kushi LH, Doyle C, McCullough M, Rock CL, Demark-Wahnefried W, Bandera EV, Gapstur S, Patel AV, Andrews K, and Gansler T (2012). American Cancer Society guidelines on nutrition and physical activity for cancer prevention. *CA: A Cancer Journal for Clinicians*. 62:30–67. [PubMed: 22237782]
17. US Department of Veteran Affairs 2016 HIV/AIDS primary care manual 2011; Available from: <https://www.hiv.va.gov/provider/manual-primary-care/index.asp>. Access date: December 15, 2018.
18. Park B, Kong S-Y, Kim J, Kim Y, Park IH, Jung S-Y, and Lee ES (2015). Health Behaviors of Cancer Survivors in Nationwide Cross-Sectional Survey in Korea: Higher Alcohol Drinking, Lower Smoking, and Physical Inactivity Pattern in Survivors with Higher Household Income. *Medicine*. 94:e1214–e1214. [PubMed: 26252280]
19. Naik H, Qiu X, Brown MC, Eng L, Pringle D, Mahler M, Hon H, Tiessen K, Thai H, Ho V, Gonos C, Charow R, Pat V, Irwin M, Herzog L, Ho A, Xu W, Jones JM, Howell D, and Liu G (2016). Socioeconomic status and lifestyle behaviours in cancer survivors: smoking and physical activity. *Current oncology (Toronto, Ont.)*. 23:e546–e555.
20. Nayak P, Paxton RJ, Holmes H, Thanh Nguyen H, and Elting LS (2015). Racial and ethnic differences in health behaviors among cancer survivors. *Am J Prev Med*. 48:729–36. [PubMed: 25998923]

21. National Research Council (US) Panel on Race, Ethnicity, and Health in Later Life; Anderson NB, Bulatao RA, Cohen B, editors. Critical Perspectives on Racial and Ethnic Differences in Health in Late Life Washington (DC): National Academies Press (US); 2004 12, Racial/Ethnic Disparities in Health Behaviors: A Challenge to Current Assumptions. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK25518/>.
22. Yaghjian L, Wolin K, Chang S-H, and Colditz G (2014). Racial disparities in healthy behaviors and cancer screening among breast cancer survivors and women without cancer: National Health Interview Survey 2005. *Cancer causes & control* : CCC. 25:605–614. [PubMed: 24634091]
23. Nunez-Rocha GM, Wall KM, Chavez-Peralta M, Salinas-Martinez AM, and Benavides-Torres RA (2013). Nutritional care, time period since diagnosis, demographics and body mass index in HIV/AIDS patients. *Rev Invest Clin*. 65:291–9. [PubMed: 24304729]
24. Friedemann-Sánchez G, Griffin JM, and Partin MR (2007). Gender differences in colorectal cancer screening barriers and information needs. *Health Expectations*. 10:148–160. [PubMed: 17524008]
25. Yager SS, Chen L, and Cheung WY (2014). Sex-based disparities in colorectal cancer screening. *Am J Clin Oncol*. 37:555–60. [PubMed: 23466582]
26. Lucero RJ, Frimpong JA, Fehlbeg EA, Bjarnadottir RI, Weaver MT, Cook C, ... & Cook RL (2017). (2017). The Relationship Between Individual Characteristics and Interest in Using a Mobile Phone App for HIV Self-Management: Observational Cohort Study of People Living With HIV. *JMIR mHealth and uHealth*. 5.
27. Nebraska D of Health and human services. eHARS (Enhanced HIV/AIDS Reporting System). 2017 <http://dhhs.ne.gov/publichealth/DataCenter/Pages/eHARS.aspx> Access date April 10, 2018
28. Centers for Disease Control and Prevention. About adult BMI 2017 https://www.cdc.gov/healthyweight/assessing/bmi/adult_bmi/index.html Access date: April 10, 2018
29. U.S. Department of Health and Human Services and US Department of Agriculture, Dietary guidelines for Americans 2015–2020. 6 ed 2015, Washington, DC: U.S. Government Printing Office.
30. Wolf AMD, Fonham ETH, Church TR, Flowers CR, Guerra CE, LaMonte SJ, Etzioni R, McKenna MT, Oeffinger KC, Shih Y-CT, Walter LC, Andrews KS, Brawley OW, Brooks D, Fedewa SA, Manassaram-Baptiste D, Siegel RL, Wender RC, and Smith RA (2018). Colorectal cancer screening for average-risk adults: 2018 guideline update from the American Cancer Society. *CA: A Cancer Journal for Clinicians*. 68:250–281. [PubMed: 29846947]
31. American Cancer Society guidelines for the early detection of cancer (2018). <https://www.cancer.org/healthy/find-cancer-early/cancer-screening-guidelines/american-cancer-society-guidelines-for-the-early-detection-of-cancer.html> Access date: April 10, 2018
32. Leeds IL and Fang SH (2016). Anal cancer and intraepithelial neoplasia screening: A review. *World journal of gastrointestinal surgery*. 8:41–51. [PubMed: 26843912]
33. Sigel K, Dubrow R, Silverberg M, Crothers K, Braithwaite S, and Justice A (2011). Cancer Screening in Patients Infected with HIV. *Curr HIV/AIDS Rep*. 8:142–52. [PubMed: 21695529]
34. Mellors JW, Munoz A, Giorgi JV, Margolick JB, Tassoni CJ, Gupta P, Kingsley LA, Todd JA, Saah AJ, Detels R, Phair JP, and Rinaldo CR (2017). Plasma Viral Load and CD4+ Lymphocytes as Prognostic Markers of HIV-1 Infection. *Annals of Internal Medicine*. 126:946–954.
35. Bauer UE, Briss PA, Goodman RA, and Bowman BA (2014). Prevention of chronic disease in the 21st century: elimination of the leading preventable causes of premature death and disability in the USA. *Lancet*. 384:45–52. [PubMed: 24996589]
36. Smith RA, Brooks D, Cokkinides V, Saslow D, & Brawley OW (2013). Cancer screening in the United States, 2013. *CA: A Cancer Journal for Clinicians*. 63:87–105.
37. Aserlind A, Maguire K, Duthely L, Wennin S, & Potter J (2017). Women Living with HIV over Age of 65: Cervical Cancer Screening in a Unique and Growing Population. *Infectious Diseases in Obstetrics and Gynecology*. 2017.
38. Bryan L, Westmaas L, Alcaraz K, and Jemal A (2014). Cigarette smoking and cancer screening underutilization by state: BRFSS 2010. *Nicotine Tob Res*. 16:1183–9. [PubMed: 24696511]
39. Mu L and Mukamal KJ (2016). Alcohol consumption and rates of cancer screening: Is cancer risk overestimated? *Cancer Causes Control*. 27:281–9. [PubMed: 26590914]

40. Mdodo R, Frazier EL, Dube SR, Mattson CL, Sutton MY, Brooks JT, and Skarbinski J (2015). Cigarette smoking prevalence among adults with HIV compared with the general adult population in the United States: cross-sectional surveys. *Ann Intern Med.* 162:335–44. [PubMed: 25732274]
41. Wells JS, Holstad MM, and Watkins Bruner D (2017). Sociodemographic Predictors of Anal Cancer Screening and Follow-up in Human Immunodeficiency Virus-Infected Individuals. *Cancer Nurs.*
42. Brooks JT, Buchacz K, Gebo KA, and Mermin J, HIV Infection and Older Americans: The Public Health Perspective, in *Am J Public Health.* 2012 p. 1516–26. [PubMed: 22698038]
43. Swendeman D, Ingram BL, and Rotheram-Borus MJ (2009). Common elements in self-management of HIV and other chronic illnesses: an integrative framework. *AIDS Care.* 21:1321–34. [PubMed: 20024709]
44. Schoenborn CA, Adams PF, Peregoy JA, (2013). Health behaviors of adults: United States, 2008–2010. *Vital Health Stat.* 10.
45. Goedert JJ, Hosgood HD, Biggar RJ, Strickler HD, and Rabkin CS (2016). Screening for Cancer in Persons Living with HIV Infection. *Trends Cancer.* 2:416–28. [PubMed: 27891533]
46. US Department of Health and Human Services. Guidelines for Prevention and Treatment of Opportunistic Infections in HIV-Infected Adults and Adolescents 2018 [Available from: https://aidsinfo.nih.gov/contentfiles/lvguidelines/glcchunk/glcchunk_343.pdf. Access date: January 10, 2019
47. Xu J and Zhou H (2017). Screening for Anal Cancer in HIV Positive Patients: Should We Make It A Standard-of-care? *J Invest Surg:*1–2.
48. US Department of Veteran Affairs. Anal Dysplasia: Primary Care of Veterans With HIV Washington DC: U.S. Department of Veterans Affairs; 2018 [Available from: <https://www.hiv.va.gov/provider/manual-primary-care/anal-dysplasia.asp>.
49. Meissner HI, Breen N, Klabunde CN, and Vernon SW (2006). Patterns of Colorectal Cancer Screening Uptake among Men and Women in the United States.
50. Chao A, Connell CJ, Cokkinides V, Jacobs EJ, Calle EE, and Thun MJ (2011). Underuse of Screening Sigmoidoscopy and Colonoscopy in a Large Cohort of US Adults. *10.2105/AJPH.* 94.10.1775.
51. Marcus JL, Chao CR, Leyden WA, Xu L, Klein DB, Horberg MA, Towner WJ, Quesenberry CP Jr., Abrams DI, Van Den Eeden SK, and Silverberg MJ (2014). Prostate cancer incidence and prostate-specific antigen testing among HIV-positive and HIV-negative men. *J Acquir Immune Defic Syndr.* 66:495–502. [PubMed: 24820107]
52. National Center for Health S, Health, United States, in Health, United States, 2016: With Chartbook on Long-term Trends in Health. 2017, National Center for Health Statistics (US): Hyattsville (MD).
53. US Department of Health and Human Services. Seeing Your Health Care Provider: Managing Your Appointments. 2017 [Available from: <https://www.hiv.gov/hiv-basics/staying-in-hiv-care/provider-visits-and-lab-test/seeing-your-health-care-provider>] Access day: January 10, 2019
54. Mutasa-Apollo T, Ford N, Wiens M, Socias ME, Negussie E, Wu P, Popoff E, Park J, Mills EJ, and Kanfers S (2017). Effect of frequency of clinic visits and medication pick-up on antiretroviral treatment outcomes: a systematic literature review and meta-analysis. *Journal of the International AIDS Society.* 20:21647. [PubMed: 28770599]
55. Ferrante JM, Ohman-Strickland P, Hahn KA, Hudson SV, Shaw EK, Crosson JC, and Crabtree BF (2008). Self-report versus Medical Records for Assessing Cancer-Preventive Services Delivery. *Cancer Epidemiol Biomarkers Prev.* 17:2987–94. [PubMed: 18990740]
56. Simons JS, Wills TA, Emery NN, and Marks RM (2015). Quantifying alcohol consumption: Self-report, transdermal assessment, and prediction of dependence symptoms. *Addict Behav.* 50:205–12. [PubMed: 26160523]
57. Norman A, Belloc R, Bergstrom A, and Wolk A (2001). Validity and reproducibility of self-reported total physical activity--differences by relative weight. *Int J Obes Relat Metab Disord.* 25:682–8. [PubMed: 11360151]

Table 1.

Characteristics of study participants, by sex (Number [%])

Characteristic	Total (n=517)	Male (n=313)	Female (n=199)	p ^a
Age				
<50 years	235 (45.5 %)	144 (46.2 %)	87 (43.7 %)	0.59
50 years	281 (54.5 %)	168 (53.9 %)	112 (56.3 %)	
Race				
Hispanic	86 (16.8 %)	65 (20.8 %)	21 (10.6 %)	0.01
Non-Hispanic, White	98 (19.1 %)	64 (20.5 %)	34 (17.1 %)	
Non-Hispanic, Black	308 (60.2 %)	172 (55.0 %)	136 (68.3 %)	
Non-Hispanic, Other	20 (3.9 %)	12 (3.8 %)	8 (4.0 %)	
Education				
Less than High-school	170 (33.3 %)	85 (27.2 %)	85 (42.9 %)	<0.0001
High-school or equivalent	164 (32.1 %)	99 (31.6 %)	65 (32.8 %)	
More than High-school	177 (34.6 %)	129 (41.2 %)	48 (24.2 %)	
Employment Status				
Unemployed	130 (26.2 %)	87 (28.5 %)	43 (22.4 %)	0.03
Unable to work/ Disabled	252 (50.7 %)	140 (45.9 %)	112 (58.3 %)	
Employed	115 (23.1 %)	78 (25.6 %)	37 (19.3 %)	
Marital Status				
Married/ Long term partner	99 (19.4 %)	55 (17.6 %)	44 (22.2 %)	0.2
Not Married/ Other	412 (80.6 %)	258 (82.4 %)	154 (77.8 %)	
Health Insurance				
Insured	503 (99.2 %)	304 (98.7 %)	194 (100.0 %)	0.14
Un-Insured	4 (0.8 %)	4 (1.3 %)	0 (0.0 %)	
Sexual Orientation				
Heterosexual	285 (57.8 %)	111 (36.4 %)	174 (92.6 %)	<0.0001
Homosexual	157 (31.9 %)	152 (49.8 %)	5 (2.7 %)	
Other	51 (10.3 %)	42 (13.8 %)	9 (4.7 %)	
CD4 T-Cell Categories				
0–199 cells/ml	41 (8.2 %)	28 (9.3 %)	13 (6.7 %)	0.01

Characteristic	Total (n=517)	Male (n=313)	Female (n=199)	P ^a
200–349 cells/ml	83 (16.7 %)	60 (20.0 %)	21 (10.9 %)	
350+ cells/ml	374 (75.1 %)	212 (70.7 %)	159 (82.4 %)	
Anti-Retroviral Medication Use				
Not Using	18 (3.6 %)	9 (3.0 %)	9 (4.7 %)	0.31
Using	483 (96.4 %)	296 (97.1 %)	182 (95.3 %)	
Hepatitis C test status				
Not Done/Do not know	126 (25.2 %)	71 (23.3 %)	55 (28.21 %)	0.29
Done/Positive result	110 (22.0 %)	73 (23.9 %)	37 (19.0 %)	
Done/Negative result	264 (52.8 %)	161 (52.8 %)	103 (52.8 %)	

Note: percentages are calculated from non-missing data

^aP-value for Chi-square test or Fisher's exact test

Table 2.

Prevalence of healthy behaviors in the study population, by sex and years since HIV diagnosis (Number [%])

Behavior categories	Male (n=313)				Female (n=199)				p ^a for comparison by sex
	Years since HIV diagnosis				Years since HIV diagnosis				
	Total	13 years (n=167)	>13 years n=141)	p ^a	Total	13 years (n=105)	>13 years (n=93)	p ^a	
BMI ^b									
Underweight	5 (1.9 %)	3 (2.0 %)	2 (1.9 %)	0.02	8 (4.5 %)	5 (5.1 %)	3 (3.8 %)	0.74	<0.0001
Normal Weight	109 (41.8 %)	74 (49.7 %)	34 (31.5 %)		30 (17.0 %)	14 (14.3 %)	16 (20.3 %)		
Overweight	89 (34.1 %)	42 (28.2 %)	45 (41.7 %)		39 (22.0 %)	23 (23.5 %)	16 (20.3 %)		
Obese	58 (22.2 %)	30 (20.1 %)	27 (25.0 %)		100 (56.5 %)	56 (57.1 %)	44 (55.7 %)		
Alcohol consumption ^c									
No consumption	96 (30.7 %)	44 (26.4 %)	50 (35.5 %)	0.20	96 (48.2 %)	43 (41.0 %)	52 (55.9 %)	0.07	<0.001
Below the recommended limit	191 (61.0 %)	107 (64.1 %)	81 (57.5 %)		86 (43.2 %)	50 (47.6 %)	36 (38.7 %)		
Above recommended limit	26 (8.3 %)	16 (9.6 %)	10 (7.1 %)		17 (8.5 %)	12 (11.4 %)	5 (5.4 %)		
Smoking									
Never Smoker	83 (27.2 %)	45 (27.8 %)	35 (25.4 %)	0.89	57 (30.7 %)	30 (30.3 %)	27 (31.4 %)	0.94	0.40
Former Smoker	69 (22.6 %)	37 (22.8 %)	32 (23.2 %)		33 (17.7 %)	17 (17.2 %)	16 (18.6 %)		
Current Smoker	153 (50.2 %)	80 (49.4 %)	71 (51.5 %)		96 (51.6 %)	52 (52.5 %)	43 (50.0 %)		
Frequency of physical activity									
Never	33 (10.7 %)	17 (10.4 %)	16 (11.5 %)	0.64	29 (14.7 %)	16 (15.2 %)	13 (14.3 %)	0.69	0.53
Rarely	91 (29.6 %)	46 (28.1 %)	42 (30.2 %)		60 (30.5 %)	35 (33.3 %)	25 (27.5 %)		
Sometimes	126 (40.9 %)	73 (44.5 %)	52 (37.4 %)		73 (37.1 %)	38 (36.2 %)	34 (37.4 %)		
Often	58 (18.8 %)	28 (17.1 %)	29 (20.9 %)		35 (17.8 %)	16 (15.2 %)	19 (20.9 %)		

Abbreviations: BMI – Body Mass Index

Note: percentages are calculated from non-missing data.

^ap-value for Chi-square test or Fisher's exact test.^bBMI was categorized as underweight (<18.5 kg/m²), normal weight (18.5–24.9 kg/m²), overweight (25.0–29.9 kg/m²) and obese (≥30.0 kg/m²) [28].

Alcohol consumption during the past six months defined as no alcohol consumption, alcohol consumption below the recommended limit of 14 drinks per week if male or 7 drinks per week if female and above the recommended limit [29].

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 3.

Age-adjusted prevalence of healthy behaviors in the study population, by sex and years since HIV diagnosis (Number [%])

Behavior categories	Male (n=307)				Female (n=198)				p ^d for comparison by sex
	Years since HIV diagnosis				Years since HIV diagnosis				
	Total	13 years (n=166)	>13 years (n=141)	p ^a	Total	13 years (n=105)	>13 years (n=93)	p ^a	
BMI ^d									
Underweight	5 (1.8%)	3 (1.7%)	2 (1.6%)	0.53	8 (4.3%)	5 (4.9%)	3 (1.6%)	0.56	<0.0001
Normal Weight	107 (40.8%)	73 (44.9%)	34 (27.7%)		29 (17.2%)	13 (13.7%)	16 (27.7%)		
Overweight	83 (34.0%)	40 (30.1%)	43 (36.2%)		36 (21.0%)	22 (23.8%)	14 (36.2%)		
Obese	56 (23.4%)	29 (23.3%)	27 (34.5%)		99 (57.5%)	55 (57.5%)	44 (34.5%)		
Alcohol Consumption ^c									
No consumption	94 (30.8%)	44 (29.5%)	50 (35.9%)	0.91	95 (48.2%)	43 (40.1%)	52 (57.4%)	0.05	<0.001
Below the recommended limit	188 (61.2%)	107 (62.0%)	81 (57.7%)		86 (43.6%)	50 (48.0%)	36 (36.7%)		
Above recommended limit	25 (8.0%)	15 (8.5%)	10 (6.4%)		17 (8.2%)	12 (11.9%)	5 (5.9%)		
Smoking Categories									
Never Smoker	80 (26.9%)	45 (27.6%)	35 (29.1%)	0.57	57 (32.6%)	30 (31.8%)	27 (33.2%)	0.81	0.40
Former Smoker	69 (22.3%)	37 (23.4%)	32 (18.2%)		33 (18.3%)	17 (16.7%)	16 (9.1%)		
Current Smoker	150 (50.8%)	79 (49.0%)	71 (52.7%)		95 (49.1%)	52 (51.5%)	43 (57.7%)		
Frequency of Physical Activity									
Never	32 (10.8%)	16 (10.4%)	16 (9.9%)	0.85	29 (14.8%)	16 (14.5%)	13 (14.0%)	0.69	0.46
Rarely	86 (28.9%)	45 (28.7%)	41 (32.7%)		58 (31.0%)	34 (34.9%)	25 (27.2%)		
Sometimes	124 (42.0%)	72 (44.4%)	52 (39.6%)		71 (36.9%)	37 (35.0%)	34 (32.2%)		
Often	55 (18.4%)	27 (16.5%)	28 (17.7%)		33 (17.3%)	156 (15.6%)	19 (26.6%)		

Note: percentages are calculated from non-missing data.

^aP-value for Mantel-Haenszel test.^bBMI was categorized as underweight (<18.5 kg/m²), normal weight (18.5–24.9 kg/m²), overweight (25.0–29.9 kg/m²) and obese (≥30.0 kg/m²) [28].

Alcohol consumption during the past six months defined as no alcohol consumption, alcohol consumption below the recommended limit of 14 drinks per week if male or 7 drinks per week if female and above the recommended limit [29].

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 4.

Prevalence of cancer screening practices in the study population, by sex and years since HIV diagnosis (Number [%])

Cancer Screening	Male				Female				p ^d for comparison by sex
	Years since HIV diagnosis				Years since HIV diagnosis				
	Total	13 years	>13 years	p ^a	Total	13 years	>13 years	p ^a	
Anal Pap-smear									
Never	173 (64.8 %)	101 (68.2 %)	72 (60.5 %)	0.41	90 (50.0 %)	48 (49.5 %)	42 (50.60 %)	0.96	<0.0001
Within past 3 years	71 (26.6 %)	36 (24.3 %)	35 (29.4 %)		84 (46.7 %)	46 (47.4 %)	38 (45.78 %)		
More than 3 years	23 (8.6 %)	11 (7.4 %)	12 (10.1 %)		6 (3.3 %)	3 (3.1 %)	3 (3.61 %)		
Colonoscopy									
Never	54 (36.2 %)	28 (49.1 %)	26 (28.3 %)	0.02	46 (46.5 %)	22 (46.8 %)	24 (46.2 %)	0.97	0.28
Within past 3 years	66 (44.3 %)	18 (31.6 %)	48 (52.2 %)		39 (39.4 %)	18 (38.3 %)	21 (40.4 %)		
More than 3 years	29 (19.5 %)	11 (19.3 %)	18 (19.6 %)		14 (14.1 %)	7 (14.9 %)	7 (13.5 %)		
Prostate									
Never	58 (38.9 %)	26 (46.4 %)	32 (34.4 %)	0.30	-	-	-	-	NA
Within past 3 years	73 (49.0 %)	25 (44.6 %)	48 (51.6 %)		-	-	-	-	
More than 3 years	18 (12.1 %)	5 (8.9 %)	13 (14.0 %)		-	-	-	-	
Cervical Pap-smear									
Never	-	-	-	-	15 (7.9 %)	7 (7.0 %)	8 (8.9 %)	0.50	NA
Within past 3 years	-	-	-	-	160 (84.2 %)	87 (87.0 %)	73 (81.1 %)		
More than 3 years	-	-	-	-	15 (7.9 %)	6 (6.0 %)	9 (10.0 %)		
Mammogram									
Never	-	-	-	-	20 (12.7 %)	6 (7.7 %)	14 (17.5 %)	0.11	NA
Within past 3 years	-	-	-	-	119 (75.3 %)	64 (82.1 %)	55 (68.8 %)		
More than 3 years	-	-	-	-	19 (12.0 %)	8 (10.3 %)	11 (13.8 %)		

Abbreviations: NA – not applicable

Note: percentages are calculated from non-missing data and cancer screenings groups are restricted to specified age ranges according to American Cancer Society guidelines, 2015 [25]; anal cancer (anal Pap-smear); no age restriction; colon cancer (colonoscopy) and prostate cancer screening: 50 years and older; breast cancer (mammogram); 40 years and older and cervical cancer (cervical Pap-smear); 21 years and above.

P-value for Chi-square test or Fisher's exact test.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 5.

Age-adjusted prevalence of cancer screening practices in the study population, by sex and years since HIV diagnosis (Number [%])

Cancer Screening	Male				Female				p ^d for comparison by sex
	Years since HIV diagnosis				Years since HIV diagnosis				
	Total	13 years	>13 years	p ^d	Total	13 years	>13 years	p ^d	
Anal Pap-smear									
Never	169 (64.9%)	98 (68.5%)	71 (53.6%)	0.55	89 (50.4%)	48 (48.5%)	41 (49.5%)	0.92	<0.0001
Within past 3 years	70 (26.7%)	35 (23.7%)	35 (38.5%)		84 (46.1%)	46 (48.5%)	38 (46.9%)		
More than 3 years	22 (8.4%)	11 (7.8%)	11 (7.9%)		6 (3.4%)	3 (3.1%)	3 (3.6%)		
Colonoscopy									
Never	54 (37.6%)	28 (50.8%)	26 (30.6%)	0.03	46 (44.6%)	22 (46.1%)	24 (43.6%)	0.97	0.53
Within past 3 years	66 (44.6%)	18 (31.5%)	48 (70.5%)		39 (40.0%)	18 (37.6%)	21 (40.2%)		
More than 3 years	28 (17.8%)	11 (17.7%)	17 (18.7%)		14 (15.3%)	7 (16.2%)	7 (13.6%)		
Prostate									
Never	58 (39.5%)	26 (47.7%)	32 (36.2%)	0.37	-	-	-	-	NA
Within past 3 years	73 (48.9%)	25 (43.2%)	48 (70.0%)		-	-	-		
More than 3 years	17 (11.5%)	5 (9.1%)	12 (13.5%)		-	-	-		
Cervical Pap-smear									
Never	-	-	-		14 (7.3%)	7 (6.3%)	7 (7.3%)	0.79	NA
Within past 3 years	-	-	-		160 (84.3%)	87 (87.0%)	73 (82.4%)		
More than 3 years	-	-	-		15 (8.4%)	6 (6.8%)	9 (10.3%)		
Mammogram									
Never	-	-	-		20 (12.4%)	6 (7.1%)	14 (17.6%)	0.12	NA
Within past 3 years	-	-	-		119 (75.5%)	64 (82.9%)	55 (68.8%)		
More than 3 years	-	-	-		19 (12.1%)	8 (10.9%)	11 (13.6%)		

Abbreviations: NA-not applicable

Note: percentages are calculated from non-missing data and cancer screenings groups are restricted to specified age ranges according to American Cancer Society guidelines, 2015 [25]; anal cancer (anal Pap-smear): no age restriction; colon cancer (colonoscopy) and prostate cancer screening: 50 years and older; breast cancer (mammogram): 40 years and older and cervical cancer (cervical pap-smear): 21 years and above.

P-value for Mantel-Haenszel test.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript