Published in final edited form as:

Psychopharmacology (Berl). 2019 July ; 236(7): 2027–2037. doi:10.1007/s00213-019-05191-6.

Persistent increase of I.V. cocaine self-administration in a subgroup of C57BL/6J male mice after social defeat stress

Danielle T. Arena¹, Herbert E. Covington III¹, Joseph F. DeBold¹, and Klaus A. Miczek^{1,2}

¹Dept. of Psychology, Tufts University, Medford, MA 02144

²Dept. of Neuroscience, Tufts University, Boston, MA 02111

Abstract

Rationale: Stressful life experiences can persistently increase the motivation for, and consumption of, intensely rewarding stimuli, like cocaine over time. In rodents, intermittent versus continuous exposure to social stress engender opposing changes to reward-related behavior, as measured by consumption of sucrose and cocaine.

Objective: The present study examines if the effects of intermittent versus continuous social stress on cocaine self-administration in mice parallel those seen in rats.

Methods: Both forms of social stress involve a brief daily physical confrontation with an aggressive resident for 10 consecutive days. Continuous social stress involves constant visual and olfactory exposure to an aggressive resident via habitation in a protected portion of the resident's home cage, while exposure to an aggressive resident during intermittent social stress is limited to a single, physical encounter per day. Implementing a femoral vein catheterization method for the first time in mice, we determined divergent changes to intravenous cocaine self-administration.

Results: Modestly increased cocaine self-administration after intermittent social stress was confirmed. In a subset of animals, continuous social stress in mice substantially increased cocaine self-administration and sucrose intake. By stark contrast, another subpopulation had substantial attenuation of cocaine self-administration and sucrose intake after continuous social stress.

Conclusions: Bimodal divergence in responding for rewarding stimuli including cocaine after social stress experience likely reflects two opposing forms of coping to continuous social stress that promote either a sensitization or attenuation of reward seeking.

Keywords

Cocaine; C57BL/6J mice; Self-administration; Social defeat stress; Intermittent versus chronic stress; Sucrose preference; Social interaction

Communicating Author Klaus A. Miczek; Klaus.Miczek@tufts.edu.

Conflict of Interest The authors declare that they have no conflict of interest.

Introduction

Coping with stress is necessary for survival, but uncontrollable stressful events can greatly exacerbate the onset of psychopathologies like drug abuse (Kleber and Gawin 1984; Der-Avakian et al. 2006; Sapolsky 2015). For example, stressful life experiences can facilitate the acquisition of cocaine use, and prompt cocaine craving or relapse in patients (Karlsgodt et al. 2003; Sinha 2008). Empirical attempts to model substance use disorders sensitive to stress experiences in animal models have examined the initiation and persistent increase of intense drug-taking, as well as reinstatement of drug-seeking after abstinence (Marinelli et al. 1996; Goeders 2002; Miczek et al. 2008, Han et al. 2017). Most behavioral and physiological findings from translational models of social stress-increased cocaine self-administration are established in non-human primates and rats, and less in mice (Morgan et al. 2002; Czoty et al. 2004; Miczek et al. 2011; Shimamoto et al. 2015). While long-term intravenous drug self-administration in mice is methodologically challenging, novel catheterization techniques enable the study of more prolonged and intense drug consumption, resulting in more clinically relevant models of substance use disorders (Thomsen and Caine 2005; Han et al. 2015). This study explores how intermittent (ISDS) and continuous social defeat stress (CSDS) alter cocaine taking in mice, in order to determine how the intensity of social stress episodes alters the ensuing patterns of behavioral and neural plasticity.

Although the intensity, frequency and probability of defensive-submissive reactions to aggression differ among strains, mice are a reliably pugnacious species (Parmigiani and Brain 1983; Guillot and Chapouthier 1996; Miczek et al. 2001). Social stress in rodents, like humans, is characterized by increased circulating glucocorticoids, immune suppression, persistent dysregulation of cardiovascular function and hyperthermia (Henry and Cassel 1969; Björkqvist 2001; de Groot et al. 1999; Miczek et al. 2008). Like other types of stress, *intermittent* social stress in rats causes sensitization to psychomotor stimulation, hastens acquisition of cocaine self-administration and augments cocaine taking during a 24-hour unlimited access binge (Miczek and Mutschler 1996; Tidey and Miczek 1997; Covington and Miczek 2001; Covington et al. 2008). Likewise, pulses of non-contingent footshock not only increase rates of cocaine self-administration, but also facilitate maintenance of high response rates at lower doses of cocaine in rats (Goeders and Guerin 1994).

The effects of social defeat on cocaine reward in mice varies based on the age at which the stress is experienced (Montagud-Romero et al. 2015; Rodriguez-Arias et al. 2017), as well as the sex of the animal (Kikusui et al. 2005), and the nature of the social defeat stress. Overall, episodic stress intensifies responding for cocaine, particularly during the acquisition and maintenance of drug taking, which also includes a leftward shift in responding for intravenous unit doses (Han et al. 2015). Rodent models of *continuous* social stress often cause a robust depressive-like phenotype, which is characterized by social avoidance and blunted reward processing (Kudryavtseva et al. 1991; Berton et al. 2006). Rats show attenuated cocaine seeking and taking after continuous social stress, but these effects have yet to be confirmed in mice (Miczek et al. 2011; Shimamoto et al. 2015).

Experience with intermittent and continuous social defeat stress either exacerbates or attenuates cocaine self-administration and alcohol intake, respectively (Miczek et al. 2011; Norman et al. 2015; Han et al. 2015). Here, we implemented femoral vein catheterization in order to delineate persistent changes in intravenous cocaine self-administration in C57BL/6J mice. Our main goal was to extend our rat studies showing divergent effects of intermittent and continuous social defeat stress on subsequent cocaine self-administration to mice (Miczek et al. 2011; Shimamoto et al. 2015). The current results provide a basis for characterizing adaptive processes in neurobiological systems after *intermittent* and *continuous* social stress and for identifying individuals who are especially vulnerable to the potentiating effects of social stress on drug taking.

Materials and Methods

Subjects

Experimental Animals—C57BL/6J male mice (Jackson Labs, Bar Harbor, ME) weighing 25–27g upon arrival were individually housed in clear polycarbonate cages ($28 \times 17 \times 14$ cm) with stainless steel wire lids and lined with pine shavings within a temperature-controlled mouse vivarium ($21 \pm 1^\circ\text{C}$, 30 to 40% humidity) that was kept on a reversed 12-hour photocycle (lights off 0600 hour). Mice were adapted to the facilities for at least 7 days before experiments. Animals were given standard rodent chow and water *ad libitum* (Lab Diet 5001 Rodent Diet, PMI Nutrition International, Brentwood, MO, USA). All housing conditions were maintained throughout the experiment (social defeat, sucrose preference and cocaine intravenous self-administration) with the exception of the chronic social defeat protocol (adapted from Kudryavtseva et al. 1991). All experimental procedures were reviewed and approved by the Tufts Institutional Animal Care and Use Committee, following the NIH Guide for the Care and Use of Laboratory Animals, 8th Edition (National Research Council 2011).

Residents—A separate group of male CFW mice (Charles River, Wilmington, MA) were pair-housed with females in clear polycarbonate cages ($28 \times 17 \times 14$ cm) for 1 month upon arrival at 8 weeks of age and served as aggressive stimulus mice (residents). Each resident was chosen for consistent aggressive behavior during confrontations with an intruder mouse (Miczek and O'Donnell 1978).

Drugs

Cocaine hydrochloride was obtained from the Research Technology Branch of the National Institute on Drug Abuse and dissolved in sterile 0.9% saline.

Experimental Design

Upon arrival, mice were randomly assigned to three groups: intermittent or continuous social defeat stress or non-stressed controls. The mice in both stress groups were socially defeated for ten consecutive days (Day 1–10). Seven to ten days after the last defeat, all animals were examined for social interaction and sucrose preference and then began cocaine self-administration.

Social Defeat Stress

Intermittent Episodic Social Defeat—Intermittently stressed mice were defeated using the resident-intruder protocol as illustrated and described previously (Miczek and O'Donnell 1978; Miczek et al. 1982; Yap and Miczek 2007) on days 1 through 10. Each social defeat episode took place in the room adjacent to where they were housed. Experimentally stressed mice were placed in a perforated cage inside the home cage of an aggressive male resident to allow unrestricted visual, auditory and olfactory contact with protection from injury. After 5 minutes of social threat in the perforated cage, stressed mice were removed from the perforated cage, and returned to the resident home cage where attack by the resident occurred. Immediately following defeat (defined as having received 30 bites within a maximum of 5 minutes), intruder mice were then placed back in the perforated cage for another 5 minutes in the aggressor's home cage. Intruder mice were then returned to the adjacent room where they were singly housed until their next defeat on the subsequent day. Intruder mice were exposed to a novel resident during each daily defeat. During each defeat episode, the latency to attack, number of bites received and total duration of each encounter was recorded.

Continuous Social Defeat—Continuously stressed mice were defeated as described previously (adapted from Miczek and O'Donnell 1978; Kudryavtseva et al. 1991). Male resident mice were allowed 24 hours to establish territory on one half of a clear polycarbonate cage (46 × 24 × 16 cm), divided lengthwise by a perforated divider with a stainless steel wire lid and pine shavings as bedding (Golden et al. 2011). Similar to the intermittent stress protocol, the intruder mouse was placed directly within the resident aggressor's home cage compartment. After 5 minutes of defeat (approximately 30 bites), the intruder was transferred across the perforated divider to the opposite compartment and housed there for the remainder of the day. Resident aggressors remained in their home cage throughout the 10-day defeat period and intruder mice were exposed to a different resident aggressor daily to prevent habituation.

Sucrose Preference

To examine sucrose preference, 50 ml bottles filled with either 1% sucrose (w/v) in tap water, or tap water alone. Bottles were introduced 4 hours into the dark photoperiod and fluid levels were noted at 4, 6 and 8 hours over two consecutive days into the dark photoperiod. The position of the tubes was changed, and bottles reweighed, after the initial 24h. To control for unintentional fluid loss and evaporation, bottle measurements were also recorded from an empty cage, and these values were subtracted from each mouse's intake. Sucrose consumption was calculated as grams of sucrose consumed (g) and as a percent of total fluid intake (sucrose + H₂O) averaged over the two days of testing.

Social Interaction Testing

Seven days after the last day of social defeat, social behavior was evaluated using a 3-chamber apparatus (85 × 30 × 35 cm) (Moy et al. 2004; Zanettini et al. 2010; Newman et al. 2016). The socially defeated or non-stressed male habituated to the central chamber of the 3-chamber apparatus for a 2.5-minute period. An unfamiliar resident aggressor stimulus mouse

was placed in a wire mesh cage in either the right or left chamber. Following this habituation period, the doors on either side of the central chamber were lifted, allowing the experimental male to move freely between the central chamber, the chamber with the resident stimulus male, and a third chamber with an empty stimulus cage during a 2.5-minute social approach test. EthoVision XT software (Noldus, Wageningen, the Netherlands) tracked the male and recorded the duration of time it spent within each of the three chambers.

Surgical Procedures

The day following sucrose preference testing (Day 23), intermittently defeated and control mice were permanently implanted with an indwelling catheter (Silastic silicon tubing, ID 0.012 in, OD 0.025 in) into the right jugular or femoral vein under a combination of ketamine (100 mg/kg i.p.) and xylazine (10 mg/kg i.p.) anesthesia (Han et al. 2015). The distal end of the catheter was passed subcutaneously from the insertion point to the area between the scapula where it exited through a small incision and was affixed to a small plastic pedestal (Plastics One, Roanoke, VA). After surgery, mice were allowed 4 days to recover. Catheters were flushed daily with 0.05ml of heparinized saline (20 IU/ml) daily to prevent clotting.

Cocaine self-administration

For cocaine self-administration, a panel with cue lights and 2 nose-poke operanda was inserted vertically into the home cage of the mouse (Miczek and de Almeida 2001). The top of the panel held a house light for cage illumination and a counterbalanced arm holding a liquid swivel (Instech Laboratories, Plymouth Meeting, PA). A 3 ml syringe in a computer-controlled, pump was connected to the swivel and pedestal by PE20 tubing. A green cue light distinguished the operanda (the “active” hole had a light present, while the “inactive” hole remained unlit). The active side remained constant throughout the experiment and was counterbalanced across subjects. A response on the active side triggered an I.V. infusion of cocaine and the stimulus light was then deactivated for 20 seconds (indicating the post-infusion time-out). All responses and infusions were recorded automatically using a computer interface and software from Med Associates (St. Albans, VT).

Daily 4 hour sessions—Following recovery from surgery, mice were trained to self-administer cocaine (0.3 mg/kg/infusion) according to a fixed ratio 1 (FR1) schedule of reinforcement (Han et al. 2015). Two 2-hour time blocks of access were separated by a 10-minute interval in which all cue and house lights were extinguished to signal unavailability of drug. Each access block terminated after the delivery of 50 infusions or after 2 hours of access. Mice acquired and were maintained on 0.3 mg/kg/infusion for 7 days. Thereafter, mice were tested according to an FR1 schedule with descending unit doses (0.1, 0.03, .01, 0.003, 0.001 mg/kg/infusion) receiving each dose for two consecutive days. Depending on cocaine self-administration during acquisition, mice with continuous stress experience were separated into one of two groups for all subsequent analyses. Responding for cocaine during training sessions after CSDS produced two subgroups that are represented by non-overlapping modes. During days 2–7 of acquisition, animals either took less than 10 infusions per session most often, or self-administered more than 90 infusions per session

most often (Figure 7). The group of “high responders” took an average of 23 mg/kg/session on days 2–7 of acquisition, while the “low responders” took an average of 3 mg/kg/session.

All intermittently defeated and contemporaneous control animals were catheterized via the jugular vein, while all continuously defeated and contemporaneous controls were catheterized via the femoral vein. A separate subset of animals void of SDS experience were catheterized via the jugular or femoral vein to determine differences in cocaine acquisition based on catheterization method.

Statistical Analysis

Data were analyzed using Prism version 7.0 (Graphpad Software Inc.). Two-way repeated measures ANOVAs were used to analyze repeated observations of social interaction and sucrose preference after either ISDS, CSDS or daily handling, cocaine self-administration after ISDS and in nonstressed controls to test for effects of catheterization method. Two-way repeated measures ANOVAs were also used to test for the effects of ISDS, CSDS and route of catheterization on cocaine self-administration. All significant results were followed by post hoc analyses with Bonferroni’s corrections for multiple comparisons. In the cocaine self-administration after CSDS, Kruskal-Wallis tests were used to analyze the relationship between control animals, those in the “high responders”, and those in the “low responders” group across days of acquisition and descending doses of drug. Pearson’s correlation coefficient (r) was generated to determine the relationship between sucrose and cocaine consumption after CSDS. The criterion for statistical significance was $p < 0.05$.

Results

Intravenous drug self-administration in mice: femoral versus jugular catheterization

By catheterizing a subset of stress-naïve mice in either the jugular or femoral vein, we confirmed that acquisition and maintenance of cocaine self-administration on an FR1 schedule of responding at 0.3 mg/kg/infusion was insignificantly affected by the route of catheterization. The benefit of femoral vein catheterization in mice, as compared to jugular, was evident by considerably extended longevity of catheters (Figure 2). A log-rank (Mantel-Cox) test revealed a significant effect of catheterization route on catheter longevity ($X^2_{(1, 28)} = 18.37$; $p < 0.001$).

CSDS produces social deficits

Using a 3-chamber apparatus, we examined time spent in each of the three chambers when the stimulus resident male was present. When tested 10 days after the last social defeat, animals with a history of CSDS showed social deficits when compared to those with a history of ISDS or daily handling. A repeated measures two-way ANOVA revealed a significant effect of chamber ($F_{2, 267} = 77.11$; $p < 0.001$) and a significant interaction between stress condition and chamber ($F_{4, 267} = 4.81$; $p < 0.001$). There was no significant main effect of stress. Post hoc analyses revealed significant attenuation of time spent in the chamber with the stimulus male by mice with a history of CSDS when compared to controls ($p < 0.05$) (Figure 3).

CSDS produces two distinct sucrose-preferring phenotypes

Although ISDS had minimal effects on subsequent consumption of and preference for 1% sucrose during a 48-hour preference test, CSDS aggregated animals into two distinct subpopulations. There was a significant, direct relationship between the amount of cocaine self-administered during acquisition and the amount of sucrose consumed during the preference test in mice that underwent CSDS ($r=0.602$, $p<0.05$; Figure 8). Based on cocaine intake during days 2–7 of acquisition, animals with a history of CSDS were either placed in the “high responders” or “low responders” group for analysis of sucrose consumption and preference.

A repeated measures two-way ANOVA revealed both a significant effect of fluid type ($F_{3, 166}=169.4$; $p<0.001$) and a significant interaction between stress condition and fluid type ($F_{3, 166}=4.22$; $p<0.01$) for 24-hour sucrose and water consumption during a 2-bottle test. There was no significant main effect of stress. Post hoc analyses revealed significantly increased consumption of 1% sucrose in mice with a history of CSDS assigned to the high responders group when compared to controls ($p<0.05$). To examine the effects of sucrose preference over water, a one-way ANOVA revealed a significant effect of fluid type ($F_{3, 83}=66.79$; $p<0.001$). Post hoc analyses found significant attenuation of sucrose preference in mice with a history of CSDS assigned to the low responders group ($p<0.001$), as well as ISDS ($p<0.001$) compared to controls (Figure 4).

Intermittent SDS modestly increases cocaine self-administration

For analysis of cocaine acquisition after previous experience with ISDS, a repeated measures two-way ANOVA revealed a significant effect of day ($F_{6, 138}=6.91$; $p<0.001$), as well as a significant effect of stress ($F_{2, 23}=6.38$; $p<0.05$). There was no significant interaction of stress and day of acquisition. Post hoc analyses found significantly increased cocaine self-administration on day one of acquisition after ISDS as compared to controls ($p<0.001$; Figure 5A).

Responding for varying doses of cocaine after ISDS was analyzed using a repeated measures two-way ANOVA to reveal a significant effect of stress ($F_{1, 72}=4.93$; $p<0.05$). There was no significant main effect of cocaine dosage or interaction of stress and cocaine dosage (Figure 5B).

Continuous SDS persistently increases or attenuates cocaine self-administration

Acquisition of cocaine self-administration after previous experience with CSDS was analyzed as a whole group using a repeated measures two-way ANOVA to reveal a significant effect of day ($F_{1, 140}=10.37$; $p<0.01$). There was no significant effect of stress and no significant interaction of stress and day of acquisition (Fig 6A).

Responding for varying doses of cocaine after CSDS was also analyzed as a whole group using a repeated measures two-way ANOVA to reveal a significant effect of cocaine dosage ($F_{5, 114}=6.82$; $p<0.001$). There was no significant main effect of stress or interaction of stress and cocaine dosage (Figure 5C).

Previous experience with CSDS aggregated mice into two distinct cocaine-taking phenotypes during days 2–7 of acquisition. Approximately half of the animals (n=6) clustered around a group mean of 3 mg/kg/session, while the other half of the animals (n=7) had a group mean of 23 mg/kg/session. Although responding for cocaine on day 1 was not indicative of subsequent cocaine taking, no member of the “low responders” took more infusions/session than the minimum number of infusions/session taken by any member of the “high responders” within days 2–7 of acquisition (Figure 7). Since cocaine self-administration data were clustered based upon the behavior itself, Kruskal-Wallis tests were run to compare cocaine self-administration between stress conditions. Responding for cocaine during acquisition after CSDS was analyzed by a Kruskal–Wallis test to reveal significant differences in the mean number of self-administered cocaine infusions across acquisition among the three conditions ($H=15.65$, $p=0.001$; Figure 6B). Results of Kruskal-Wallis testing were also significant for effect of cocaine dose ($H=12.32$, $p=0.001$; Figure 6D).

Discussion

The present results confirm increased intravenous cocaine taking in mice exposed to intermittent social defeat stress. We extend previous findings by comparing *continuous* to *intermittent* social stress followed by subsequent cocaine self-administration. We found that these two histories of social defeat had divergent effects on patterns of cocaine self-administration. Experience with continuous social defeat stress aggregated C57Bl/6J mice into two distinct subpopulations upon examination of subsequent cocaine intake during daily sessions. In fact, one subpopulation self-administered a mean of 23 mg/kg/session, whereas a second subpopulation averaged 3 mg/kg/session. The maximal number of infusions taken by any member of the low responders never exceeded the minimum number of infusions taken by any member of the high responders (Figure 7). These patterns of cocaine taking were closely correlated with those for sucrose preference, a putative measure of anhedonia (Willner et al. 1987; Berton et al. 2006; Krishnan et al. 2007). Preference for sucrose over water revealed overlap with patterns of cocaine taking in continuously defeated mice. Specifically, the amount of sucrose consumed during the preference test predicted the amount of cocaine self-administered during acquisition (Figure 8). After experiencing continuous SDS, mice showed either increased or attenuated intake of both sucrose and cocaine when compared to animals with a history of ISDS, or those with no social defeat experience. Taken together, the correlation in patterns of consumption for cocaine and sucrose suggests two hypotheses. First, chronic social stress may result in divergent neuroplastic processes to engender either increased consummatory behavior or an anhedonic-like phenotype. Second, anhedonia may present itself across behavior under the control of several highly reinforcing stimuli.

The results from the current set of experiments extend previous findings regarding the effects of exposure to continuous SDS. Continuous SDS in mice has produced phenotypes that translate to symptoms indicative of anhedonia in humans (Björkqvist 2001). Continuous SDS effectively blunts reward processing as measured by reduced intake of sucrose in both rats and mice (Willner et al. 1987; Rygula et al. 2005; Macedo et al. 2018), and attenuated cocaine self-administration in both male and female rats (Miczek et al. 2011; Shimamoto et

al. 2015). Remarkably, continuous SDS engendered differentiated intake of both sucrose and cocaine, with approximately half of the mice increasing intake (Figure 8). It remains to be determined how the intermittency, intensity and controllability of stress alters the motivation for, and consumption of, rewarding stimuli, such as sucrose and cocaine. While continuous SDS consistently produced animals with social deficits, it resulted in two starkly contrasting phenotypes of sucrose and cocaine self-administration. These data may indicate a vulnerability to cocaine use based upon the individual's response to stress experience (Charney 2004; Krishnan et al. 2007).

The translational value of stress-increased cocaine use in animal models depends heavily on drug access conditions (Ahmed and Koob 1998; Kawa et al. 2016; Ahmed et al. 2018). Until now, jugular catheters were limited in patency preventing the study of persistent and intense drug intake in mouse models of intravenous drug self-administration. The approach to the femoral vein extends the patency of catheters by several weeks and increases the time to accumulate drug intake. Allowing for long-term drug self-administration in mice enables the study of higher and longer rates of consumption, motivation, and more translationally relevant intermittent access conditions (Mantsch et al. 2007; Zimmer, Oleson and Roberts 2012). Earlier work demonstrated that the pharmacokinetics of cocaine depend not on the route of intravenous catheterization for the study of self-administered drugs, as long as the distance from the insertion site to the heart is properly accounted for (Collins and Kantak, 2002; Czoty et al. 2004; Nader and Reboussin 1994). Minor changes to the pharmacodynamics of intravenous cocaine self-administration will alter susceptibility and sensitization of the drug (Samaha and Robinson, 2002), and the current set of experiments in mice confirms there is an insignificant difference in responding for cocaine based upon the method of catheterization. Limited patency of jugular catheters has hindered our ability to study drug self-administration in the same mouse both before and after SDS, but femoral catheterization techniques render this approach feasible (Figure 2).

Individuals are known to vary considerably in their behavioral, neural and endocrine responses to social stressors (Wingfield and Sapolsky 2003; de Boer et al. 2017; Wood and Valentino 2017). In line with bimodal population distributions, two points emerge from the overall results of our study. First, ten days of continuous social defeat stress significantly reduced social interactions in all of our B6 mice. Beyond this general deficit in social behavior, which is an important measure of stressor vulnerability, our chronically stressed mice did however diverge, with regards to reward sensitivity. As described above, either an *increase* or *decrease* in cocaine and sucrose taking was phenotypically expressed after chronic social stress. Thus, our data indicate that subpopulations exist beyond a distinct depressive-like phenotype, including completely opposite responses to reward sensitivity. This evidence may further explain the biological complexities of affective syndromes, which often occur with different outcomes to antidepressant treatment, even in patients with the same diagnosis (Labonté et al. 2017).

Second, intermittently stressed mice, when compared to non-stressed controls, generated significantly more responding for cocaine under the current limited access self-administration conditions. However, elevated cocaine taking by *intermittently* stressed mice was far less than the amount self-administered by high-responding (HC) *chronically* stressed

mice. Taken together, these data indicate that social stress can gradually lead to the development of a distinct reward seeking phenotype, whether it be intensely augmented, or robustly diminished. Specifically, brief intermittent stress modestly increases cocaine taking, while more intense stress exposures lead to a divergence in cocaine reward, either further increasing, or conversely, completely compromising drug taking. In line with our behavioral data, molecular results have indicated a *recruitment* of many molecular neural mechanisms during the process of developing a *resiliency* to chronic social stress (Krishnan et al. 2007; Hultman et al. 2018; Lorsch et al. 2018). Future experiments are warranted to determine whether or not neural adaptations are associated with differential patterns of cocaine taking across a range of individuals after chronic, intense social stress experiences.

Acknowledgments

The authors would like to thank J. Thomas Sopko for his assistance during manuscript preparation.

Funding This work was funded by NIDA grant DA031734 to KAM

References

- Ahmed SH, Badiani A, Miczek KA, Müller CP (2018) Non-pharmacological factors that determine drug use and addiction. *Neuroscience & Biobehavioral Reviews*
- Ahmed SH, Koob GF (1998) Transition from moderate to excessive drug intake: change in hedonic set point. *Science* 282(5387), 298–300 [PubMed: 9765157]
- Berton O, McClung CA, DiLeone RJ, Krishnan V, Renthal W, Russo SJ, Graham D, Tsankova NM, Bolanos CA, Rios M, Monteggia LM (2006) Essential role of BDNF in the mesolimbic dopamine pathway in social defeat stress. *Science* 311(5762), 864–868 [PubMed: 16469931]
- Björkqvist K (2001) Social defeat as a stressor in humans. *Physiology & behavior* 73(3), 435–442 [PubMed: 11438372]
- Charney DS (2004) Psychobiological mechanisms of resilience and vulnerability: implications for successful adaptation to extreme stress. *Am J Psychiatry* 161(2):195–216 [PubMed: 14754765]
- Collins SL, Kantak KM (2002) Neuronal nitric oxide synthase inhibition decreases cocaine self-administration behavior in rats. *Psychopharmacology* 159(4), 361–369 [PubMed: 11823888]
- Covington HE, Miczek KA (2001) Repeated social-defeat stress, cocaine or morphine. *Psychopharmacology* 158(4), 388–398 [PubMed: 11797060]
- Covington HE, Tropea TF, Rajadhyaksha AM, Kosofsky BE, Miczek KA (2008) NMDA receptors in the rat VTA: a critical site for social stress to intensify cocaine taking. *Psychopharmacology* 197(2), 203–216 [PubMed: 18097654]
- Czoty PW, Morgan D, Shannon EE, Gage HD, Nader MA (2004) Characterization of dopamine D 1 and D 2 receptor function in socially housed cynomolgus monkeys self-administering cocaine. *Psychopharmacology* 174(3), 381–388 [PubMed: 14767632]
- de Boer SF, Buwalda B, Koolhaas JM (2017) Untangling the neurobiology of coping styles in rodents: towards neural mechanisms underlying individual differences in disease susceptibility. *Neuroscience & Biobehavioral Reviews* 74, 401–422 [PubMed: 27402554]
- de Groot J, van Milligen FJ, Moonen-Leusen BW, Thomas G, Koolhaas JM (1999) A single social defeat transiently suppresses the anti-viral immune response in mice. *Journal of neuroimmunology* 95(1–2), 143–151 [PubMed: 10229124]
- Der-Avakian A, Bland ST, Schmid MJ, Watkins LR, Spencer RL, Maier SF (2006) The role of glucocorticoids in the uncontrollable stress-induced potentiation of nucleus accumbens shell dopamine and conditioned place preference responses to morphine. *Psychoneuroendocrinology* 31(5), 653–663 [PubMed: 16584846]
- Goeders NE (2002) The HPA axis and cocaine reinforcement. *Psychoneuroendocrinology* 27(1–2), 13–33 [PubMed: 11750768]

- Goeders NE, Guerin GF (1994) Non-contingent electric footshock facilitates the acquisition of intravenous cocaine self-administration in rats. *Psychopharmacology* 114(1), 63–70 [PubMed: 7846208]
- Golden SA, Covington HE, Berton O, Russo SJ (2011) A standardized protocol for repeated social defeat stress in mice. *Nature protocols* 6(8), 1183 [PubMed: 21799487]
- Guillot PV, Chapouthier G (1996) Intermale aggression and dark/light preference in ten inbred mouse strains. *Behavioural brain research* 77(1–2), 211–213 [PubMed: 8762172]
- Han X, Albrechet-Souza L, Doyle MR, Shimamoto A, DeBold JF, Miczek KA (2015) Social stress and escalated drug self-administration in mice II. Cocaine and dopamine in the nucleus accumbens. *Psychopharmacology* 232(6), 1003–1010 [PubMed: 25216798]
- Han X, DeBold JF, Miczek KA (2017) Prevention and reversal of social stress-escalated cocaine self-administration in mice by intra-VTA CRFR1 antagonism. *Psychopharmacology* 234(18), 2813–2821 [PubMed: 28698920]
- Harris AZ, Atsak P, Bretton ZH, Holt ES, Alam R, Morton MP, Abbas AI, Leonardo ED, Bolkan SS, Hen R, Gordon JA (2018) A novel method for chronic social defeat stress in female mice. *Neuropsychopharmacology* 43(6):1276–1283 [PubMed: 29090682]
- Henry JP, Cassel JC (1969) Psychosocial factors in essential hypertension recent epidemiologic and animal experimental evidence. *American Journal of Epidemiology* 90(3), 171–200 [PubMed: 4899564]
- Holly EN, Shimamoto A, DeBold JF, Miczek KA (2012) Sex differences in behavioral and neural cross- sensitization and escalated cocaine taking as a result of episodic social defeat stress in rats. *Psychopharmacology* 224(1), 179–188 [PubMed: 22926005]
- Hultman R, Ulrich K, Sachs BD, Blount C, Carlson DE, Ndubuiuzi N, Bagot RC, Parise EM, Vu MAT, Gallagher NM, Wang J (2018). Brain-wide Electrical Spatiotemporal Dynamics Encode Depression Vulnerability. *Cell* 173(1), 166–180 [PubMed: 29502969]
- Karlsgodt KH, Lukas SE, Elman I (2003) Psychosocial stress and the duration of cocaine use in non-treatment seeking individuals with cocaine dependence. *The American journal of drug and alcohol abuse* 29(3), 539–551 [PubMed: 14510039]
- Kawa AB, Bentzley BS, Robinson TE (2016) Less is more: prolonged intermittent access cocaine self-administration produces incentive-sensitization and addiction-like behavior. *Psychopharmacology* 233(19–20), 3587–3602 [PubMed: 27481050]
- Kleber HD, Gawin FH (1984) The spectrum of cocaine abuse and its treatment. *J Clin Psychiatry* 45:18–23
- Krishnan V, Han MH, Graham DL, Berton O, Renthal W, Russo SJ, Ghose S (2007) Molecular adaptations underlying susceptibility and resistance to social defeat in brain reward regions. *Cell* 131(2), 391–404 [PubMed: 17956738]
- Kudryavtseva NN, Bakshtanovskaya IV, Koryakina LA (1991) Social model of depression in mice of C57BL/6J strain. *Pharmacology Biochemistry and Behavior* 38(2), 315–320
- Parmigiani S, Brain PF (1983) Effects of residence, aggressive experience and intruder familiarity on attack shown by male mice. *Behavioural Processes* 8(1), 45–57 [PubMed: 24923607]
- Macedo GC, Morita GM, Domingues LP, Favoretto CA, Suchecki D, Quadros IMH (2018) Consequences of continuous social defeat stress on anxiety-and depressive-like behaviors and ethanol reward in mice. *Hormones and behavior* 97, 154–161 [PubMed: 29056427]
- Mantsch JR, Cullinan WE, Tang LC, Baker DA, Katz ES, Hoks MA, Ziegler DR (2007) Daily cocaine self-administration under long-access conditions augments restraint-induced increases in plasma corticosterone and impairs glucocorticoid receptor-mediated negative feedback in rats. *Brain research* 1167, 101–111 [PubMed: 17689506]
- Marinelli M, Le Moal M, Piazza PV (1996) Acute pharmacological blockade of corticosterone secretion reverses food restriction-induced sensitization of the locomotor response to cocaine. *Brain research* 724(2), 251–255 [PubMed: 8828576]
- Miczek KA, Mutschler NH (1996) Activational effects of social stress on IV cocaine self-administration in rats. *Psychopharmacology* 128(3), 256–264 [PubMed: 8972545]

- Miczek KA, de Almeida RM (2001) Oral drug self-administration in the home cage of mice: alcohol-heightened aggression and inhibition by the 5-HT 1B agonist anpirtoline. *Psychopharmacology* 157(4), 421–429 [PubMed: 11605102]
- Miczek KA, Thompson ML, Shuster L (1982) Opioid-like analgesia in defeated mice. *Science* 215(4539), 1520–1522 [PubMed: 7199758]
- Miczek KA, Maxson SC, Fish EW, Faccidomo S (2001) Aggressive behavioral phenotypes in mice. *Behavioural brain research* 125(1–2), 167–181 [PubMed: 11682108]
- Miczek KA, Yap JJ, Covington HE (2008) Social stress, therapeutics and drug abuse: preclinical models of escalated and depressed intake. *Pharmacology & therapeutics* 120(2), 102–128 [PubMed: 18789966]
- Miczek KA, Nikulina EM, Shimamoto A, Covington HE (2011) Escalated or suppressed cocaine reward, tegmental BDNF, and accumbal dopamine caused by episodic versus continuous social stress in rats. *Journal of Neuroscience* 31(27), 9848–9857 [PubMed: 21734276]
- Miczek KA, O'Donnell JM (1978) Intruder-evoked aggression in isolated and nonisolated mice: effects of psychomotor stimulants and L-dopa. *Psychopharmacology* 57(1), 47–55 [PubMed: 26933]
- Montagud-Romero S, Aguilar MA, Maldonado C, Manzanedo C, Miñarro J, Rodríguez-Arias M (2015) Acute social defeat stress increases the conditioned rewarding effects of cocaine in adult but not in adolescent mice. *Pharmacology Biochemistry and Behavior* 135, 1–12
- Morgan D, Brebner K, Lynch WJ, Roberts DCS (2002) Increases in the reinforcing efficacy of cocaine after particular histories of reinforcement. *Behavioural pharmacology* 13(5), 389–396 [PubMed: 12394415]
- Moy SS, Nadler JJ, Perez A, Barbaro RP, Johns JM, Magnuson TR, Piven J, Crawley JN (2004) Sociability and preference for social novelty in five inbred strains: an approach to assess autistic-like behavior in mice. *Genes, Brain and Behavior* 3(5), 287–302
- Nader MA, Reboussin DM (1994) The effects of behavioral history on cocaine self-administration by rhesus monkeys. *Psychopharmacology* 115(1–2), 53–58 [PubMed: 7862912]
- Norman KJ, Seiden JA, Klickstein JA, Han X, Hwa LS, DeBold JF, Miczek KA (2015) Social stress and escalated drug self-administration in mice I. Alcohol and corticosterone. *Psychopharmacology* 232(6), 991–1001 [PubMed: 25242256]
- Newman EL, Gunner G, Huynh P, Gachette D, Moss SJ, Smart TG, Rudolph U, DeBold JF, Miczek KA (2016) Effects of Gabra2 Point Mutations on Alcohol Intake: Increased Binge-Like and Blunted Chronic Drinking by Mice. *Alcoholism: Clinical and Experimental Research* 40(11), 2445–2455
- Labonté B, Engmann O, Purushothaman I, Menard C, Wang J, Tan C, Scarpa JR, Moy G, Loh YH, Cahill M, Lorsch ZS (2017) Sex-specific transcriptional signatures in human depression. *Nature medicine* 23(9), 1102
- Lorsch ZS, Loh YHE, Purushothaman I, Walker DM, Parise EM, Salery M, Cahill ME, Hodes GE, Pfau ML, Kronman H, Hamilton PJ (2018) Estrogen receptor α drives pro-resilient transcription in mouse models of depression. *Nature communications* 9(1), 1116
- Rodríguez-Arias M, Montagud-Romero S, Rubio-Araiz A, Aguilar MA, Martín-García E, Cabrera R, Maldonado R, Porcu F, Colado MI, Miñarro J (2017) Effects of repeated social defeat on adolescent mice on cocaine-induced CPP and self-administration in adulthood: integrity of the blood–brain barrier. *Addiction biology* 22(1), 129–141 [PubMed: 26374627]
- Rygula R, Abumaria N, Flügge G, Fuchs E, Rütther E, Havemann-Reinecke U (2005) Anhedonia and motivational deficits in rats: impact of chronic social stress. *Behavioural brain research* 162(1), 127–134 [PubMed: 15922073]
- Samaha AN, Li Y, Robinson TE (2002) The rate of intravenous cocaine administration determines susceptibility to sensitization. *Journal of Neuroscience* 22(8), 3244–3250 [PubMed: 11943825]
- Sapolsky RM (2015) Stress and the brain: individual variability and the inverted-U. *Nature neuroscience* 18(10), 1344 [PubMed: 26404708]
- Shimamoto A, Holly EN, Boyson CO, DeBold JF, Miczek KA (2015) Individual differences in anhedonic and accumbal dopamine responses to chronic social stress and their link to cocaine self-administration in female rats. *Psychopharmacology* 232(4), 825–834 [PubMed: 25178816]

- Sinha R (2008) Chronic stress, drug use, and vulnerability to addiction. *Annals of the New York Academy of Sciences* 1141(1), 105–130 [PubMed: 18991954]
- Takahashi A, Chung JR, Zhang S, Zhang H, Grossman Y, Aleyasin H, Flanigan ME, Pfau ML, Menard C, Dumitriu D, Hodes GE (2017) Establishment of a repeated social defeat stress model in female mice. *Scientific Reports* 7(1), 12838 [PubMed: 28993631]
- Thomsen M, Caine SB (2005) Chronic intravenous drug self-administration in rats and mice. *Current Protocols in Neuroscience* 32(1), 9–20
- Tidey JW, Miczek KA (1997) Acquisition of cocaine self-administration after social stress: role of accumbens dopamine. *Psychopharmacology* 130(3), 203–212 [PubMed: 9151353]
- Willner P, Towell A, Sampson D, Sophokleous S, Muscat R (1987) Reduction of sucrose preference by chronic unpredictable mild stress, and its restoration by a tricyclic antidepressant. *Psychopharmacology* 93(3), 358–364 [PubMed: 3124165]
- Wingfield JC, Sapolsky RM (2003) Reproduction and resistance to stress: when and how. *Journal of neuroendocrinology* 15(8), 711–724 [PubMed: 12834431]
- Wood SK, Valentino RJ (2017) The brain norepinephrine system, stress and cardiovascular vulnerability. *Neuroscience & Biobehavioral Reviews* 74, 393–400 [PubMed: 27131968]
- Yap JJ, Miczek KA (2007) Social defeat stress, sensitization, and intravenous cocaine self-administration in mice. *Psychopharmacology* 192(2), 261–273 [PubMed: 17297635]
- Zanettini C, Carola V, Iacono LL, Moles A, Gross C, d'Amato FR (2010) Postnatal handling reverses social anxiety in serotonin receptor 1A knockout mice. *Genes, Brain and Behavior* 9(1), 26–32
- Zimmer BA, Oleson EB, Roberts DC (2012) The motivation to self-administer is increased after a history of spiking brain levels of cocaine. *Neuropsychopharmacology* 37(8), 1901 [PubMed: 22453139]

Fig 1:
Experimental Timeline.

Fig 2:
Longevity of either jugular (n=17) or femoral (n=11) vein catheterization in non-stressed controls. Data are expressed as percent of catheter survival over time.

Fig 3:

Three chamber compartment testing with an unfamiliar, stimulus resident male mouse 10 days after social defeat. Data are expressed as mean; ** $p < 0.01$ vs. controls.

Fig 4: Twenty-four-hour consumption and preference for 1.0% sucrose solution versus water in a 2- bottle preference test by intermittently stressed mice (I; n=36), continuously stressed mice placed into the high responders group (HC; n=7), continuously stressed mice placed into the low responders group (LC; n=6), and controls (C; n=38). All values are mean $\pm$ SEM. *= p<0.05, ***=p<0.001 compared to the relevant control group.

Fig 5:

A. Acquisition of cocaine self-administration (0.3 mg/kg/inf) after intermittent SDS (n=14, filled squares) and contemporaneous controls (n=12, open circles) B. Responding for various doses of intravenous cocaine in mice with a history of intermittent social defeat (n=6) and contemporaneous controls (n = 9). Individual data were averaged over 2 sessions. Data are expressed as mean; * $p < 0.05$ vs. controls.

Fig 6:

A. Acquisition of cocaine self-administration (0.3 mg/kg/inf) after continuous social defeat (n=13, black squares) and contemporaneous controls (n=9, open circles). B. Acquisition of cocaine self-administration (0.3 mg/kg/inf) after continuous social defeat (high responders: n= 7, dark grey squares; low responders: n= 6, light grey triangles) and contemporaneous controls (n=9, open circles) C. Responding for various doses of intravenous cocaine in mice with a history of continuous social defeat (n=9, black squares) and contemporaneous controls (n=6, open circles). D. Responding for various doses of intravenous cocaine in mice with a history continuous social defeat (high responders: n= 5, dark grey squares; low responders: n= 4, light grey triangles) and contemporaneous controls (n=6, open circles). Data from each dose on the dose-effect curves were averaged over 2 sessions. Data are expressed as mean $\pm$ SEM.

Fig 7:

A. In animals with a history of continuous SDS (high responders: n= 7, dark grey bars; low responders: n= 6, light grey bars), the maximum number of cocaine infusions/session taken by any member of the low responders between days 2–7 of acquisition never exceeded the minimum number of cocaine infusions taken by high responders on corresponding days. B. The number of cocaine infusions across days 2–7 of acquisition formed a continuous distribution in animals with no history of SDS (controls: n= 9), unlike the bimodal distribution produced after CSDS.

Fig 8:

Twenty-four-hour intake of 1% sucrose during a preference test is predictive of cocaine consumption during acquisition in mice with a history of continuous SDS [(n=17) ($r=0.34$; $p=0.014$)]. Open circles denote nonstressed controls (n=4), light grey squares denote animals placed in the low responders group (n=6) and dark grey squares denote high responders (n=7).