Published in final edited form as:

J Steroid Biochem Mol Biol. 2011 October ; 127(1-2): 108–117. doi:10.1016/j.jsbmb.2010.11.007.

Bridging Epidemiology and Model Organisms to Increase Understanding of Endocrine Disrupting Chemicals and Human Health Effects

Tracey J. Woodruff

Program on Reproductive Health and the Environment, Department of Obstetrics, Gynecology, and Reproductive Sciences, University of California, San Francisco

Abstract

Concerning temporal trends in human reproductive health has prompted concern about the role of environmental mediated risk factors. The population is exposed to chemicals present in air, water, food and in a variety of consumer and personal care products and multiple chemicals are found human populations around the globe. Recent reviews find that endocrine disrupting chemicals (EDCs) can adversely affect reproductive and developmental health. However, there are still many knowledge gaps. This paper reviews some of the key scientific concepts relevant to integrating information from human epidemiologic and model organisms to understand the relationship between EDC exposure and adverse human health effects. Additionally, areas of new insights which influence the interpretation of the science are briefly reviewed, including: enhanced understanding of toxicity pathways; importance of timing of exposure; contribution of multiple chemical exposures; and low dose effects. Two cases are presented, thyroid disrupting chemicals and anti-androgens chemicals, which illustrate how our knowledge of the relationship between EDCs and adverse human health effects is strengthened and data gaps reduced when we integrate findings from animal and human studies

Keywords

endocrine disrupting chemicals; thyroid hormones; anti-androgens; pregnancy; susceptible periods of development; epidemiology; model organisms

Introduction

Concerning temporal trends in human reproductive health has prompted concern about the role of environmental mediated risk factors. Studies report increases in reproductive diseases and decline in reproductive function since the mid-20th century among certain locations and

Address correspondence to: Tracey J. Woodruff, Program on Reproductive Health and the Environment, University of California, San Francisco, 1330 Broadway St., Suite 1100, Oakland, CA 94612, woodrufft@obgyn.ucsf.edu, Phone: 510.986.8942; Fax: 510.986.8960.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

populations (primarily in the developed world), previously reviewed and illustrated in Table 1 [1].

The relatively short time frame over which decline in reproductive health and function has been observed cannot be explained by genetic changes. Environmental chemicals have been identified as one of the potential risk factors that may be contributing to observed changes in reproductive health [2-4]. Over roughly the same period, manufacture and use of both natural and synthetic chemicals has increased by over 20 fold [5]. In the US, there are approximately 87,000 chemical substances registered for use in commerce as of 2006, and about 3,000 chemicals manufactured or imported in excess of 1 million pounds each. [6].

The population is exposed to chemicals present in air, water, food and in a variety of consumer and personal care products. In the United States, nationally representative samples of population through the National Health and Nutrition Examination Survey find that every individual has measured levels of multiple environmental chemicals in his/her body [7, 8]. There are similar findings from studies in Europe [9], and populations in the Arctic far from pollution sources [10]. Thus it is expected that all human populations are exposed to some level of synthetic chemicals.

Previous studies have demonstrated environmental chemicals can adversely impact human health. Poisoning incidents with mercury in Japan and polychlorinated biphenyl (PCB) in Taiwan, produced neurological, reproductive, and developmental effects, even when the mother was asymptomatic [11-13]. Occupational exposures to 1,2-dibromo-3-chloropropane (DBCP), produced male infertility [14]. Science has evolved to study health effects of lower level exposures among the population. For example, reviews of the epidemiological evidence find that contemporary exposures to PCBs are associated with a decrease in semen quality, specifically reduced sperm motility [2] and prenatal exposures to methylmercury and PCBs at contemporary concentrations can increase risk of neurological deficits in children [15-18].

Environmental contaminants can adversely affect reproductive health through diverse biological mechanisms. Historically, much of the scientific inquiry focused on genotoxic or mutagenic chemicals and cancer effects [19, 20]. Since the latter part of the 20th century there has increased emphasis on an important class of chemicals called endocrine disrupting chemicals (EDCs) that interfere with the production, release, transport, metabolism, binding, action, or elimination of natural hormones in the body and are responsible for maintenance of homeostasis and regulation of developmental processes and their potential effects on reproductive and developmental effects [2]. Recent reviews and scientific consensus statements find that “the evidence for adverse reproductive outcomes (infertility, cancers, malformations) from exposure to endocrine disrupting chemicals is strong, and there is mounting evidence for effects on other endocrine systems, including thyroid, neuroendocrine, obesity and metabolism, and insulin and glucose homeostasis.” [2].

However, there are still many gaps in our understanding of the relationship between EDC exposures and reproductive and developmental effects. First, the number of chemicals that have been evaluated for effects on human health remains limited [6]. Second, many reproductive and developmental health conditions, such as female reproductive effects (e.g.

fibroids, endometriosis), male reproductive effects (e.g. cryptorchidism, prostate cancer), and childhood diseases (e.g. obesity, cancer), which are likely to be influenced by EDC exposures have not been fully evaluated for their contribution. As new efforts are initiated to address knowledge gaps the links between EDCs and reproductive and development outcomes, understanding how findings from human and model organisms can each contribute to filling the gaps is critical. Findings from human observational studies provide direct evidence of the relationship between exposure to environmental chemicals and subsequent adverse health effects. However, limitations in the epidemiologic data, including scientific and ethical, require the use of findings from mammalian and other model organisms to complement human epidemiologic evidence. This paper reviews some of the key scientific concepts relevant to integrating information from human epidemiologic and model organisms to understand the relationship between EDC exposure and adverse human health effects. This is followed by two cases studies: first on thyroid disrupting chemicals and second on anti-androgens chemicals, illustrating how the interplay between the findings from two disciplines can enhance our understanding of the relationship between exposure and disease.

Key Concepts and Emerging Science

Sources of Scientific Information

Our understanding of potentially harmful effects of exposure to environmental contaminants comes from a variety of sources including wildlife life studies, *in vitro* and *in vivo* toxicology studies, epidemiologic studies, and clinical evidence. The most extensive information comes from animal studies, which are a preferable method for assessing the potential for adverse human effects and for developing strategies for prevention of harmful exposures. Environmental contaminants are not intended for human use, and it is unethical to knowingly expose humans to these chemicals under experimental conditions to assess for harmful effects.

The role of model organisms—Animals have long been used to understand the effects of chemical exposure on human reproduction and development [21]. One of the first reproductive and developmental studies, from 1919, used rats to study the effects alcohol on fetal development, before it was identified in humans [22, 23]. The reliability of experimental animal data for reproductive and developmental health has been well established and presently, there is no example of a chemical agent that has adversely affected human reproduction or development but has not caused the same or similar adverse effects in animal models [24]. Model organisms studies have many advantages, including ability to control the experimental setting to enhance ability to observe chemical effects, and shorter time from exposure to disease [25]. Multiple reviews have concluded that there is concordance of developmental and reproductive effects and that humans are as sensitive or more sensitive than the most sensitive animal species [23, 26] [23, 24, 26-29]. Given that there is general conservation of biologic function across animal species, including humans; animal studies provide important insights into potential human harm [30].

However, limitations in the design of traditional toxicological studies have limited their utility for studying reproductive or developmental outcomes. These include: use of insensitive strains; exposures that do not cover sensitive periods of development; exposures to single chemicals at high doses rather than the mixture of low doses of chemicals more often seen by the public; and a focus on overt disease endpoints. While animal studies continue to provide important information to predict human harm, epidemiologic and clinical studies provide complimentary sources of information.

The role of epidemiology—Human observational studies of environmental chemicals provide the most direct evidence of the relationship between exposure and increased risk of adverse health outcomes, and are often the basis of assessments about harmful effects of EDCs. Studies are typically designed to evaluate whether the change in the risk factor of interest, or the chemical exposure, is related to the change in the incidence or prevalence of the disease of study, while at the same time accounting for factors that may influence that relationship. Model animal studies tend to narrow their methodological scope to increase their ability to detect presence of an effect from chemical exposure, while human observational studies necessarily include the factors which can influence the exposure/effect relationship as they are often impossible to control. These include pre-existing exposure to other potentially influential chemicals, biological variability (e.g. disease status, age, gender), and typically lower levels of exposure. This makes the interpretation of the results of observational human studies challenging because these factors that influence risk (e.g. age, disease status, co-exposures), can obscure the signal for a particular chemical/outcome relationship. Features of model organisms and epidemiology are illustrated in Table 2.

Emerging Scientific Themes and their Implications

Since the latter part of the 20th century, the evolution of our understanding about the role of EDCs in human health has been accompanied by increasing insights into additional factors which influence the relationship between chemical exposure and risk of adverse health effects. These include: enhanced understanding of toxicity pathways; importance of timing of exposure; contribution of multiple chemical exposures; and low dose effects. The growth in the science in each of these areas is influencing study designs in epidemiology and model organisms, the interpretation of the data from each, and the need to integrate knowledge from both the fields. Each of these areas and their contribution to the field is discussed below.

Biological pathways—Advances in molecular biology, computational biology and information sciences is transforming our ability to interrogate and understand disease etiology on multiple levels, which in turn requires advancing our approaches to assessing potential risks from chemical exposures [31, 32]. We are faced with expanding information at the genetic level, such as through genome wide association studies [33], epigenetic assessments [34], and metabolomics [35], providing unprecedented information about the pathways from exposure to overt disease. Concurrently, the technological capacity to test multiple chemicals and their effects on biological pathways rapidly through high throughput vitro assays expands the amount of knowledge available for interpretation [36]. The advancement of scientific ability to evaluate early perturbations in the disease process

increases the opportunity to identify potentially harmful chemicals using early biological markers of disease [4], and is a feature which can be integrated across the interpretation data from human and model organism studies.

Periods of Susceptibility—Human development is vulnerable to biological disruption particularly when these changes occur during critical windows of development, which typically occur periconceptually, during pregnancy, infancy, childhood, puberty, pregnancy and lactation. Perturbations from chemical exposures can lead to important functional deficits and increased risks of disease and disability in infants, children and across the entire span of human life [4, 30, 37]. The prenatal period has become a particular area of focus of environmental chemical exposure for two primary reasons. First, over the past 60 years it has become clear that the placenta does not protect the fetus from damaging chemicals, [38], that chemicals can be found in the fetus, and in some cases may be biomagnified. For example, analysis of second trimester amniotic fluid samples from 51 women found the presence of at least one environmental contaminant. [39], and for mercury, fetal exposures to environmental contaminants may be higher than maternal. [40-42]. Second, while the effects of prenatal exposure on immediate outcomes, such as birth defects or childhood illness has been relatively well known [15], newer science find that *in utero* disturbances from the external environment can increase risk of adult disease (in particular the central nervous system, the cardiovascular system, the endocrine system and the immune system), referred to as the Developmental Basis of Adult Disease/Dysfunction [2].

Chemical background, exposure to mixtures and biomonitoring—Numerous animal studies now find that co-exposures to chemicals can enhance risks of adverse health effects compared to risks from exposure to individual chemicals, even if the chemicals act through different mechanisms of action [43-46]. The National Academy of Sciences has recently recommended that chemical exposures that can affect the same common adverse outcome should be considered, and in many cases are additive [47].

Studies of exposure pathways show the population is in constant contact with environmental chemicals in air, food, water and consumer products [48-51]. Now we have extensive biomonitoring data (measurements of chemicals in biological matrices), which show ubiquitous exposure to many EDCs in the general population found in homes, communities and workplaces [8]. Further, new analysis of CDC population-based biomonitoring data among pregnant women in the U.S. found virtually all pregnant women have body burdens of a number of EDCs in their body include certain pesticides, perchlorate, Bisphenol-A (BPA), phthalates, perfluorochemicals (PFCs), polychlorinated biphenyls (PCBs) and polybrominated diphenyl ethers (PBDEs) (Table 3).

Biological background—Background health status, as influenced by age, pre-existing disease, genetics, and other factors, can influence the effect of chemical exposure on subsequent health risks. For example, individuals with preexisting immune suppression, such as organ transplant patients, those who are HIV positive, and those at early or late lifestages, might experience disproportionately higher risk of adverse health effects due to chemical exposure compared to more healthy individuals. A vast number of biological processes that are ongoing throughout a lifetime can contribute to disease etiology. For

example, neurological deficits and cancer, are relatively prevalent in the population, and have multiple additional factors that contribute to their disease etiology. Thus, we would expect that other ongoing biological processes are already occurring that can contribute independently to disease risk, and EDC exposure would be in addition to these ongoing biological processes [52].

Multiple Mechanisms of Action—Exposure to a chemical can influence disease processes through multiple mechanisms of action. Hence, studies that focus on a particular pathway or individual biological event may provide an incomplete picture of the range of effects produced by a particular chemical [53].

Effects at Low Doses—Traditionally, the evaluation of noncancer effects from chemical assumes that there is an exposure below which exposures are anticipated to not contribute to increase risk of disease (referred to as a “threshold”). However, scientific evidence now finds that adverse health effects can occur from exposures at common population levels (e.g. lead and mercury), and threshold cannot be identified in the human population. Further, background chemical exposures, pre-existing vulnerabilities in the population, such as from age and pre-existing disease, can contribute to population background risks such that practically, a threshold will not exist in the population and we expect risks even at low exposures [52].

Implications and Case Studies

Our increasing appreciation of the influence of biological background (i.e. age and disease status) and chemical background to EDC and disease etiology comes from integrating findings from human and model organisms studies. In turn, this base of knowledge guides the interpretation of insights from studies from both disciplines. The following case studies illustrate the importance of each and how scientific concepts aid in interpreting and expanding our knowledge base.

Thyroid hormone perturbations and neurodevelopmental outcomes—This case study of thyroid hormone (TH) perturbations during pregnancy and neurological outcomes illustrates how human studies provided initial evidence of the link between exposure and outcome that was later enhanced by studies in model organisms.

Thyroid Hormone Overview: THs, thyroxine (T₄) and triiodothyronine (T₃), are essential to neurological development. Insufficient levels of TH during development can lead to mild to severe cognitive impairment, neurobehavioral disorders, hypomyelination and attendant physical impairments [54, 55].

First insights into effects of TH decrements during pregnancy on neurological development comes from studies during the first half of the 20th century of human populations with severe endemic goiter, caused by insufficient TH, where children were found to be increased risk of mental retardation and other neurocognitive deficits [56]. While clinically relevant decrements in T₄ during pregnancy were recognized as harmful, later epidemiological studies observed neurobehavioral delays in children born to women with mild to moderately low levels of T₄, not indicative of maternal hypothyroidism, during the first trimester of

pregnancy [57-59] [57, 59-61]. Other epidemiologic studies found that small deficits in circulating levels of TH are associated with decreased cognitive performance at various times during development and adulthood [62-68].

Period of susceptibility: The fetal thyroid gland does not begin to concentrate iodide until the 12th gestational week [69] and does not produce significant quantities of TH until about week 20 [69]. Thus, until about the 20th week the fetus is entirely dependent on maternal THs [70], and T4 decrements during this time period can have a greater impact of subsequent neurological risks.

While human studies have shown the importance of the relationship between maternal TH and neurocognitive development in the child, mechanistic studies of the role of chemical exposures in TH changes has been an important complement to the field. They have enhanced our understanding of the mechanisms involved, the changing TH levels during fetal development, role of multiple chemical exposures, and the importance of different THs to neurodevelopment.

The effects of TH on the developing brain are directly related to serum concentrations of T₄ in laboratory animal studies, with animal studies also finding that the developing brain is sensitive to small reductions in serum TH [71-79]. For example, declining serum T₄ has been shown to be associated with linear declines in the number of oligodendrocytes and increases in the number of astrocytes in white matter of rat pups [80], and effects on the developing hippocampus [77].

Multiple Chemical Exposures.: A number of environmental chemicals are capable of disrupting TH levels, often through different mechanisms (illustrated in Table 4) [81, 82]. For example, perchlorate inhibits the uptake of iodine, resulting in decreased synthesis of TH. PCBs and the pesticide acetochlor activate enzymes in the liver that increase excretion of TH, thus reducing circulating levels of TH [83]. Crofton et al. evaluated to--rats exposed to a mixture of 18 TDCs (dioxins, dibenzofurans and PCBs), at doses comparable to human exposures, for effects on serum T₄ [43]. Components of this mixture affect T₄ through two different mechanisms of action: the dioxins, dibenzofurans and dioxin-like PCBs activate one set of liver enzymes, and the non-dioxin-like PCBs activate a separate set of liver enzymes. Crofton et al. found the mixture had a dose-additive effect on T₄ at environmentally-relevant doses and a 2-3 fold greater than dose-additive effect on T₄ at higher doses [43], demonstrating that exposures to TDCs acting on different mechanisms can have cumulative effects.

Metrics of TH disruption: Clinical sciences first relies on changes in TSH to indicate thyroid related condition [84]. However, there are situations where serum T₄ and T₃ levels may not be associated with serum TSH as they are under normal conditions, such as in the case of liver disease, or if there are defects in the proteins responsible for transporting T₄ and T₃ in the circulation or in the receptors mediating negative feedback of T₄ on TSH [85, 86]. Likewise, there are chemicals, such as PCBs, that can cause a decrease in serum total and free T₄ in animals without causing a concomitant increase in serum TSH [87, 88].

Biological Background.: Surveillance of the human population finds significant portions are already at risk from additional T₄ decrements. About one third of U.S. women, including pregnant women, have low iodine intake (iodine is required for TH synthesis) [89]). In addition, between 1999-2002, 7.3% of the U.S. population aged 12 years and older reported that they had thyroid disease or were taking thyroid medication [90]. In addition, among women receiving TH replacement medication for treatment of hypothyroidism, 14% had TSH and T₄ measurements indicating continued hypothyroidism [90]. Pregnancy causes an increased demand on the thyroid gland and hypothyroidism is more than twice as common among pregnant women than among non-pregnant women ages 12-49 [90].

Conclusions Regarding Thyroid Data: A number of environmental chemicals are capable of disrupting TH levels, including decreasing circulating levels of T₄. Compensatory mechanisms are likely insufficient to counteract the potential adverse health effects of these T₄ decrements, and thus the fetus is at increased risk for neurological effects from exposure to low levels of TDCs. This conclusion is supported by complimentary findings from the animal and human literature. First, findings in both animals and humans indicate that even small T₄ decrements can have adverse neurodevelopmental consequences in the offspring. Second, fetuses and infants do not have stored TH, and thus have limited capacity to respond to TH decrements during critical stages of development. Third, human evidence find a substantial prevalence of TH and iodine insufficiency in the population of U.S. women, putting them an increased risk of exposure to chemicals that affect TH levels. Fourth, animal studies find that exposure to TDCs can act together to affect TH levels and that human studies show people are currently exposed to multiple TDCs.

Anti-Androgens and male reproductive development—A number of different chemicals can affect androgen action. Phthalates are a relatively well characterized class of chemicals in this group and provide a case example of how model organisms played a pivotal role in our understanding of the effects on anti-androgenic substances on male reproductive development that lead to complimentary human studies, significantly increasing our overall knowledge in this field.

Phthalates are a group of industrial chemicals used as plasticizers, and as solubilizing and stabilizing agents. They are found in numerous products including personal care products, medical devices, various plastics include flooring, polyvinyl chloride type plastics and tubings [8]. It was discovered in 2000 that phthalates were also ubiquitously measured in people through reports from the first CDC national biomonitoring report and in publication [91].

Effects of Exposure to Anti-Androgens During Fetal Life: Proper male reproductive tissue development is dependent on a transient peak in testosterone levels during fetal development. Disruption to androgen activity during this critical developmental window can result in a number of male reproductive tract abnormalities, including retained female structures, such as nipples (in animals), and malformations of male reproductive structures, such as hypospadias (an abnormal location of the urethral opening) and cryptorchidism [53].

Evolution of our understanding of critical windows and anti-androgens: Early studies of phthalates focused on adult exposures and primarily effects other than reproductive, though some standard reproductive studies noted effects on male rat testicles [92]. Differential effects by age were noted in early studies, with early ages being more susceptible, though it was not until later studies when effects specific to the fetal exposures were identified through multigeneration study of DBP in the rat [93]. Follow up studies in rats identified a critical window of exposure during fetal developmental [94-98]. Additionally, studies identified a sensitive period of exposure during gestational days 15-17 for dibutyl phthalate exposures in the rat, about gestational weeks 8 to 14 in humans [99].

Phthalates appear to interfere with androgen activity by inhibiting testosterone synthesis. Specifically, phthalates with four to six carbon side chains interfere with the production of testosterone by inhibiting the uptake of cholesterol into the mitochondria by Steroidogenic Acute Regulatory protein (StAR) protein and by inhibiting some, but not all, of the enzymes in the steroidogenic pathway [100]. These phthalates have been most consistently shown to reduce testosterone production in animal research to date, though other phthalates may also be important [101, 102].

Studies in rats have demonstrated that a phthalate-induced decrease in testosterone predictably results in a syndrome of anti-androgenic reproductive abnormalities characterized by: malformations of the epididymis, vas deferens, seminal vesicles, prostate, hypospadias, cryptorchidism and testicular injury; permanent changes (feminization) in the retention of nipples and areolae (sexually dimorphic structures in rodents); and demasculinization of the growth of the perineum, resulting in a reduced anogenital distance [98]. This constellation of effects has been referred to as the “phthalate syndrome” and, as with other anti-androgenic chemicals, the severity of effects increases with the dose [99].

The spectrum of effects seen in phthalate syndrome parallels a spectrum of human diseases that has been termed “testicular dysgenesis syndrome” (TDS): infertility, cryptorchidism, hypospadias, and testicular cancer, which has been hypothesized to have a fetal origin associated with male development of the testis [103].

Role of human observational studies: In 2000, with publication of the first CDC biomonitoring results in the peer reviewed literature, Blount et al. showed that there was widespread exposure in the US population to a number of different phthalates, finding metabolites of DBP, DEP and BzBP in over 95% of the population [91]. They also found that women of reproductive age had significantly higher levels of metabolites of DBP than other age and gender groups. While phthalates had been measured in the various species since the 1970s, including deep-sea jellyfish [104], fish caught in various parts of North America [105], and bovine tissue [106], (this was the first measurements in humans of wide spread common exposure). Further, the higher exposures among women of reproductive age combined with findings in animals suggested that male reproductive development could be at risk.

Given ubiquitous exposure in the human population, identified male reproductive effects, and analytic ability to measure body burdens, human epidemiologic studies could evaluate

potential health effects at exposure levels experienced by the general population, which are lower than the doses in the animal studies. Several epidemiological studies have evaluated the relationship between adult exposures and male reproductive effects, in particular semen quality – with some studies finding an association between increased phthalate levels and decreases in semen quality in males [107], (though not in every study [108]).

Other human studies evaluated prenatal exposures, previously identified by animal studies as a sensitive period for exposure. In particular, the study by Swan et al. found associations between certain prenatal phthalate exposure and male developmental reproductive outcomes, in particular shortened ano-genital distance [109]. This study was notable for several reasons. First, it used a marker of male reproductive effects reflecting feminization that was common in toxicological studies, but not applied in epidemiologic studies – ano-genital distance. Subsequent human studies have observed ano-genital distance to be sexual dimorphic and reflective of penile length and growth in males [110, 111]. This novel application was possible only by cross fertilization across model organisms and human observation studies. Other studies also identified effects from developmental exposures including changes in reproductive hormone levels in male infants exposed to phthalates in breast milk [112].

Exposure to Mixtures of Androgen Receptor Antagonists and Androgen Synthesis

Disruptors: Studies find that exposure to a mixture of chemicals with both similar and different mechanisms of action results in dose-additive effects [46, 113]. Rats exposed to a mixture of androgen receptor antagonists, vinclozolin, procymidone, and flutamide, at doses that would not have caused hypospadias alone, resulted in over 50% of animals with hypospadias [113]. A second study analyzed the effects of prenatal exposure to a mixture of seven different anti-androgens (phthalates and pesticides) with differing mechanisms of action (i.e., androgen receptor antagonist or inhibition of androgen synthesis) [46]. All combinations produced cumulative, dose additive outcomes in the androgen-dependent tissues. Swan found that a summary measure of phthalates, integrating exposure to five phthalate metabolites associated with ano-genital distance, was more predictive of the ano-genital distance than the individual metabolite concentrations [114].

Conclusion Regarding Anti-androgen Data: The prenatal exposure of males to anti-androgenic chemicals illustrates how findings from animal and human studies together enhance our understanding of effects: 1) animal studies identified critical periods of development sensitive to exposures; 2) animal and human studies propose that perturbations early in the pathway of male reproductive development result in a wide array of permanent and irreversible adverse outcomes; and 3) animal studies show that exposure to different chemicals that affect male reproductive development can have a cumulative effect even if through different mechanisms of action.

Conclusions

The complexity of human disease requires that we draw broadly upon available information to understand the potential impacts of EDCs. Our knowledge of the EDC/disease relationship is strengthened and data gaps reduced when we integrate findings from animal

and human studies. Our evolving understanding of the relationship between exposure and disease and the factors that can influence risk, including chemical background exposures, disease status and susceptibility can put a portion of the population at risk from small, incremental exposures, as illustrated in Table 5. Thus, future research efforts should build upon these findings to improve the utility of new research and to contribute to the efforts to understand, and prevent, exposures to harmful chemicals.

References

- [1]. Woodruff TJ, Schwartz J, Giudice LC, Research agenda for environmental reproductive health in the 21st century, *J Epidemiol Community Health* 64(4) (2010) 307–310. [PubMed: 20348330]
- [2]. Diamanti-Kandarakis E, Bourguignon JP, Giudice LC, Hauser R, Prins GS, Soto AM, Zoeller RT, Gore AC, Endocrine-disrupting chemicals: an Endocrine Society scientific statement, *Endocr Rev* 30(4) (2009) 293–342. [PubMed: 19502515]
- [3]. Euling SY, Selevan SG, Pescovitz OH, Skakkebaek NE, Role of environmental factors in the timing of puberty, *Pediatrics* 121 Suppl 3 (2008) S167–171. [PubMed: 18245510]
- [4]. Woodruff TJ, Carlson A, Schwartz JM, Giudice LC, Proceedings of the Summit on Environmental Challenges to Reproductive Health and Fertility: Executive summary, *Fertil Steril* 89(2 Suppl) (2008) e1–e20. [PubMed: 18308046]
- [5]. Federal Reserve Board, G.17 - Industrial Production and Capacity Utilization, Industrial Production for Chemicals (NAICS=325), The Federal Reserve Board Data Download Program, 2008 p. <http://www.federalreserve.gov/datadownload/Download.aspx?rel=G17&series=4374e4378caec4376beaa4874d28776fa28768c28771e&filetype=sheet&l&label=include&lavout=seriescolumn&from=28701/28701/21919&to=28712/28731/22009>.
- [6]. U.S. EPA, What is the TSCA Chemical Substance Inventory, US Environmental Protection Agency, 2006.
- [7]. National Center for Environmental Health, Third National Report on Human Exposure to Environmental Chemicals, US Department of Health and Human Services, Atlanta, GA, 2005.
- [8]. CDC, Fourth National Report on Human Exposure to Environmental Chemicals Centers for Disease Control and Prevention, National Center for Environmental Health, Atlanta, GA, 2009.
- [9]. Porta M, Puigdomènech E, Ballester F, Selva J, Ribas-Fitó N, Llop S, López T, Monitoring concentrations of persistent organic pollutants in the general population: The international experience, *Environment International* 34(4) (2008) 546–561. [PubMed: 18054079]
- [10]. Arctic Monitoring and Assessment Programme (AMAP), AMAP Assessment 2009: Human Health in the Arctic, Oslo, 2009.
- [11]. Guo YL, Lambert GH, Hsu CC, Hsu MM, Yucheng: health effects of prenatal exposure to polychlorinated biphenyls and dibenzofurans, *Int Arch Occup Environ Health* 77(3) (2004) 153–158. [PubMed: 14963712]
- [12]. Harada M, Congenital Minamata disease: intrauterine methylmercury poisoning, *Teratology* 18(2) (1978) 285–288. [PubMed: 362594]
- [13]. Harada M, Minamata disease: methylmercury poisoning in Japan caused by environmental pollution, *Crit Rev Toxicol* 25(1) (1995) 1–24. [PubMed: 7734058]
- [14]. Teitelbaum DT, The toxicology of 1,2-dibromo-3-chloropropane (DBCP): a brief review, *Int J Occup Environ Health* 5(2) (1999) 122–126. [PubMed: 10330513]
- [15]. Stillerman KP, Mattison D, Giudice L, Woodruff TJ, Environmental exposures and adverse pregnancy outcomes: a review of the science, *Reprod Sci* 15(7) (2008) 631–650. [PubMed: 18836129]
- [16]. Lederman SA, Jones RL, Caldwell KL, Rauh V, Sheets SE, Tang D, Viswanathan S, Becker M, Stein JL, Wang RY, Perera FP, Relation between cord blood mercury levels and early child development in a World Trade Center cohort, *Environ Health Perspect* 116(8) (2008) 1085–1091. [PubMed: 18709170]

- [17]. Patandin S, Lanting CI, Mulder PGH, Boersma ER, Sauer PJJ, Weisglas-Kuperus N, Effects of environmental exposure to polychlorinated biphenyls and dioxins on cognitive abilities in Dutch children at 42 months of age, *J. Pediatr* 134(1) (1999) 33–41. [PubMed: 9880446]
- [18]. Stewart PW, Lonky E, Reihman J, Pagano J, Gump BB, Darvill T, The relationship between prenatal PCB exposure and intelligence (IQ) in 9-year-old children, *Environ Health Perspect* 116(10) (2008) 1416–1422. [PubMed: 18941588]
- [19]. Clewell HJ, Crump KS, Quantitative estimates of risk for noncancer endpoints, *Risk Anal* 25(2) (2005) 285–289. [PubMed: 15876204]
- [20]. Rall DP, Thresholds?, *Environ Health Perspect* 22 (1978) 163–165. [PubMed: 648484]
- [21]. Holson JF, Desesso JM, Jacobson CF, Farr CH, Appropriate use of animal models in the assessment of risk during prenatal development: an illustration using inorganic arsenic, *Teratology* 62(1) (2000) 51–71. [PubMed: 10861633]
- [22]. Arlitt A, The effect of alcohol on the intelligent behavior of the white rat and its progeny, *Psychol. Monog* 26 (1919) 1–50.
- [23]. Kimmel CA, Holson JF, Hogue CJ, Carlo G, Reliability of Experimental Studies for Predicting Hazards to Human Development, NCTR Technical Report for Experiment No. 6015, Jefferson, AR, 1984.
- [24]. Nemec MD, Kaufman LE, Stump DG, Lindstrom P, Varsho BJ, Holson JF, Significance, Reliability, and Interpretation of Developmental and Reproductive Toxicity Study Findings, *Developmental Reproductive Toxicology: A Practical Approach*, Informa Healthcare, 2006.
- [25]. Zapponi GA, Marcello I, Carere A, Prevention, ethics and science: lessons from Lorenzo Tomatis, *Ann Ist Super Sanita* 44(1) (2008) 8–12. [PubMed: 18469370]
- [26]. National Research Council, Scientific Frontiers in Developmental Toxicology and Risk Assessment, National Academy Press, Washington, DC, 2000.
- [27]. Hemminki K, Vineis P, Extrapolation of the evidence on teratogenicity of chemicals between humans and experimental animals: chemicals other than drugs, *Teratog Carcinog Mutagen* 5(4) (1985) 251–318. [PubMed: 2866606]
- [28]. Newman LM, Johnson EM, Staples RE, Assessment of the Effectiveness of Animal Developmental Toxicity Testing for Human Safety, *Reproductive Toxicology* 7(4) (1993) 359–390. [PubMed: 8400625]
- [29]. Nisbet ICT, Karch NJ, Chemical Hazards to Human Reproduction, Noyes Data Corp., Park Ridge, NJ, 1983.
- [30]. Crain DA, Janssen SJ, Edwards TM, Heindel J, Ho SM, Hunt P, Iguchi T, Juul A, McLachlan JA, Schwartz J, Skakkebaek N, Soto AM, Swan S, Walker C, Woodruff TK, Woodruff TJ, Giudice LC, Guillette LJ Jr., Female reproductive disorders: the roles of endocrine-disrupting compounds and developmental timing, *Fertil Steril* 90(4) (2008) 911–940. [PubMed: 18929049]
- [31]. Collins FS, Gray GM, Bucher JR, Toxicology. Transforming environmental health protection, *Science* 319(5865) (2008) 906–907. [PubMed: 18276874]
- [32]. National Research Council, Toxicity Testing in the 21st Century: A Vision and Strategy, The National Academies Press, Washington, DC, 2007.
- [33]. National Institutes of Health, Genome-Wide Association Studies (GWAS), 2010.
- [34]. Weinhold B, Epigenetics: the science of change, *Environ Health Perspect* 114(3) (2006) A160–167. [PubMed: 16507447]
- [35]. Guyton KZ, Kyle AD, Aubrecht J, Cogliano VJ, Eastmond DA, Jackson M, Keshava N, Sandy MS, Sonawane B, Zhang L, Waters MD, Smith MT, Improving prediction of chemical carcinogenicity by considering multiple mechanisms and applying toxicogenomic approaches, *Mutat Res* 681(2-3) (2009) 230–240. [PubMed: 19010444]
- [36]. USEPA, ToxCast, 2010.
- [37]. Grandjean P, Bellinger D, Bergman A, Cordier S, Davey-Smith G, Eskenazi B, Gee D, Gray K, Hanson M, van den Hazel P, Heindel JJ, Heinzow B, Hertz-Picciotto I, Hu H, Huang TT, Jensen TK, Landrigan PJ, McMillen IC, Murata K, Ritz B, Schoeters G, Skakkebaek NE, Skerfving S, Weihe P, The faroes statement: human health effects of developmental exposure to chemicals in our environment, *Basic Clin Pharmacol Toxicol* 102(2) (2008) 73–75. [PubMed: 18226057]

- [38]. Rusyniak DE, Furbee RB, Pascuzzi R, Historical neurotoxins: what we have learned from toxins of the past about diseases of the present, *Neurol Clin* 23(2) (2005) 337–352. [PubMed: 15757788]
- [39]. Foster W, Chan S, Platt L, Hughes C, Detection of endocrine disrupting chemicals in samples of second trimester human amniotic fluid, *J Clin Endocrinol Metab* 85(8) (2000) 2954–2957. [PubMed: 10946910]
- [40]. Barr DB, Bishop A, Needham LL, Concentrations of xenobiotic chemicals in the maternal-fetal unit, *Reprod Toxicol* 23(3) (2007) 260–266. [PubMed: 17386996]
- [41]. Rollin HB, Rudge CV, Thomassen Y, Mathee A, Odland JO, Levels of toxic and essential metals in maternal and umbilical cord blood from selected areas of South Africa--results of a pilot study, *J Environ Monit* 11(3) (2009) 618–627. [PubMed: 19280040]
- [42]. Stern AH, Smith AE, An assessment of the cord blood:maternal blood methylmercury ratio: implications for risk assessment, *Environ Health Perspect* 111(12) (2003) 1465–1470. [PubMed: 12948885]
- [43]. Crofton KM, Craft ES, Hedge JM, Gennings C, Simmons JE, Carchman RA, Carter WH Jr., DeVito MJ, Thyroid-hormone-disrupting chemicals: evidence for dose-dependent additivity or synergism, *Environ Health Perspect* 113(11) (2005) 1549–1554. [PubMed: 16263510]
- [44]. Gray LE Jr., Wilson VS, Stoker T, Lambright C, Furr J, Noriega N, Howdeshell K, Ankley GT, Guillette L, Adverse effects of environmental antiandrogens and androgens on reproductive development in mammals, *Int J Androl* 29(1) (2006) 96–104; discussion 105–108. [PubMed: 16466529]
- [45]. Rajapakse N, Silva E, Kortenkamp A, Combining xenoestrogens at levels below individual no-observed-effect concentrations dramatically enhances steroid hormone action, *Environ Health Perspect* 110(9) (2002) 917–921. [PubMed: 12204827]
- [46]. Rider CV, Furr J, Wilson VS, Gray LE, A mixture of seven antiandrogens induces reproductive malformations in rats, *International Journal of Andrology* 31(2) (2008) 249–262. [PubMed: 18205796]
- [47]. National Research Council, Phthalates and Cumulative Risk Assessment, in: N.A.o. Science (Ed.), Washington D.C., 2008.
- [48]. EWG, National Tap Water Quality Database, Environmental Working Group, Washington, DC, 2005.
- [49]. NLM, Household Products Database, National Institutes of Health, National Library of Medicine, Bethesda, MD.
- [50]. U.S. Department of Agriculture, Pesticide Data Program; Annual Summary, Calendar Year 2006, Washington, DC, 2007.
- [51]. Dougherty CP, Henricks Holtz S, Reinert JC, Panyacosit L, Axelrad DA, Woodruff TJ, Dietary exposures to food contaminants across the United States, *Environ Res* 84(2) (2000) 170–185. [PubMed: 11068931]
- [52]. National Academy of Science, Science and Decisions: Advancing Risk Assessment, National Research Council Committee on Improving Risk Analysis Approaches Used by the U.S. EPA, Washington, DC, 2008.
- [53]. Woodruff TJ, Zeise L, Axelrad DA, Guyton KZ, Janssen S, Miller M, Miller GG, Schwartz JM, Alexeeff G, Anderson H, Birnbaum L, Bois F, Cogliano VJ, Crofton K, Euling SY, Foster PM, Germolec DR, Gray E, Hattis DB, Kyle AD, Luebke RW, Luster MI, Portier C, Rice DC, Solomon G, Vandenberg J, Zoeller RT, Meeting report: moving upstream-evaluating adverse upstream end points for improved risk assessment and decision-making, *Environ Health Perspect* 116(11) (2008) 1568–1575. [PubMed: 19057713]
- [54]. van der Sluijs Veer L, Kempers MJ, Last BF, Vulsma T, Grootenhuis MA, Quality of life, developmental milestones, and self-esteem of young adults with congenital hypothyroidism diagnosed by neonatal screening, *J Clin Endocrinol Metab* 93(7) (2008) 2654–2661. [PubMed: 18460566]
- [55]. Zoeller RT, Crofton KM, Mode of action: developmental thyroid hormone insufficiency--neurological abnormalities resulting from exposure to propylthiouracil, *Crit Rev Toxicol* 35(8-9) (2005) 771–781. [PubMed: 16417044]

- [56]. de Escobar GM, Obregon MJ, del Rey FE, Maternal thyroid hormones early in pregnancy and fetal brain development, *Best Pract Res Clin Endocrinol Metab* 18(2) (2004) 225–248. [PubMed: 15157838]
- [57]. Haddow JE, Palomaki GE, Allan WC, Williams JR, Knight GJ, Gagnon J, O'Heir CE, Mitchell ML, Hermos RJ, Waisbren SE, Faix JD, Klein RZ, Maternal thyroid deficiency during pregnancy and subsequent neuropsychological development of the child, *N Engl J Med* 341(8) (1999) 549–555. [PubMed: 10451459]
- [58]. Morreale de Escobar G, Obregón MJ, Escobar del Rey F, Is neuropsychological development related to maternal hypothyroidism or to maternal hypothyroxinemia?, *J. Clin. Endocrinol. Metab* 85 (2000) 3975–3987. [PubMed: 11095417]
- [59]. Pop VJ, Brouwers EP, Vader HL, Vulsma T, van Baar AL, de Vijlder JJ, Maternal hypothyroxinaemia during early pregnancy and subsequent child development: A 3-year follow-up study, *Clin Endocrinol (Oxf)* 59(3) (2003) 282–288. [PubMed: 12919150]
- [60]. Haddow JE, Subclinical hypothyroidism and pregnancy outcomes, *Obstet Gynecol* 106(1) (2005) 198; author reply 198–199. [PubMed: 15994654]
- [61]. Pop VJ, Kuijpers JL, van Baar AL, Verkerk G, van Son MM, de Vijlder JJ, Vulsma T, Wiersinga WM, Drexhage HA, Vader HL, Low maternal free thyroxine concentrations during early pregnancy are associated with impaired psychomotor development in infancy, *Clin Endocrinol (Oxf)* 50(2) (1999) 149–155. [PubMed: 10396355]
- [62]. Glinoer D, Rovet J, Gestational hypothyroxinemia and the beneficial effects of early dietary iodine fortification, *Thyroid* 19(5) (2009) 431–434. [PubMed: 19415991]
- [63]. Heyerdahl S, Oerbeck B, Congenital hypothyroidism: developmental outcome in relation to levothyroxine treatment variables, *Thyroid* 13(11) (2003) 1029–1038. [PubMed: 14651787]
- [64]. LaFranchi SH, Austin J, How should we be treating children with congenital hypothyroidism?, *J Pediatr Endocrinol Metab* 20(5) (2007) 559–578. [PubMed: 17642417]
- [65]. Oerbeck B, Sundet K, Kase BF, Heyerdahl S, Congenital hypothyroidism: influence of disease severity and L-thyroxine treatment on intellectual, motor, and school-associated outcomes in young adults, *Pediatrics* 112(4) (2003) 923–930. [PubMed: 14523187]
- [66]. Rovet J, Simic N, The role of transient hypothyroxinemia of prematurity in development of visual abilities, *Semin Perinatol* 32(6) (2008) 431–437. [PubMed: 19007682]
- [67]. Selva KA, Harper A, Downs A, Blasco PA, Lafranchi SH, Neurodevelopmental outcomes in congenital hypothyroidism: comparison of initial T4 dose and time to reach target T4 and TSH, *J Pediatr* 147(6) (2005) 775–780. [PubMed: 16356430]
- [68]. Simic N, Asztalos EV, Rovet J, Impact of neonatal thyroid hormone insufficiency and medical morbidity on infant neurodevelopment and attention following preterm birth, *Thyroid* 19(4) (2009) 395–401. [PubMed: 19355829]
- [69]. Morreale de Escobar G, Obregon MJ, Escobar del Rey F, Is neuropsychological development related to maternal hypothyroidism or to maternal hypothyroxinemia?, *J Clin Endocrinol Metab* 85(11) (2000) 3975–3987. [PubMed: 11095417]
- [70]. Morreale de Escobar G, The role of thyroid hormone in fetal neurodevelopment, *Journal of Pediatric Endocrinology and Metabolism* 13 (2001) 1453–1462.
- [71]. Gilbert ME, Paczkowski C, Propylthiouracil (PTU)-induced hypothyroidism in the developing rat impairs synaptic transmission and plasticity in the dentate gyrus of the adult hippocampus, *Brain Res Dev Brain Res* 145(1) (2003) 19–29. [PubMed: 14519490]
- [72]. Gilbert ME, Alterations in synaptic transmission and plasticity in area CA1 of adult hippocampus following developmental hypothyroidism, *Brain Res Dev Brain Res* 148(1) (2004) 11–18. [PubMed: 14757514]
- [73]. Gilbert ME, Sui L, Dose-dependent reductions in spatial learning and synaptic function in the dentate gyrus of adult rats following developmental thyroid hormone insufficiency, *Brain Res* 1069(1) (2006) 10–22. [PubMed: 16406011]
- [74]. Gilbert ME, Sui L, Developmental exposure to perchlorate alters synaptic transmission in hippocampus of the adult rat, *Environ Health Perspect* 116(6) (2008) 752–760. [PubMed: 18560531]

- [75]. Gilbert ME, Sui L, Walker MJ, Anderson W, Thomas S, Smoller SN, Schon JP, Phani S, Goodman JH, Thyroid hormone insufficiency during brain development reduces parvalbumin immunoreactivity and inhibitory function in the hippocampus, *Endocrinology* 148(1) (2007) 92–102. [PubMed: 17008398]
- [76]. Goodman JH, Gilbert ME, Modest Thyroid Hormone Insufficiency During Development Induces a Cellular Malformation in the Corpus Callosum: A Model of Cortical Dysplasia, *Endocrinology* (2007).
- [77]. Royland JE, Parker JS, Gilbert ME, A genomic analysis of subclinical hypothyroidism in hippocampus and neocortex of the developing rat brain, *Journal of neuroendocrinology* 20(12) (2008) 1319–1338. [PubMed: 19094080]
- [78]. Sui L, Anderson WL, Gilbert ME, Impairment in short-term but enhanced long-term synaptic potentiation and ERK activation in adult hippocampal area CA1 following developmental hypothyroidism, *Toxicol Sci* (In Press) (2005).
- [79]. Sui L, Gilbert ME, Pre- and postnatal propylthiouracil-induced hypothyroidism impairs synaptic transmission and plasticity in area CA1 of the neonatal rat hippocampus, *Endocrinology* 144(9) (2003) 4195–4203. [PubMed: 12933695]
- [80]. Sharlin DS, Tighe D, Gilbert ME, Zoeller RT, The Balance between Oligodendrocyte and Astrocyte Production in Major White Matter Tracts Is Linearly Related to Serum Total Thyroxine, *Endocrinology* 149(5) (2008) 2527–2536. [PubMed: 18276755]
- [81]. Howdeshell KL, A model of the development of the brain as a construct of the thyroid system, *Environ. Health Perspect* 110 (2002) 337–348.
- [82]. Brucker-Davis F, Effects of environmental synthetic chemicals on thyroid function, *Thyroid* 8(9) (1998) 827–856. [PubMed: 9777756]
- [83]. Crofton KM, Thyroid disrupting chemicals: mechanisms and mixtures, *Int J Androl* 31(2) (2008) 209–223. [PubMed: 18217984]
- [84]. American Thyroid Association, *Thyroid Function Tests*, 2010.
- [85]. Burman KD, Commentary: Discordant measurements of serum triiodothyronine (T₄), thyroxine (T₃) and thyroid-stimulating hormone (TSH), *Clinical chemistry* 54(7) (2008) 1246. [PubMed: 18593966]
- [86]. Targher G, Montagnana M, Salvagno G, Moghetti P, Zoppini G, Muggeo M, Lippi G, Association between serum TSH, free T₄ and serum liver enzyme activities in a large cohort of unselected outpatients, *Clin Endocrinol (Oxf)* 68(3) (2008) 481–484. [PubMed: 17941901]
- [87]. Klaassen CD, Hood AM, Effects of microsomal enzyme inducers on thyroid follicular cell proliferation and thyroid hormone metabolism, *Toxicol Pathol* 29(1) (2001) 34–40. [PubMed: 11215682]
- [88]. Hood A, Hashmi R, Klaassen CD, Effects of microsomal enzyme inducers on thyroid-follicular cell proliferation, hyperplasia, and hypertrophy, *Toxicol Appl Pharmacol* 160(2) (1999) 163–170. [PubMed: 10527915]
- [89]. Caldwell KL, Jones R, Hollowell JG, Urinary iodine concentration: United States National Health And Nutrition Examination Survey 2001-2002, *Thyroid* 15(7) (2005) 692–699. [PubMed: 16053386]
- [90]. Aoki Y, Belin RM, Clickner R, Jeffries R, Phillips L, Mahaffey KR, Serum TSH and Total T₄ in the United States Population and Their Association With Participant Characteristics: National Health and Nutrition Examination Survey (NHANES 1999-2002), *Thyroid* 17(12) (2007) 1211–1223. [PubMed: 18177256]
- [91]. Blount BC, Silva MJ, Caudill SP, Needham LL, Pirkle JL, Sampson EJ, Lucier GW, Jackson RJ, Brock JW, Levels of seven urinary phthalate metabolites in a human reference population, *Environ Health Perspect* 108(10) (2000) 979–982.
- [92]. Gray TJB, Butterworth KR, Gaunt IF, Grasso P, Gangolli SD, Short-term toxicity study of di-(2-ethylhexyl) phthalate in rats, *Food and Cosmetics Toxicology* 15(5) (1977) 389–399. [PubMed: 598790]
- [93]. Wine RN, Li L-H, Barnes LH, Gulati DK, Chapin RE, Reproductive Toxicity of Di-n-Butylphthalate in a Continuous Breeding Protocol in Sprague-Dawley Rats, *Environ Health Perspect* 105(1) (1997) 102–107. [PubMed: 9074889]

- [94]. Mylchreest E, Cattley RC, Foster PM, Male reproductive tract malformations in rats following gestational and lactational exposure to Di(n-butyl) phthalate: an antiandrogenic mechanism?, *Toxicol Sci* 43(1) (1998) 47–60. [PubMed: 9629619]
- [95]. Mylchreest E, Foster PM, DBP exerts its antiandrogenic activity by indirectly interfering with androgen signaling pathways, *Toxicol Appl Pharmacol* 168(2) (2000) 174–175.
- [96]. Mylchreest E, Sar M, Cattley RC, Foster PM, Disruption of androgen-regulated male reproductive development by di(n-butyl) phthalate during late gestation in rats is different from flutamide, *Toxicol Appl Pharmacol* 156(2) (1999) 81–95. [PubMed: 10198273]
- [97]. Mylchreest E, Sar M, Wallace DG, Foster PM, Fetal testosterone insufficiency and abnormal proliferation of Leydig cells and gonocytes in rats exposed to di(n-butyl) phthalate, *Reprod. Toxicol* 16 (2002) 19–28. [PubMed: 11934529]
- [98]. Mylchreest E, Wallace DG, Cattley RC, Foster PM, Dose-dependent alterations in androgen-regulated male reproductive development in rats exposed to Di(n-butyl) phthalate during late gestation, *Toxicol Sci* 55(1) (2000) 143–151. [PubMed: 10788569]
- [99]. National Research Council, *Phthalates and Cumulative Risk Assessment: The Task Ahead*, National Academies Press, Washington, D.C., 2008.
- [100]. Barlow NJ, Phillips SL, Wallace DG, Sar M, Gaido KW, Foster PM, Quantitative changes in gene expression in fetal rat testes following exposure to di(n-butyl) phthalate, *Toxicol Sci* 73(2) (2003) 431–441. [PubMed: 12700402]
- [101]. Gray LE Jr., Ostby J, Furr J, Price M, Veeramachaneni DN, Parks L, Perinatal exposure to the phthalates DEHP, BBP, and DINP, but not DEP, DMP, or DOTP, alters sexual differentiation of the male rat, *Toxicol Sci* 58(2) (2000) 350–365. [PubMed: 11099647]
- [102]. Liu K, Lehmann KP, Sar M, Young SS, Gaido KW, Gene expression profiling following in utero exposure to phthalate esters reveals new gene targets in the etiology of testicular dysgenesis, *Biol Reprod* 73(1) (2005) 180–192. [PubMed: 15728792]
- [103]. Toppari J, Virtanen H, Skakkebaek N, Environmental contaminants and the reproductive and fertility effects in the male: Possible role of fetal exposure, in: Woodruff TJ, Janssen SJ, Guillelte LJ Jr., Giudice LC (Eds.), *Environmental Impacts on Reproductive Health and Fertility*, Cambridge Press, Cambridge, UK, 2010.
- [104]. Morris RJ, Phthalic acid in the deep sea jellyfish Atolla, *Nature* 227(5264) (1970) 1264. [PubMed: 4393820]
- [105]. Mayer FL, Stalling DL, Johnson JL, PHTHALATE ESTERS AS ENVIRONMENTAL CONTAMINANTS, *Nature* 238(5364) (1972) 411–413. [PubMed: 4626399]
- [106]. Taborsky RG, Isolation studies on a lipoidal portion of the bovine pineal gland, *Journal of Agricultural and Food Chemistry* 15(6) (1967) 1073–1076.
- [107]. Hauser R, Meeker JD, Duty S, Silva MJ, Calafat AM, Altered semen quality in relation to urinary concentrations of phthalate monoester and oxidative metabolites, *Epidemiology* 17(6) (2006) 682–691. [PubMed: 17003688]
- [108]. Jonsson BA, Richthoff J, Rylander L, Giwercman A, Hagmar L, Urinary phthalate metabolites and biomarkers of reproductive function in young men, *Epidemiology* 16(4) (2005) 487–493. [PubMed: 15951666]
- [109]. Swan SH, Main KM, Liu F, Stewart SL, Kruse RL, Calafat AM, Mao CS, Redmon JB, TERNAND CL, Sullivan S, Teague JL, Decrease in anogenital distance among male infants with prenatal phthalate exposure, *Environ Health Perspect* 113(8) (2005) 1056–1061. [PubMed: 16079079]
- [110]. Sathyanarayana S, Beard L, Zhou C, Grady R, Measurement and correlates of ano-genital distance in healthy, newborn infants, *International Journal of Andrology* 33(2) 317–323. [PubMed: 20132349]
- [111]. Thankamony A, Ong KK, Dunger DB, Acerini CL, Hughes IA, Anogenital Distance from Birth to 2 Years: A Population Study, *Environ Health Perspect* 117(11) (2009) 1786–1790. [PubMed: 20049133]
- [112]. Main KM, Mortensen GK, Kaleva MM, Boisen KA, Damgaard IN, Chellakooty M, Schmidt IM, Suomi AM, Virtanen HE, Petersen DV, Andersson AM, Toppari J, Skakkebaek NE, Human breast milk contamination with phthalates and alterations of endogenous reproductive hormones

- in infants three months of age, *Environ Health Perspect* 114(2) (2006) 270–276. [PubMed: 16451866]
- [113]. Christiansen S, Scholze M, Axelstad M, Boberg J, Kortenkamp A, Hass U, Combined exposure to anti-androgens causes markedly increased frequencies of hypospadias in the rat, *Int J Androl* (2008 (epub)).
- [114]. Swan SH, Environmental phthalate exposure in relation to reproductive outcomes and other health endpoints in humans, *Environ Res* 108(2) (2008) 177–184. [PubMed: 18949837]
- [115]. Bray F, Richiardi L, Ekblom A, Pukkala E, Cuninkova M, Moller H, Trends in testicular cancer incidence and mortality in 22 European countries: continuing increases in incidence and declines in mortality, *Int J Cancer* 118(12) (2006) 3099–3111. [PubMed: 16395710]
- [116]. Shah MN, Devesa SS, Zhu K, McGlynn KA, Trends in testicular germ cell tumours by ethnic group in the United States, *Int J Androl* 30(4) (2007) 206–213; discussion 213–204. [PubMed: 17708751]
- [117]. U.S. Environmental Protection Agency, American's Children and the Environment, Washington, DC, 2008.
- [118]. Centers for Disease Control and Prevention, Prevalence of autism spectrum disorders - Autism and Developmental Disabilities Monitoring Network, United States, 2006, *MMWR Surveill Summ* 58(10) (2009) 1–20.
- [119]. Pastor PN, Reuben CA, Diagnosed attention deficit hyperactivity disorder and learning disability: United States, 2004–2006, *Vital Health Stat* 10(237) (2008) 1–14.
- [120]. Paulozzi LJ, International trends in rates of hypospadias and cryptorchidism, *Environ Health Perspect* 107(4) (1999) 297–302. [PubMed: 10090709]
- [121]. Vu LT, Nobuhara KK, Laurent C, Shaw GM, Increasing prevalence of gastroschisis: population-based study in California, *J Pediatr* 152(6) (2008) 807–811. [PubMed: 18492521]
- [122]. Harris KB, Pass KA, Increase in congenital hypothyroidism in New York State and in the United States, *Mol Genet Metab* 91(3) (2007) 268–277. [PubMed: 17512233]
- [123]. Swan SH, Hertz-Picciotto I, Reasons for infecundity, *Fam Plann Perspect* 31(3) (1999) 156–157. [PubMed: 10379434]
- [124]. Brett K, Fecundity in 2002 NSFG women 15–24 years of age (personal communication), National Center for Health Statistics, Hyattsville, MD, 2008.
- [125]. Davidoff MJ, Dias T, Damus K, Russell R, Bettgowda VR, Dolan S, Schwarz RH, Green NS, Petrini J, Changes in the gestational age distribution among U.S. singleton births: impact on rates of late preterm birth, 1992 to 2002, *Semin Perinatol* 30(1)(2006) 8–15. [PubMed: 16549207]
- [126]. Dahlstrom BL, Engh ME, Bukholm G, Oian P, Changes in the prevalence of pre-eclampsia in Akershus County and the rest of Norway during the past 35 years, *Acta Obstet Gynecol Scand* 85(8) (2006) 916–921. [PubMed: 16862468]
- [127]. Getahun D, Nath C, Ananth CV, Chavez MR, Smulian JC, Gestational diabetes in the United States: temporal trends 1989 through 2004, *Am J Obstet Gynecol* 198(5) (2008) 525 e521–525. [PubMed: 18279822]
- [128]. Akslaede L, Sorensen K, Petersen JH, Skakkebaek NE, Juul A, Recent decline in age at breast development: the Copenhagen Puberty Study, *Pediatrics* 123(5) (2009) e932–939. [PubMed: 19403485]
- [129]. Jorgensen N, Askund C, Carlsen E, Skakkebaek NE, Coordinated European investigations of semen quality: results from studies of Scandinavian young men is a matter of concern, *Int J Androl* 29(1) (2006) 54–61; discussion 105–108. [PubMed: 16466524]
- [130]. Carlsen E, Giwercman A, Keiding N, Skakkebaek NE, Evidence for decreasing quality of semen during past 50 years, *Bmj* 305(6854) (1992) 609–613. [PubMed: 1393072]
- [131]. Andersson A, Jensen TK, Petersen JH, Joergensen T, Borch-Johnsen K, Skakkebaek NE, Trends in Leydig cell function in Danish men, *Human Reproduction* 20(suppl. 1) (2005) i26.
- [132]. Perheentupa A, Laatikainen T, Vierula M, Skakkebaek NE, Andersson AM, Toppari T, Clear Birth Cohort Effect in Serum Testosterone and SHBG Levels in Finnish Men, *Endocrine Society Meeting* 2006, 2006.
- [133]. Woodruff T, Zota A, Schwartz J, Environmental chemicals in pregnant women in the US: NHANES 2003–2004, *Environ Health Perspect* ((in review)).

- [134]. Woodruff TJ, Janssen SJ, Guillette LJ Jr., Giudice LC, Environmental Impacts on Reproductive Health and Fertility, Cambridge University Press, New York, NY, 2010.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 1:

Examples of recent trends in select reproductive disease, disorders and function¹.

Reproductive Diseases/Disorders	Increase	Period	Location	Reference
Testicular cancer	1– 6%	1953 - 1999	Europe	[115]
Testicular cancer	60%	1973 - 2003	USA	[116]
Certain childhood cancers	20 – 24%	1976 - 2005	USA	[117]
Autism	57%	2002 - 2006	USA	[118]
Attention Deficit Hyperactivity Disorder	3% per year	1997 - 2006	USA	[119]
Birth defects:				
<i>Cryptorchidism</i>	200%	1970 - 1993	USA	[120]
<i>Gastroschisis</i>	300%	1978 - 2005	California	[121]
<i>Congenital hypothyroidism</i>	138%	1987 - 2003	New York	[122]
Reproductive Function		Time	Location	Ref.
Reported difficulty conceiving and maintaining pregnancy				
<i>All ages</i>	60% more women	1982; 2002	USA	[123, 124]
<i><25 years old</i>	200% more women	1982; 2002	USA	[123, 124]
Prematurity	2.9% shorter gestation	1992 - 2002	USA	[125]
Pre-eclampsia	19-36%	1968-2002	Norway	[126]
Gestational Diabetes	122%	1989-2004	USA	[127]
Premature puberty:				
<i>Age at onset of breast development</i>	1 – 2 years younger	1940 - 1994	USA, Denmark	[3] [128]
<i>Age at onset of menstruation</i>	2.5 – 4 months younger	1940 - 1994	USA	[3]
Sperm count	~1% decline per year	1931 - 1994	Western countries	[129, 130]
Serum testosterone	1% decline per year	1987 - 2004	Boston, USA	[131, 132]

¹Updated from Woodruff et al. [1]

Table 2:
Comparing experimental features of human epidemiology studies and experimental animal studies.

Human Epidemiologic Data	Experimental Animal Data
Exposure Issues	
Exposure chemical(s) may be unclear or difficult to estimate, individual may not remember what they were exposed to or unaware of exposure	Controlled dosage of chemical(s) is clear
Reflects real world setting of multiple exposures, but can make it challenging for interpreting effects of individual chemical	Often only exposed to individual chemical, does not account for exposure to other chemicals that may affect same adverse health outcome
Duration of exposure often unknown	Controlled duration
Exposure may cover multiple windows of susceptibility or may not include specific exposure timing for critical windows	Controlled exposure can be applied for known or suspected periods of susceptibility
Experimental Design	
Generally make use of existing data, which often cross-sectional and/or retrospective	Study design is experimental
Confounding factors may be present (diet, disease, smoking, etc)	Confounding factors are reduced through experimental design (controlled exposures and diet) and use of healthy animals
Genetic variability	Known genetics: selection of homogeneous or heterogeneous strains.
Indirect determination of outcomes (endpoints). Come from birth certificates, medical records, for example	Direct endpoints studied
Interpretation and use of results	
No need for extrapolation to human populations and usually not for high to low dose	Need to extrapolate to humans and often from high to low dose
Human studies can take many years	Studies can be performed in short periods of time, usually less than 2 years

Table 3:

Percentage of U.S. Pregnant Women with Select EDCs Measured in their Body

Chemical analyte	Percent of U.S. pregnant women with detectable levels of the analyte
PFCs (serum; µg/L) (N=76)	
PFOA	99
PFOS	99
PBDEs (serum; ng/g lipid) (N=75)	
BDE-47	99
BDE-99	87
BDE-100	99
BDE-153	100
PCBs (serum; ng/g lipid) (N=75)	
PCB-118	100
PCB-138 & 158	100
PCB-153	100
PCB-180	96
Organochlorine (OC) pesticides (serum; ng/g lipid) (N=71)	
DDT	62
DDE	100
Hexachlorobenzene	100
Environmental phenols (urine; µg/L) (N=86)	
Bisphenol-A	96
Triclosan	87
Benzophenone-3	100
Phthalates (urine; µg/L) (N=91)	
Mono-n-butyl phthalate (MnBP)	99
Mono-ethyl phthalate (MEP)	100
Mono-benzyl phthalate (MBzP)	100
Mono-isobutyl phthalate (MiBP)	99
Perchlorate (urine; µg/L) (N=89)	
Perchlorate	100

* Based on analysis of representative sample of U.S. population by the U.S. Centers for Disease Control and Prevention, National Health and Nutrition Examination Survey 2003-2004 adapted from Woodruff et al. [133].

Table 4:

Classes, mechanisms of action, and effects of thyroid disrupting chemicals on thyroid hormone homeostasis *

Class	Mechanism	Effects on THs	Chemicals
Iodine Transport	Competition/block of NIS	Decreased thyroidal synthesis of T3 & T4	Perchlorate, chlorate, bromate, Nitrates, thiocyanate
Synthesis Inhibitors	Inhibition of TPO	Decreased thyroidal synthesis of T3 & T4	Methimazole, propylthiouraea, amitrole, mancozeb, soy isoflavones, benzophenone 2, 1-methyl-3-propyl-imidazole-2-thione
Transport Disruption	Altered binding to serum transport proteins	unknown	Hydroxyl-PCBs, EMD 49209; pentachlorophenol
Enhanced Hepatic Catabolism	Upregulation of glucuronylsyltransferases or sulfotransferases (via CAR/PRX or AhR)	Increased biliary elimination of T3 & T4	Acetochlor, phenobarbital, 3-methylcolanthrene, PCBs, 1-methyl-3-propyl-imidazole-2-thione
Enhanced Cellular Transport	Upregulation of OATPs or MCT transporters via CAR/PRX or AhR	Increased biliary elimination of T3 & T4	TCPOBOP, pregnenolone-16 α -carbonitrile, TCDD, rifampicin, phenobarbital, oltipraz
Sulfo-transferases	Inhibition of sulfotransferases (SULTs)	Decrease sulfation of THs	Hydroxylated PCBs, triclosan, pentachlorophenol
Deiodinases	Inhibition or upregulation of deiodinases	Decreased peripheral synthesis of T3	FD&C Red Dye #3, propylthiouracil, PCBs, octyl-methoxycinnamate
TR agonists & antagonists	Direct or indirect alterations in TR-TRE binding	Altered activation of TH dependent gene transcription	TetrabromoBisphenol A, Bisphenol A, HydroxyPCBs

* Reprinted from Crofton 2005 [43]

Table 5:
Features of the human population that can influence risks from exposure to environmental contaminants

Feature	Definition	Example	Implications
Chemical Background	Concurrent or preexisting body burdens of environmental chemicals	Over 90% of the U.S. population has measurable levels of plasticizers, pesticides, flame retardants and perfluorinated chemicals	Exposure to an individual chemical in addition to other exposures can increase risk of subsequent disease
Biological Background	Health status, as influenced by age, pre-existing disease, genetics, and other intrinsic biological factors.	About 10% of adults are deficient in thyroid hormone, low thyroid hormone during pregnancy can cause neurodevelopment harm in children.	Chemical exposure has greater effect on people with pre-existing disease, genetic predisposition, etc.
Population Variability and Defining Normal	An individual may have smaller range of normal physiological function than would be observed over the whole population	Individual variability in normal thyroid hormone levels is smaller than the range of normal in the population	Chemical exposures which perturb physiological systems within the "normal" population range may still have effects at the individual level.
Small individual effects versus population effects	Chemical exposure may produce a small increase in risk or effect at the level of the individual but result in large shifts in effect at the population level.	Observed prenatal phthalate levels slightly decreases the anogenital distance in malebabies	US population exposure to phthalates is ubiquitous, so small changes can shift the population distribution of the measurement, resulting in increased effects for a segment of the population.
Periods of Susceptibility	Exposure during vulnerable periods of development can pose a risk of irreversible effects, both in the short and long term, and diminished capacity for recovery.	Exposure to solvents during pregnancy can result in miscarriage or preterm delivery or can cause permanent neurological damage to the fetus. (eg. Fetal alcohol syndrome).	Exposures during susceptible periods can increase risk of permanent adverse effects that may manifest early or not until later in life.

Reprinted from Woodruff and Janssen (permission required) [134]