Published in final edited form as:

Birth Defects Res C Embryo Today. 2013 March ; 99(1): 45–49. doi:10.1002/bdrc.21029.

Risk Communication and Decision Tools for Children's Health Protection

Patrice Sutton¹ and Tracey J. Woodruff^{*,1}

¹Program on Reproductive Health and the Environment, Department of Obstetrics, Gynecology, and Reproductive Sciences, University of California, San Francisco, California, USA

Abstract

Scientific discovery linking the environment to beneficial and adverse health children's health outcomes is rapidly expanding leading scientists and health professionals to call for timely action to prevent harm and secure benefits. A robust method to synthesize what is known about the environmental drivers of health is a foundational step to making the science actionable by individuals and decision-makers. To meet this need, a methodology called the Navigation Guide was crafted by a collaboration of 22 clinical and environmental health scientists. The Navigation Guide proceeds from methods of research synthesis used in clinical settings but accounts for differences between environmental and clinical health sciences related to the evidence-base and decision-context. The methodology can be used to develop evidence profiles that provide simple, transparent summaries, such as practice guidelines or other evidence-based recommendations for prevention. Establishing proof-of-concept of the method is underway. Development of the Navigation Guide is extremely timely as it coincides with growing recognition of the need for updated methods in risk assessment. The costs in 2008 to the U.S. healthcare system for treatment of childhood illnesses linked to toxic environmental exposures is conservatively estimated to be over \$76 billion, and it is anticipated that U.S. healthcare policy decisions will increasingly rely on systematic reviews of the evidence. The Navigation Guide is poised to provide a methodological bridge to link healthcare decision-making to efforts to reduce toxic environmental exposures. The institutionalization of the Navigation Guide would provide a concrete mechanism for linking science to action to protect children's health.

Introduction

Scientific discovery linking the environment to beneficial and adverse health children's health outcomes is rapidly expanding. Studies find that individually and together, the social, built and nutritional environment, and physical and chemical agents are key drivers of child health.(Burke and Miller, 2011; Cheadle et al., 2010; Guilloteau et al., 2009; Kordas, 2010; Marmot and Bell, 2009; Morello-Frosch and Shenassa, 2006; Morello-Frosch et al., 2011; Ren et al., 2010; Schneider et al., 2001; Weiss and Bellinger, 2006; Williams and Sternthal, 2010; Wright, 2010; Wright et al., 2010; Wright, 2008) Science over the last decade spotlights the importance of environmental exposures that occur during periods of

*Correspondence to: tracey.woodruff@ucsf.edu.

development, i.e., in utero, and during infancy, childhood and adolescence. This is because these periods are windows of increased vulnerability to environmental contributors to health. (Crain et al., 2008; Grandjean et al., 2008; Newbold and Heindel, 2010; Woodruff et al., 2010) Moreover, the consequences of developmental exposures can extend across the lifetime of individuals and potentially across generations. (Olden et al., 2011) In recognition of the science linking environmental exposures during periods of development to a myriad of health outcomes that can manifest from infancy to old age, leading scientists and health professionals have called for timely action to prevent harm and secure benefits. (Diamanti-Kandarakis et al., 2009; Grandjean et al., 2008; Reuben and President's Cancer Panel, 2010; UCSF Program on Reproductive Health and the Environment, 2012; Woodruff et al., 2008)

Communication and decision tools for preventing exposure to environmental chemicals need to encompass not only individual action but also action at policy levels (Lubick, 2011). While action at the individual level can potentially reduce exposure to some toxic chemicals (Ji et al., 2010; Lu et al., 2006; Rudel et al., 2011), and informed consumer-purchasing patterns can send a signal to the marketplace to help drive societal change (Bailin et al., 2008; Layton, 2011), individuals alone can do little to impact many exposures, such as air and water pollution. Further, decisions on the individual level about avoiding toxic exposures are also often affected by external factors that limit making healthier choices (Adler and Stewart, 2009).

Moreover, a formidable barrier to prevention is that in their efforts to take action on the science, clinicians, parents and policy makers are confronted with a plethora of information, some contradictory, about environmental and chemical risks to children's health. They want and need evidence-based information to inform what they do to prevent exposures and promote health. Better decision tools are needed to evaluate the soundness of available information, synthesize it and translate it into public-friendly prevention strategies for health professionals, parents and decision-makers.

Taking Action to Prevent Harm and Secure Benefits: The Role of Research Synthesis

Clinicians, health impacted populations, advocates and policy makers generally lack the capacity to efficiently, systematically and transparently incorporate the meaning of the science into their decision-making. The science linking the environment to health is voluminous, of variable quality, complex, and largely unfamiliar to decision-makers in clinical and policy arenas. Keeping abreast of its meaning can be overwhelming. An unrelenting deluge of bits and pieces of information can serve to mask rather than amplify early warning signals and undermine our capacity to act wisely. A robust method to synthesize what is known about the environmental drivers of health is a necessary foundational step to making the science actionable by individuals and decision-makers.

Methods in Research Synthesis

Common features of the many and varied methods currently used by environmental health scientists are that they are expert opinion-based and do not provide weight of evidence

summaries for non-cancer effects. Historically, similar methods prevailed in the clinical field which largely relied on a system of expert reviews on which to base treatment decisions. (Rennie and Chalmers, 2009) However, starting in the 1970s, the role of expert reviews began to be questioned, and systematic approaches that harness expertise to a rigorous, transparent, and explicit methodology to evaluate a clearly formulated question were advanced. Empirical evidence in clinical health sciences demonstrates the superiority of systematic reviews for patient outcomes. (Rennie and Chalmers, 2009) A landmark paper by Antman and colleagues compared expert opinion-based recommendations for treatment of myocardial infarction published in scientific reviews and clinical textbooks to statistical analyses of the combined results of randomized controlled trials. (Antman et al., 1992) This research documented the lack of timely incorporation of experimental evidence into expert-based recommendations with some reviewers not even mentioning effective therapies, and others recommending therapies proven to be ineffective or even dangerous.

Methods to synthesize the science into evidence-based decisions have been developed and validated in clinical arenas. (GRADE Working Group, 2011; Montori and Guyatt, 2008) But due to differences between environmental and clinical health sciences related to the evidence-base and decision-context these methods are not seamlessly applicable to environmental exposures.

In the clinical setting, *in vivo*, *in vitro*, and human experimental evidence combined with an analysis of risks and benefits inform human exposure decisions prior to the entry of substances into the marketplace (Figure 1). The “gold standard” of evidence for clinical risk benefit decisions is a well-conducted randomized control trial. Systematic reviews in the clinical sciences proceed from this evidence and context.

In stark contrast, population exposure to exogenous substances in the environment typically occurs before regulatory scrutiny of a compound and in the absence of risk–benefit analysis, because of the current regulatory structure for governing manufactured chemicals. Ethical considerations virtually preclude experimental human data from the environmental health evidence stream, so we must rely on *in vitro* and *in vivo* studies for early warnings of adverse effects and on human observational studies to assess the nature and extent of the damage. (Zapponi et al., 2008)

To bridge this gap between the evidence streams and decision contexts in clinical and environmental health sciences, the University of California, San Francisco Program on Reproductive Health and the Environment undertook an interdisciplinary collaboration to craft an evidence-based medicine methodology to evaluate environmental contaminants and their potential effects on reproductive and developmental health. The result is the Navigation Guide, the product of a yearlong collaboration of 22 clinical and environmental health scientists and/or practitioners, from governmental and non-governmental organizations in the U.S. and Europe. (Woodruff et al., 2011) The Navigation Guide proceeds from methods in the clinical sciences but accounts for the differences in evidence and decision context described above.

Overview of the Navigation Guide

The Navigation Guide methodology involves 4 steps (Figure 2). Applying the methodology requires the expertise of toxicologists, epidemiologists, statisticians, industrial hygienists, clinicians and other potential scientific expertise. The method also provides for the incorporation of the expertise of non-scientists, including health-impacted populations and their advocates, for example, in framing a meaningful study question and incorporating local knowledge, values and preferences into recommendations for prevention.

While such expertise is incorporated into every step, it is not an “expert-opinion” methodology. What is a fundamental shift from existing methods of expert review in environmental health science is that each of the 4 steps is conducted in a thorough, consistent and transparent manner, every “judgment” is documented, and that the science is clearly delineated from values, preferences, costs and/or other considerations. In short, the rationale for a decision is traceable, reproducible, comprehensible and therefore testable over time.

Government agencies can use the Navigation Guide methodology to craft evidence-based statements regarding the relationship between an environmental exposure and health (Steps 1–3). The first case study of steps 1–3 of the Navigation Guide methodology was completed in 2013 by a collaboration of scientists at USEPA, UCSF, and Johns Hopkins. The results established proof-of-concept for applying a systematic and transparent method of review in environmental health to link the science to timely, prevention-oriented action. The case study addressed the question, “What is the impact of developmental exposure to perfluorooctanoic acid (PFOA) on fetal growth?”

Government agencies called on to make risk management decisions can also apply Step 4 of the Navigation Guide to grade the strength of recommendations for prevention. Decision-tools need to distinguish the “strength of the evidence” from the “strength of the recommendation” to make transparent the values and preferences that underlie risk management decisions. This distinction also allows for incorporating the recommendations made by the National Academy of Sciences (NAS) in *Science and Decisions* to identify and protect against chemicals that can harm human health before entering into a long process to establish numerical levels of risk and to cull those decisions that are not sensitive to the resolution of uncertainty from the risk assessment process, i.e., decisions for which additional information would have little or no value added to support the decision. (National Research Council, 2008) Professional societies, healthcare organizations, and other potential guideline developers working with toxicologists can use the Navigation Guide to craft consistent and timely recommendations to improve patient, and ultimately population, health outcomes.

Conclusion

Providing clinicians, parents and policy-makers with evidence-based information about the environmental drivers of children’s health is key to prevention. The *Navigation Guide* is a method to develop evidence profiles that provide simple, transparent summaries, such as

practice guidelines or other evidence-based recommendations for prevention. Development of the Navigation Guide methodology is extremely timely as it coincides with growing recognition of the need for updated methods in risk assessment (National Research Council, 2008; National Research Council (U.S.) Committee on Improving Risk Analysis Approaches Used by the U.S. EPA et al., 2009) and related calls for government agencies such as the USEPA to incorporate systematic approaches into decision-making (2011; Charnley, 2011; Hopkinson, 2011; Lovell, 2011; Vitter and Inhofe, June 30, 2011). Beginning in 2012, systematic methods of research synthesis in environmental health science have been advanced by the National Toxicology Program Office of Health Assessment and Translation (OHAT) in its development of literature-based evaluations to reach conclusions about potential human health hazards and to examine the state of the science. The output from an OHAT evaluation can vary including, but not limited to: NTP Monograph, state-of-the-science workshop report, or peer-reviewed journal publication. (National Toxicology Program, 2013)

Having a method that understands the opportunities and challenges of applying current clinical approaches to environmental health is also critical. While not yet on the radar of healthcare decision-makers, the costs in 2008 to the U.S. healthcare system for treatment of childhood illnesses linked to toxic environmental exposures is conservatively estimated to be over \$76 billion (Trasande and Liu, 2011). At the same time, it is anticipated that U.S. healthcare policy decisions will increasingly rely on systematic reviews of the evidence; the recent healthcare reform legislation allocated \$1.1 billion dollars for comparative effectiveness research (Department of Health and Human Services, 2009). The Navigation Guide is poised to provide a methodological bridge to link healthcare decision-making to efforts to reduce toxic environmental exposures.

Realization of this goal could transform the science by making it actionable in a timely manner. Conversely, it would also shed light on the consequences of our inaction that now finds refuge under a mounting pile of scientific publications. In 1961, Sir Bradford Hill emphasized that “strong evidence” does not imply “crossing every ‘t’, and swords with every critic, before we act.” (Hill, 1965). He proposed differential standards of evidence for different actions, a recommendation echoed by the NAS a half-century later in *Science and Decisions*. The institutionalization of the Navigation Guide would provide a concrete mechanism for linking science to action to protect children’s health.

References

- 2011 GOP Senators Broaden EPA Science Attacks To Criticize NAAQS Process. Risk Policy Report: InsideEPA.
- Adler NE, Stewart J. 2009 Reducing obesity: motivating action while not blaming the victim. *Milbank Q* 87(1):49–70. [PubMed: 19298415]
- Antman EM, Lau J, Kupelnick B, Mosteller F, Chalmers TC. 1992 A comparison of results of meta-analyses of randomized control trials and recommendations of clinical experts. Treatments for myocardial infarction. *JAMA* 268(2):240–248. [PubMed: 1535110]
- Bailin PS, Byrne M, Lewis S, Liroff R. 2008 Public Awareness Drives Market for Safer Alternatives: Bisphenol A Market Analysis Report. Falls Church, VA: Investor Environmental Health Network.
- Burke MG, Miller MD. 2011 Practical Guidelines for Evaluating Lead Exposure in Children with Mental Health Conditions. *Postgrad Med* 123(1).

- Charnley G 2011 Testimony for the Review of EPA's Integrated Risk Information System (IRIS) In: House Science Space and Technology Subcommittee on Investigations and Oversight, editor. Washington, D.C.
- Cheadle A, Samuels SE, Rauzon S, Yoshida SC, Schwartz PM, Boyle M, Beery WL, Craypo L, Solomon L. 2010 Approaches to measuring the extent and impact of environmental change in three California community-level obesity prevention initiatives. *Am J Public Health* 100(11):2129–2136. [PubMed: 20935262]
- Crain DA, Janssen SJ, Edwards TM, Heindel J, Ho SM, Hunt P, Iguchi T, Juul A, McLachlan JA, Schwartz J, Skakkebaek N, Soto AM, Swan S, Walker C, Woodruff TK, Woodruff TJ, Giudice LC, Guillette LJ. 2008 Female reproductive disorders: the roles of endocrine-disrupting compounds and developmental timing. *Fertil Steril* 90(4):911–940. [PubMed: 18929049]
- Department of Health and Human Services. 2009 Recovery Act Allocates \$1.1 Billion for Comparative Effectiveness Research.
- Diamanti-Kandarakis E, Bourguignon JP, Giudice LC, Hauser R, Prins GS, Soto AM, Zoeller RT, Gore AC. 2009 Endocrine-disrupting chemicals: an Endocrine Society scientific statement. *Endocr Rev* 30(4):293–342. [PubMed: 19502515]
- GRADE Working Group. 2011 Frequently Asked Questions.
- Grandjean P, Bellinger D, Bergman A, Cordier S, Davey-Smith G, Eskenazi B, Gee D, Gray K, Hanson M, van den Hazel P, Heindel JJ, Heinzow B, Hertz-Picciotto I, Hu H, Huang TT, Jensen TK, Landrigan PJ, McMillen IC, Murata K, Ritz B, Schoeters G, Skakkebaek NE, Skerfving S, Weihe P. 2008 The faroes statement: human health effects of developmental exposure to chemicals in our environment. *Basic & clinical pharmacology & toxicology* 102(2):73–75. [PubMed: 18226057]
- Guilloteau P, Zabielski R, Hammon H, Metges C. 2009 Adverse effects of nutritional programming during prenatal and early postnatal life, some aspects of regulation and potential prevention and treatments. *Journal of physiology and pharmacology : an official journal of the Polish Physiological Society* 60:17–35. [PubMed: 19996479]
- Hill AB. 1965 The Environment and Disease: Association or Causation? *Proc R Soc Med* 58:295–300. [PubMed: 14283879]
- Hopkinson J 2011 Alliance Plans Panel To Address Key Scientific Issues In Risk Assessment Risk Policy Report.
- Ji K, Lim Kho Y, Park Y, Choi K. 2010 Influence of a five-day vegetarian diet on urinary levels of antibiotics and phthalate metabolites: A pilot study with “Temple Stay” participants. *Environ Res* 110(4):375–382. [PubMed: 20227070]
- Kordas K 2010 Iron, Lead, and Children's Behavior and Cognition. *Annu Rev Nutr* 30:123–148. [PubMed: 20433349]
- Layton L 2011 Wal-Mart bypasses federal regulators to ban controversial flame retardant. *The Washington Post*.
- Lovell A 2011 GAO Reviewing EPA Changes To Chemical Risk Assessment Process. *InsideEPA*.
- Lu C, Toepel K, Irish R, Fenske RA, Barr DB, Bravo R. 2006 Organic diets significantly lower children's dietary exposure to organophosphorus pesticides. *Environ Health Perspect* 114(2):260. [PubMed: 16451864]
- Lubick N 2011 Advising Parents in the Face of Scientific Uncertainty: An Environmental Health Dilemma. *Environ Health Perspect* 119(10).
- Marmot MG, Bell R. 2009 Action on health disparities in the United States. *JAMA: The Journal of the American Medical Association* 301(11):1169. [PubMed: 19293419]
- Montori VM, Guyatt GH. 2008 Progress in evidence-based medicine. *JAMA* 300(15):1814–1816. [PubMed: 18854545]
- Morello-Frosch R, Shenassa ED. 2006 The environmental “riskscape” and social inequality: implications for explaining maternal and child health disparities. *Environ Health Perspect* 114(8): 1150–1153. [PubMed: 16882517]
- Morello-Frosch R, Zuk M, Jerrett M, Shamasunder B, Kyle AD. 2011 Understanding the cumulative impacts of inequalities in environmental health: implications for policy. *Health Aff (Millwood)* 30(5):879–887. [PubMed: 21555471]

- National Research Council. 2008 Phthalates and cumulative risk assessment: the task ahead. Washington, D.C.: National Academies Press xix, 188 p. p.
- National Research Council (U.S.) Committee on Improving Risk Analysis Approaches Used by the U.S. EPA, National Research Council (U.S.) Board on Environmental Studies and Toxicology, National Research Council (U.S.) Division on Earth and Life Studies. 2009 Science and decisions: advancing risk assessment. Washington, D.C.: National Academies Press xviii, 403 p. p.
- National Toxicology Program. 2013 Office of Health Assessment and Translation (OHAT) Evaluation Process In: Program NT, editor. Online.
- Newbold R, Heindel J. 2010 Developmental exposures and implications for early and latent disease In: Woodruff TJ JS GJL, Giudice LC, editor. Environmental Impacts on Reproductive Health and Fertility. Cambridge, UK: Cambridge University Press p 92–102.
- Olden K, Freudenberg N, Dowd J, Shields AE. 2011 Discovering how environmental exposures alter genes could lead to new treatments for chronic illnesses. *Health Aff (Millwood)* 30(5):833–841. [PubMed: 21555469]
- Ren C, Park SK, Vokonas PS, Sparrow D, Wilker E, Baccarelli A, Suh HH, Tucker KL, Wright RO, Schwartz J. 2010 Air pollution and homocysteine: more evidence that oxidative stress-related genes modify effects of particulate air pollution. *Epidemiology* 21(2):198–206. [PubMed: 20110814]
- Rennie D, Chalmers I. 2009 Assessing authority. *JAMA* 301(17):1819–1821. [PubMed: 19417202]
- Reuben SH, President's Cancer Panel 2010 Reducing Environmental Cancer Risk: What We Can Do Now. Bethesda: National Cancer Institute. Report nr 1437934218.
- Rudel RA, Gray JM, Engel CL, Rawsthorne TW, Dodson RE, Ackerman JM, Rizzo J, Nudelman JL, Brody JG. 2011 Food Packaging and Bisphenol A and Bis(2-Ethylhexyl) Phthalate Exposure: Findings from a Dietary Intervention. *Environ Health Perspect*.
- Schneider JS, Lee M, Anderson D, Zuck L, Lidsky T. 2001 Enriched environment during development is protective against lead-induced neurotoxicity. *Brain Res* 896(1–2):48–55. [PubMed: 11277972]
- Trasande L, Liu Y. 2011 Reducing the staggering costs of environmental disease in children, estimated at \$76.6 billion in 2008. *Health Aff (Millwood)* 30(5):863–870. [PubMed: 21543421]
- UCSF Program on Reproductive Health and the Environment. 2012 Professional Statements Database. San Francisco.
- Vitter D, Inhofe J. 6 30, 2011 Letter to Jackson Lisa P, Administrator USEPA. WA, DC.
- Weiss B, Bellinger DC. 2006 Social ecology of children's vulnerability to environmental pollutants. *Environ Health Perspect* 114(10):1479–1485. [PubMed: 17035129]
- Williams DR, Sternthal M. 2010 Understanding Racial-ethnic Disparities in Health. *J Health Soc Behav* 51(1 suppl):S15–S27. [PubMed: 20943580]
- Woodruff TJ, Carlson A, Schwartz JM, Giudice LC. 2008 Proceedings of the summit on environmental challenges to reproductive health and fertility: executive summary. *Fertil Steril* 89(2 Suppl):e1–e20. [PubMed: 18308046]
- Woodruff TJ, Janssen SJ, Guillette LJ Jr, Giudice LC. 2010 Environmental impacts on reproductive health and fertility. New York, NY: Cambridge University Press.
- Woodruff TJ, Sutton P, The Navigation Guide Work Group. 2011 An Evidence-Based Medicine Methodology To Bridge The Gap Between Clinical And Environmental Health Sciences. *Health Aff (Millwood)* 30(5):931–937. [PubMed: 21555477]
- Wright RJ. 2010 Perinatal stress and early life programming of lung structure and function. *Biol Psychol*.
- Wright RJ, Visness CM, Calatroni A, Grayson MH, Gold DR, Sandel MT, Lee-Parritz A, Wood RA, Kattan M, Bloomberg GR, Burger M, Togias A, Witter FR, Sperling RS, Sadovsky Y, Gern JE. 2010 Prenatal maternal stress and cord blood innate and adaptive cytokine responses in an inner-city cohort. *Am J Respir Crit Care Med* 182(1):25–33. [PubMed: 20194818]
- Wright RO. 2008 Neurotoxicology: What can context teach us? *The Journal of pediatrics* 152(2):155. [PubMed: 18206679]
- Zapponi GA, Marcello I, Carere A. 2008 Prevention, ethics and science: lessons from Lorenzo Tomatis. *Ann Ist Super Sanita* 44(1):8–12. [PubMed: 18469370]

Figure 1.
Streams of Evidence for Chemical Toxicity Assessment in Clinical and Environmental Health Sciences

Figure 2.
Overview of the Navigation Guide Methodology
Source: Woodruff TJ, Sutton P. The navigation guide work group. 2011.