CORRESPONDENCE

Acute Confusional States in Hospital

by PD Dr. med. Norbert Zoremba, Ph.D. and Prof. Dr. med. Mark Coburn in issue 7/2019

Take into Account Postoperative Cognitive Dysfunction

The article successfully combines the topic of hospital treatment with the risk of developing postoperative delirium (POD) (1). The authors point out that rather than being a transient disease (*“Durchgangssyndrom”*, transitory psychotic syndrome), POD is associated with increased mortality, prolonged hospitalization, and poorer outcome. The article advocates structured and consistent implementation of validated testing procedures as well as preventive and therapeutic approaches with non-pharmacological procedures. The authors conclude that adequate prevention of delirium, timely diagnosis, identification of precipitating factors, and rapid initiation of causal treatments are critical to the success of the treatment (1).

Our working group on the prevention of POD wishes to add that postoperative cognitive dysfunction (POCD) can also trigger persistent cognitive dysfunction after delirium (2). POCD can last up to two years postoperatively. POD is therefore an important risk factor for POCD. The preoperative presence of somatic diseases and subclinical delirious symptoms is an important risk factor for POD and POCD. Its prevention requires subclinical delirious symptoms and somatic diseases to be identified preoperatively, and then to implement compensatory measures. Our own investigations have shown this (3). For instance, one important somatic disease was preoperative urinary tract infection. However, prevention of delirium in the general hospital is only one aspect of adequately caring for people with cognitive impairment in the general hospital. Consistent gerontopsychiatric care for nursing home residents with dementia can reduce the risks of re-hospitalization and of delirium (4). In addition to delirium prevention in hospital, outpatient care for patients with dementia must also be strengthened as an interface to the hospital. The present work encourages this.

DOI: 10.3238/arztebl.2019.0376a

References

1. Zoremba N, Coburn M: Acute confusional states in hospital. *Dtsch Arztebl Int* 2019; 116: 101–6.
2. Kratz T, Diefenbacher A: Kognitive Akut- und Langzeitfolgen intensivmedizinischer Behandlung. *Nervenarzt* 2016; 87: 246–52.
3. Kratz T, Heinrich M, Schlauf E, Diefenbacher A: The prevention of postoperative confusion—a prospective intervention with psychogeriatric liaison on surgical wards in a general hospital. *Dtsch Arztebl Int* 2015; 112: 289–96.
4. Kirchen-Peters S, Diefenbacher A: Gerontopsychiatrische Konsiliar- und Liaisondienste – eine Antwort auf die Herausforderung Demenz? *Z Gerontol Geriatr* 2014; 47: 595–604.

Prof. Dr. med. Torsten Kratz

Prof. Dr. med. Albert Diefenbacher, MBA

Evangelisches Krankenhaus Königin Elisabeth Herzberge gGmbH,

Abteilung für Psychiatrie, Psychotherapie und Psychosomatik
Berlin, Germany
T.Kratz@keh-berlin.de

Conflict of interest statement

The authors declare that no conflict of interest exists.

In Reply:

The aim of our article was to give an overview of the current state of knowledge of delirium in the hospital. In particular, diagnosis, strategies for prevention of delirium, and delirium treatment are essential for the outcome of the patient (1). The statements in our article refer to both internal medicine patients and operative patients, regardless of patient age or pre-existing conditions, and are intended to raise awareness of the problem of delirium in the hospital (2).

We fully support the statements of Professors Kratz and Diefenbacher about postoperative delirium and postoperative cognitive deficits as well as about indications of patients with dementia. However, these patient groups are a special collective that does not reflect the full range of patients in a hospital. Our task must be to ensure targeted delirium prevention, early delirium detection, and rapid delirium treatment in an extensive manner. This is the only way to avoid delirium during hospitalization, or to treat it in a timely manner, for patients with and without cognitive impairment from all specialist disciplines, thereby avoiding a protracted or permanent cognitive impairment (3, 4).

DOI: 10.3238/arztebl.2019.0376b

References

1. Avelino-Silva TJ, Campora F, Curiati JAE, et al.: Prognostic effects of delirium motor subtypes in hospitalized older adults: a prospective cohort study. *PLoS One* 2018; 13: e0191092.
2. Zoremba N, Coburn M: Acute confusional states in hospital. *Dtsch Arztebl Int* 2019; 116: 101–6.
3. Devlin JW, Skrobik Y, Gelinas C, et al.: Clinical practice guidelines for the prevention and management of pain, agitation/sedation, delirium, immobility, and sleep disruption in adult patients in the ICU. *Crit Care Med* 2018; 46: e825–73.
4. Aldecoa C, Bettelli G, Bilotta F, et al.: European Society of Anaesthesiology evidence-based and consensus-based guideline on postoperative delirium. *Eur J Anaesthesiol* 2017; 34: 192–214.

PD Dr. med. Norbert Zoremba PhD

Klinik für Anästhesiologie, operative Intensivmedizin und Schmerztherapie
Sankt Elisabeth Hospital Gütersloh, Germany
norbert.zoremba@sankt-elisabeth-hospital.de

Prof. Dr. med. Mark Coburn

Klinik für Anästhesiologie, Universitätsklinikum RWTH Aachen, Germany

Conflict of interest statement

PD Dr. Zoremba and Prof. Coburn have received congress fee and travel cost reimbursement, as well as lecture honoraria, from Orion Pharma and MD Horizonte GmbH