CORRESPONDENCE

The Perioperative Care of Older Patients

Time for a New, Interdisciplinary Approach

by Dr. med. Cynthia Olotu, Prof. Dr. med. Arved Weimann, Prof. Dr. med. Christian Bahrs, Prof. Dr. med. Wolfgang Schwenk, Prof. Dr. med. Martin Scherer, and Prof. Dr. med. Rainer Kiefmann in issue 5/2019

Multicenter Study

Regarding this valuable presentation about the need for perioperative geriatric care, we would like to point out that timely initiation of individual preventive measures through multicomponent intervention can reduce the prevalence and incidence of delirium in hospitals (1, 2). To date, appropriately structured concepts have only rarely been used in Germany. To prevent delirium and postoperative cognitive dysfunction (POCD), the multicenter study PAWEL was launched in 2017, funded by the Innovation Fund of the Federal Joint Committee (G-BA, *Gemeinsamer Bundesausschuss*) (3, 4). For this, more than 1200 patients from seven hospitals were recruited, screened pre-admission and post-discharge, and randomly assigned to a daily multimodal intervention.

In addition to evaluating the perioperative delirium prevalence, the project focuses on developing a risk score for preoperative detection of delirium risk and for implementing a three-level, multi-professional training concept for care providers. A multicenter delirium study will be carried out for the first time in Europe, on a large German cohort, which will assess the prevalence of POCD at 2, 6, and 12 months post-operatively as well as its range of influence by non-pharmacological measures.

Special attention will be paid to the health economic evaluation (in cooperation with the University of Duisburg-Essen and the public health insurance company *Allgemeine Ortskrankenkasse* [AOK] Baden-Württemberg), in which costs of potential consequences of delirium will be recorded. The care structure analysis aims to show that cross-sectoral intervention with multi-professional training and daily prevention based on the delirium scores is cost-effective for service providers, that the improved quality of results has no additional costs, and that care needs and requirements are lower than with standard treatment. The results are expected by the end of 2020 and should provide the basis for German guidelines on the prevention and management of delirium and POCD, which is often insufficiently defined. Further, they will be entered into the quality contracts of the Institute for Quality Assurance and Transparency in Health Care (*Institut für Qualitätssicherung und Transparenz im Gesundheitswesen*; IQTIG) with the health care insurance providers, thereby improving the quality of treatment for high-risk patients. Finally, the results will make it

easier to weigh indications for elective surgery as well as any improvements in quality of life that it might bring.

DOI: 10.3238/arztebl.2019.0373a

References

1. Olotu C, Weimann A, Bahrs C, Schwenk W, Scherer M, Kiefmann R: The perioperative care of older patients—time for a new, interdisciplinary approach. *Dtsch Arztebl Int* 2019; 116: 63–9.
2. Siddiqi N, Harrison JK, Clegg A, et al.: Interventions for preventing delirium in hospitalised non-ICU patients. *Cochrane Database Syst Rev* 2016; 3: CD005563.
3. Gemeinsamer Bundesausschuss – Innovationsausschuss: PAWEL-Studie. <https://innovations-fonds.g-ba.de/projekte/versorgungsforschung/pawel-patientensicher-heit-wirtschaftlichkeit-und-lebensqualitaet-reduktion-von-delirrisiko-und-postoperativer-kognitiver-dysfunktion-pocd-nach-elektivoperationen-im-alter.16> (last accessed on 26 February 2019).
4. Sánchez A, Thomas C, Deeken F, et al.: Patient safety, cost-effectiveness, and quality of life: reduction of delirium risk and postoperative cognitive dysfunction after elective procedures in older adults—study protocol for a stepped-wedge cluster randomized trial (PAWEL Study). *Trials* 2019; 20: 71.

PD Dr. med. Christine Thomas

Klinik für Psychiatrie und Psychotherapie für Ältere
Krankenhaus Bad Cannstatt, Klinikum Stuttgart, Germany
c.thomas@klinikum-stuttgart.de

Patricia Sabbah, MA; MSc.

Klinik für Psychiatrie und Psychotherapie für Ältere
Krankenhaus Bad Cannstatt, Klinikum Stuttgart, Germany

Prof. Dr. med. Dr. phil. Michael Rapp

Professur Sozial- und Präventivmedizin, Universität Potsdam, Germany

Prof. Dr. med. Gerhard Eschweiler

Geriatrisches Zentrum Universitätsklinikum Tübingen, Germany

Conflict of interest statement

Prof. Rapp has received consultant honoraria from Lilly, Malteser Deutschland GmbH, and Project GmbH, authorship and lecture fees from Dr. Willmar Schwabe GmbH, lecture fees from the Forum für medizinische Fortbildung FormF GmbH, Akademie Heiligenfeld GmbH, Labor 28 GmbH, Alzheimer Gesellschaft Brandenburg, Wichern Krankenhaus GmbH, DKH Henriettenstiftung GmbH, the Gemeindepsychiatrischen Verbund (Community Mental Health Group), the Altenhilfe GmbH – Friederich von Bodelschwingh Klinik, the Ruppiner Kliniken GmbH, and the Klinikum Quakenbrück.

The remaining authors declare that no conflict of interest exists.

Disregard for Competence

Reading the review on the particular challenges of perioperative medicine for older patients, one could easily reach the conclusion that neither departments of geriatrics nor geriatric expertise exist in acute care hospitals in the Federal Republic of Germany (1). At no point is it noted that, across the country, day-in and day-out, qualified geriatricians and geriatric teams are

virtually everywhere in many hospitals, occupying themselves with concrete geriatric problems in the acute inpatient setting. Topics such as preoperatively restricted mobility, undernutrition, questions of early rehabilitation, and questions of delirium prophylaxis and treatment are addressed here competently. The authors very correctly state that patients can receive the optimal benefits only once all physicians and therapists know and implement the concept. The key points of geriatric work have been firmly in place for many years, especially with respect to the benefits of early rehabilitation; nonetheless, the authors themselves do not seem to know these points, even though they were in contact with the appropriate specialist society, according to the acknowledgments. In my view, the competence in geriatrics that has accumulated and been painstakingly developed over 20 years in the Federal Republic of Germany has not earned such a disregard.

DOI: 10.3238/arztebl.2019.0373b

Reference

1. Olotu C, Weimann A, Bahrs C, Schwenk W, Scherer M, Kieffmann R: The perioperative care of older patients—time for a new, interdisciplinary approach. *Dtsch Arztebl Int* 2019; 116: 63–9.

PD Dr. med. H. Burkhardt

II. Medizinische Klinik
Sektion Altersmedizin
UMM Universitätsmedizin Mannheim, Germany
geriatrie@umm.de

Conflict of interest statement

The author declares that no conflict of interest exists.

Preoperative Screening Required

The article successfully links the topics of our increasingly aging population, an increase in the older patient group in the general hospital setting, and an increased postoperative risk (1). The authors point out that needs-based care for older people requires not only interdisciplinary cooperation but also adaptation and optimization of hospital procedures. The existing perioperative treatment pathways are not yet sufficiently implemented in daily life. We are grateful that postoperative delirium (POD) has been taken up in this context. POD is a relevant risk factor for the outcome, need for care, and care home admission of dementia patients.

The article advocates POD screening until the fifth postoperative day. Our working group on the prevention of POD would like to add that preoperative delirium screening should be carried out intensively. Our own investigations have shown that, depending on the physical and cognitive status of older persons, pre-existing subclinical delirium is an important risk factor for POD and its consequences (2). About 10%–20% of those aged over 70 are already delirious at hospital admission (3). We propose to include delirium screening in preoperative assessment (in addition to frailty, cognitive impairment, osteoporosis, risk of falling, undernutrition, and polypharmacy).

Further, we would like to point out that, in addition to POD, postoperative cognitive dysfunction (POCD)

can also occur. POCD can last up to two years postoperatively. POD is an important risk factor for POCD, as are the presence of somatic diseases and subclinical delirious symptoms preoperatively (4).

The perception of POD as a highly relevant risk factor for the later quality of life of older people should receive the same importance as other preoperative diagnoses and treatments, therefore it should be included in the preoperative information for older people. The present work encourages this.

DOI: 10.3238/arztebl.2019.0374a

References

1. Olotu C, Weimann A, Bahrs C, Schwenk W, Scherer M, Kieffmann R: The perioperative care of older patients—time for a new, interdisciplinary approach. *Dtsch Arztebl Int* 2019; 116: 63–9.
2. Kratz T, Heinrich M, Schlauß E, Diefenbacher A: Preventing postoperative delirium: a prospective intervention with psychogeriatric liaison on surgical wards in a general hospital. *Dtsch Arztebl Int* 2015; 112: 289–96.
3. Levkoff S, Cleary P, Liptzin B: Epidemiology of delirium: an overview of research issues and findings. *International Psychogeriatrics* 1991; 3: 149–67.
4. Kratz T, Diefenbacher A: Kognitive Akut- und Langzeitfolgen intensivmedizinischer Behandlung. *Nervenarzt* 2016; 87: 246–52.

Prof. Dr. med. Torsten Kratz

Prof. Dr. med. Albert Diefenbacher, MBA

Evangelisches Krankenhaus Königin Elisabeth Herzberge gGmbH
Abteilung für Psychiatrie, Psychotherapie und Psychosomatik
Berlin, Germany
T.Kratz@keh-berlin.de

Conflict of interest statement

The authors declare that no conflict of interest exists.

In Reply:

Postoperative delirium is one of the most common and devastating complications of older patients. Kratz and Diefenbacher rightly emphasize that delirium can also be present preoperatively. Patients with preoperative delirium present with conspicuous findings in cognitive testing, which is recommended as part of the preoperative assessment. In this case, considering the possibility of preoperative delirium and then carrying out delirium screening is a sensible additional step. It should be taken into account that the aspects of “fluctuation” and “acute onset” according to ICD-10 may be difficult to assess here. Similar to Kratz and Diefenbacher, Thomas also emphasizes the high relevance of postoperative cognitive dysfunction (POCD), which can develop from delirium. In our article, we address the serious consequences of delirium, without however calling POCD by its name. Nonetheless, this clinical picture is explicitly included here, as are all cognitive changes that negatively affect quality of life and daily functioning of patients but that do not meet criteria for POCD. In general, diagnosis of POCD is based solely on psychometric test results, and patient-relevant factors such as impact on independence and daily life activities are often not included. Furthermore, outside the specialties of anesthesiology and surgery, there is a continuing demand for redefinition of the concept of POCD (1).

We (the authors) are eagerly following the promising PAWEL study by Thomas *et al.*. Prevention of delirium is an essential therapeutic goal of perioperative geriatric medicine. Implementation of appropriate measures often fails in practice due to time and personnel resources. The health economic evaluation of later costs of delirium—which are difficult to capture in monetary terms—gives this study an outstanding importance. Hopefully, this will show that delirium prevention leads to an extra financial burden only in the short term, and that it can save on costs to health care as well as to society over the long term.

Finally, it should be emphasized that our demands for perioperative geriatric medicine by no means represents a disregard of the geriatric expertise, as feared by Burkhardt. The expertise of geriatric departments is indispensable for perioperative care of older patients. However, given the number of affected patients and the numerous hospitals with no geriatric departments, presenting and assisting every older patient at risk is not possible. Gerontological “basic

knowledge” is too often lacking in our clinical daily practice. Therefore, our demands were aimed primarily at sensitizing the non-geriatrician care provider to the special features of older patients—so that perioperative geriatric medicine can be practiced even when no geriatrician is present.

DOI: 10.3238/arztebl.2019.0374b

References

1. Evered L, Silbert B, Knopman DS, et al.: Recommendations for the nomenclature of cognitive change associated with anaesthesia and surgery-2018. *Br J Anaesth* 2018; 121: 1005–12.
2. Olotu C, Weimann A, Bahrs C, Schwenk W, Scherer M, Kiefmann R: The perioperative care of older patients—time for a new, interdisciplinary approach. *Dtsch Arztebl Int* 2019; 116: 63–9.

On behalf of the authors

Dr. med. Cynthia Olotu
Universitätsklinikum Hamburg-Eppendorf
Klinik und Poliklinik für Anästhesiologie
Hamburg, Germany
c.olotu@uke.de

Conflict of interest statement

The author declares that no conflict of interest exists.