Ascending Aortic Aneurysm by Cardiac CT Angiography

Raveen Pal, MD; Ambarish Gopal, MD; Matthew J. Budoff, MD

Division of Cardiology, Department of Medicine, University of California, Irvine, California (Pal);

Division of Cardiology, Los Angeles Biomedical Research Institute at Harbor-UCLA, Torrance,

California (Budoff); Division of Cardiology, Loma Linda University Medical Center, Loma Linda,

California (Gopal)

ABSTRACT

Thoracic aortic aneurysm (TAA) is an increasingly frequent and potentially life-threatening medical diagnosis. TAA is defined as a localized dilatation of the aorta to more than 50% of baseline. Patients are most commonly asymptomatic at the time of diagnosis. Diagnosis is made incidentally on chest radiography or echocardiography. Diagnosis is then confirmed by echocardiography, left ventricular angiography, cardiac computed tomography or cardiac magnetic resonance. Cardiac computed tomography angiography (CTA) is an excellent imaging modality for this purpose as it allows simultaneous analysis of the coronary artery tree, which is useful in determining the most suitable procedure for each patient.

Thoracic aortic aneurysm (TAA) is an increasingly frequent and potentially life-threatening medical diagnosis. TAA is defined as a localized dilatation of the aorta to more than 50% of baseline. TAAs are classified in the same manner as aortic dissections. The Stanford system divides TAAs into 2 categories: Stanford Type A, which involves the ascending and/or transverse aorta, and Stanford Type B, which involves the descending aorta distal to the subclavian.

The etiology of TAAs includes risk factors for atherosclerosis (hypertension, smoking history, age, male gender), connective tissue disorders (Marfan's syndrome, Ehlers-Danlos syndrome), infections, trauma, or idiopathic causes.¹

Patients are most commonly asymptomatic at the time of diagnosis. Diagnosis is made incidentally on chest radiography or echocardiography. If there are presenting symptoms, these may include chest, back or flank pain or heart failure.

Three main treatment options include: (1) medical management of asymptomatic TAAs that are <6 cm in diameter. Care includes β -blocker treatment and routine yearly surveillance with imaging; (2) endovascular stent placement; (3) open surgical repair.

Complications of TAAs include aortic regurgitation, aortic leak, aortic rupture, and death. After aneurysmal size exceeds 6 cm, the risk for rupture and dissection is greater than 6.9% per year, and risk of death is 11.8% per year.² If a TAA ruptures, the risk of death is anywhere from 50% to 100%.^{3,4} Hence, it is extremely important to identify patients at high risk of rupture and offer corrective

surgery or endovascular stenting. The most important determinant of risk of rupture is size of aneurysm. Current guidelines propose that TAAs be scheduled for elective repair when they reach >6 cm in diameter, unless the patient has Marfan's syndrome or a bicuspid aortic valve, in which case the size limit is 5 cm.⁵ Other indications for repair include aneurysms >5 cm which have been rapidly expanding (>1 cm per y), symptomatic aneurysms regardless of size, associated severe aortic regurgitation, evidence of dissection, history of trauma, and a history consistent with mycotic aneurysm.

As stated, most cases are diagnosed incidentally by a finding of mediastinal enlargement on chest radiograph. Diagnosis is then confirmed by echocardiography, left ventricular angiography, cardiac computed tomography or cardiac magnetic resonance. Cardiac computed tomography angiography (CTA) is an excellent imaging modality for this purpose as it allows simultaneous analysis of the coronary artery tree, which is useful in determining the most suitable procedure for each patient. It is important to make measurements orthogonal to the imaging plane, as axial data has been demonstrated to overestimate thoracic aortic diameters. Typically, measurements made at both the level of the pulmonary artery and 10 millimeters above the left main coronary artery will help discern the normal diameters; measures are also made at the maximum diameter. Normal diameters should be age and gender based, although common clinical practice is to describe all aortas >4 mm as dilated.⁶

We present images from an asymptomatic patient with a Type A ascending aortic aneurysm (Figures 1

Figure 1. A, B, and C show anterior to lateral projections of three-dimensional volume rendered images that demonstrate the ascending aortic aneurysm which involves dilatation of the aortic root and extends to the beginning of the transverse aorta.

Figure 2. A and B show two-dimensional coronal and axial views using CTA showing the ascending aortic aneurysm extending from the root of the aorta to the transverse aorta (A is a coronal projection and B is an axial slice through the area of interest demonstrating the maximum diameter of 7.5 cm). Abbreviation: CTA = computed tomography angiography.

and 2). This case demonstrates the ability to accurately assess an incidentally discovered aortic aneurysm with CTA. It is imperative that physicians who interpret

cardiac CT, evaluate the ascending and descending aorta when performing these studies. This patient was subsequently scheduled for surgery given the location and size of the aneurysm (7.5 cm) and associated high risk of rupture.

References

1. Khalil A, Tarik T, Porembka D. Aortic pathology: aortic trauma, debris, dissection, and aneurysm. *Crit Care Med.* 2007;35(8 suppl): S392–S400.
2. Davies RR, Goldstein LJ, Coady MA, et al. Yearly rupture or dissection rates for thoracic aortic aneurysms: simple prediction based on size. *Ann Thorac Surg.* 2002;73:17–27; discussion, 27–28.
3. Johansson G, Markstrom U, Swedenborg J. Ruptured thoracic aortic aneurysms: a study of incidence and mortality rates. *J Vasc Surg.* 1995;21:985–988.
4. Bickerstaff LK, Pairolero PC, Hollier LH. Thoracic aortic aneurysms: a population-based study. *Surgery.* 1982;92:1103–1108.
5. Bonow RO, Carabello BA, Chatterjee K, et al. American College of Cardiology/American Heart Association Task Force on Practice Guidelines. *J Am Coll Cardiol.* 2008;52:e1–e142.
6. Mao SS, Ahmadi N, Shah B, et al. Normal thoracic aorta diameter on cardiac computed tomography in healthy asymptomatic adult: impact of age and gender. *Acad Rad.* 2008;15:827–834.