Reviews

A Systematic Review of Gender Differences in Mortality after Coronary Artery Bypass Graft Surgery and Percutaneous Coronary Interventions

CATHERINE KIM, M.D., M.P.H., RITA F. REDBERG, M.D., M.SC., TIANA PAVLIC, M.D., KIM A. EAGLE, M.D.

Division of General Internal Medicine, Departments of Medicine and Obstetrics & Gynecology, University of Michigan, Ann Arbor, Michigan, USA

Summary

Gender differences exist in outcomes, particularly early mortality, for percutaneous interventions (PCI) and coronary artery bypass graft surgery (CABG). Better understanding of this issue may target areas for improvement for all patients undergoing revascularization. Therefore, we summarized the evidence on gender differences in PCI and CABG outcomes, particularly early mortality, and mediators of this difference. Using the key terms “women” or “gender,” “revascularization,” “coronary artery bypass,” “angioplasty,” “stent,” and “coronary intervention,” we searched MEDLINE from 1985 to 2005 for all randomized controlled trials (RCTs) and registries reporting outcomes by gender. Bibliographies and the Web sites of cardiology conferences were also reviewed. The literature was examined to identify gender differences in outcomes and mediators of these differences. We identified 23 studies reporting outcomes by gender for CABG and 48 studies reporting outcomes by gender for PCI. The majority of studies noted greater in-hospital mortality in women than in men, with mortality differences resolving with longer follow-up. Early mortality differences were reduced but not consistently eliminated after adjustment for comorbidities, procedural characteristics, and body habitus. Power to detect gender

differences after multivariate adjustment was limited by declining mortality rates and small sample size. Gender was an independent risk factor for complications after both CABG and PCI. Women experience greater complications and early mortality after revascularization. Future exploration is needed of gender differences in quality of care and benefit from combinations of stenting and antiplatelet, and anticoagulant medications in order to optimize treatment.

Key words: women, coronary artery bypass graft, percutaneous coronary intervention, outcomes, gender, review

Clin. Cardiol. 2007; 30: 491–495.

© 2007 Wiley Periodicals, Inc.

Introduction

In the United States in 2002, 373,000 men and 142,000 women underwent CABG and 802,000 men and 402,000 women underwent a PCI, and these numbers are increasing.¹ Women may experience poorer outcomes, particularly greater early mortality, than men. This may be related to women’s less favorable risk profile. However, it is unclear whether additional factors contribute to the difference. Further understanding of these associations may identify opportunities to improve coronary care.

In this review, we summarize the status of gender differences in morbidity and mortality after CABG and PCI. We also summarize potential mediators of these differences. Finally, we present evidence on gender differences in specific aspects of periprocedural care that may affect outcomes, such as use of stents and adjunctive medications.

Methods

To identify randomized trials, we reviewed MEDLINE references published between January 1, 1985, and March 31, 2005, in the subset of randomized controlled trials and human subjects with these search terms: (women OR gender) AND coronary disease AND (revascularization OR coronary artery bypass OR stent) OR angioplasty OR coronary intervention) which yielded 1,725 articles. To identify registries, we re-

This work was supported by a grant from the Walter and Esther Hewlett Fund for Cardiovascular Research in Women (KA), the American Diabetes Association (CK), the Robert Wood Johnson Foundation (RR), and the Flight Attendant Medical Research Institute (RR).

Address for reprints:

Catherine Kim, M.D., M.P.H.

Division of General Internal Medicine

Departments of Medicine and Obstetrics Gynecology

University of Michigan

Ann Arbor, MI 48109, USA

e-mail: cathkim@umich.edu

Received: April 18, 2006

Accepted: May 24, 2006

Published online in Wiley InterScience

(www.interscience.wiley.com).

DOI:10.1002/clc.20000

© 2006 Wiley Periodicals, Inc.

viewed MEDLINE references in the same period in the subset of human subjects with the addition of "registry," which yielded 441 articles. We also reviewed reference lists and web pages of cardiology conferences for articles not yet published. Finally, we reviewed summaries published by the American Heart Association, the European Heart Society, and the Cochrane Collaboration. We excluded studies that did not report outcomes separately by gender or did not report on outcomes by procedure. We excluded studies that only reported outcomes by acute coronary syndrome (ACS) type, such as non-ST-elevation myocardial infarction (NSTEMI) or STEMI, and those that did not report on mortality or on a composite endpoint including mortality. Due to the small numbers of women enrolled, lack of reporting by gender, and the relatively small number of studies, we excluded studies examining only excimer laser angioplasty, directional atherectomy, cutting balloon angioplasty, extraction atherectomy, vascular brachytherapy, and saphenous vein graft interventions.

We identified 23 studies reporting outcomes by gender for CABG and 48 studies reporting outcomes by gender for PCI; in several instances, more than one report came from the same database, and in these instances the largest dataset was reported upon.

Results

Gender Differences in Early Mortality

Unadjusted in-hospital mortality rates from PCI registries, PCI trials, and CABG studies were higher in women than men. The majority of studies demonstrated similar late mortality between men and women. Several explanations have been offered for the increased early mortality in women: greater number of comorbidities and degree and type of vessel involvement, smaller coronary vessel size and body habitus, delayed intervention, less frequent use of arterial grafts, and racial and socioeconomic differences. Adjustment for these factors eliminated the gender difference in some studies, but not in others.

Comorbidities, anatomy, and early mortality: Women undergoing PCI and CABG were on average older than men and more often had diabetes, hypertension, congestive heart failure, and severe noncardiac disease. These factors, particularly age, account for much of the gender difference in early mortality in studies of both PCI² and CABG.³ Paradoxically, studies have suggested that gender differences are greater in younger than in older patients.^{4,5} Diabetes may also have a greater adverse impact on women than on men.^{6,7} Lesion type and total number of lesions may not be independently associated with outcome after adjustment for other risk factors,⁸ but acknowledgment of degree of vessel involvement (anatomy) as well as clinical risk factors may be required to account for gender differences.⁹ Of note, several PCI^{4,8,10,11} and CABG^{12,13} series still demonstrated gender differences in early mortality after adjustment for clinical and anatomical factors.

Smaller coronary arteries and body habitus: Since women tend to have smaller coronary arteries than men, investigators

have explored vessel size as a mediator of women's higher mortality after CABG and PCI. Presumably, smaller vessel size would be related to lower angiographic success and/or greater periprocedural complications due to technical difficulties. However, the relationship between vessel size and outcomes is inconsistent, with some studies showing differences^{3,14} and others showing no differences.^{7,15,16}

When information on vessel size is not available, markers for body habitus have sometimes been used as surrogates. In one CABG registry, the correlation between body surface area (BSA) and coronary artery size was 0.9,¹⁷ although other studies have found correlations as low as 0.1.^{18,19} As with studies adjusting for reference vessel size, studies examining markers of body habitus and mortality are inconsistent. Several CABG^{5,20–22} and PCI^{23–25} studies have demonstrated associations between gender and mortality after adjustment for body habitus, while other CABG^{3,17,26} and PCI^{11,19,27–31} studies have not.

One explanation for the conflicting results may be that gender differences in mortality are more pronounced with larger BSA, and different studies may enroll patients with different ranges of body habitus.^{22,32,33} In addition, Trabattoni *et al.* found that restenosis rates after PCI were higher in women, despite adjustment for BMI and risk factor differences. Of interest is the fact that when they also considered final balloon size, the magnitude of the gender difference was decreased.³⁴ Final balloon size, not BMI, was predictive of restenosis regardless of gender.

Early intervention and referral: Some studies have shown that women have longer times to PCI in the setting of ischemia,^{27,35–37} although the effect on gender mortality differences was minimal. When disease is not as severe or more diffuse, or morbidity is higher from related procedures after early intervention such as unplanned CABG, men may benefit more than women from early intervention.

Timing of referral for CABG may affect CABG outcomes. Studies have suggested that women may be referred for CABG less often than men, and possibly later in the course of disease.^{37–41} It is difficult to determine if referral bias before catheterization leads to increased clinical risk in women who do undergo CABG, but the contribution to mortality appears to be relatively small.

Women may be referred less for elective CABG than men but may undergo emergent CABG more often, either as the result of PCI complications or ACS.⁴² In turn, emergent CABG is associated with greater mortality than nonemergent CABG. Stents may be more likely to close in small vessels; Watanabe *et al.* found that women were more likely to undergo same-admission CABG after stent placement, presumably because of complications or unsatisfactory angiographic results.⁴³ However, more recent reports suggest that emergent CABG may not be as important a factor in gender differences in mortality.^{11,16}

Graft type and off-pump technology: Since arterial grafts have higher patency rates, widespread use of arterial grafts instead of saphenous vein grafts should reduce CABG mortality. Women have lower rates of arterial, specifically internal tho-

racic or mammary artery (IMA) use than men,^{15, 22, 33, 44} even after adjustment for other factors such as age, nonelective surgery, and extent of disease.^{15, 45} Thus, lower use of IMAs could be a potential mediator of gender differences in outcomes. However, the independent relationship with early mortality is inconsistent, with some CABG series demonstrating an association^{15, 26, 33} and others finding none.^{22, 45–47} Similarly, the use of off-pump revascularization may improve CABG outcomes, particularly periprocedural neurological events,⁴⁸ but it is unclear whether this effect is due to avoidance of cardiopulmonary bypass and/or other aspects of healthcare delivery, such as surgical skill.

Income and race: Women have lower income than men, although the effect of income as a mediator of gender differences in revascularization outcomes has not been widely studied. In a Canadian registry, adjustment for women's lower income reduced but did not eliminate the 1-year gender differences in mortality after CABG.²⁶ In an analysis of patients undergoing PCI or CABG at the Cleveland Clinic, lower median family income was associated with higher odds of receiving PCI instead of CABG, but income was not associated with mortality after either procedure.⁴⁹

The interaction of race, gender, and outcomes has not been widely studied. Of note, although racial and gender differences have been documented for revascularization, these have not been associated with similar differences in mortality.⁵⁰ Hartz *et al.* noted a significant interaction between race and gender in the performance of CABG, but not in CABG mortality.³² In another registry, black women were at lower risk for a composite endpoint of death, MI, and revascularization at 1 year than black men, with no gender difference among white patients.¹⁶

Adjunctive periprocedural therapies: Glycoprotein inhibitors: Without exception, randomized trials of periprocedural therapies are at least partially funded by the companies that profit from these technologies. With this in mind, randomized trials of glycoprotein IIb/IIIa inhibitors in conjunction with PCI have generally found benefit in both men and women, usually by a composite endpoint.^{51, 53} Gender differences may vary with the specific glycoprotein inhibitor, although the differences are small and the data are not consistent.^{54, 56} In contrast, the effect of IIb/IIIa inhibitors may differ for CABG, with minimal benefit on mortality.⁵⁷

Adjunctive periprocedural therapies: Stents: Studies of outcomes after coronary stenting in new device trials or registries suggest improved outcomes in both men and women when compared with past studies. As with glycoprotein inhibitors, firmer conclusions regarding the benefits of new technologies are limited by power, lack of reporting by gender, or both. Consequently, recent reductions in periprocedural mortality have made gender differences more difficult to detect, but it is unclear how much of this is attributable to the new technologies versus other improvements in care or patient selection.⁵⁸ Despite overall reductions in periprocedural mortality,⁵⁹ some studies suggest that women may not benefit as much as men from revascularization with drug-eluting stents.^{8, 60}

Conclusions

Gender differences in outcomes are greatest for in-hospital mortality. These differences are diminished after adjustment for risk factors such as age and diabetes. Correction for body habitus further diminishes the gender difference, although it is unclear how body habitus influences mortality. It is unclear how lower referral rates for women are related to their higher risk profile and increased adverse outcomes. Finally, it is unknown whether the uneven benefit of newer therapies, particularly drug-eluting stents and glycoprotein inhibitors, is attributable to gender differences in comorbidity patterns or other factors.

The American Heart Association's 2005 Scientific Statement notes that women have increased in-hospital mortality after both elective and primary PCI compared with men in the majority of reports, but that such differences are smaller after adjustment for women's greater age, smaller BSA, and comorbidities at the time of presentation.⁵⁹ Whether such differences are still significant after adjustment is unclear, since approximately half of the studies demonstrated persistently elevated adjusted odds ratios with the lower bounds of the confidence interval exceeding 1.⁵⁹

The American Heart Association and the American College of Cardiology, in collaboration with the American Association for Thoracic Surgery and the Society of Thoracic Surgeons, note that in-hospital CABG mortality is on average greater in women than in men, and sometimes, but not always, this disparity was reduced after adjustment for use of IMA grafts, age, and BSA. However, the joint statement noted that CABG compared with medical therapy still appeared to benefit women in appropriately selected patients.⁴⁷ The European Society of Cardiology guidelines do not make specific recommendations or statements regarding gender. No Cochrane Collaboration meta-analyses or statements on PCI or CABG have examined gender-specific outcomes.

Knowledge of the reasons for gender differences in outcomes after coronary revascularization can optimize therapy for all patients. The relative importance of factors in mediating these gender differences, particularly early mortality, requires studies that are adequately powered to conduct multivariate risk adjustment, study of the interactions between therapy and overall risk factor patterns, and closer inspection of processes of care, particularly medication use, surrounding revascularization. It is essential that clinical trials and registries report gender-specific outcomes, as it is likely that predictors of risk, including gender, will continue to evolve as procedure type and patient risk profiles change. It is also important to make a careful assessment of the influence of funding sources, as studies of newer technologies are heavily reliant upon the companies benefiting from these technologies.

References

1. American Heart Association. *Heart Disease and Stroke Statistics—2005 Update*. Dallas, TX: American Heart Association, 2005

2. Antoniucci D, Valenti R, Moschi G, Migliorini A, Trapani M, et al.: Sex-based differences in clinical and angiographic outcomes after primary angioplasty or stenting for acute myocardial infarction. *Am J Cardiol* 2001;87:289–293
3. Fisher L, Kennedy J, Davis K, Maynard C, Fritz J, et al.: Association of sex, physical size, and operative mortality after coronary artery bypass in the Coronary Artery Surgery Study (CASS). *J Thorac Cardiovasc Surg* 1982;84:334–341
4. Cowley M, Mullin S, Kelsey S, Kent K, Gruentzig A, et al.: Sex differences in early and long-term results of coronary angioplasty in the NHLBI PTCA Registry. *Circulation* 1985;71(1):90–97
5. Vaccarino V, Abramson J, Veledar E, Weintraub W: Sex differences in hospital mortality after coronary artery bypass surgery. Evidence for a higher mortality in younger women. *Circulation* 2002;105:1176–1181
6. Mehilli J, Kastrati A, Bollwein H, Dibra A, Schuhlen H, et al.: Gender and restenosis after coronary artery stenting. *Eur Heart J* 2003;24:1523–1530
7. Mehilli J, Kastrati A, Dirschinger J, Bollwein H, Neumann F, et al.: Differences in prognostic factors and outcomes between women and men undergoing coronary artery stenting. *J Am Med Assoc* 2000;284(14):1799–1805
8. Jackevicius C, Pinto R, Daly P, Mackie K, Dzavik V: Routine use of glycoprotein IIb/IIIa inhibitor therapy is associated with an improved in-hospital outcome after percutaneous coronary intervention: Insights from a large, prospective, single-centre registry. *Can J Cardiol* 2005;21(1):27–32
9. Alfonso F, Hernandez R, Banaños C, Fernandez-Ortiz A, Escaned J, et al.: Initial results and long-term clinical and angiographic outcome of coronary stenting in women. *Am J Cardiol* 2000;86:1380–1383
10. Ellis S, Myler R, King S, Douglas JJ, Topol E, et al.: Causes and correlates of death after unsupported coronary angioplasty: Implications for use of angioplasty and advanced support techniques in high-risk settings. *Am J Cardiol* 1991;68:1447–1451
11. Malenka D, Wennberg D, Quinton H, O'Rourke D, McGrath P, et al.: Gender-related changes in the practice and outcomes of percutaneous coronary interventions in Northern New England from 1994 to 1999. *J Am Coll Cardiol* 2002;40(12):2092–2101
12. Iyer V, Russell W, Leppard P, Craddock D: Mortality and myocardial infarction after coronary artery surgery. A review of 12,003 patients. *Med J Aust* 1993;159:166–170
13. Weintraub W, Wenger N, Jones E, Craver J, Guyton R: Changing clinical characteristics of coronary surgery patients: Differences between men and women. *Circulation* 1993;88(part 2):79–86
14. O'Connor N, Morton J, Birkmeyer J, Olmstead E, O'Connor G, for the Northern New England Cardiovascular Disease Study Group: Effect of coronary artery diameter in patients undergoing coronary bypass surgery. *Circulation* 1996;93(4):652–655
15. Mickleborough L: Gender differences in quality of distal vessels: Effect on results of coronary artery bypass grafting. *J Thorac Cardiovasc Surg* 2003;126:950–958
16. Robertson T, Kennard E, Mehta S, Popma J, Carrozza J, et al.: Influence of gender on in-hospital clinical and angiographic outcomes and on one-year follow-up in the New Approaches to Coronary Intervention (NACI) Registry. *Am J Cardiol* 1997;80(10A):26K–39K
17. Woods S, Noble G, Smith J, Hassefeld K: The influence of gender in patients undergoing coronary artery bypass graft surgery: An eight-year prospective hospitalized cohort study. *J Am Coll Surg* 2003;196:428–434
18. Sheifer S, Canos M, Weinfurt K, Arora U, Mendelsohn F, et al.: Sex differences in coronary artery size assessed by intravascular ultrasound. *Am Heart J* 2000;139:649–653
19. Cantor W, Miller J, Hellkamp A, Kramer J, Peterson E, et al.: Role of target vessel size and body surface area on outcomes after percutaneous coronary interventions in women. *Am Heart J* 2002;144:297–302
20. Hogue C, Barzilai B, Pieper K, Coombs L, DeLong E, et al.: Sex differences in neurological outcomes and mortality after cardiac surgery: A Society of Thoracic Surgery National Database Report. *Circulation* 2001;103:2133–2137
21. Christakis G, Weisel R, Buth K, Fremes S, Rao V, et al.: Is body size the cause for poor outcomes of coronary artery bypass operations in women? *J Thorac Cardiovasc Surg* 1995;110(5):1344–1356
22. Edwards F, Carey J, Grover F, Bero J, Hartz R: Impact of gender on coronary bypass operative mortality. *Ann Thorac Surg* 1998;66(1):125–131
23. Clayton T, Pocock S, Henderson R, Poole-Wilson P, Shaw T, et al.: Do men benefit more than women from an interventional strategy in patients with unstable angina or non-ST elevation myocardial infarction? The impact of gender in the RITA-3 Trial. *Eur Heart J* 2004;25:1641–1650
24. Kelsey S, James M, Holubkov A, Holubkov R, Cowley M, et al.: Results of percutaneous transluminal coronary angioplasty in women. 1985–1986 National Heart, Lung, and Blood Institute's Coronary Angioplasty Registry. *Circulation* 1993;87:720–727
25. Lansky A, Mehran R, Dangas G, Desai K, Costantini-Ortiz C, et al.: New-device angioplasty in women: Clinical outcome and predictors in a 7,372-patient registry. *Epidemiology* 2002;13:S46–S51
26. Guru V, Fremes S, Tu J: Time-related mortality for women after coronary artery bypass graft surgery: A population-based study. *J Thorac Cardiovasc Surg* 2004;127:1158–1165
27. Lansky A, Pietras C, Costa R, Tsuchiya Y, Brodie B, et al.: Gender differences in outcomes after primary angioplasty versus primary stenting with and without abciximab for acute myocardial infarction. *Circulation* 2005;111:1611–1618
28. Peterson E, Lansky A, Kramer J, Anstrom K, Lanzilotta M: Effect of gender on the outcomes of contemporary percutaneous coronary intervention. *Am J Cardiol* 2001;88:359–364
29. Bell M, Holmes DJ, Berger P, Garratt K, Bailey K, et al.: The changing in-hospital mortality of women undergoing percutaneous transluminal coronary angioplasty. *J Am Med Assoc* 1993;269(16):2091–2095
30. Arnold A, Mick M, Piedmonte M, Simpfordorfer C: Gender differences for coronary angioplasty. *Am J Cardiol* 1994;74(1):18–21
31. Weintraub W, Wenger N, Kosinski A, Douglas JJ, Liberman H, et al.: Percutaneous transluminal coronary angioplasty in women compared with men. *J Am Coll Cardiol* 1994;24(1):81–90
32. Hartz R, Rao A, Plomondon M, Grover F, Shroyer A: Effects of race, with or without gender, on operative mortality after coronary artery bypass grafting: A study using the Society of Thoracic Surgeons National Database. *Ann Thorac Surg* 2001;71:512–520
33. Abramov D, Tamariz M, Sever J, Chrstakis G, Bhatnagar G, et al.: The influence of gender on the outcome of coronary artery bypass surgery. *Ann Thorac Surg* 2000;70:800–806
34. Trabattini D, Bartorelli A, Montorsi P, Fabbiochi F, Loaldi A, et al.: Comparison of outcomes in women and men treated with coronary stent implantation. *Cathet Cardiovasc Intervent* 2003;58:20–28
35. Stone G, Grines C, Browne K, Marco J, Rothbaum D, et al.: Comparison of in-hospital outcome in men versus women treated by either thrombolytic therapy or primary coronary angioplasty for acute myocardial infarction. *Am J Cardiol* 1995;75:987–992
36. Vakili B, Kaplan R, Brown D: Sex-based differences in early mortality of patients undergoing primary angioplasty for first acute myocardial infarction. *Circulation* 2001;104:3034–3038
37. Blomkalns A, Chen A, Hochman J, Peterson E, Trynosky K, et al.: Gender disparities in the diagnosis and treatment of non-ST segment elevation acute coronary syndromes. Large-scale observations from the CRUSADE National Quality Improvement Initiative. *J Am Coll Cardiol* 2005;45(6):832–837
38. Khan S, Nessim S, Gray R, Czer L, Chaux A, et al.: Increased mortality of women in coronary artery bypass surgery: Evidence for referral bias. *Ann Intern Med* 1990;112:561–567
39. Sadanandan S, Cannon C, Gibson C, Murphy S, DiBattiste P, et al.: A risk score to estimate the likelihood of coronary artery bypass surgery during the index hospitalization among patients with unstable angina and non-ST-segment elevation myocardial infarction. *J Am Coll Cardiol* 2004;44(4):799–803
40. Thompson C, Humphries K, Gao M, Galbraith P, Norris C, et al.: Revascularization use and survival outcomes after cardiac catheterization in British Columbia and Alberta. *Can J Cardiol* 2004;20(14):1417–1423
41. Chandra N, Ziegelstein R, Rogers W, Tiefenbrunn A, Gore J, et al.: Observations of the treatment of women in the United States with myocardial infarction: A report from the National Registry of Myocardial Infarction-I. *Arch Intern Med* 1998;158(9):981–988
42. Seshadri N, Whitlow P, Acharya N, Houghtaling P, Blackstone E, et al.: Emergency coronary artery bypass surgery in the contemporary percutaneous coronary intervention area. *Circulation* 2002;106(18):2346–2350
43. Watanabe C, Maynard C, Ritchie J: Comparison of short-term outcomes following coronary artery stenting in men versus women. *Am J Cardiol* 2001;88:848–852
44. Jacobs A, Kelsey S, Brooks M, Faxon D, Chaitman B, et al.: Better outcome for women compared with men undergoing coronary revascularization. *Circulation* 1998;98:1279–1285
45. Aldea G, Gaudiani J, Shapira O, Jacobs A, Weinberg J, et al.: Effect of gender on postoperative outcomes and hospital stays after coronary artery bypass grafting. *Ann Thorac Surg* 1999;67(4):1097–1103
46. Davierwala P, Maganti M, Yau T: Decreasing significance of left ventricular dysfunction and reoperative surgery in predicting coronary artery bypass grafting-associated mortality: A twelve-year study. *J Thorac Cardiovasc Surg* 2003;126(5):1335–1344
47. Eagle K, Guyston R, Davidoff R, Edwards F, Ewy G, et al.: ACC/AHA 2004 Guideline Update for Coronary Artery Bypass Graft Surgery.

- American College of Cardiology Foundation and the American Heart Association Available at: www.americanheart.org. Accessed April, 2005
48. Mack M, Brown P, Kugelmass A, Battaglia S, Tarkington L, et al.: Current status and outcomes of coronary revascularization 1999 to 2002: 148,396 surgical and percutaneous procedures. *Ann Thorac Surg* 2004;77(3): 761–766
 49. Brener S, Lytle B, Casserly I, Schneider J, Topol E, et al.: Propensity analysis of long-term survival after surgical or percutaneous revascularization in patients with multivessel coronary artery disease and high-risk features. *Circulation* 2004;109:2290–2295
 50. Anand S, Xie C, Mehta S, Franzosi M, Joyner C, et al.: For the CURE Investigators: Differences in the management and prognosis of women and men who suffer from acute coronary syndromes. *J Am Coll Cardiol* 2005;46(10):1845–1851
 51. The CAPTURE Investigators: Randomised placebo-controlled trial of abciximab before and during coronary intervention in refractory unstable angina: The CAPTURE study. *Lancet* 1997;349:1429–1435
 52. The EPIC Investigators: Use of a monoclonal antibody directed against the platelet glycoprotein IIb/IIIa receptor in high-risk coronary angioplasty. *N Engl J Med* 1994;330(14):956–961
 53. The EPILOG Investigators: Platelet glycoprotein IIb/IIIa receptor blockade and low-dose heparin during percutaneous coronary revascularization. *N Engl J Med* 1997;336:1689–1696
 54. Topol E, Moliterno D, Herrmann H, Powers E, Grines C, et al.: Comparison of two platelet glycoprotein IIb/IIIa inhibitors, tirofiban and abciximab, for the prevention of ischemic events with percutaneous coronary revascularization. *N Engl J Med* 2001;344:1888–1894
 55. Fernandes L, Tcheng J, O'Shea J, Weinber B, Lorenz T, et al.: Is glycoprotein IIb/IIIa antagonism as effective in women as in men following percutaneous coronary intervention? *J Am Coll Cardiol* 2002;40(6):1085–1091
 56. Iakovou I, Dangas G, Mehran R, Lansky A, Kobayashi Y, et al.: Gender differences in clinical outcome after coronary artery stenting with use of glycoprotein IIb/IIIa inhibitors. *Am J Cardiol* 2002;89:976–979
 57. Roffi M, Mukherjee D, Chew D, Bhatt D, Cho L, et al.: Lack of benefit from intravenous platelet glycoprotein IIb/IIIa receptor inhibition as adjunctive treatment for percutaneous interventions of aortocoronary bypass grafts: A pooled analysis of five randomized clinical trials. *Circulation* 2002;106(24):3063–3067
 58. Vogt A, Engel H, Glunz H, Sattelberger U, Reil G, et al.: Early results of coronary angioplasty despite more complex interventions (Registry of the German Community Hospitals 1993–2000). *Am J Cardiol* 2002;90:1008–1009
 59. Lansky A, Hochman J, Ward P, Mintz G, Fabunmi R, et al.: Percutaneous coronary intervention and adjunctive pharmacotherapy in women. A statement for Healthcare Professionals from the American Heart Association. *Circulation* 2005;111:940–953
 60. Lemos P, Serruys P, van Domburg R, Saia F, Arampatzis C, et al.: Unrestricted utilization of sirolimus-eluting stents compared with conventional bare stent implantation in the “real world.” *Circulation* 2004;109:190–195