A Plate-based Cytotoxicity Assay for the Assessment of Rat Placental Natural Killer Cell Cytolytic Function

Cedar H. Baik¹, Mallory Geer¹, Olivia K. Travis², Denise C. Cornelius^{1,2}

¹Department of Emergency Medicine, University of Mississippi Medical Center

²Department of Experimental Therapeutics and Pharmacology, University of Mississippi Medical Center

Abstract

It is well known that decidual natural killer (NK) cells play a critical role in establishment and maintenance of normal pregnancy. Recent studies have demonstrated an altered population of circulating and decidual NK cells in women who suffer from adverse pregnancy complications such as recurrent miscarriage and preeclampsia. Studies from our group have shown that hypertension in pregnancy is associated with an increased population of activated NK cells in the placenta based on the expression of surface activation markers. This manuscript provides a detailed protocol to assess the cytotoxic function of NK cells isolated from placentas in a preeclampsia-like animal model of surgically induced placental ischemia. The following steps are described in detail: generation of single cell suspension, NK cell isolation, ex vivo stimulation, effector:target cell co-culture, and the cytotoxicity assay.

Keywords

Immunology and Infection; Issue 148; cytotoxicity; natural killer cell; Lymphoprep; lactate dehydrogenase; immunology; inflammation; hypertension

Introduction

Preeclampsia is a hypertensive disorder of pregnancy characterized by fetal growth restriction, end organ damage and chronic immune activation. Chronic immune activation in women with preeclampsia leads to increased circulating and placental inflammatory cytokines, an imbalance in CD4⁺ T Cells populations, and an increased population of activated Natural Killer (NK) cells¹. Studies recently published by our lab demonstrate a role for NK cells in causing some of the pathophysiology associated with preeclampsia in the Reduced Uterine Perfusion Pressure (RUPP) rat model of preeclampsia. Using flow cytometry to measure surface expression of activation markers on NK cells, an increased

Correspondence to: Denise C. Cornelius at dcornelius@umc.edu.

Disclosures

No conflicts of interest, financial or otherwise, are declared by the authors.

Video Link

The video component of this article can be found at <https://www.jove.com/video/58961/>

population of activated NK cells in the circulation and placentas of RUPP rats compared to normal pregnant (NP) rats was observed².

To confirm the flow cytometry observations, functional studies to assess the cytotoxic activity of NK cells isolated from the placentas of NP and RUPP rats were performed. There are several methods available for the assessment of cytotoxic function of cytotoxic CD8⁺ T cells and NK cells. The gold standard for functional cytotoxic analysis is the chromium release assay³. Other developed protocols utilized include flow cytometry⁴, image cytometry⁵, calcein release⁶, and most recently bioluminescence⁷. This video will provide a detailed protocol on using the well-established lactate dehydrogenase (LDH) release assay to measure cytotoxic function of NK cells using a commercially available LDH cytotoxicity assay kit.

Protocol

All protocols were approved by the Institutional Animal Care and Use Committee at the University of Mississippi Medical Center. The care and handling of the animals were in accord with the National Institutes of Health guidelines for ethical animal treatment.

1. Lymphocyte Cell Isolation from Placentas

1. Remove one placenta from the rat uterus (gestation day 19) and place in 10 mL of ice-cold PBS⁸.
2. Place one placenta on a 100 μ m filter and sit in a Petri dish containing 13.5 mL of RPMI and 1.5 mL of FBS (total volume is 15 mL). Use the flat side of a syringe plunger to push the placenta through the filter into the Petri dish.
3. Prepare three 15 mL conical tubes for each tissue. Add 3 mL of density gradient medium (see **Table of Materials**) to each tube, then carefully overlay 5 mL of homogenized placenta into each tube.
4. Centrifuge for 25 min at $300 \times g$ at room temperature (RT) with **no brake**. Collect the **thin white buffy layer** with a transfer pipette.

NOTE: After centrifugation, 3 layers are visible, the red RPMI at the top, the white buffy layer in the middle, and the clear density gradient medium layer at the bottom. Use a transfer pipette to pull up the white buffy layer from the tube. Combine buffy layer from all tubes of the same placenta into 1 tube.

5. Add 10 mL of RPMI to combined buffy layers. Centrifuge for 10 min, $300 \times g$, at 4 °C and discard supernatant.

2. Isolation of Natural Killer Cells

1. Resuspend cell pellet in 50 μ L of ice-cold PBS
2. Add biotin-labeled CD3 antibody to pelleted cells according to the manufacturer's protocol and mix well with a pipette. Place the tube in a tube rotator and incubate 20 min at 4 °C.

3. Add 1 mL of RPMI, centrifuge for 10 min at $400 \times g$ and 4°C , and discard supernatant.
4. Resuspend pellet in 1 mL of RPMI and combine with 150 μL of magnetic beads in a 1.5 mL microcentrifuge tube. Place the microcentrifuge tube in a tube rotator and rotate while incubating for 30 min at 4°C .
NOTE: Pull out release buffer at this time and allow to reach RT in the biosafety cabinet.
5. Place the tubes in the magnet for 1 min. Collect supernatant and save CD3⁻ cell population in a 15 mL tube on ice.
NOTE: This is the CD3⁻ population of cells.
6. Remove the tube from the magnet and add 1 mL of RPMI. Mix cells and beads 5 times with a pipette.
7. Repeat step 2.5.
8. Centrifuge the CD3⁻ population of cells for 10 min at $400 \times g$ and 4°C and discard supernatant. Resuspend the cell pellet in 50 μL of ice-cold PBS
9. Add biotin-labeled CD161a antibody to CD3⁻ cells according to the manufacturer's protocol and mix well. Place tube in a tube rotator and incubate for 20 min at 4°C .
10. Add 1 mL of RPMI, centrifuge for 10 min at $400 \times g$ and 4°C , and discard supernatant. Resuspend pellet in 1 mL of RPMI and combine with 150 μL of magnetic beads in a 1.5 mL microcentrifuge tube.
11. Place microcentrifuge tube in a tube rotator and rotate while incubating for 30 min at 4°C .
12. Place the tubes in a magnet for 1 min. Collect supernatant and discard CD3⁻/CD161a⁻ cells.
13. Remove tube from magnet and add 1 mL of RPMI. Mix cells and beads 5 times with a pipette.
14. Repeat step 2.12.
15. Remove microcentrifuge tubes from the magnet and add 1 mL of RT Release buffer. Place tube on tube rotator and rotate while incubating for 15 min at RT.
16. Place tubes in magnet for 1 min. Collect supernatant in a new 15 mL conical tube on ice.
NOTE: This is the population of NK cells.
17. Remove tube from magnet and add 1 mL of RT RPMI. Mix cells and beads 5 times with pipette.
18. Repeat step 2.16, placing supernatant in the same tube. Mix well and take a 20 μL sample to count cells

NOTE: Keep tube on ice.

19. Centrifuge CD3⁻/CD161a⁺ cells for 10 min, $400 \times g$ at 4 °C, then remove supernatant. Resuspend the cells in RPMI (10% FBS, 1% Pen/Strep, 2 ng/ mL IL-2) and seed at a concentration of 3×10^5 cells/well in a 6-well plate in 2.5 mL of NK Cell Activation Media. Incubate cells for 48 h at 37 °C, 5% CO₂ in a humidified incubator.

3. Cytotoxicity Assay: Retrieving NK Cells or YAC1 from Culture or Passing Cells

NOTE: All steps must be conducted under the hood. All cell tubes must be kept on ice at all times.

1. Use a glass serological pipet to collect YAC1 cells and media from flask, place in a 50 mL tube on ice, and mix well. Take 20 µL to count cells.
2. Spin the YAC1 cells for 10 min at $300 \times g$ and 4 °C. Count cells while these are spinning.
3. Add trypsin/EDTA to each well in a NK cell 6-well plate. Tap the plate and place in incubator.
4. After cells have incubated with trypsin/EDTA for ~5 min at 37 °C, scrape the plate/flask with a sterile plate scraper. Add 1 mL of NK Cell Media to each well.
5. Collect cells and media with serological pipet and collect in a 15 mL centrifuge tube. Take 20 µL to count cells.
6. Spin the NK cells for 10 min, $400 \times g$ at 4 °C. Count the sample of cells from step 3.5 during this centrifugation.

NOTE: View the culture plates under a microscope before discarding them to make sure there are no more cells adhered to the bottom of the wells. Since the experiment uses NK cells and YAC1 cells, be sure to count each one **separately**.

7. Count cells and resuspend at the concentrations determined in optimization trials to test for the appropriate number of target:effector ratio. This will be achieved by re-suspending the pellet in their corresponding media to make the following cell concentrations: YAC1 at 4×10^5 cells/mL and NK cells at 2×10^7 cells/mL.

4. Cytotoxicity Assay: Assay Protocol

1. Use a round bottom, culture treated 96-well plate to set up the plate as suggested in Table 1. This table shows the experimental controls and 3 sets of NK experimental columns. This can be expanded to a total of 10 NK experimental columns in a 96-well plate.
2. Centrifuge the assay plate at $250 \times g$ for 4 min to be certain that the effector and target cells are in contact. Incubate the 96-well plate for 5 hours in a humidified chamber incubator at 37 °C, 5% CO₂ to achieve ample contact between target and effector cells and target cell lysis by effector cells.

NOTE: The protocol can be paused here.

3. 45 min prior to harvesting supernatants, add 10 μ L of 10x Lysis Solution to the Target Cell Maximum LDH Release wells (Wells 1E, 1F, 1G, and 1H) and place the plate back in the humidified chamber.
4. After the incubation time is completed, centrifuge the plate at $250 \times g$ for 4 min. Using a multichannel pipettor, transfer 50 μ L aliquots from all wells to a fresh 96-well flat-bottom assay plate.
NOTE: Do not touch the bottom of the wells so that cells are not transferred into the fresh assay plate. Do not use a cultured treated plate as the fresh assay plate.
5. **Make Assay Reagent.**
 1. After Assay Buffer has reached room temperature, add 12 mL of Assay Buffer to one Substrate Mix bottle. Invert and shake gently until completely dissolved. 1 bottle is enough for two 96-well plates.
NOTE: Assay Buffer should be protected from light while being thawed. Mix immediately prior to use.
6. Add 50 μ L of Assay Reagent to all wells in the assay plate from step 4.6. Cover the plate with foil to protect it from light and incubate for 30 min at room temperature.
NOTE: Newly made Assay Reagent should be stored in a freezer. Reagent is good for approximately 8 weeks.
7. Add 50 μ L of Stop Solution to each well. Read plate within 1 hour after adding Stop Solution and record the absorbance at 490 nm. Representative data is shown in Table 2.

5. Calculation of Results

1. Calculate the average absorbance value from the Culture Medium Background wells and subtract from all absorbance values for Experimental, Target Cell Spontaneous LDH Release and Effector Cell Spontaneous LDH Release wells.
2. Calculate the average absorbance values for the Volume Correction Control wells and subtract from the absorbance values acquired for the Target Cell Maximum LDH Release Control wells.
3. Use the corrected values from Steps 5.1 and 5.2 in the following formula to calculate percent cytotoxicity for each effector:target well.

$$\% \text{ Cytotoxicity} = \frac{\text{Experimental} - \text{Effector Spontaneous} - \text{Target Spontaneous}}{\text{Target Maximum} - \text{Target Spontaneous}} \times 100$$

Representative Results

Placental NK cells obtained from NP and RUPP rats were incubated for 5 h with target cells in their respective medias at a ratio of 50:1 (NK:target). Absorbance was recorded at 490 nm

and the raw data is shown in Table 2. The average absorbance of the Culture Medium Background and the Volume Correction control wells were calculated. These averages were subtracted from the appropriate wells indicated in the manufacturer's protocol and are represented in Table 3. The corrected values were then used to obtain the % cytotoxicity using the protocol provided by the manufacturer (Table 4).

Discussion

There are a number of important key notes to consider for optimal results. The sterility of the cells utilized is very important. After collection of the placenta, it is important that preparation and isolation of the NK cells are performed under sterile conditions in a biosafety cabinet. Furthermore, because all cells release LDH upon cellular damage, care should be taken to obtain a high viability of NK cells after isolation and during the co-culture process. Too much spontaneous LDH release from the NK cells can often result in negative, unusable data

There are some aspects of this protocol that will be specific to the user and their needs. For example, there are various methods for isolation of mononuclear cells from placenta in addition to the methods in this article, such as mechanical and enzymatic disaggregation. Additionally, it is up to the user to determine the optimal target cells number and effector:target ratio that will be suitable for their experiment. In our hands, 10,000 target cells and an effector:target ratio of 50:1 was determined to be optimal for our experiments. These numbers may change depending on the target cells chosen and on the tissue from which the NK cells are isolated. Other investigators have utilized this method to assess NK function of cells isolated from liver and spleen⁹. Therefore, this method can be employed in various areas of research, not just in studies of preeclampsia, as we have chosen to use it.

The gold standard for cytotoxicity assessment is the chromium release assay⁵. While this method is reliable and reproducible, the most obvious limitation is the use of radioactive exposure risk to personnel and radioactive disposal. Other, non-radioactive cytotoxicity assays such as the flow cytometry based and calcien release assays are reported to have comparable results to CRA^{6,10,11}. However, flow-cytometry-based assays require additional specialized training of personnel in flow cytometry to run the assay. The LDH release assay doesn't require any specialized training, as this test only requires simple pipetting to perform the assay. Additionally, while the LDH release assay can be utilized using standard spectroscopy or fluorescent detection, the calcien release and flow-cytometry-based methods require equipment capable of detecting fluorescence. Limitations of the LDH release assay include an underestimation of cell death due to incomplete release of LDH from damaged cells.

In this assay, the measurement of LDH released into the media occurs by an enzymatic reaction in which iodonitrotetrazolium (a tetrazolium salt) is converted into formazan (red in color). The absorbance is measured by standard spectroscopy and the amount of damaged cells in the culture is proportional to the intensity of the red color. This assay allows for accurate determination of damaged or injured cells in a sample. Finally, this colorimetric

assay can also be used with other cell types to assay cytotoxicity occurring via cell-mediated as well as chemical-mediated mechanisms.

Acknowledgments

This work was supported by the National Heart Lung and Blood Institute of the National Institutes of Health under grant R00HL130456, the National Institute of General Medical Sciences of the National Institutes of Health under award P20GM104357, and by the Mississippi INBRE, funded by an Institutional Development Award (IDeA) from the National Institutes of General Medical Sciences of the National Institutes of Health under grant number P20GM103476. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

References

1. Cornelius DC, Cottrell J, Amaral LM, & LaMarca B Inflammatory Mediators: A causal link to hypertension during pregnancy- Studies in Preeclampsia. *British Journal of Pharmacology*. (2018).
2. Elfarra J et al. Natural killer cells mediate pathophysiology in response to reduced uterine perfusion pressure. *Clinical Science*. 131 (23), 2753–2762, (2017). [PubMed: 29042488]
3. Brunner KT, Mauel J, Cerottini JC, Chapuis B Quantitative assay of the lytic action of immune lymphoid cells on 51-Cr-labelled allogeneic target cells in vitro; inhibition by isoantibody and by drugs. *Immunology*. 14 (2), 181–196 (1968). [PubMed: 4966657]
4. Kim GG, Donnenberg VS, Donnenberg AD, Gooding W, Whiteside TL A novel multiparametric flow cytometry-based cytotoxicity assay simultaneously immunophenotypes effector cells: comparisons to a 4 h 51Cr-release assay. *Journal of Immunological Methods*. 325 (1–2), 51–66, (2007). [PubMed: 17617419]
5. Somanchi SS, McCulley KJ, Somanchi A, Chan LL, Lee DA A Novel Method for Assessment of Natural Killer Cell Cytotoxicity Using Image Cytometry. *PLoS One*. 10 (10), e0141074, (2015). [PubMed: 26492577]
6. Neri S, Mariani E, Meneghetti A, Cattini L, Facchini A Calcein-acetyoxymethyl cytotoxicity assay: standardization of a method allowing additional analyses on recovered effector cells and supernatants. *Clinical and Diagnostic Laboratory Immunology*. 8 (6), 1131–1135, (2001). [PubMed: 11687452]
7. Karimi MA et al. Measuring cytotoxicity by bioluminescence imaging outperforms the standard chromium-51 release assay. *PLoS One*. 9(2), e89357, (2014). [PubMed: 24586714]
8. Caporossi C, Nogueira PL, Marques JC, Assis RM, Aguilar-Nascimento JE Validation of the gastroschisis experimental model and the influence of the mother's diet enriched with glutamine in the fetal morphology. *Acta Cirúrgica Brasileira*. 29 (3), 158–165, (2014). [PubMed: 24626727]
9. Lv LH et al. Functional distinction of rat liver natural killer cells from spleen natural killer cells under normal and acidic conditions in vitro. *Hepatobiliary and Pancreatic Diseases International*. 11 (3), 285–293 (2012). [PubMed: 22672823]
10. Flieger D et al. A novel non-radioactive cellular cytotoxicity test based on the differential assessment of living and killed target and effector cells. *Journal of Immunological Methods*. 180 (1), 1–13 (1995). [PubMed: 7897241]
11. Jang YY et al. An improved flow cytometry-based natural killer cytotoxicity assay involving calcein AM staining of effector cells. *Annals of Clinical Laboratory Science*. 42 (1), 42–49 (2012). [PubMed: 22371909]

Table 1:**Assay Plate Layout.**

TCS = Target cell Spontaneous LDH Release (50 μ L target cells + 50 μ L target cell medium); TCM = Target Cell Maximum LDH Release (50 μ L target cells + 50 μ L target cell medium); VCC = Volume Correction Control (50 μ L target cell medium + 50 μ L NK Cell medium); CMB = Culture Medium Background Control (50 μ L target cell medium + 50 μ L NK Cell medium); ECSR = Effector Cell Spontaneous Release (50 μ L NK cells + 50 μ L NK cells medium); EW = Experimental wells (50 μ L target cells + 50 μ L NK cells).

1	2	3	4	5
TCS	VCC	ECSR-1	ECSR-2	ECSR-3
TCS	VCC	ECSR-1	ECSR-2	ECSR-3
TCS	VCC	ECSR-1	ECSR-2	ECSR-3
TCS	VCC	ECSR-1	ECSR-2	ECSR-3
TCM	CMB	EW-1	EW-2	EW-3
TCM	CMB	EW-1	EW-2	EW-3
TCM	CMB	EW-1	EW-2	EW-3
TCM	CMB	EW-1	EW-2	EW-3

Table 2:
Raw absorbance data of an assay plate containing the experimental controls in columns 1 and 2 and experimental wells to measure the cytotoxicity of placental NK cells from NP and RUPP rats against YAC1 target cells.

1	2	3	4	5
		NP	RUPP	NP
0.615	0.484	0.473	0.463	0.471
0.586	0.484	0.458	0.474	0.469
0.61	0.486	0.464	0.459	0.469
0.578	0.495	0.478	0.458	0.482
0.605	0.55	0.524	0.526	0.543
0.576	0.504	0.519	0.502	0.528
0.565	0.512	0.515	0.513	0.531
0.563	0.581	0.526	0.508	0.519

Table 3.**Corrected Absorbance.**

Absorbance of wells after subtraction of average absorbance of control wells according to the manufacturer's protocol. Column 1, Wells 1–4, are TCS absorbances values - the average CMB absorbance value. Column 1, Wells 5–8, are the TCM absorbance values - the average VCC absorbance value.

1	2	3	4	5
		NP	RUPP	NP
0.12775		−0.01425	−0.02425	−0.01625
0.09875		−0.02925	−0.01325	−0.01825
0.12275		−0.02325	−0.02825	−0.01825
0.09075		−0.00925	−0.02925	−0.00525
0.06825		0.03675	0.03875	0.05575
0.03925		0.03175	0.01475	0.04075
0.02825		0.02775	0.02575	0.04375
0.02625		0.03875	0.02075	0.03175

Table 4.**Cytotoxicity Calculations.**

Average percent cytotoxicity of replicate samples of placental NK cells isolated from NP and RUPP rats.

	NP	RUPP	NP
	128.9916	108.8235	93.69748
	63.44538	118.9076	66.80672
	75.92593	72.75132	64.28571
	66.27907	63.17829	83.33333
Average % Cytotoxicity	83.66049	90.91518	77.03081