Letters (continued from page 328)

References

1. Simpson RJ, Podolak R, Mangano CA, Foster JR, Dalldorf FG: Vagal syncope during recurrent pulmonary embolism. *J Am Med Assoc* 1983;249:390-393
2. Sievert H, Babic UU, Hausdorf G, Schneider M, Höpp HW, Pfeiffer D, Pfisterer M, Friedli B, Urban P: Transcatheter closure of atrial

- septal defect and patent foramen ovale with the ASDOS device (a multi-institutional European trial). *Am J Cardiol* 1998;82:1405-1413
3. Meacham RR III, Headley AS, Bronze MS, Lewis JB, Rester MM: Impending paradoxical embolism. *Arch Intern Med* 1998;158:438-448

Evolution of the Diagnosis of Acute Myocardial Infarction

C. R. Conti: *Clin Cardiol* 1999;22:163-164

To the Editor:

I enjoyed the Editor's Note "Evolution of the Diagnosis of Acute Myocardial Infarction," especially because it recapitulates many of my own experiences. I believe it stops short on the evolution of our understanding of the electrocardiogram, however. Several hospitals in our area have followed the suggestions of Ronald Selvester by adapting a 15-lead electrocardiogram for all chest pain patients.¹ We are convinced that the added leads have markedly improved our diagnostic accuracy and believe that it is time to once again expand the usefulness of this fantastic tool. We believe that somewhere between 20 and 30% of the patients with acute myocardial infarction will have their infarct identified by this simple expansion of the three extra leads and recognition of their utility.

M.H. Ellestad, M.D.
Director of Research MHI
Clinical Professor of Medicine UCI

Long Beach Memorial Medical Center
Long Beach, California

Reference

1. Ahmad J, Selvester RHS, Jaleel QM, Ellestad MH: Improved identification of acute myocardial infarction with 16-lead ECG. In *Electrocardiology '97* (Bacharova L and Macfarlane PW, Eds.). Singapore: World Scientific Publishing Co. Pte. Ltd., 1997

Author's reply:

I agree with Dr. Ellestad's letter.

C. Richard Conti, M.D., M.A.C.C.
Editor-in-Chief, *Clinical Cardiology*
Professor of Medicine
Division of Cardiology
Department of Medicine
University of Florida
Gainesville, Florida

Intrapericardial Therapeutics and Diagnostics (IPTD): Potential Advantages, Recent Advances, Experimental Direct Therapy of Cardiac Diseases and Arrhythmias

Clin Cardiol 1999;22(suppl 1):I-1-I-42

To the Editor:

I enjoyed reading the supplement to *Clinical Cardiology* entitled "Intrapericardial Therapeutics and Diagnostics" and edited by Dr. David Spodick. Certainly it is exciting to contemplate the pericardial space as a readily accessible route for therapeutic agents that could act directly on the heart surface and the subepicardium.

Drinker and Field¹ considered the pericardial space to be rather "inactive," but subsequently various workers, including our group,² were able to show considerable and brisk absorption from the pericardial space of various substances, including blood.

I have two concerns about using the pericardial space as a therapeutic access route, neither of which necessarily presents an insurmountable problem. Under certain circumstances absorption from this space may be quite brisk, both via the surface of the heart and from parietal pericardial venous and lymphatic drainage channels. The lymphatic routes that we³ and others have estab-

lished include the usual myocardial drainage pathway to the right lymphatic duct and the internal mammary chains bilaterally. Therefore, systemic effects of instilled agents would merit consideration. The other concern is the possibility of stimulating fibrotic reactions or causing bleeding into the pericardial space that could lead to adhesive or constrictive pericarditis.

Albert J. Miller, M.D.
Professor of Clinical Medicine (Cardiology)
Northwestern University Medical School
Chicago, Illinois

Reference

1. Drinker CK, Field ME: Absorption from the pericardial cavity. *J Exper Med* 1931;53:143-150
2. Miller AJ: *The Lymphatics of the Heart*. New York: Raven Press, 1982
3. Miller AJ, DeBoer A, Pick R, Palmer A, Van Pelt L: The lymphatic drainage of the pericardial space in the dog. *Lymphology* 1988; 21:227-233