Long-term outcomes of total hip arthroplasty in patients younger than 55 years: a systematic review of the contemporary literature

Xin Yu Mei, MD
Ying Jia Gong, MD
Oleg Safir, MD
Allan Gross, MD
Paul Kuzyk, MD, MASc

Accepted Oct. 15, 2018; Published online
June 1, 2019

Correspondence to:

X. Mei
Department of Orthopaedic Surgery
Rm 476A Mount Sinai Hospital
600 University Ave
Toronto ON M5G 1X5
xmei@qmed.ca

DOI: 10.1503/cjs.013118

Background: Total hip arthroplasty (THA) is increasingly performed in younger patients despite the lack of comprehensive assessment of long-term outcomes. We systematically reviewed the contemporary literature to assess the 1) indications, 2) implant selection and long-term survivorship, 3) complication and reoperation rates and 4) radiographic and functional outcomes of primary THA in patients younger than 55 years.

Methods: We searched the Embase and MEDLINE databases for English-language articles published between 2000 and 2018 that reported outcomes of primary THA in patients younger than 55 years with a minimum follow-up duration of 10 years.

Results: Thirty-two studies reporting on 3219 THA procedures performed in 2434 patients met our inclusion criteria. The most common preoperative diagnoses were avascular necrosis (1044 [32.4%]), osteoarthritis (870 [27.0%]) and developmental dysplasia of the hip (627 [19.5%]). Modular implants (3001 [93.2%]), cementless fixation (2214 [68.8%]) and metal-on-polyethylene bearings (1792 [55.7%]) were frequently used. The mean 5- and 10-year survival rates were 98.7% and 94.6%, respectively. Data on survival beyond 10 years were heterogeneous, with values of 27%–99.5% at 10–14 years, 59%–84% at 15–19 years, 70%–77% at 20–24 years and 60% at 25–30 years. Rates of dislocation, deep infection and reoperation for any reason were 2.4%, 1.2% and 16.3%, respectively. The mean Harris Hip Score improved from 43.6/100 to 91.0/100.

Conclusion: Total hip arthroplasty in patients younger than 55 years provides reliable outcomes at up to 10 years. Future studies should evaluate the outcomes of THA in this population at 15–20 years' follow-up.

Contexte : On effectue de plus en plus d'arthroplasties totales de la hanche (ATH) chez des patients qui ne sont pas âgés, malgré l'absence d'évaluation exhaustive des issues à long terme. Nous avons procédé à une revue systématique de la littérature récente pour analyser 1) les indications, 2) la sélection des implants et la survie à long terme, 3) les taux de complications et de réintervention, et 4) les résultats radiographiques et fonctionnels des ATH primaires chez les patients de moins de 55 ans.

Méthodes : Nous avons interrogé les bases de données Embase et MEDLINE pour recenser les articles de langue anglaise publiés entre 2000 et 2018 qui faisaient état des issues d'ATH primaires chez des patients de moins de 55 ans suivis pendant au moins 10 ans.

Résultats : Trente-deux études portant sur 3219 ATH effectuées chez 2434 patients répondaient à nos critères d'inclusion. Les diagnostics préopératoires les plus fréquents étaient la nécrose avasculaire (1044 [32,4%]), l'arthrose (870 [27,0%]) et la dysplasie développementale de la hanche (627 [19,5%]). Les implants modulaires (3001 [93,2%]), la fixation non cimentée (2214 [68,8%]) et le couple métal-polyéthylène (1792 [55,7%]) ont été fréquemment utilisés. Les taux de survie moyens à 5 et à 10 ans étaient de 98,7 % et de 94,6 %, respectivement. Les données sur la survie au-delà de 10 ans étaient hétérogènes, allant de 27 % à 99,5 % après 10 à 14 ans, de 59 % à 84 % après 15 à 19 ans, de 70 % à 77 % après 20 à 24 ans et de 60 % après 25 à 30 ans. Les taux de dislocation, d'infection profonde et de réintervention, toutes causes confondues, étaient de 2,4 %, de 1,2 % et de 16,3 %, respectivement. Le score de Harris moyen s'est amélioré, passant de 43,6/100 à 91,0/100.

Conclusion : L'arthroplastie totale de la hanche chez les patients de moins de 55 ans donne des résultats fiables pour les 10 premières années après l'intervention. Les prochaines études devraient évaluer les issues de l'arthroplastie de la hanche dans cette population après 15 à 20 ans de suivi.

Total hip arthroplasty (THA) reliably decreases pain and improves function and quality of life in patients with advanced hip disease at up to 25–30 years of follow-up.^{1,2} Despite early concerns over prosthetic longevity in patients with higher activity levels, improvements in implant design and surgical technique have led to increased demand for THA in younger, active patients.^{3,4} Kurtz and colleagues³ reviewed the American National Inpatient Sample database from 2006 and reported the proportion of primary THA procedures performed annually in patients younger than 55 years to be about 21%, with a projected rise to 28% by 2030. This proportion is slightly lower outside the United States, with rates of 11.9%, 6.4% and 13.2% reported by the 2014 Canadian,⁵ United Kingdom⁶ and Australian⁷ joint replacement registries, respectively.

Earlier studies showed high revision rates following Charnley low-friction arthroplasty in younger patients compared to older cohorts,^{8–10} with the main modes of failure being aseptic loosening and wear-induced osteolysis.^{11,12} Inferior implant survival in younger patients has been attributed to higher activity levels as well as a higher proportion of patients with inflammatory arthritis and congenital hip disease as their preoperative diagnosis.¹³ Despite these challenges, recent innovations including cementless fixation and alternative bearing surfaces have shown considerable promise in addressing many previous limitations of THA in younger patients.

Given these technological advances, we performed a systematic review of the contemporary literature with the aim of assessing the 1) indications, 2) implant selection and long-term survivorship, 3) rates of complication and reoperation and 4) radiographic and functional outcomes of primary THA in patients younger than 55 years of age.

METHODS

Literature search and study selection

We performed a literature search according to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines. Eligible studies were identified through a systematic search of MEDLINE and Embase databases from inception to April 2018. Database search terms included “total hip arthroplasty,” “total hip replacement,” “younger than 55,” “younger than 50,” “younger than 40,” “younger than 30,” “less than 55,” “less than 50,” “less than 40,” “less than 30” and “young patient.” We reviewed the bibliographies of all retrieved studies for relevant articles.

Two authors (X.Y.M. and Y.J.G.) independently screened the titles and abstracts of articles identified through the search strategy for eligibility. The predetermined inclusion criteria were 1) reporting of primary THA outcomes, 2) series of at least 20 patients, 3) mean follow-

up of at least 10 years, 4) all patients younger than 55 years at the time of surgery and 5) reporting of implant selection and survivorship, complications and functional outcomes. Articles were restricted to those that were published in full and written in English. Studies were excluded if they 1) were published before 2000, 2) were conference abstracts, case reports, reviews or surgical technique articles, 3) did not adequately report implant selection and survivorship or 4) reported outcomes of hemiarthroplasty, hip resurfacing, metal-on-metal bearings or revision THA. When multiple studies reported on the same study population, the article with the longest follow-up duration was selected, and the other studies were removed. Studies that passed the initial title and abstract screening were reviewed in full with the use of the same eligibility criteria. Any disagreement between authors was resolved by consensus.

Data extraction and quality assessment

Two authors (X.Y.M. and Y.J.G.) independently extracted the relevant data from each study and recorded them in an Excel document. Data collected included study design and period; patient age, sex and body mass index; mean follow-up duration and proportion of patients lost to follow-up; preoperative diagnosis and surgical approach; and implant selection, type of fixation and bearing surface. Outcome measures included implant survivorship at 5 years, 10 years and final follow-up; rates of dislocation, deep infection, other complications and reoperation; radiographic assessment; and functional outcome scores. Disagreements between reviewers were resolved by consensus.

Two authors (X.Y.M. and Y.J.G.) independently assessed the methodological quality of eligible studies. The Cochrane Collaboration’s tool for assessing risk of bias in randomized trials¹⁴ was used for randomized controlled trials, and the Methodological Index for Non-Randomized Studies (MINORS) was used for nonrandomized studies.¹⁵ Review Manager (RevMan) version 5.3 (Cochrane Collaboration) was used to construct the risk of bias summary. Any disagreement between reviewers was resolved by consensus. Agreement between reviewers on individual MINORS items was measured with the Cohen κ .¹⁶

Statistical analysis

We used descriptive statistics to summarize patient demographic characteristics, implant selection and outcome measures. We calculated weighted means for all interval and ratio data. Stratification by implant fixation was not performed owing to inconsistent reporting of outcomes and the small number of studies using cemented and hybrid fixation. A p value < 0.05 was considered statistically significant. We performed all statistical analysis using SPSS version 24.0 software (IBM Corp.).

RESULTS

The initial search yielded 435 potentially relevant articles after exclusion of duplicates (Fig. 1). After review of the titles and abstracts, 384 articles were excluded. Twenty-three additional articles were excluded after full-text review, leaving 32 articles eligible for inclusion in the systematic review.

Patient demographic characteristics

A total of 3219 primary THA procedures performed in 2434 patients between 1969 and 2006 were included. The year of surgery was 1970–1980 for 171 hips (5.3%), 1980–

1990 for 650 (20.2%), 1990–2000 for 1044 (32.4%) and after 2000 for 495 (15.4%); 9 studies reporting on 859 procedures (26.7%) had dates of surgery spanning more than 1 decade and thus could not be sorted into these intervals. A total of 777 procedures (24.1%) were performed after 1985. Of the 2434 patients, 1144 (47.0%) (standard deviation [SD] 17.2, range 20.5–85.7) were women. The mean age was 42.0 years (SD 7.4 yr, range 17.9–52.5 yr), and the mean body mass index was 26.6 (SD 2.7, range 22–29.6). The mean follow-up duration was 15.5 years (SD 5.9 yr, range 10–28.4 yr); 185 patients (7.6% [SD 8.1, range 0–39.1]) were lost to follow-up. The most common preoperative diagnoses were avascular necrosis (1044 cases

Fig. 1. Flow diagram showing study selection.

[32.4%]), primary/secondary osteoarthritis (870 [27.0%]) and developmental dysplasia of the hip (627 [19.5%]) (Table 1). A summary of the patient demographic characteristics is presented in Table 2, with full details available in Appendix 1 (available at cansurg.ca/013118-a1).

Surgical technique and implant selection

A variety of surgical approaches including anterolateral, posterolateral, posterior and transtrochanteric were used. Modular implants were used in 3001 hips (93.2%) and monoblock implants in 218 (6.8%). Implant fixation was cementless in 2214 hips (68.8%), hybrid in 540 (16.8%) and cemented in 465 (14.4%). Among the 2399 cementless femoral stems described, the fixation design was metaphyseal-fitting in 1544 cases (64.4%), metaphyseal-diaphyseal-junction-fitting in 366 (15.3%), diaphyseal-fitting in 223 (9.3%) and screwed into the femoral canal in 137 (5.7%); the fixation design was not reported in 129 cases (5.4%). The bearing surface used was metal-on-conventional-polyethylene in 1792 hips (55.7%), ceramic-on-ceramic in 748 (23.2%), ceramic-on-conventional-polyethylene in 530 (16.5%), metal-on-highly-cross-linked polyethylene in 147 (4.6%) and ceramic-on-highly-cross-linked polyethylene in 2 (0.1%). The femoral head diameter ranged from 22 to 36 mm. Two studies reported use of acetabular autografts.^{40,43} Full surgical and implant details are presented in Appendix 1.

Implant survivorship, complications and reoperation

The 5- and 10-year revision-free implant survival rates were 98.7% (SD 1.5%, range 95%–100%) and 94.6% (SD 5.5%, range 78.1%–100%), respectively. In studies

with mean follow-up beyond 10 years, reported revision-free survival rates were 27%–99.5% (16 studies) at 10–14 years, 59%–84% (2 studies) at 15–19 years, 70%–77% (2 studies) at 20–24 years and 60% (1 study) at 25–30 years. Rates of dislocation, deep infection and reoperation for any reason were 2.4% (SD 2.5%, range 0%–10.9%), 1.2% (SD 1.5%, range 0%–7%) and 16.3% (SD 13.6%, range 0%–63.8%), respectively. Aseptic loosening was the most common reason for reoperation. A summary of implant survivorship and complications is presented in Table 2, with full details available in Appendix 1.

Radiographic assessment and functional outcome

Nonprogressive acetabular and femoral radiolucent lines were observed in 418 (13.0%) (SD 18.4%, range 0%–85.2%) and 225 (7.0%) (SD 13.8%, range 0%–90.9%) components, respectively. Progressive radiolucency was not commonly reported. Functional outcome at final follow-up was reported with the use of the Harris Hip Score or the Merle d'Aubigné Score in all but 1 study, which used the Western Ontario and McMaster Universities Osteoarthritis Index.³⁴ The Harris Hip Score was reported in 26 studies, with a mean postoperative score of 91.0/100 (SD 4.8, range 81–98) and a mean improvement of 47.4 points (SD 4.6, range 32.5–53). The Merle d'Aubigné Score was reported in 7 studies, with a mean postoperative score of 16.0/18 (SD 1.6, range 10.5–17.1) and a mean improvement of 7.1 points (SD 1.5, range 5.6–8.7). Other reported outcome scores included the Hip Disability and Osteoarthritis Outcome Score, Oxford Hip Score and modified UCLA Activity Score.

Study quality

The level of evidence was I in 3 studies^{17–19} and IV in 29 studies.^{20–48} The risk of bias summary for each included randomized trial and the interrater agreement for each MINORS item are presented in Table 3 and Table 4, respectively. The overall risk of bias of the randomized trials was low. The majority of included case series were of low methodological quality. Agreement between the reviewers was considered satisfactory for all items. The mean MINORS global score was 11.0/16 (range 7–14).

DISCUSSION

We found that a large proportion of younger patients underwent THA for avascular necrosis and osteoarthritis secondary to congenital, developmental or traumatic anatomic abnormalities. This result is in keeping with the current literature.⁴⁹ These patients often present with major structural abnormalities such as proximal femoral deformity and femoral head collapse that increase the complexity of THA. Furthermore, patients who have undergone

Table 1. Preoperative diagnosis in patients younger than 55 years of age who underwent total hip arthroplasty

Diagnosis	No. (%) of hips n = 3219
Avascular necrosis	1044 (32.4)
Osteoarthritis (primary/secondary)	870 (27.0)
Developmental dysplasia of hip	627 (19.5)
Inflammatory arthropathy	267 (8.3)
Rheumatoid arthritis	92 (2.9)
Unclassified inflammatory arthropathy	70 (2.2)
Ankylosing spondylitis	58 (1.8)
Juvenile idiopathic arthritis	47 (1.5)
Post trauma	105 (3.3)
Congenital dislocation of hip	39 (1.2)
Slipped capital femoral epiphysis	36 (1.1)
Tumour	33 (1.0)
Legg-Calvé-Perthes disease	32 (1.0)
Septic arthritis	21 (0.6)
Osteochondritis dissecans	16 (0.5)
Still disease	2 (0.1)
Other/not reported	127 (3.9)

Table 2 (part 1 of 2). Summary of patient demographic characteristics, implant selection, survivorship and outcomes

Study	No. of patients	No. of hips	Age, yr, mean (range)	Length of follow-up, yr, mean (range)	Implant fixation	Bearing surface	Implant survival, %			Outcome; no. (%) of hips		
							5 yr	10 yr	Final	Dislocation	Deep infection	Reoperation
Choi et al., ²⁷ 2017	17	20	36.2 (21–40)	11 (10–13.5)	Cementless, hybrid	Metal-on-HXLPE	95	95	95 (11 yr)	1 (5.0)	0 (0.0)	1 (5.0)
Philippot et al., ³⁹ 2017	114	137	41 (18–50)	21.9 (3.3–30.9)	Cementless	MoP	NR	NR	77 (21.9 yr)	15 (10.9) (intra-prosthetic)	1 (0.7)	44 (32.1)
Schmoulders et al., ⁴¹ 2017	77	81	48 (30–50)	13.5 (9.7–16.9)	Cementless	CoP	NR	96.8	93 (13.5 yr)	1 (1.2)	0 (0.0)	6 (7.4)
Stambough et al., ⁴⁴ 2016	72	75	41.2 (17–50)	10 (8.2–11.9)	Cementless	Metal-on-HXLPE	96	92	92 (10 yr)	1 (1.3)	1 (1.3)	5 (6.7)
Kim et al., ¹⁸ 2016	200	400	52.5 (26–54)	11.8 (10–13)	Cementless	CoC	99.5	99.5	99.5 (11.8 yr)	0 (0.0)	0 (0.0)	2 (0.5)
Kim et al., ¹⁹ 2016	171	342	48 (21–50)	26.1 (25–27)	Hybrid, cementless	MoP	NR	NR	78.5 cup, 95.5 stem (26.1 yr)	5 (1.5)	1 (0.3)	73 (21.3)
McLaughlin et al., ³⁷ 2016	67	82	36.4 (20–49)	25 (20–29)	Cementless	MoP	NR	NR	90 stem (27 yr)	0 (0.0)	3 (3.7)	9 (11.0)
Kim et al., ³⁵ 2014	70	88	45.6 (19–49)	28.4 (27–29)	Cementless	MoP	NR	NR	66 cup, 90 stem (28.4 yr)	2 (2.3)	0 (0.0)	39 (44.3)
Babovic et al., ²¹ 2013	50	54	38.9 (15–50)	10.4 (NR)	Cementless	Metal-on-HXLPE 52, ceramic-on-HXLPE 2	NR	NR	98.1 (10.4 yr)	1 (1.9)	0 (0.0)	1 (1.9)
Chana et al., ²⁵ 2013	100	110	45.8 (20–55)	11.5 (10–13.5)	Cementless	CoC	NR	96.5	96.5 (11.5 yr)	1 (0.9) (late traumatic)	0 (0.0)	4 (3.6)
Schmitz et al., ⁴⁰ 2013	48	69	25 (16–29)	11.5 (7–23)	Cemented	MoP	NR	86	75 (15 yr)	2 (2.9)	4 (5.8)	16 (23.2)
Yoon et al., ⁴⁸ 2012	62	75	24 (18–30)	11.4 (10–13.4)	Cementless	CoC	NR	98.9	98.9 (10 yr)	1 (1.3) (with fractured liner)	0 (0.0)	1 (1.3)
Biernard et al., ²² 2011	80	93	44 (16–50)	12.3 (9.8–15.5)	Cementless	MoP	NR	84	84 (12.3 yr)	6 (6.5)	0 (0.0)	17 (18.3)
Faldini et al., ³⁰ 2011	28	34	47 (44–50)	12 (10–14)	Cementless	MoP 27, CoC 7	100	100	100 (12 yr)	1 (2.9)	0 (0.0)	0 (0.0)
Hsu et al., ³³ 2011	62	80	38.6 (16–49)	10.1 (10–12.3)	Hybrid	CoC	NR	96.3	96.3 (10 yr)	0 (0.0)	0 (0.0)	3 (3.8)
Kim et al., ¹⁷ 2011	78 79	109 110	43.4 (21–50) 46.8 (21–49)	18.4 (17–19) 18.4 (16–19)	Hybrid, cementless	MoP	NR	93.6	85.5 (20 yr)	3 (1.4)	3 (1.4)	38 (17.4)
Pakvis et al., ³⁸ 2011	131	158	42 (18–50)	13.2 (10–18)	Cementless	CoP 100, MoP 58	NR	98	80 (14 yr)	7 (4.4)	3 (1.9)	39 (24.7)
Boyer et al., ²³ 2010	69	76	39 (NR)	10 (7–15)	Hybrid, cementless	CoC	NR	92	92 (10 yr)	3 (3.9)	1 (1.3)	5 (6.6)
Burston et al., ²⁴ 2010	44	54	39.5 (18–50)	12.5 (10–17)	Cemented, hybrid	MoP	NR	NR	79.2 (12.5 yr)	4 (7.1)	2 (3.6)	14 (25.0)
Flecher et al., ³¹ 2010	212	233	42.6 (20–50)	10 (5–16)	Cementless	CoP	NR	96.7	87 (15 yr)	6 (2.6)	4 (1.7)	23 (9.9)
Wroblewski et al., ⁴⁷ 2010	26	35	17.9 (12–19)	15.6 (2.3–34)	Cemented	MoP 25, CoP 10	NR	NR	59 (15.6 yr)	0 (0.0)	1 (2.9)	16 (45.7)
Akbar et al., ²⁰ 2009	59	70	35 (22–40)	14 (10–16)	Cementless	CoP	100	100	86 (14 yr)	0 (0.0)	0 (0.0)	5 (7.1)
Lewthwaite et al., ³⁶ 2008	101	123	42 (NR)	12.5 (10–17)	Cemented	MoP	NR	94.4	92.6 (12.5 yr)	1 (0.8)	1 (0.8)	13 (10.6)
Utting et al., ⁴⁵ 2008	53	70	40 (19–49)	13.6 (12–16)	Hybrid	MoP	NR	NR	84 cup/liner (stem NR) (16 yr)	2 (2.9)	1 (1.4)	13 (18.6)
Wangen et al., ⁴⁶ 2008	42	47	25 (15–30)	13 (10–16)	Cementless	MoP	NR	NR	51 (13 yr)	3 (6.4)	0 (0.0)	24 (51.1)
Singh et al., ⁴² 2004	33	38	42 (22–49)	10 (5.3–14.2)	Hybrid, cementless	CoP 36, MoP 2	NR	NR	90.5 cemented cup, 96 cementless cup, 100 stem (12 yr)	0 (0.0)	0 (0.0)	4 (10.5)

Table 2 (part 2 of 2). Summary of patient demographic characteristics, implant selection, survivorship and outcomes

Study	No. of patients	No. of hips	Age, yr, mean (range)	Length of follow-up, yr, mean (range)	Implant fixation	Bearing surface	Implant survival, %			Outcome; no. (%) of hips		
							5 yr	10 yr	Final	Dislocation	Deep infection	Reoperation
Keener et al., ³⁴ 2003	42	57	42 (18–49)	25.7 (25–30)	Cemented	MoP	NR	NR	60 (30 yr)	1 (1.8)	4 (7.0)	22 (38.6)
Crowther et al., ²⁸ 2002	44	56	37 (22–49)	11 (9–14)	Cementless	MoP	NR	NR	87.5 (11 yr)	0 (0.0)	1 (1.8)	7 (12.5)
Chiu et al., ²⁶ 2001	33	47	28.8 (17–39)	14.9 (6.9–21.1)	Cemented	MoP	97.8	84.5	27 (15 yr)	2 (4.3)	1 (2.1)	30 (63.8)
Duffy et al., ²⁹ 2001	72	82	32 (17–39)	10.3 (10–14)	Cementless	MoP	96.3	78.1	78.1 (10 yr)	1 (1.2)	2 (2.4)	26 (31.7)
Garcia-Cimbrelo et al., ³² 2000	58	67	32.4 (18–39)	21.7 (5–25)	Cemented	MoP	98.5	88	70 (24 yr)	3 (4.5)	0 (0.0)	19 (28.4)
Smith et al., ⁴³ 2000	40	47	41 (21–50)	18.2 (17–20)	Cemented	MoP	NR	96.8	96.8 (10 yr)	1 (2.1)	0 (0.0)	6 (12.8)

CoC = ceramic-on-ceramic; CoP = ceramic-on-conventional-polyethylene; HXLPE = highly cross-linked polyethylene; MoP = metal-on-conventional-polyethylene; NR = not reported.

Table 3. Risk of bias for each included randomized trial¹⁴

Study	Random sequence generation (selection bias)	Allocation concealment (selection bias)	Blinding of participants and personnel (performance bias)	Blinding of outcome assessment (detection bias)	Incomplete outcome data (attrition bias)	Selective reporting (reporting bias)	Other bias
Kim et al. ¹⁷	+	+	?	+	+	+	?
Kim et al. ¹⁸	+	+	?	+	+	+	?
Kim et al. ¹⁹	+	+	?	+	+	+	?

+ = low risk of bias; ? = unclear risk of bias.

Table 4. Interreviewer agreement on Methodological Index for Non-Randomized Studies items¹⁵

Item	κ coefficient*
1. A clearly stated aim	0.862
2. Inclusion of consecutive patients	0.759
3. Prospective collection of data	0.603
4. Endpoints appropriate to the aim of the study	0.755
5. Unbiased assessment of the study endpoint	0.683
6. Follow-up period appropriate to the aim of the study	1.000
7. Loss to follow-up less than 5%	0.861
8. Prospective calculation of the study size	0.896

*A value > 0.4 is considered satisfactory.

previous surgery may have retained hardware, extensive scar tissue, heterotopic ossification or limb length discrepancy that require a greater degree of preoperative planning than routine primary THA in older patients with osteoarthritis.⁴⁹ Although THA had historically been performed in younger patients for rheumatoid arthritis, only 8.3% of hips in our review had inflammatory arthritis of any kind as their preoperative diagnosis. The decreasing demand for THA in patients with inflammatory arthritis likely reflects advances in disease-modifying antirheumatic drugs, leading to improved management of these conditions.⁵⁰

The predominance of modular, cementless implants in our review is also consistent with current practice patterns in North America and Europe. Modular implants have gained substantial popularity as they allow surgeons to independently adjust offset, version and limb length both

preoperatively and intraoperatively.⁵¹ In addition, modular head and liner exchange can be a less invasive revision option in cases of instability and eccentric polyethylene wear with well-fixed femoral and acetabular components.⁵² Although modularity is associated with risk of trunnion corrosion, adverse local tissue reactions and component fracture, the only modularity-related complications in our review were 4 cases of fractured ceramic liner.

The inconsistent survivorship of cemented implants in younger patients has led to a preference for cementless fixation in this population.^{53,54} We found that a variety of cementless femoral component designs were used, with metaphyseal-fitting stems being the most common. Metaphyseal-fitting stems are thought to not only increase proximal load transfer to reduce stress shielding and thigh pain, but also preserve diaphyseal bone for future revision,^{55–57} thus making them a popular choice in younger patients, who are expected to outlive their implants. Success with metaphyseal-fitting stems has furthered interest in shorter stem designs for additional bone preservation.^{18,58} In a within-patient randomized trial, Kim and colleagues¹⁸ reported the outcomes of 200 patients who underwent bilateral THA and were randomly allocated to receive conventional cementless stems in one hip and short cementless stems in the contralateral hip. Those authors found decreased stress shielding in the short stems but no difference in survivorship or functional outcome between the implants after a mean follow-up duration of 10.8 years.

The most commonly reported bearings in our review were metal-on-conventional-polyethylene, ceramic-on-

ceramic and ceramic-on-conventional-polyethylene. This may be related to our inclusion of only studies with long-term follow-up. Although highly cross-linked polyethylene has been shown to improve wear rates compared to conventional polyethylene,^{59,60} long-term data showing a clear increase in clinical survivorship are not yet available. The superior wear resistance of ceramic-on-ceramic bearings offers potential reduction of wear-induced osteolysis in comparison to metal-on-conventional-polyethylene bearings, although no study to date has shown a significant difference in reoperation rates. There have also been concerns over the risk of ceramic component fracture, chipping on insertion and squeaking.⁶¹ The incidence of ceramic-related complications in our review was relatively low, with 4 cases (0.3%) of fractured ceramic liners and 5 cases (0.4%) of squeaking. Ceramic-on-conventional-polyethylene bearings have also shown excellent wear resistance in the long term,^{62,63} although studies comparing these bearings with metal-on-conventional-polyethylene bearings have shown mixed results.⁶⁴⁻⁶⁷ We excluded metal-on-metal bearings from our review owing to widespread concerns over increased metal ion levels and adverse local tissue reactions to metallic wear debris.

The overall revision-free survival rate was 98.7% at 5 years and 94.6% at 10 years. These values are similar to those reported for THA in older patients. Mäkelä and colleagues⁶⁸ reviewed the Nordic Arthroplasty Registry for all primary THA procedures performed in patients older than 55 years between 1995 and 2011. Using the outcome of revision for any reason, they reported 10-year survivorship rates of 91.8%, 90.0% and 92.2% for cementless, hybrid and cemented components, respectively, in patients aged 55–64 years. Similarly, Hailer and colleagues⁶⁹ reviewed the Swedish Hip Arthroplasty Register from 1992 to 2007 and reported 10-year survivorship rates of 85% and 94% for cementless and cemented THA components, respectively. Survivorship beyond 10 years was inconsistently reported because the final follow-up periods varied greatly between studies. Therefore, we were able to present only the range of final survivorship rates reported in the included studies. This reflects the current paucity of studies with more than 15 years' follow-up, which is an inherent limitation of newer technology. Only 2 studies^{26,46} showed implant survivorship rates less than 75% at 10–15 years. Chiu and colleagues²⁶ reported acetabular component survivorship of 27% at 15 years in 47 Charnley low-friction arthroplasty procedures performed using early cementing techniques. They found inadequate cement mantle around 27 components (57%) that correlated strongly with subsequent aseptic loosening and emphasized the importance of good cementing technique. Wangen and colleagues⁴⁶ reported acetabular component survivorship of 51% at 13 years in 47 cement-

less THA procedures using hydroxyapatite-coated hemispherical cups in patients younger than 30 years. They attributed the high rates of cup loosening to the tendency for dissolution and resorption of the hydroxyapatite coating on the acetabular side, which had been observed in previous studies.^{70,71}

Rates of dislocation (2.4%) and deep infection (1.2%) in our review are comparable to values reported in large registry studies (1.9% and 1.3%, respectively).^{72,73} Rates of periprosthetic fracture were also relatively low in comparison to those in the literature (0.3% v. 1%), likely secondary to superior bone quality in younger patients.⁷⁴ The overall reoperation rate was high, at 16.3%, although many operations were late revisions for aseptic loosening more than 10 years after THA.

Radiographic evaluation of implant stability was performed with the DeLee and Charnley⁷⁵ and Gruen⁷⁶ systems for acetabular and femoral components, respectively. Progressive radiolucency was rarely reported, which minimized the risk of a large number of impending aseptic failures. Substantial improvements and high postoperative Harris Hip and Merle d'Aubigné scores suggest that most patients achieved satisfactory functional outcome.

Limitations

This study has several limitations. First, our review was restricted to English-language studies published after 1999 that had a mean follow-up duration of 10 years or more. Although having these strict inclusion criteria allowed us to focus on contemporary, long-term outcomes that are critical to counselling patients preoperatively, they may have resulted in exclusion of studies that contribute substantially to the literature on THA in younger patients. For instance, our minimum follow-up cut-off may have led to underrepresentation of highly cross-linked polyethylene in our study owing to its relative newness. Second, most of the included studies were small, retrospective case series of low methodological quality and were heterogeneous with respect to preoperative diagnosis, surgical technique and implant selection. This limited our ability to perform a meta-analysis and reflects the current lack of prospective, standardized, multicentre data in the orthopedic literature. Third, the paucity of studies with 15 years' follow-up limited our ability to draw conclusions about implant survivorship beyond 10 years. Finally, there was considerable variability in outcome reporting between studies. For instance, many studies did not differentiate between primary and secondary osteoarthritis. Moreover, 6 different clinical outcome scores were used among the included studies, with 11 studies reporting only postoperative scores. These inconsistencies not only limited statistical analysis but may also have led to misclassification bias.

CONCLUSION

Our study provides important trends and data that should help surgeons counsel younger patients undergoing THA. The most common preoperative diagnoses appear to be avascular necrosis, osteoarthritis and developmental dysplasia of the hip. Modular, cementless implants with metal-on-conventional-polyethylene bearings were used in most cases, with high survivorship seen at up to 10 years. Rates of dislocation and infection were comparable to those with THA in older patients, and good functional outcomes were routinely achieved. Future studies should aim to evaluate the outcome of THA in this population at 15–20 years' follow-up.

Affiliations: From the Department of Orthopaedic Surgery, Mount Sinai Hospital, Toronto, Ont. (Mei, Safir, Gross, Kuzyk); and the Department of Family Medicine, Queen's University, Kingston, Ont. (Gong).

Competing interests: None declared by X.Y. Mei and Y.J. Gong. O. Safir declares paid consultancies with Zimmer Biomet and DePuy Synthes, speaker fees from Zimmer Biomet, and research support from DePuy Synthes in the last 2 years. A. Gross declares speaker fees from Zimmer, and paid consultancies with Zimmer and Intellijoint within the last 2 years. P. Kuzyk declares research support from DePuy Synthes.

Contributors: X.Y. Mei, O. Safir, A. Gross and P. Kuzyk designed the study. X.Y. Mei acquired and analyzed the data, which Y.J. Gong also analyzed. X.Y. Mei wrote the article, which all authors reviewed and approved for publication.

References

- Callaghan JJ, Bracha P, Liu SS, et al. Survivorship of a Charnley total hip arthroplasty. A concise follow-up, at a minimum of thirty-five years, of previous reports. *J Bone Joint Surg Am* 2009;91:2617-21.
- Klapach AS, Callaghan JJ, Goetz DD, et al. Charnley total hip arthroplasty with use of improved cementing techniques: a minimum twenty-year follow-up study. *J Bone Joint Surg Am* 2001;83-a:1840-8.
- Kurtz SM, Lau E, Ong K, et al. Future young patient demand for primary and revision joint replacement: national projections from 2010 to 2030. *Clin Orthop Relat Res* 2009;467:2606-12.
- Maradit Kremers H, Larson DR, Crowson CS, et al. Prevalence of total hip and knee replacement in the United States. *J Bone Joint Surg Am* 2015;97:1386-97.
- Hip and knee replacements in Canada, 2014–2015: Canadian Joint Replacement Registry annual report*. Ottawa: Canadian Institute for Health Information; 2017.
- Age and gender of primary hip replacement patients (graph). *11th annual report 2014: National Joint Registry for England, Wales, Northern Ireland and the Isle of Man*. Hemel Hempstead (UK): National Joint Registry; 2015. Available: www.njrreports.org.uk/hips-primary-procedures-patient-characteristics/H06v4NJR?reportid=24DFC886-2530-4EEE-9327-999776161705&defaults=DC__Reporting_Period__Date_Range=%222015%7CNJR2014%22J__Filter__Calendar__Year=%22MAX%22,R__Filter__Gender=%22All%22,H__Filter__Joint=%22Hip%22 (accessed 2019 Apr. 24).
- Hip, knee and shoulder arthroplasty: annual report 2014*. Adelaide (Australia): Australian Orthopaedic Association National Joint Replacement Registry; 2014.
- Dorr LD, Kane TJ 3rd, Conaty JP. Long-term results of cemented total hip arthroplasty in patients 45 years old or younger. A 16-year follow-up study. *J Arthroplasty* 1994;9:453-6.
- Chandler HP, Reineck ET, Wixson RL, et al. Total hip replacement in patients younger than thirty years old. A five-year follow-up study. *J Bone Joint Surg Am* 1981;63:1426-34.
- Sharp DJ, Porter KM. The Charnley total hip arthroplasty in patients under age 40. *Clin Orthop Relat Res* 1985;201:51-6.
- Homesley HD, Minnich JM, Parvizi J, et al. Total hip arthroplasty revision: a decade of change. *Am J Orthop* 2004;33:389-92.
- Kurtz S, Mowat F, Ong K, et al. Prevalence of primary and revision total hip and knee arthroplasty in the United States from 1990 through 2002. *J Bone Joint Surg Am* 2005;87:1487-97.
- Schreurs BW, Hannink G. Total joint arthroplasty in younger patients: Heading for trouble? *Lancet* 2017;389:1374-5.
- Higgins JP, Altman DG, Gotzsche PC, et al.; Cochrane Bias Methods Group; Cochrane Statistical Methods Group. The Cochrane Collaboration's tool for assessing risk of bias in randomised trials. *BMJ* 2011;343:d5928.
- Slim K, Nini E, Forestier D, et al. Methodological index for non-randomized studies (MINORS): development and validation of a new instrument. *ANZ J Surg* 2003;73:712-6.
- McHugh ML. Interrater reliability: the kappa statistic. *Biochem Med* 2012;22:276-82.
- Kim YH, Kim JS, Park JW, et al. Comparison of total hip replacement with and without cement in patients younger than 50 years of age: the results at 18 years. *J Bone Joint Surg Br* 2011;93:449-55.
- Kim YH, Park JW, Kim JS. Ultrashort versus conventional anatomic cementless femoral stems in the same patients younger than 55 years. *Clin Orthop Relat Res* 2016;474:2008-17.
- Kim YH, Park JW, Kim JS, et al. Twenty-five- to twenty-seven-year results of a cemented vs a cementless stem in the same patients younger than 50 years of age. *J Arthroplasty* 2016;31:662-7.
- Akbar M, Aldinger G, Krahmer K, et al. Custom stems for femoral deformity in patients less than 40 years of age: 70 hips followed for an average of 14 years. *Acta Orthop* 2009;80:420-5.
- Babovic N, Trousdale RT. Total hip arthroplasty using highly cross-linked polyethylene in patients younger than 50 years with minimum 10-year follow-up. *J Arthroplasty* 2013;28:815-7.
- Biemond JE, Pakvis DF, Van Hellemond GG, et al. Long-term survivorship analysis of the cementless Spotorno femoral component in patients less than 50 years of age. *J Arthroplasty* 2011;26:386-90.
- Boyer P, Hutten D, Loriaut P, et al. Is alumina-on-alumina ceramic bearings total hip replacement the right choice in patients younger than 50 years of age? A 7- to 15-year follow-up study. *Orthop Traumatol Surg Res* 2010;96:616-22.
- Burston BJ, Yates PJ, Hook S, et al. Cemented polished tapered stems in patients less than 50 years of age: a minimum 10-year follow-up. *J Arthroplasty* 2010;25:639-99.
- Chana R, Facek M, Tilley S, et al. Ceramic-on-ceramic bearings in young patients: outcomes and activity levels at minimum ten-year follow-up. *Bone Joint J* 2013;95-b:1603-9.
- Chiu KY, Ng TP, Tang WM, et al. Charnley total hip arthroplasty in Chinese patients less than 40 years old. *J Arthroplasty* 2001;16:92-101.
- Choi WK, Kim JJ, Cho MR. Results of total hip arthroplasty with 36-mm metallic femoral heads on 1st generation highly cross linked polyethylene as a bearing surface in less than forty-year-old patients: minimum ten-year results. *Hip Pelvis* 2017;29:223-7.
- Crowther JD, Lachiewicz PF. Survival and polyethylene wear of porous-coated acetabular components in patients less than fifty years old: results at nine to fourteen years. *J Bone Joint Surg Am* 2002;84:729-35.

29. Duffy GP, Berry DJ, Rowland C, et al. Primary uncemented total hip arthroplasty in patients < 40 years old: 10- to 14-year results using first-generation proximally porous-coated implants. *J Arthroplasty* 2001;16(Suppl 1):140-4.
30. Faldini C, Miscione MT, Chehrassan M, et al. Congenital hip dysplasia treated by total hip arthroplasty using cementless tapered stem in patients younger than 50 years old: results after 12-years follow-up. *J Orthop Traumatol* 2011;12:213-8.
31. Flecher X, Pearce O, Parratte S, et al. Custom cementless stem improves hip function in young patients at 15-year followup. *Clin Orthop Relat Res* 2010;468:747-55.
32. Garcia-Cimbrelo E, Cruz-Pardos A, Cordero J, et al. Low-friction arthroplasty in patients younger than 40 years old: 20- to 25-year results. *J Arthroplasty* 2000;15:825-32.
33. Hsu JE, Kinsella SD, Garino JP, et al. Ten-year follow-up of patients younger than 50 years with modern ceramic-on-ceramic total hip arthroplasty. *Semin Arthroplasty* 2011;22:229-33.
34. Keener JD, Callaghan JJ, Goetz DD, et al. Twenty-five-year results after Charnley total hip arthroplasty in patients less than fifty years old: a concise follow-up of a previous report. *J Bone Joint Surg Am* 2003;85:1066-72.
35. Kim YH, Park JW, Park JS. The 27 to 29-year outcomes of the PCA total hip arthroplasty in patients younger than 50 years old. *J Arthroplasty* 2014;29:2256-61.
36. Lewthwaite SC, Squires B, Gie GA, et al. The Exeter™ universal hip in patients 50 years or younger at 10–17 years' followup. *Clin Orthop Relat Res* 2008;466:324-31.
37. McLaughlin JR, Lee KR. Total hip arthroplasty with an uncemented tapered femoral component in patients younger than 50 years of age: a minimum 20-year follow-up study. *J Arthroplasty* 2016;31:1275-8.
38. Pakvis D, Biemond L, Van Hellemond, et al. A cementless elastic monoblock socket in young patients: a ten to 18-year clinical and radiological follow-up. *Int Orthop* 2011;35:1445-51.
39. Philippot R, Neri T, Boyer B, et al. Bousquet dual mobility socket for patient under fifty years old. More than twenty year follow-up of one hundred and thirty one hips. *Int Orthop* 2017;41:589-94.
40. Schmitz MW, Busch VJ, Gardeniers JW, et al. Long-term results of cemented total hip arthroplasty in patients younger than 30 years and the outcome of subsequent revisions. *BMC Musculoskel Disord* 2013;14:37.
41. Schmolders J, Amvrazis G, Pennekamp PH, et al. Thirteen year follow-up of a cementless femoral stem and a threaded acetabular cup in patients younger than fifty years of age. *Int Orthop* 2017;41:39-45.
42. Singh S, Trikha SP, Edge AJ. Hydroxyapatite ceramic-coated femoral stems in young patients. A prospective ten-year study. *J Bone Joint Surg Br* 2004;86:1118-23.
43. Smith SE, Estok DM 2nd, Harris WH. 20-year experience with cemented primary and conversion total hip arthroplasty using so-called second-generation cementing techniques in patients aged 50 years or younger. *J Arthroplasty* 2000;15:263-73.
44. Stambough JB, Pashos G, Bohnenkamp FC, et al. Long-term results of total hip arthroplasty with 28-millimeter cobalt–chromium femoral heads on highly cross-linked polyethylene in patients 50 years and less. *J Arthroplasty* 2016;31:162-7.
45. Utting MR, Raghuvanshi M, Amirfeyz R, et al. The Harris–Galante porous-coated, hemispherical, polyethylene-lined acetabular component in patients under 50 years of age: a 12- to 16-year review. *J Bone Joint Surg Br* 2008;90:1422-7.
46. Wangen H, Lereim P, Holm I, et al. Hip arthroplasty in patients younger than 30 years: excellent ten to 16-year follow-up results with a HA-coated stem. *Int Orthop* 2008;32:203-8.
47. Wroblewski BM, Purbach B, Siney PD, et al. Charnley low-friction arthroplasty in teenage patients: the ultimate challenge. *J Bone Joint Surg Br* 2010;92:486-8.
48. Yoon HJ, Jeong JY, Kang SY, et al. Alumina-on-alumina THA performed in patients younger than 30 years: a 10-year minimum followup study. *Clin Orthop Relat Res* 2012;470:3530-6.
49. Polkowski GG, Callaghan J, Mont M, et al. Total hip arthroplasty in the very young patient. *J Am Acad Orthop Surg* 2012;20:487-97.
50. Hekmat K, Jacobsson L, Nilsson JA, et al. Decrease in the incidence of total hip arthroplasties in patients with rheumatoid arthritis — results from a well defined population in south Sweden. *Arthritis Res Ther* 2011;13:R67.
51. Duwelius PJ, Burkhart B, Carnahan C, et al. Modular versus non-modular neck femoral implants in primary total hip arthroplasty: Which is better? *Clin Orthop Relat Res* 2014;472:1240-5.
52. Walmsley DW, Waddell JP, Schemitsch EH. Isolated head and liner exchange in revision hip arthroplasty. *J Am Acad Orthop Surg* 2017;25:288-96.
53. Dorr LD, Kane ITJ, Pierce Conaty J. Long-term results of cemented total hip arthroplasty in patients 45 years old or younger: a 16-year follow-up study. *J Arthroplasty* 1994;9:453-6.
54. Dorr LD, Lueckett M, Conaty JP. Total hip arthroplasties in patients younger than 45 years. A nine- to ten-year follow-up study. *Clin Orthop Relat Res* 1990;260:215-9.
55. Mallory TH, Head WC, Lombardi AV Jr. Tapered design for the cementless total hip arthroplasty femoral component. *Clin Orthop Relat Res* 1997;344:172-8.
56. Renkawitz T, Francesco SS, Grifka J, et al. A new short uncemented, proximally fixed anatomic femoral implant with a prominent lateral flare: design rationals and study design of an international clinical trial. *BMC Musculoskelet Disord* 2008;9:147.
57. Saito J, Aslam N, Tokunaga K, et al. Bone remodeling is different in metaphyseal and diaphyseal-fit uncemented hip stems. *Clin Orthop Relat Res* 2006;451:128-33.
58. Kim YH, Park JW, Kim JS. Is diaphyseal stem fixation necessary for primary total hip arthroplasty in patients with osteoporotic bone (Class C bone)? *J Arthroplasty* 2013;28:139-46.e1.
59. Glyn-Jones S, Thomas GE, Garfield-Roberts P, et al. The John Charnley Award: highly crosslinked polyethylene in total hip arthroplasty decreases long-term wear: a double-blind randomized trial. *Clin Orthop Relat Res* 2015;473:432-8.
60. Kuzyk PR, Saccone M, Sprague S, et al. Cross-linked versus conventional polyethylene for total hip replacement: a meta-analysis of randomised controlled trials. *J Bone Joint Surg Br* 2011;93:593-600.
61. Kamath AF, Prieto H, Lewallen DG. Alternative bearings in total hip arthroplasty in the young patient. *Orthop Clin North Am* 2013;44:451-62.
62. Urban JA, Garvin KL, Boese CK, et al. Ceramic-on-polyethylene bearing surfaces in total hip arthroplasty. Seventeen to twenty-one-year results. *J Bone Joint Surg Am* 2001;83-a:1688-94.
63. Wroblewski BM, Siney PD, Fleming PA. Low-friction arthroplasty of the hip using alumina ceramic and cross-linked polyethylene. A ten-year follow-up report. *J Bone Joint Surg Br* 1999;81:54-5.
64. Nakahara I, Nakamura N, Nishii T, et al. Minimum five-year follow-up wear measurement of longevity highly cross-linked polyethylene cup against cobalt–chromium or zirconia heads. *J Arthroplasty* 2010;25:1182-7.

65. Stilling M, Nielsen KA, Soballe K, et al. Clinical comparison of polyethylene wear with zirconia or cobalt–chromium femoral heads. *Clin Orthop Relat Res* 2009;467:2644–50.
66. Hernigou P, Bahrami T. Zirconia and alumina ceramics in comparison with stainless-steel heads. Polyethylene wear after a minimum ten-year follow-up. *J Bone Joint Surg Br* 2003;85:504–9.
67. Epinette JA, Manley MT. No differences found in bearing related hip survivorship at 10–12 years follow-up between patients with ceramic on highly cross-linked polyethylene bearings compared to patients with ceramic on ceramic bearings. *J Arthroplasty* 2014;29:1369–72.
68. Mäkelä KT, Matilainen M, Pulkkinen P, et al. Failure rate of cemented and uncemented total hip replacements: register study of combined Nordic database of four nations. *BMJ* 2014;348:f7592.
69. Hailer NP, Garellick G, Kärrholm J. Uncemented and cemented primary total hip arthroplasty in the Swedish Hip Arthroplasty Register: evaluation of 170,413 operations. *Acta Orthop* 2010;81:34–41.
70. Lazarinis S, Kärrholm J, Hailer NP. Increased risk of revision of acetabular cups coated with hydroxyapatite: a Swedish Hip Arthroplasty Register study involving 8,043 total hip replacements. *Acta Orthop* 2010;81:53–9.
71. Reikeras O, Gunderson RB. Failure of HA coating on a gritblasted acetabular cup: 155 patients followed for 7–10 years. *Acta Orthop Scand* 2002;73:104–8.
72. Meek RM, Allan DB, McPhillips G, et al. Epidemiology of dislocation after total hip arthroplasty. *Clin Orthop Relat Res* 2006;447:9–18.
73. Dale H, Skråmm I, Løwer HL, et al. Infection after primary hip arthroplasty: a comparison of 3 Norwegian health registers. *Acta Orthop* 2011;82:646–54.
74. Della Rocca GJ, Leung KS, Pape HC. Periprosthetic fractures: epidemiology and future projections. *J Orthop Trauma* 2011;25 (Suppl 2):S66–70.
75. DeLee JG, Charnley J. Radiological demarcation of cemented sockets in total hip replacement. *Clin Orthop Relat Res* 1976;121:20–32.
76. Gruen TA, McNeice GM, Amstutz HC. “Modes of failure” of cemented stem-type femoral components: a radiographic analysis of loosening. *Clin Orthop Relat Res* 1979;141:17–27.

CJS's top viewed articles*

1. **Research questions, hypotheses and objectives**
Farrugia et al.
Can J Surg 2010;53:278–81
2. **Nonsurgical treatment of chronic anal fissure: nitroglycerin and dilatation versus nifedipine and botulinum toxin**
Tranqui et al.
Can J Surg 2006;49:41–5
3. **Clinical practice guideline: management of acute pancreatitis**
Greenberg et al.
Can J Surg 2016;59:128–40
4. **Blinding: Who, what, when, why, how?**
Karanicolas et al.
Can J Surg 2010;53:345–8
5. **Complications associated with laparoscopic sleeve gastrectomy for morbid obesity: a surgeons' guide**
Sarkosh et al.
Can J Surg 2013;56:347–52
6. **Surgical approach in primary total hip arthroplasty: anatomy, technique and clinical outcomes**
Petis et al.
Can J Surg 2015;58:128–39
7. **Tracheostomy: from insertion to decannulation**
Engels et al.
Can J Surg 2009;52:427–33
8. **Pharmacological management of postoperative ileus**
Zeinali et al.
Can J Surg 2009;52:153–7
9. **Granulomatous mastitis including breast tuberculosis and idiopathic lobular granulomatous mastitis**
Bakaris et al.
Can J Surg 2006;49:427–30
10. **Characteristics of highly successful orthopedic surgeons: a survey of orthopedic chairs and editors**
Klein et al.
Can J Surg 2013;56:192–8

*Based on page views on PubMed Central of research, reviews, commentaries and discussions in surgery. Updated July. 16, 2019.