Published in final edited form as:

Angew Chem Int Ed Engl. 2019 July 08; 58(28): 9434–9438. doi:10.1002/anie.201903308.

Ligand-Controlled Regiodivergent Enantioselective Rhodium-Catalyzed Alkene Hydroboration

Andrew J. Bochat, Veronika M. Shoba, James M. Takacs*

Department of Chemistry and Nebraska Center for Integrated Biomolecular Communication, University of Nebraska-Lincoln, 807 Hamilton Hall, Lincoln, NE, 68588-0304 (USA)

Abstract

Regiocontrol in the rhodium-catalyzed boration of vinyl arenes is typically dominated by the presence of the conjugated aryl substituent. However, small differences in TADDOL-derived chiral monophosphite ligands can override this effect and direct rhodium-catalyzed hydroboration of β -aryl and -heteroaryl methylenes by pinacolborane to produce either chiral primary or tertiary borated products selectively. The regiodivergent behavior is coupled with enantiodivergent addition of the borane. The nature of the TADDOL backbone substituents and that of the phosphite moiety function synergistically to direct the sense and extent of regioselectivity and enantioinduction. Twenty substrates undergo each reaction mode with regioselectivities reaching greater than 20:1 and enantiomer ratios reaching up to 98:2. A variety of subsequent transformations illustrate the potential utility of each product.

Entry for the Table of Contents

Regiodivergent with enantioselectivity included! Small changes in the structure of the TADDOL-derived chiral monophosphite ligand direct the rhodium-catalyzed hydroboration of methylenes to give either primary or tertiary chiral boronic esters.

Keywords

regiodivergent; enantiodivergent; rhodium-catalyzed hydroboration; catalysis

The enantioselective preparation of chiral boronic esters is of current interest due to their synthetic utility through a diverse set of stereospecific transformations^[1] and the growing appreciation that boronic acid derivatives may hold significant potential in medicinal

* jtakacs1@unl.edu.

Dedicated to Professor David A. Evans

Supporting information and the ORCID identification number(s) for the author(s) of this article can be found under:

chemistry.^[2] Considerable progress has been realized in the development of regioselective asymmetric hydroboration^{[3],[4]} and asymmetric borometallation^[5] reactions, most exploiting substrate control through use of activated alkenes or directing groups. Catalysts that override inherent substrate bias to selectively produce either of two regioisomers offer greater synthetic utility. In addition to several recent examples of non-stereocontrolled borylative difunctionalizations via regiodivergent borocupration,^[6] the ligand-controlled regiodivergent palladium-catalyzed hydroboration of terminal alkynes serves to illustrate the potential. Prabhu^[7] reported a bulky *N*-heterocyclic carbene ligand promotes regioselective boropalladation to deliver boron to the terminus; protonolysis leads to **1** (Figure 1). In contrast, tricyclohexylphosphine effects delivery of boron to the internal position leading to the regioisomeric product **2**.

Examples of chiral catalyst-controlled regiodivergent and enantioselective reactions are less common.^{[8],[9]} Herein, we report that relatively modest structural changes to a chiral TADDOL-monophosphite ligand results in a switch in both regioselectivity and the sense of enantioselectivity in the catalytic asymmetric hydroboration (CAHB) of β -aryl methylidene substrates **3** by pinacolborane (pinBH). This methodology provides access to either rhodium-catalyzed γ -boration at the unsubstituted terminus leading to the primary organoboronic ester **4** (*si*-face addition favored)^[10] or β -boration leading to the tertiary organoboronic ester **5** (*re*-face addition favored)^[11] using ligands (*R,R*)-**T1a** or (*R,R*)-**T2b**, respectively.

To probe the influence of structural changes in the ligand on the γ - to β -boration ratio (i.e., **4:5**) and the levels of asymmetric induction (i.e., *re:si* face addition), a series of TADDOL-derived phosphite ligands were evaluated under a standard set of screening conditions for the CAHB of **3a** (Figure 2). The selected TADDOL-derived ligands varied in the substituents at two positions. The **T1-T6** series of phosphites differ with respect to the aryl appendages on the TADDOL backbone; for example, the aryl appendages are 3,5-dimethylphenyl groups in the **T1** series, 4-methylphenyl groups in **T2**, etc. Next, each TADDOL backbone was evaluated as its phenyl phosphite (i.e., **T1a-T6a**, R = C₆H₅) and as its pentafluorophenyl phosphite (i.e., **T1b-T6b**, R = C₆F₅); for example, **T1a** is the TADDOL-phenylphosphite bearing 3,5-dimethylphenyl appendages and **T1b** is the corresponding pentafluorophenyl phosphite.

Phenylphosphite **T1a** (Ar = 3,5-Me₂C₆H₃) displays a significant preference for enantioselective γ -boration giving predominantly **4a** (6:1 γ : β); the major product reflects addition of pinBH to the *si*-face of the substrate (96:4 er). In contrast, the reaction run using phenylphosphite **T2a** (Ar = 4-MeC₆H₄) shows a slight preference for β -boration (1:1.1 γ : β), and the major regioisomer **5a** arises via predominant addition of pinBH to the *re*-face (93:7 er). The results show that the aryl substituents appended to the TADDOL backbone have a significant effect on the sense of enantioinduction and regioselectivity. The nature of the phosphite substituent also plays a significant role. Compared to the phenylphosphites **T1a-T3a**, the corresponding pentafluorophenyl phosphites **T1b-T3b** give enhanced levels of β -boration. Among the latter, **T2b** (Ar = 4-MeC₆H₄) gives the highest level (1:10 γ : β) and enantioselectivity favoring *re*-face addition (97:3 er).^[12] Ligands **T4-T6** generally are less selective than the other structural analogs examined. The 4-^tBuC₆H₄-TADDOL derivative

T5b is a curious case. It affords the β -boration product **5a** with moderate regioselectivity (1:4 γ : β), but in contrast to other β -selective ligands (e.g., **T2b**), pinBH adds to the *si*-face of the alkene (96:4 er).

Having identified the γ , *si*-face selective ligand **T1a** and β , *re*-face selective ligand **T2b**, a series of methylidenes were examined to probe the substrate scope of each.^[13] Figure 3A summarizes the results obtained using each ligand for CAHB/oxidation of 4-substituted aryl methylidene derivatives **3b-j**. Halogenated derivatives generally work well affording predominantly **4b-d** using **T1a** and predominantly **5b-d** using **T2b**. Yields are in the range of 64–76% after oxidation with enantiomer ratios (er) up to 98:2. The methyl benzoate derivative **3e**, 4-trifluoromethylphenyl derivative **3f**, and 4-methylphenyl derivative **3g** undergo γ -boration using the Rh-**T1a** catalyst to afford **4e** (70%, 94:6 er), **4f** (76%, 94:6 er), and **4g** (63%, 98:2 er), respectively. The more electron rich aromatic derivatives, 4-methoxyphenyl **3h** and 4-(dimethylamino)phenyl **3i**, afford their respective γ -borylated products **4h** (57%, 95:5 er) and **4i** (57%, 94:6 er) albeit with lower regiocontrol. The Boc-protected 4-aminophenyl derivative **3j** exhibits good regioselectivity, but the yield of **4j** (52%, 93:7 er) is again moderate.

The β -boration results obtained using the **T2b** catalyst for substrates **3b-f** are like those described above for the **T1a** γ -boration catalyst; yields for **5b-f** range from 64–78% with enantiomer ratios from 96:4 to 98:2 er. The two catalyst systems differ to a greater extent with substrates bearing electron donating substituents. In contrast to the moderate yields but good enantioselectivity using the **T1a** catalyst, **3g-i** (especially the 4-methoxyphenyl and 4-(dimethylamino)phenyl derivatives **3h** and **3i**) give somewhat higher yields but exhibit attenuated enantioselectivity for β -borylated **5g** (66%, 91:9 er), **5h** (76%, 86:14 er) and **5i** (60%, 80:20 er) using **T2b**.

Figure 3B illustrates the results obtained with several more highly functionalized aryl derivatives including several methylidenes bearing heteroaryl substituents. Using the **T1a**-catalyst, the 3-methoxyphenyl (**3k**), 3-chlorotolyl (**3l**), 3,4-methylenedioxyphenyl (**3m**), and indole derivative (**3n**) afford **4k-n** in 60–75% yield and up to 97:3 er. The thienyl and benzofuran derivatives **3o-q** give γ -borylated products in more moderate yields (50–59%) due to higher levels of β -boration for **4o** and **4q**, and higher levels of the reduction product for **4p**. The enantioselectivity is also lower for these substrates, ranging from 90:10 er for **4o** to 65:35 er for **4q**. In contrast, the 3- and 2-thienyl derivatives are efficient substrates in the **T2b**-catalyzed reaction affording **5o** (85%, 98:2 er) and **5p** (71%, 96:4 er), respectively. The benzofuran derivative **3q** provides good regio- and enantioselectivity for **5q** (40%, 94:6 er); the moderate yield reflects competing reduction in this case.^[4a, 14] The yields, regio- and enantioselectivities obtained for **5k** (60%, 94:6) and **5l** (65%, 95:5) are similar to those observed for the corresponding γ -borylated products. However, substrates bearing multiple donor groups can give surprising results; **5k** and **5l**, for example, form with the opposite sense of stereoreduction (see SI).

Several other limitations are found (Figure 3C). Substrates bearing *ortho*-substituted aryl groups (e.g., **3r** and **3s**) do not exhibit regio- or enantioswitching. Both catalyst systems provide almost exclusively γ -borylated products with little asymmetric induction. Unlike aryl

substituted methylenes **3a-s**, the alkyl-substituted methylene **3t** undergoes β -boration using the **T1a**-catalyst to give **5t** (85%, 95:5 er);^[4a] less than 5% of the γ -borylated product is formed. β -Boration of **3t** giving **5t** (58%, 92:8 er) remains the major pathway using ligand **T2b**, although the percentage of γ -boration increases to 32%.

The synthetic versatility of enantioselective, regiodivergent CAHB is demonstrated by the selected conversions of **4a** and **5a** to **6-21** (Figure 4).^[15] Both the primary and tertiary pinacol boronic esters are converted to the corresponding cesium^[16] or potassium^[17] trifluoroborate salts, **6a** and **7a** respectively, in 97–98% yield. Similarly, boronic acids **6b** and **7b** and the corresponding *N*-methyliminodiacetic acid (MIDA) esters^[18] **6c** and **7c** are readily accessible (62–72%). The one-carbon homologated boronic esters **8** (75%) and **9** (55%) are obtained via the Matteson protocol.^[19] In the case of the tertiary boronic ester **5a**, the reaction proceeds with stereoretention to afford (*R*)-**9** (98:2 er). Hydrolysis of the primary boronic ester **4a** to the boronic acid, followed by Raney Nickel cleavage of N-O bond affords chiral oxaborolane **10** (71%). Reduction of the oxime C-N double bond in **5a** with NaCNBH₃ affords **11** (77%); the reduction works equally as well for **4a** (results not shown, see SI).

In addition to forming boron derivatives **6–11**, other transformations highlight the flexible and diverse ways in which **4a** and **5a** can be elaborated. Using Aggarwal's modification of the Zweifel olefination,^[20] the primary boronic ester **4a** is converted to ketone **12** (57%). For the tertiary boronic ester **5a**, rhodium-catalyzed 1,2-addition to 4-nitrobenzaldehyde followed by DMP oxidation affords ketone **13** (59%, 96:4 er).^[21] Ketones **12** and **13** undergo acid-catalyzed cyclizations to the chiral dihydrooxazine **14** (39%) and chiral isoxazoline **15** (53%, 96:4 er), respectively.

The transition-metal-free sp^2 - sp^3 cross-couplings of electron-rich aromatic systems convert **4a** and **5a** to the thienyl derivatives **16** (70%) and **17** (84%), respectively.^[22] Treatment of the primary boronic ester **4a** with bromobenzene under typical palladium-catalyzed Suzuki cross-coupling conditions^[5c] affords **18**, a product that is not readily accessible using the transition-metal-free procedure employed for thienylation. Protodeboration of the tertiary boronic ester **5a** affords **19** (90%, 98:2 er).^[23] In our hands, the common amination procedure^[24] is only successful for the primary boronic ester **4a**; a moderate yield of the chiral 1,3-aminoalcohol derivative **20** (43%) is obtained. Oxidation of C-B bond in the tertiary boronic ester **5a** followed by N-O bond cleavage gives chiral 1,2-diol **21** (86%, 99:1 er).

In summary, the CAHB of styrene and related vinyl arenes (e.g., β -methylstyrene or indene derivatives) often leads to formation of the chiral secondary benzylic borated products. The observed regioselectivity is usually attributed to the formation of an intermediate η^3 -complex with the aryl substituent.^[25] However, aryl methylenes (e.g., α -methylstyrene) generally give the primary boronic ester. The chiral, tertiary benzylic regioisomer is not formed, presumably due to hindrance in forming the corresponding η^3 -complexed intermediate. We find that rhodium-catalyzed hydroboration of β -aryl methylenes affords either the chiral primary or tertiary boronic ester depending on the TADDOL-derived phosphite ligand employed. Twenty substrates are shown to undergo regioselective reaction

using each catalyst system. Regioselectivities reach levels greater than 20:1 and enantiomer ratios up to 98:2 are obtained. Conversions of **3a** to **6-21** illustrate the potential synthetic utility of regiodivergent CAHB.

In terms of the mechanism, the aryl substituents appended to the TADDOL backbone as well as the nature of the arylphosphite moiety act synergistically to affect the sense of regioselectivity and enantioinduction. This appears to be due, at least in part, to an electronic effect because the pentafluorophenyl substituent is not unique in promoting β -boration. For example, the corresponding **T2** ligand bearing either a (4-trifluoromethyl)phenyl or a 4-cyanophenyl phosphite substituent also exhibit regioswitching, although the levels of regio- and enantioselectivity are not as high. Deuterium labeling experiments with **T1a** and **T2b** show that addition of H (D) is rapid and reversible, leading to competitive H/D-exchange in the substrate using either regiodivergent catalyst.^[26] Several hypotheses that might account for the coupled regio- and enantiodivergent CAHB, effected by ostensibly similar catalyst systems, are currently under investigation.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

Funding from the NIH National Institutes of General Medical Sciences (R01-GM100101) is gratefully acknowledged.

References

- [1]. a) Matteson DS, J. Org. Chem 2013, 78, 10009; [PubMed: 23875690] b) Sandford C, Aggarwal VK, Chem. Commun 2017, 53, 5481.
- [2]. a) Das BC, Thapa P, Karki R, Schinke C, Das S, Kambhampati S, Banerjee SK, Veldhuizen PV, Verma A, Weiss LM, Evans T, Future Med. Chem 2013, 5, 653; [PubMed: 23617429] b) Diaz DB, Yudin AK, Nat. Chem 2017, 9, 731; [PubMed: 28754930] c) Antonio Abad G, Alexey R, Erik A-J, Jose GT-F, Curr. Med. Chem 2018, 25, 1;d) Mereddy GR, Chakradhar A, Rutkoski RM, Jonnalagadda SC, J. Organomet. Chem 2018, 865, 12.
- [3]. For recent examples, see a) Chen J, Xi T, Lu Z, Org. Lett 2014, 16, 6452; [PubMed: 25438006] b) Chen J, Xi T, Ren X, Cheng B, Guo J, Lu Z, Org. Chem. Front 2014, 1, 1306;c) Zhang L, Zuo Z, Wan X, Huang Z, J. Am. Chem. Soc 2014, 136, 15501; [PubMed: 25325782] d) Hu N, Zhao G, Zhang Y, Liu X, Li G, Tang W, J. Am. Chem. Soc 2015, 137, 6746; [PubMed: 25938599] e) Wang Z, He X, Zhang R, Zhang G, Xu G, Zhang Q, Xiong T, Zhang Q, Org. Lett 2017, 19, 3067. [PubMed: 28590749]
- [4]. For related publications, see a) Shoba VM, Thacker NC, Bochat AJ, Takacs JM, Angew. Chem. Int. Ed 2016, 55, 1465; Angew. Chem. 2016, 128, 1487;b) Chakrabarty S, Takacs JM, J. Am. Chem. Soc 2017, 139, 6066; [PubMed: 28414243] c) Chakrabarty S, Takacs JM, ACS Catal. 2018, 8, 10530. [PubMed: 31134137]
- [5]. a) Matsuda N, Hirano K, Satoh T, Miura M, J. Am. Chem. Soc 2013, 135, 4934; [PubMed: 23495912] b) Jia T, Cao P, Wang B, Lou Y, Yin X, Wang M, Liao J, J. Am. Chem. Soc 2015, 137, 13760; [PubMed: 26458555] c) Chen B, Cao P, Yin X, Liao Y, Jiang L, Ye J, Wang M, Liao J, ACS Catal. 2017, 7, 2425;d) Logan KM, Brown MK, Angew. Chem. Int. Ed 2017, 56, 851; Angew. Chem. 2017, 129, 869.
- [6]. a) Sakae R, Hirano K, Miura M, J. Am. Chem. Soc 2015, 137, 6460; [PubMed: 25951059] b) Su W, Gong T-J, Lu X, Xu M-Y, Yu C-G, Xu Z-Y, Yu H-Z, Xiao B, Fu Y, Angew. Chem. Int. Ed 2015, 54, 12957; Angew. Chem. 2015, 127, 13149;c) Su W, Gong T-J, Zhang Q, Zhang Q, Xiao

- B, Fu Y, ACS Catal. 2016, 6, 6417;d) Sardini SR, Brown MK, J. Am. Chem. Soc 2017, 139, 9823; [PubMed: 28699339] e) Smith KB, Brown MK, J. Am. Chem. Soc 2017, 139, 7721; [PubMed: 28581713] f) Jia T, He Q, Ruscoe RE, Pulis AP, Procter DJ, Angew. Chem. Int. Ed 2018, 57, 11305; Angew. Chem. 2018, 130, 11475.
- [7]. Ojha DP, Prabhu KR, Org. Lett 2016, 18, 432. [PubMed: 26760916]
- [8]. For the use of enantiomeric catalysts with chiral substrates to effect regiodivergent catalytic reactions, see: Funken N, Zhang Y-Q, Gansäuer A, Chem. - Eur. J 2017, 23, 19. [PubMed: 27813183]
- [9]. For examples of regiodivergent, enantioselective, metal-catalyzed reactions, see: a) Dalton DM, Rappé AK, Rovis T, Chem. Sci 2013, 4, 2062; [PubMed: 23671790] b) Cahard E, Male HPJ, Tissot M, Gaunt MJ, J. Am. Chem. Soc 2015, 137, 7986; [PubMed: 26090564] c) Richmond E, Khan IU, Moran J, Chem. - Eur. J 2016, 22, 12274; [PubMed: 27461524] d) Xu G, Zhao H, Fu B, Cang A, Zhang G, Zhang Q, Xiong T, Zhang Q, Angew. Chem. Int. Ed 2017, 56, 13130; Angew. Chem. 2017, 129, 13310; e) Zhou H, Wei Z, Zhang J, Yang H, Xia C, Jiang G, Angew. Chem. Int. Ed 2017, 56, 1077; Angew. Chem. 2017, 129, 1097; f) Liang L, Niu H-Y, Xie M-S, Qu G-R, Guo H-M, Org. Chem. Front 2018, 5, 3148; g) Xu S, Zhang Z-M, Xu B, Liu B, Liu Y, Zhang J, J. Am. Chem. Soc 2018, 140, 2272. [PubMed: 29303569]
- [10]. The absolute configuration for (R)-4a was determined by conversion to (R)-2,3-diphenylpropanol; see SI.
- [11]. The absolute configuration for (R)-5a was determined by conversion to (R)-2-phenylpropane-1,2-diol; see SI.
- [12]. Other phenyl phosphites also afford predominantly β -boration. For example, the (4-trifluoromethyl)phenyl phosphite T2c (Ar = 4-MeC₆H₄) gives (1:8 γ : β) with enantioselectivity favoring re-face addition (92:8 er), and the 4-cyanophenyl phosphite T2d (Ar = 4-MeC₆H₄) is similar.
- [13]. The metal-to-ligand ratio (Rh:L) employed varies on a case-by-case basis to maximize enantioselectivity, see SI.
- [14]. Competing alkene reduction typically accounts for low yields of boronic ester; see Shoba VM, Takacs JM, J. Am. Chem. Soc 2017, 139, 5740. [PubMed: 28394591]
- [15]. In scaling up to 2.65 mmol of 3a, a reduced catalyst (0.5 mol%) and 1:1 Rh:L ratio afforded: 4a (70%, 97:3 er) using T1a and 5a (81% 93:7 er) using T2b.
- [16]. a) Lennox AJJ, Lloyd-Jones GC, Angew. Chem. Int. Ed 2012, 51, 9385; Angew. Chem. 2012, 124, 9519.
- [17]. Bagutski V, Ros A, Aggarwal VK, Tetrahedron 2009, 65, 9956.
- [18]. Lee SJ, Gray KC, Paek JS, Burke MD, J. Am. Chem. Soc 2008, 130, 466. [PubMed: 18081295]
- [19]. a) Sadhu KM, Matteson DS, Organometallics 1985, 4, 1687; b) Thomas SP, French RM, Jheengut V, Aggarwal VK, Chem. Rec 2009, 9, 24. [PubMed: 19243084]
- [20]. a) Sonawane RP, Jheengut V, Rabalakos C, Larouche-Gauthier R, Scott HK, Aggarwal VK, Angew. Chem. Int. Ed 2011, 50, 3760; Angew. Chem. 2011, 123, 3844.
- [21]. a) Ros A, Aggarwal VK, Angew. Chem. Int. Ed 2009, 48, 6289; Angew. Chem. 2009, 121, 6407.
- [22]. Bonet A, Odachowski M, Leonori D, Essafi S, Aggarwal VK, Nat. Chem 2014, 6, 584. [PubMed: 24950327]
- [23]. Nave S, Sonawane RP, Elford TG, Aggarwal VK, J. Am. Chem. Soc 2010, 132, 17096. [PubMed: 21080646]
- [24]. Mlynarski SN, Karns AS, Morken JP, J. Am. Chem. Soc 2012, 134, 16449. [PubMed: 23002712]
- [25]. a) Collins BSL, Wilson CM, Myers EL, Aggarwal VK, Angew. Chem. Int. Ed 2017, 56, 11700; Angew. Chem. 2017, 129, 11860.
- [26]. Chakrabarty S, Palencia H, Morton MD, Carr RO, Takacs JM, Chem. Sci 2019, DOI: 10.1039/c8sc05613g.

Prior Work by Prabhu: Regiodivergent Hydroboration **This Work: Regiodivergent Enantioselective Hydroboration**

Figure 1.
Regiodivergent hydroboration of alkenes.

Figure 2.

Comparison of phenyl and pentafluorophenyl phosphites for a series of TADDOL backbone derivatives. Notes: the structures shown correlate to the major enantiomer from HPLC, except the product from **T5b** which arises from the opposite sense of π -facial selectivity.

Figure 3A

X =	F	Cl ^a	Br ^a	CO ₂ Me ^b	CF ₃ ^a	CH ₃	OMe	NMe ₂	N(H)Boc
	4b 74%, 97:3 er (<5% β)	4c 65%, 98:2 er (<5% β)	4d 75%, 97:3 er (<5% β)	4e 70%, 94:6 er (<5% β)	4f 76%, 94:6 er (<5% β)	4g 63%, 98:2 er (8% β)	4h 57%, 95:5 er (19% β)	4i 57%, 94:6 er (14% β)	4j 52%, 93:7 er (5% β)
	5b 73%, 96:4 er (13% γ)	5c 64%, 98:2 er (<5% γ)	5d 76%, 96:4 er (8% γ)	5e 78%, 96:4 er (<5% γ)	5f 75%, 98:2 er (7% γ)	5g 66%, 91:9 er (18% γ)	5h 76%, 86:14 er (5% γ)	5i^c 60%, 80:20 er (5% γ)	5j 65%, 94:6 er (5% γ)

Figure 3B

Figure 3C

Figure 3A-C.

Substrate scope: **A**. 4-Substituted phenyl derivatives; **B**. 3-Substituted phenyl and heteroaromatic derivatives; **C**. *ortho*-Substituted aryl and alkyl derivatives. Reported yields and enantiomer ratios are for the isolated alcohol after oxidation. **DG** = (Me₂C=N-O). Notes: ^a used 1:1 Rh:L ratio; ^b yield of boronic ester before oxidation with NaBO₃-H₂O; ^c Compounds **5i**, **5k**, and **5l** exhibited an opposite sign of optical rotation and order of elution on chiral HPLC; see SI.

Figure 4.

Versatility of the primary and tertiary boronic esters **4a** and **5a** [DG = (Me₂C=N-O)].

Reaction Conditions: (a) CsF or KF, MeCN/H₂O; (b) BCl₃, DCM; (c) BCl₃, DCM; MIDA, DMSO; (d) LiCH₂Cl, -78 °C Et₂O; (e) BCl₃, DCM; Raney Ni, H₂, MeOH/THF; (f) NaBH₃CN, MeOH/HCl; (g) LiC(OEt)=CH₂, THF, -78 °C; I₂; NaOMe, MeOH; (h) KF, MeCN/H₂O; O₂NC₆H₄CHO, 5 mol% [Rh(cod)Cl]₂, dioxane, 80 °C; DMP, DCM; (i) **12** or **13**, HCl/H₂O/MeOH, 40 °C; (j) thiophene, nBuLi, THF, -78 °C; NBS; (k) PhBr, 15 mol% Pd(OAc)₂, 15 mol% BINAP, KOH, THF/H₂O, 100 °C; (l) TBAF, toluene; (m) MeON(H)Li, THF, 60 °C; Boc₂O; (n) H₂O₂, NaOH, MeOH/H₂O; Raney Ni, H₂