Published in final edited form as:

Prev Med. 2019 June ; 123: 84–90. doi:10.1016/j.ypmed.2019.03.001.

Does optimal parenting style help offspring maintain healthy weight into mid-life?

Ying Chen, ScD^{1,2}, Ichiro Kawachi, MD, Ph.D³, Lisa F. Berkman, Ph.D^{2,3,4,5}, Claudia Trudel-Fitzgerald, Ph.D^{3,6}, Laura D. Kubzansky, Ph.D^{3,6}

¹Human Flourishing Program, Institute for Quantitative Social Science, Harvard University;

²Department of Epidemiology, Harvard T.H. Chan School of Public Health;

³Department of Social and Behavioral Sciences, Harvard T.H. Chan School of Public Health;

⁴Department of Global Health and Population, Harvard T.H. Chan School of Public Health;

⁵Center for Population and Development Studies, Harvard University;

⁶Lee Kum Sheung Center for Health and Happiness, Harvard T.H. Chan School of Public Health.

Abstract

An authoritative parenting style is generally associated with healthier body weight in children and adolescents. However, whether the protective effect of an authoritative style on offspring body weight may persist into adulthood has seldom been investigated. In this study we examined the longitudinal association between parenting style and body mass index (BMI) change in mid-life. Longitudinal data from the Midlife in the United States Study (N=3,929) were analyzed using generalized estimating equations, adjusting for a range of relevant covariates. Parenting styles were assessed at phase I (1995–1996) using items measuring parental warmth and control, while BMI was assessed at phases I and II (2004–2006). Four parenting styles were derived following prior research: authoritative, authoritarian, permissive, and uninvolved styles. Compared to an authoritative style, an authoritarian style was associated with 14% higher increase in the standardized BMI change score ($\beta=0.14$, 95% confidence interval: 0.03, 0.26). While there was suggestive evidence that an uninvolved versus authoritative style might also be associated with greater BMI increase, we found no differences between a permissive and authoritative style. This study suggested that the protective effect of an authoritative parenting style on offspring body weight may persist well into mid-life, particularly as compared to the authoritarian style and possibly the uninvolved style. Such work may reinforce the importance of a public health focus on improving parenting practices and suggest the value of implementing parenting programs, as one strategy for increasing the likelihood that individuals can maintain healthy weight well into adulthood.

Corresponding author: Ying Chen, ScD, 129 Mt Auburn Street, Cambridge, MA 02138. Phone: 617-945-9001. yingchen@fas.harvard.edu.

Conflict of Interest: The authors declare no conflict of interest.

Keywords

Parenting styles; Parental warmth; Parental control; Body weight; Lifecourse

INTRODUCTION

Obesity has been identified as a public health crisis worldwide (NCD Risk Factor Collaboration, 2017). In addition to targeting individual-level risk factors for obesity such as unhealthy diet and sedentary lifestyle, the Institute of Medicine has called for a population-based integrative approach for obesity prevention that also considers modifiable factors in the broader social environment (Committee on Accelerating Progress in Obesity Prevention, 2012).

Family represents individuals' immediate social surroundings and is potentially a critical source of social support that influences health (Alvarez et al., 2016). In particular, parenting practices may shape offspring health and well-being over the lifecourse (Britto et al., 2017). One of the most widely-studied parenting practices is parenting style. Based on levels of parental warmth and parental control, prior researchers (Maccoby and Martin, 1983) identified four general parenting styles: the authoritative (high in both warmth and control), authoritarian (low in warmth and high in control), permissive (high in warmth and low in control) and uninvolved style (low in both warmth and control). Empirical evidence generally suggests that the authoritative style is associated with better offspring health and well-being compared to other parenting styles (Pinquart, 2016; Pinquart, 2017a; Pinquart, 2017b).

Recent research, though mostly cross-sectional, has begun to consider parenting styles in relation to childhood obesity. A number of reviews have suggested that an authoritative style is associated with healthier body weight and weight-related behaviors in children and adolescents, as compared to other parenting styles (Sleddens et al., 2011; Sokol et al., 2017; Vollmer and Mobley, 2013). In comparison, only a handful of studies have explored whether the protective effects of an authoritative style on maintaining healthy body weight may persist beyond adolescence, but the limited evidence does suggest a lingering beneficial influence in young adulthood (Fuemmeler et al., 2012). However, to our knowledge whether the association may extend into middle or late adulthood has never been examined. From a lifecourse perspective, parental influences may shape offspring health not only in childhood but also well into adulthood (Holt-Lunstad et al., 2017). Children often model dietary and exercising behaviors from their parents, and such behavioral patterns may persist into adulthood and exert long-term health effects (Sokol et al., 2017). In fact, obese children are at substantially higher risk of staying obese when they grow up into adults (Simmonds et al., 2016).

Prior studies on parenting styles and offspring body weight often considered a limited range of confounders, generally including factors such as demographic characteristics and socioeconomic status [SES] (Sokol et al., 2017). Other factors may also have important linkages to parenting styles and body weight but few studies have included them. For example, childhood familial characteristics such as family structure (Sokol et al., 2017),

parental abuse (Danese and Tan, 2014), residential stability (Anderson et al., 2014; Jones, 2015) and family religiousness (Bornstein et al., 2017; Goeke-Morey and Cummings, 2017) have been linked with either parenting or children's risk of obesity. However, whether they may confound the association of parenting style with offspring body weight is understudied.

This study examined the longitudinal association between parenting styles and offspring body weight over a 9-year follow-up in mid-life, controlling for a wide range of childhood family environment characteristics as potential confounders (e.g., family SES, family structure, parental abuse, residential stability, family religiousness). Parenting styles were recalled when offspring were middle-aged. Because parenting styles were retrospectively reported, as a sensitivity analysis we additionally adjusted for a number of adulthood characteristics (e.g., adulthood SES, depression, chronic health conditions) that may affect how parenting styles were recalled (Widom et al., 2004). We hypothesized that the authoritative parenting style would be associated with healthier body weight in mid-life, compared to other parenting styles.

METHODS

Study Sample

Data were from the Midlife in the United States (MIDUS) study. MIDUS was initiated in 1994–1995 to study health and well-being in mid-life. At the first wave (MIDUS I), 7,108 non-institutionalized individuals aged between 25 to 74 years across the United States were enrolled through a random selection process. Participants were invited to participate in a phone interview, and then received a self-administered questionnaire (SAQ). A second wave of the study (MIDUS II) took place in 2004–2005, which followed up 70% (N=4,963) of the original participants. Details of the MIDUS recruitment and follow-up procedures were reported elsewhere (Brim et al., 2004; Shaw et al., 2004).

Because parenting styles and body weight were only assessed in the SAQ, the analytic sample for the present study was drawn from respondents who completed the SAQ at both waves (N=3,929; 569 of them either siblings or twins, and we adjusted for potential clustering by sibling status in the analyses). This study was approved by the institutional review board (IRB) at the authors' institution. The original MIDUS study was approved by the IRB at participating institutions, and all participants provided informed consent (Radler, 2014).

Measures

Independent Variables

Parental warmth. At phase I, participants recalled parental warmth during their years of growing up. A six-item Parental Affection Scale (Rossi, 2001) (Table S1) was used to assess maternal and paternal warmth separately (e.g., “How much love and affection did your mother/father give you?”). Response categories ranged from 1 (a lot) to 4 (not at all). When appropriate, items were reverse coded so that a higher score reflected greater warmth. Maternal and paternal warmth scores were calculated for participants with valid data on at least half of the scale items, by averaging responses across items. Following prior research,

an overall parental warmth score was calculated by averaging the maternal and paternal warmth scores (Rothrauff et al., 2009). The scale had good internal consistency reliability in this sample ($\alpha = .89$ for the maternal scale, $\alpha = .91$ for the paternal scale). It also showed positive associations with multiple domains of well-being in prior work (Chen et al., 2018; Moran et al., 2018), providing some evidence for construct validity.

Parental control.: Participants recalled parental control during their years of growing up at phase I. A three-item Parental Control Scale (Rossi, 2001) (Table S1) was used to assess maternal and paternal control separately (e.g., “How strict was your mother/father with her/his rules for you?”). Response options ranged from 1 (a lot) to 4 (not at all). Responses were scored so that a higher score represented greater control. Maternal and paternal control scores were calculated for participants with valid data on at least half of the scale items, by averaging responses across items. Following prior research, an overall parental control score was calculated by averaging the maternal and paternal control scores (Rothrauff et al., 2009). The scale showed acceptable internal consistency reliability in this sample ($\alpha = .74$ for both the maternal and paternal scale). In prior work (Enns et al., 2002), greater paternal control measured with this scale was associated with lower risk of externalizing disorders in male children, which provided some evidence for construct validity.

Parenting style.: Four parenting style typologies were created based on distinct constellations of parental warmth and control (Maccoby and Martin, 1983). As specific cut-points for these measures have not been validated, we followed common practice in epidemiologic studies of psychosocial characteristics (Kubzansky et al., 2014) and characterized individuals scoring in the top tertile of the distribution of scores on each subscale as being distinctively high in warmth and in control (Figure S1). The authoritative style included participants scoring in the top tertile of both warmth and control (14.11%); the authoritarian style included respondents scoring in the bottom and middle tertiles of warmth but in the top tertile of control (18.07%); the permissive style included those scoring in the top tertile of warmth but in the bottom and middle tertiles of control (18.37%); and the uninvolved style included those in the bottom and middle tertiles of both warmth and control (49.45%). To evaluate the sensitivity of any associations to these particular cut-points, we also followed another categorization approach from prior literature (Rothrauff et al., 2009) by using a median split to define high versus low warmth or control (Figure S2).

Dependent variable

Body weight.: At both phases participants reported their height and weight, based on which body mass index (BMI, kg/m^2) was calculated. The self-reported BMI showed high concordance ($r=0.92$) with BMI measured by clinicians or trained staffs in a subgroup who participated in a biomarker project at phase II ($N=1,255$, the subgroup was comparable to the full sample in terms of self-reported BMI and major demographic and health-related characteristics) (Dienberg Love et al., 2010). We calculated a BMI change score (used as a continuous variable) by subtracting BMI at phase I from BMI at phase II. To minimize influence of extreme outliers, the score was winsorized at the 1st and 99th percentile (i.e., respondents with a score < the 1st percentile or > the 99th percentile were assigned the value for the 1st and 99th percentile). We standardized the BMI change score (mean=0, standard

deviation=1) and used the standardized score as the dependent variable in all models, to facilitate comparison of effect estimates across studies in which BMI may follow a different distribution (Landis, 2005).

Covariates

Demographic factors.: Demographic covariates included participant age (in years), sex (male, female), and race (white, black, other races), all self-reported at phase I.

Childhood characteristics.: All childhood characteristics were recalled at phase I. Childhood SES was assessed with the highest educational attainment of parents (less than high school, high school, some college, college degree or higher) (Miller et al., 2011). Severe parental abuse was assessed using one question from the Conflict Tactics Inventory (Straus, 1979): “During your childhood, how often did your mother/father kicked, bit, or hit you with a fist or an object, beat you up, choked, burned or scalded you”. Response options ranging from 1 (often) to 4 (never), and responses were coded so a higher score represented greater abuse. An overall parental abuse score was created by averaging the maternal and paternal abuse scores, and was used as a continuous variable (Savla et al., 2013). Two-parent family structure was assessed with a single question: “Did you live with both biological parents up until you were 16?” (yes, no). Childhood residential area was also queried (rural, small town, medium-sized town, suburbs, city, moved around). Childhood residential stability was assessed with a single question: “How many times during your childhood did you move to a totally new neighborhood or town?” Participants reporting less than 3 residential moves were considered as having residential stability (Bures, 2003). Family religiousness was also queried: “How important was religion in your home when you were growing up?”, with responses ranging from 1 (not at all important) to 4 (very important). The measure was used as a continuous variable.

Adulthood characteristics.: Adulthood covariates were assessed at phase I. Participants reported their current marital status (married, divorced/separate, widowed, never married) and their own educational attainment (less than high school, high school, some college, college degree or higher). Household income was also self-reported (in U.S. dollars, income greater than \$300,000 was recoded as \$300,000 to minimize risk of deductive disclosure), and quartiles of household income were created. Major depression (yes, no) over past year was assessed with the validated Composite International Diagnostic Interview Short Form (CIDI-SF) (Aalto-Setälä et al., 2002; Kessler et al., 1998). Participants who reported ever having or taking medication for any of the following conditions were considered as having chronic physical health conditions (yes, no): cancer, heart attack, diabetes, and stroke.

Statistical Analyses

All statistical analyses were performed in SAS 9.4 ($p < .05$ significance level, two-tailed). Chi-square tests and analysis of variance tests were used to examine distribution of participant characteristics by parenting styles.

Generalized estimating equation models (GEE) were used to examine the association between parenting styles and BMI change, adjusting for clustering by sibling status. The

base model adjusted for age, sex and race. A second model further adjusted for childhood SES. The third model additionally accounted for a range of other childhood familial factors. We performed several sensitivity analyses. Because parenting styles were retrospectively reported in mid-life, to account for the possibility that adulthood factors might influence how parenting styles were recalled, we additionally adjusted for adulthood characteristics concurrently assessed with parenting styles. We also reanalyzed the primary sets of models with parenting styles created using the median split to define high versus low levels of parental warmth and control.

Because results of the primary analyses suggested that the strongest differences were between parenting styles low in warmth (i.e., authoritarian, uninvolved style) and high in warmth (i.e., authoritative style), we performed a post-hoc analysis to explore whether effects of parenting styles were primarily driven by the dimension of parental warmth. Specifically, we reanalyzed the primary models first including parental warmth and then including parental control as the independent variable in separate models. We also conducted stratified analyses to consider effects of parental warmth within low, moderate, or high levels of parental control.

In the full analytic sample (N=3,929), 10 participants were missing data on parenting style, 301 were missing data on body weight at baseline or the follow-up, and another 261 were missing data on covariates. Complete-case analysis would result in a loss of 14.6% (n=572) of the participants. We performed a multivariate normal multiple imputation procedure (number of imputed datasets=5) to impute missing data on all variables, as it often provides more accurate estimates compared to other methods of handling missing data (Sterne et al., 2009). We also performed complete-case analysis as a sensitivity analysis.

RESULTS

Participants were predominantly white (93.48%) and slightly higher percentage female (55.48%), with the mean baseline age of 47.39 years (SD=12.43). The mean BMI increase was 1.24 kg/m² (SD=3.14) over an average of 9-years of follow-up. Participants generally reported high levels of parental warmth (mean=2.97, range: 1 to 4) and parental control (mean=3.00, range: 1 to 4). Descriptive analyses suggested participants with authoritative parents were more likely to report a two-parent family structure, high family religiousness and residential stability in childhood, and were more likely to be married and have no depression in adulthood, compared to those raised by authoritarian and permissive parents (Table 1).

Compared to those raised by authoritative parents, participants with authoritarian parents were 16% higher in the standardized score of BMI increase ($\beta=0.16$, 95% confidence interval [CI]: 0.05, 0.28), adjusting for age, sex and race (Table 2). Additionally adjusting for childhood SES ($\beta=0.16$, 95% CI: 0.05, 0.28) and other childhood covariates ($\beta=0.14$, 95% CI: 0.03, 0.26) did not change the association. Notably, the effect size of authoritative versus authoritarian parenting style was even larger than having the highest versus the lowest category of childhood family SES (i.e., parental education as college degree or higher versus less than high school). Moreover, the association remained robust in the sensitivity analysis

that further considered adulthood characteristics. To a lesser extent, the uninvolved versus authoritative style was also associated with a greater BMI increase, but the association was somewhat attenuated in the fully-adjusted model. In comparison, there was no difference in the permissive versus authoritative style in any model. Sensitivity analyses using the median split to define high versus low levels of parental warmth and control yielded similar results (Table S2). Further, the complete-case analysis also yielded somewhat attenuated but largely similar results (Table S3).

When considering individual dimensions of parenting separately, parental warmth was associated with substantially less BMI increase over time (Table 3), whereas parental control was not associated with BMI change ($p>0.05$ in all models, results not shown). However, the stratified analyses suggested that effects of parental warmth were evident only for individuals scoring in the top tertile of parental control (Table 3). Results of this post-hoc analysis were consistent with our primary analyses suggesting that the authoritative style was likely associated with healthier body weight, particularly as compared to the authoritarian style.

DISCUSSION

This is the first longitudinal study suggesting that the authoritative parenting style may exert a protective effect on offspring body weight well into mid-life, particularly as compared to the authoritarian style and also possibly the uninvolved style. However, there was no difference between the authoritative and permissive style. The post-hoc analyses on individual dimensions of parenting also indicated that parental warmth may help offspring maintain healthy body weight, only when a high level of parental control was also present.

Findings of this study are consistent with prior work in younger populations which generally suggested that the authoritative parenting style was associated with healthier body weight in children and adolescents, as compared to other parenting styles (Sokol et al., 2017). This study expands prior literature by adding evidence that the protective effect of the authoritative style may persist into mid-life. Parental warmth may provide children with a sense of emotional security and self-worth. Such experiences could facilitate the formation of developmental assets such as social integration and self-regulation, which may help increase resilience and reduce unhealthy coping strategies (e.g., binge eating) under stressful situations (Holmes, 2014). In addition, authoritative parents may also teach children healthy practices and set reasonable expectations on their behaviors. Some of these behavioral patterns including diet and exercise habits may persist into adulthood and affect one's weight trajectories (Kwon et al., 2015; Watts et al., 2018). However, effects may be tempered by influences occurring later in life. Thus, while parental influences set a developmental trajectory, social relationships developed in later life may redirect the course. For instance, as individuals transition to adolescence and adulthood, peers and partners may reshape one's behavioral perceptions and patterns (e.g., smoking, diet), and offset early parental influences (Scalici and Schulz, 2014).

Compared to the authoritative style, the elevated BMI increase associated with the authoritarian style was particularly pronounced, and this finding may be attributable to the

synergistic effects of low parental warmth and high parental control. Although the dimension of parental control was not associated with BMI change in this middle-aged sample, prior studies have separately linked low parental warmth and excessive parental control with increased risk of childhood obesity (Gartstein et al., 2018; Larsen et al., 2015; Rhee et al., 2016). Low parental warmth may result in higher emotional distress and low self-efficacy, while high parental control may lead to lack of capacity for reasoning and self-regulation. Each of these sequelae could increase risk of adopting unhealthy weight-related behaviors to cope with distress (Larsen et al., 2015; Topham et al., 2011). Somewhat unexpectedly, evidence suggesting the uninvolved versus authoritative parental style was associated with higher offspring BMI increase was modest in this study. This might be due in part to participant characteristics specific to this sample, who generally reported high levels of parental warmth and control. The limited variation might attenuate our ability to detect effects of an uninvolved style, if any.

This study has several limitations. First, parenting style and other childhood characteristics were retrospectively reported in mid-life, which may be subject to recall bias. However, there has been evidence suggesting that concerns about retrospectively reported childhood experiences may not be as significant as previously thought (Patten et al., 2015). For instance, in prior work, childhood maltreatment experiences as recalled in adulthood showed high concordance with official records such as data from Child Protective Services and medical records, suggesting reasonable validity (Pinto et al., 2014; Shin et al., 2013). In addition, the longitudinal study design and sensitivity analyses adjusting for adulthood characteristics provide reassurance that the observed associations are not entirely due to bias. Second, although this study adjusted for a wide range of childhood characteristics, residual confounding is possible by other factors for which information was not available such as childhood health status and parental mental health. Due to the lack of temporal separation in the assessments of childhood parenting style and adulthood health-related characteristics, we did not perform a causal mediation analysis (VanderWeele, 2015), and as a result it remains unclear whether parenting styles may have a direct effect on offspring adulthood weight, or only indirectly through shaping children's later life experiences. Finally, MIDUS participants did not comprise a nationally representative sample (e.g., predominantly white), thus results of this study may not be generalizable to other populations. These limitations were, however, balanced by strengths including the prospective study design, the large sample size with a 9-year follow-up, and the rigorous confounding control in this study.

The World Health Organization has called for parenting programs at the population level (World Health Organization, 2007, 2009), yet the progress on implementing large-scale parenting programs has been relatively slow (Chu et al., 2012). Such programs seek to improve general parenting styles (e.g., reducing coercive discipline practice) and specific parenting practices (e.g., improving skills in communicating about health), and have been linked to multiple better health and well-being outcomes in children including reduced risk of obesity (Gerards et al., 2012; Hubbs-Tait et al., 2016; Yap et al., 2016). This study further suggests that the beneficial effects of such programs, if any, on children's body weight may persist into mid-life, and helps strengthen the case for implementing parenting programs at the population level. Further research on parenting styles and the broader family

environment can help inform more targeted interventions, and identify resources within the family that may be targeted for improving obesity prevention and control.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

ACKNOWLEDGEMENTS

The original Midlife in the United States study was supported by the John D. and Catherine T. MacArthur Foundation Research Network on Successful Midlife Development. Follow-up data collection was supported by the National Institute on Aging (P01-AG020166). Funding agencies had no role in the data collection, analysis, or interpretation; nor were they involved in the writing or submission of this publication. We thank the staff of the clinical research centers at Georgetown University, University of Wisconsin-Madison, and University of California, Los Angeles for their support in conducting this study. CTF received a postdoctoral fellowship from the Fonds de Recherche du Québec-Santé.

REFERENCES

- Aalto-Setälä T, Haarasilta L, Marttunen M, Tuulio-Henriksson A, Poikolainen K, Aro H, Lonnqvist J, 2002 Major depressive episode among young adults: CIDI-SF versus SCAN consensus diagnoses. *Psychol Med* 32:1309–14. [PubMed: 12420900]
- Alvarez EC, Kawachi I, Romani JR, 2016 Family social capital and health – a systematic review and redirection. *Sociology of Health & Illness* 39:5–29. [PubMed: 27813121]
- Anderson S, Leventhal T, Dupéré V, 2014 Residential mobility and the family context: A developmental approach. *Journal of Applied Developmental Psychology* 35:70–78.
- Bornstein MH, Putnick DL, Lansford JE, Al-Hassan SM, Bacchini D, Bombi AS, Chang L, Deater-Deckard K, Di Giunta L, et al., 2017 ‘Mixed blessings’: parental religiousness, parenting, and child adjustment in global perspective. *Journal of Child Psychology and Psychiatry* 58:880–92. [PubMed: 28244602]
- Brim OG, Ryff CD, Kessler RC, 2004 How healthy are we? : a national study of well-being at midlife. University of Chicago Press, Chicago.
- Britto PR, Lye SJ, Proulx K, Yousafzai AK, Matthews SG, Vaivada T, Perez-Escamilla R, Rao N, Ip P, et al., 2017 Nurturing care: promoting early childhood development. *The Lancet* 389:91–102.
- Bures RM, 2003 Childhood Residential Stability and Health at Midlife. *American Journal of Public Health* 93:1144–48. [PubMed: 12835200]
- Chen Y, Kubzansky LD, VanderWeele TJ, 2018 Parental warmth and flourishing in mid-life. *Soc Sci Med* 220:65–72. [PubMed: 30396119]
- Chu JT, Farruggia SP, Sanders MR, Ralph A, 2012 Towards a public health approach to parenting programmes for parents of adolescents. *J Public Health (Oxf)* 34 Suppl 1:i41–7. [PubMed: 22363030]
- Committee on Accelerating Progress in Obesity Prevention, Food and Nutrition Board, Institute of Medicine, 2012 in: Glickman D, Parker L, Sim LJ, Del Valle Cook H, Miller EA (Eds.), Accelerating Progress in Obesity Prevention: Solving the Weight of the Nation. National Academies Press, Washington, DC.
- Danese A, Tan M, 2014 Childhood maltreatment and obesity: systematic review and meta-analysis. *Mol Psychiatry* 19:544–54. [PubMed: 23689533]
- Dienberg Love G, Seeman TE, Weinstein M, Ryff CD, 2010 Bioindicators in the MIDUS national study: protocol, measures, sample, and comparative context. *Journal of aging and health* 22:1059–80. [PubMed: 20876364]
- Enns MW, Cox BJ, Clara I, 2002 Parental bonding and adult psychopathology: results from the US National Comorbidity Survey. *Psychol Med* 32:997–1008. [PubMed: 12214798]

- Fuemmeler BF, Yang C, Costanzo P, Hoyle RH, Siegler IC, Williams RB, Ostbye T, 2012 Parenting styles and body mass index trajectories from adolescence to adulthood. *Health Psychol* 31:441–9. [PubMed: 22545979]
- Gartstein MA, Seamon E, Thompson SF, Lengua LJ, 2018 Parenting matters: Moderation of biological and community risk for obesity. *J Appl Dev Psychol* 56:21–34. [PubMed: 29910526]
- Gerards SM, Dagnelie PC, Jansen MW, van der Goot LO, de Vries NK, Sanders MR, Kremers SP, 2012 Lifestyle Triple P: a parenting intervention for childhood obesity. *BMC Public Health* 12:267. [PubMed: 22471971]
- Goeke-Morey MC, Cummings EM, 2017 Religiosity and parenting: recent directions in process-oriented research. *Current opinion in psychology* 15:7–12. [PubMed: 28813272]
- Holmes J, 2014 *John Bowlby and attachment theory*, 2nd ed. Routledge, London, UK.
- Holt-Lunstad J, Robles TF, Sbarra DA, 2017 Advancing social connection as a public health priority in the United States. *American Psychologist* 72:517. [PubMed: 28880099]
- Hubbs-Tait L, Kimble A, Hingle M, Novotny R, Fiese B, 2016 Systematic Review of Child Obesity Prevention and Treatment Trials Addressing Parenting. *The FASEB Journal* 30:1155.6–55.6. [PubMed: 26644349]
- Jones A, 2015 Residential mobility and trajectories of adiposity among adolescents in urban and non-urban neighborhoods. *J Urban Health* 92:265–78. [PubMed: 25801487]
- Kessler RC, Andrews G, Mroczek D, Ustun B, Wittchen H-U, 1998 The World Health Organization Composite International Diagnostic Interview short-form (CIDI-SF). *International Journal of Methods in Psychiatric Research* 7:171–85.
- Kubzansky LD, Winning A, Kawachi I, 2014 Affective States and Health, in: Berkman LF, Kawachi I, Glymour MM (Eds.), *Social Epidemiology: New perspectives on social determinants of global population health*. Oxford University Press, New York, pp. 320–64.
- Kwon S, Janz KF, Letuchy EM, Burns TL, Levy SM, 2015 Active lifestyle in childhood and adolescence prevents obesity development in young adulthood. *Obesity* 23:2462–69. [PubMed: 26538514]
- Landis R, 2005 Standardized regression coefficient, in: Everitt B, Howell D (Eds.), *Encyclopedia of Statistics in Behavioral Science*. John Wiley & Sons Ltd, Chichester, p. 1892.
- Larsen JK, Hermans RC, Sleddens EF, Engels RC, Fisher JO, Kremers SP, 2015 How parental dietary behavior and food parenting practices affect children's dietary behavior. Interacting sources of influence? *Appetite* 89:246–57. [PubMed: 25681294]
- Maccoby E, Martin J, 1983 Socialization in the context of the family: Parent–child interaction, in: Mussen P, Hetherington E (Eds.), *Handbook of Child Psychology*, 4th ed. Wiley, New York, pp. 1–101.
- Miller GE, Lachman ME, Chen E, Gruenewald TL, Karlamangla AS, Seeman TE, 2011 Pathways to resilience: maternal nurturance as a buffer against the effects of childhood poverty on metabolic syndrome at midlife. *Psychol Sci* 22:1591–9. [PubMed: 22123777]
- Moran KM, Turiano NA, Gentzler AL, 2018 Parental warmth during childhood predicts coping and well-being in adulthood. *J Fam Psychol* 32:610–21. [PubMed: 29708363]
- NCD Risk Factor Collaboration, 2017 Worldwide trends in body-mass index, underweight, overweight, and obesity from 1975 to 2016: a pooled analysis of 2416 population-based measurement studies in 128.9 million children, adolescents, and adults. *Lancet* 390:2627–42. [PubMed: 29029897]
- Patten SB, Wilkes TCR, Williams JVA, Lavorato DH, el-Guebaly N, Schopflocher D, Wild C, Colman I, Bulloch AGM, 2015 Retrospective and prospectively assessed childhood adversity in association with major depression, alcohol consumption and painful conditions. *Epidemiology and Psychiatric Sciences* 24:158–65. [PubMed: 24480045]
- Pinquart M, 2016 Associations of parenting styles and dimensions with academic achievement in children and adolescents: A meta-analysis. *Educational Psychology Review* 28:475–93.
- Pinquart M, 2017a Associations of parenting dimensions and styles with externalizing problems of children and adolescents: An updated meta-analysis. *Dev Psychol* 53:873–932. [PubMed: 28459276]

- Pinquart M, 2017b Associations of parenting dimensions and styles with internalizing symptoms in children and adolescents: A meta-analysis. *Marriage & Family Review* 53:613–40.
- Pinto R, Correia L, Maia Â, 2014 Assessing the reliability of retrospective reports of adverse childhood experiences among adolescents with documented childhood maltreatment. *Journal of Family Violence* 29:431–38.
- Radler BT, 2014 The Midlife in the United States (MIDUS) Series: A National Longitudinal Study of Health and Well-being. Open Health Data 2.
- Rhee KE, Jelalian E, Boutelle K, Dickstein S, Seifer R, Wing R, 2016 Warm parenting associated with decreasing or stable child BMI during treatment. *Childhood Obesity* 12:94–102. [PubMed: 26895374]
- Rossi AS, 2001 Caring and doing for others: Social responsibility in the domains of family, work, and community. University of Chicago Press.
- Rothrauff TC, Cooney TM, An JS, 2009 Remembered parenting styles and adjustment in middle and late adulthood. *Journals of Gerontology Series B: Psychological Sciences and Social Sciences* 64:137–46.
- Savla JT, Roberto KA, Jaramillo-Sierra AL, Gambrel LE, Karimi H, Butner LM, 2013 Childhood abuse affects emotional closeness with family in mid- and later life. *Child abuse & neglect* 37:388–99. [PubMed: 23369347]
- Scalici F, Schulz PJ, 2014 Influence of perceived parent and peer endorsement on adolescent smoking intentions: parents have more say, but their influence wanes as kids get older. *PLoS One* 9:e101275. [PubMed: 24991921]
- Shaw BA, Krause N, Chatters LM, Connell CM, Ingersoll-Dayton B, 2004 Emotional support from parents early in life, aging, and health. *Psychol Aging* 19:4–12. [PubMed: 15065927]
- Shin SH, Miller DP, Teicher MH, 2013 Exposure to childhood neglect and physical abuse and developmental trajectories of heavy episodic drinking from early adolescence into young adulthood. *Drug Alcohol Depend* 127:31–8. [PubMed: 22749563]
- Simmonds M, Llewellyn A, Owen CG, Woolacott N, 2016 Predicting adult obesity from childhood obesity: a systematic review and meta-analysis. *Obes Rev* 17:95–107. [PubMed: 26696565]
- Sleddens EF, Gerards SM, Thijs C, de Vries NK, Kremers SP, 2011 General parenting, childhood overweight and obesity-inducing behaviors: a review. *Int J Pediatr Obes* 6:e12–27. [PubMed: 21657834]
- Sokol RL, Qin B, Poti JM, 2017 Parenting styles and body mass index: a systematic review of prospective studies among children. *Obes Rev* 18:281–92. [PubMed: 28086262]
- Sterne JA, White IR, Carlin JB, Spratt M, Royston P, Kenward MG, Wood AM, Carpenter JR, 2009 Multiple imputation for missing data in epidemiological and clinical research: potential and pitfalls. *BMJ* 338:b2393. [PubMed: 19564179]
- Straus MA, 1979 Measuring Intrafamily Conflict and Violence: The Conflict Tactics (CT) Scales. *Journal of Marriage and Family* 41:75–88.
- Topham GL, Hubbs-Tait L, Rutledge JM, Page MC, Kennedy TS, Shriver LH, Harriett AW, 2011 Parenting styles, parental response to child emotion, and family emotional responsiveness are related to child emotional eating. *Appetite* 56:261–64. [PubMed: 21232566]
- VanderWeele T, 2015 Explanation in causal inference: methods for mediation and interaction. Oxford University Press, New York, NY.
- Vollmer RL, Mobley AR, 2013 Parenting styles, feeding styles, and their influence on child obesogenic behaviors and body weight. A review. *Appetite* 71:232–41. [PubMed: 24001395]
- Watts A, Berge JM, Loth K, Larson N, Neumark-Sztainer D, 2018 The Transmission of Family Food and Mealtime Practices From Adolescence to Adulthood: Longitudinal Findings From Project EAT-IV. *J Nutr Educ Behav* 50:141–47 e1. [PubMed: 29054340]
- Widom CS, Raphael KG, DuMont KA, 2004 The case for prospective longitudinal studies in child maltreatment research: commentary on Dube, Williamson, Thompson, Felitti, and Anda (2004). *Child Abuse Negl* 28:715–22. [PubMed: 15261466]
- World Health Organization, 2007 Helping parents in developing countries improve adolescents' health. World Health Organization, Geneva.

- World Health Organization, 2009 Preventing violence through the development of safe, stable and nurturing relationships between children and their parents and caregivers. Series of briefings on violence prevention: the evidence. World Health Organization, Geneva, Switzerland.
- Yap MBH, Morgan AJ, Cairns K, Jorm AF, Hetrick SE, Merry S, 2016 Parents in prevention: A meta-analysis of randomized controlled trials of parenting interventions to prevent internalizing problems in children from birth to age 18. *Clin Psychol Rev* 50:138–58. [PubMed: 27969003]

Participant characteristics by parenting styles in childhood (The Midlife in the United States Study 1995/1996 – 2004/2005 questionnaire wave, N=3,929)

Participant Characteristics	Full Sample N=3,929	Parenting Styles in Childhood			P-Value
		Authoritative N = 553	Authoritarian N = 708	Permissive N = 720	
Baseline age, years, mean (SD)	47.39 (12.43)	47.44 (12.96)	46.88 (11.70)	48.38 (12.97)	0.04
Male, N (%)	1,749 (44.52)	522 (47.07)	294 (38.43)	390 (45.94)	0.002
Race, N (%)					<0.001
White	3,673 (93.48)	1,015 (91.52)	708 (92.55)	802 (94.46)	
Black	148 (3.77)	68 (6.13)	30 (3.92)	24 (2.83)	
Others	108 (2.75)	26 (2.34)	27 (3.53)	23 (2.71)	
Childhood characteristics					
Highest parental education, N (%)					0.06
Less than high school	985 (25.79)	283 (26.37)	220 (29.45)	172 (21.16)	
High school	1,377 (36.05)	379 (35.32)	260 (34.81)	310 (37.48)	
Some college	301 (15.73)	177 (16.50)	107 (14.32)	138 (16.69)	
College degree or higher	857 (22.43)	234 (21.81)	160 (21.42)	204 (24.67)	
Parental abuse score, mean (SD)	1.25 (0.55)	1.13 (0.39)	1.51 (0.75)	1.08 (0.28)	<0.001
Lived with biological parents, N (%)	3,174 (80.83)	969 (87.38)	605 (79.08)	694 (81.84)	<0.001
Childhood residential area, N (%)					<0.001
Rural	952 (24.55)	313 (28.58)	200 (26.42)	203 (24.08)	
Small town	1,003 (25.86)	291 (26.58)	185 (24.44)	209 (24.79)	
Medium-sized town	458 (11.81)	123 (11.23)	75 (9.91)	97 (11.51)	
Suburbs	606 (15.63)	158 (14.43)	116 (15.32)	127 (15.07)	
City	697 (17.97)	173 (15.80)	135 (17.83)	169 (20.05)	
Moved around	162 (4.18)	37 (3.38)	46 (6.08)	38 (4.51)	
Residential stability, N (%)	2,915 (74.84)	873 (79.15)	536 (70.71)	654 (77.58)	<0.001
Family religiousness, N (%)					<0.001
Religion not at all important	168 (4.28)	13 (1.17)	44 (5.75)	26 (3.07)	
Religion not very important	603 (15.37)	86 (7.77)	122 (15.95)	112 (13.21)	
Religion somewhat important	1,380 (35.19)	327 (29.54)	266 (34.77)	317 (37.38)	

Participant Characteristics	Full Sample N=3,929	Parenting Styles in Childhood			P-Value
		Authoritative N = 553	Authoritarian N = 708	Permissive N = 720	
Religion very important	1,771 (45.16)	681 (61.52)	333 (43.53)	393 (46.34)	
Adulthood characteristics					
Marital status, N (%)					<0.001
Married	2,814 (71.64)	854 (77.08)	532 (69.54)	597 (70.32)	826 (69.06)
Divorced/separated	537 (13.67)	111 (10.02)	124 (16.21)	103 (12.13)	197 (16.47)
Widowed	166 (4.23)	37 (3.34)	32 (4.18)	47 (5.54)	50 (4.18)
Never married	411 (10.46)	106 (9.57)	77 (10.07)	102 (12.01)	123 (10.28)
Educational attainment, N (%)					0.16
Less than high school	264 (6.72)	71 (6.40)	61 (7.97)	43 (5.06)	86 (7.19)
High school	1,099 (27.97)	320 (28.85)	209 (27.32)	234 (27.56)	331 (27.68)
Some college	1,145 (29.14)	323 (29.13)	224 (29.28)	231 (27.21)	366 (30.60)
College degree or higher	1,421 (36.17)	395 (35.62)	271 (35.42)	341 (40.16)	413 (34.53)
Household income quartiles, N (%)					0.95
Bottom quartile (\$0-\$32,499)	931 (24.24)	254 (23.43)	180 (24.13)	192 (23.13)	298 (25.40)
Second quartile (\$32,500-\$58,999)	986 (25.67)	280 (25.83)	194 (26.01)	208 (25.06)	303 (25.83)
Third quartile (\$59,000-\$98,999)	965 (25.12)	269 (24.82)	191 (25.60)	213 (25.66)	292 (24.89)
Top quartile (\$99,000-\$300,000+)	959 (24.97)	281 (25.92)	181 (24.26)	217 (26.14)	280 (23.87)
Major depression, N (%)	469 (11.94)	89 (8.03)	127 (16.60)	71 (8.36)	181 (15.13)
Any chronic health condition, N (%)	593 (15.09)	164 (14.79)	117 (15.29)	140 (16.49)	170 (14.21)
					0.55

Note. Percentages refer to the proportion of individuals within each parenting style category with that characteristic. p comes from χ^2 or analysis of variance tests.

Parenting styles in childhood and BMI change in mid-life (The Midlife in the United States Study 1995/1996–2004/2005 questionnaire wave, N=3,929)

Table 2.

	Model 1 β (95% CI)	Model 2 β (95% CI)	Model 3 β (95% CI)	Model 4 β (95% CI)
Parenting styles (Authoritative as ref)				
Authoritarian style	0.16 (0.05, 0.28) **	0.16 (0.05, 0.28) **	0.14 (0.03, 0.26) *	0.12 (0.01, 0.23) *
Permissive style	0.06 (−0.06, 0.17)	0.06 (−0.05, 0.18)	0.07 (−0.05, 0.18)	0.06 (−0.06, 0.17)
Uninvolved style	0.11 (0.01, 0.20) *	0.11 (0.01, 0.20) *	0.10 (−0.00, 0.20)	0.08 (−0.02, 0.18)
Age (standardized)	−0.21 (−0.24, −0.18) ***	−0.22 (−0.25, −0.19) ***	−0.22 (−0.25, −0.19) ***	−0.21 (−0.25, −0.17) ***
Male (female as ref)	−0.09 (−0.15, −0.02) **	−0.09 (−0.15, −0.02) **	−0.09 (−0.16, −0.02) **	−0.07 (−0.13, −0.002) *
Race (White as ref)				
Black	0.14 (−0.06, 0.33)	0.12 (−0.08, 0.32)	0.11 (−0.09, 0.31)	−0.10 (−0.10, 0.30)
Others	0.08 (−0.15, 0.31)	0.09 (−0.14, 0.31)	0.08 (−0.15, 0.31)	0.05 (−0.17, 0.28)
Childhood characteristics				
Parental education (<high school as ref)				
High school		0.01 (−0.09, 0.10)	0.02 (−0.08, 0.12)	0.02 (−0.07, 0.12)
Some college		−0.01 (−0.11, 0.09)	0.00 (−0.10, 0.11)	0.02 (−0.08, 0.13)
College degree or higher		−0.14 (−0.24, −0.04) **	−0.11 (−0.22, −0.01) *	−0.08 (−0.18, 0.03)
Parental abuse score (standardized)			0.04 (0.00, 0.07) *	0.03 (−0.01, 0.06)
Lived with biological parents (no as ref)				0.04 (−0.05, 0.12)
Childhood residential area (rural as ref)				
Small town			−0.03 (−0.12, 0.06)	−0.03 (−0.12, 0.06)
Medium-sized town			−0.06 (−0.18, 0.05)	−0.07 (−0.19, 0.05)
Suburbs			−0.06 (−0.16, 0.05)	−0.07 (−0.17, 0.04)
City			−0.07 (−0.17, 0.03)	−0.08 (−0.18, 0.03)
Moved around			−0.15 (−0.35, 0.05)	−0.15 (−0.35, 0.05)
Childhood residential stability (no as ref)			−0.03 (−0.12, 0.05)	−0.04 (−0.12, 0.05)
Family religiousness (not at all as ref)				
Religion not very important			−0.16 (−0.35, 0.03)	−0.17 (−0.36, 0.02)
Religion somewhat important			−0.12 (−0.29, 0.05)	−0.12 (−0.29, 0.05)

	Model 1 β (95% CI)	Model 2 β (95% CI)	Model 3 β (95% CI)	Model 4 β (95% CI)
Religion very important			-0.12 (-0.28, 0.05)	-0.11 (-0.27, 0.05)
Adulthood characteristics				
Marital status (married as ref)				
Divorced/separated				0.22 (0.12, 0.33) ***
Widowed				0.17 (-0.01, 0.36)
Never married				0.13 (0.00, 0.25) *
Education attainment (< high school as ref)				
High school				0.05 (-0.10, 0.20)
Some college				-0.04 (-0.20, 0.12)
College degree or higher				-0.08 (-0.12, -0.05)
Household income (bottom quartile as ref)				
Second quartile				0.06 (-0.04, 0.15)
Third quartile				0.10 (0.00, 0.21) *
Top quartile				0.11 (0.01, 0.22) *
Major depression (no as ref)				0.16 (-0.28, -0.04) **
Any chronic health condition (no as ref)				-0.01 (-0.11, 0.09)

Note: Generalized estimating equations with normal distribution and identity link were used in all models to estimate the mean change in BMI (standardized score, mean=0, standard deviation=1) by parenting styles, adjusting for clustering by sibling status.

* $p < .05$,

** $p < .01$,

*** $p < .001$

Table 3.

Parental warmth in childhood and BMI change in mid-life, stratified by levels of parental control (The Midlife in the United States Study 1995/1996–2004/2005 questionnaire wave, N=3,929)

	Full Sample	Stratified by Parental Control		
		Bottom tertile	Middle tertile	Top tertile
	β (95% CI)	β (95% CI)	β (95% CI)	β (95% CI)
Model 1	−0.05 (−0.08, −0.02) **	−0.06 (−0.11, 0.00)	−0.01 (−0.07, 0.04)	−0.08 (−0.14, −0.02) **
Model 2	−0.05 (−0.08, −0.01) **	−0.05 (−0.11, 0.01)	−0.01 (−0.07, 0.05)	−0.08 (−0.14, −0.02) *
Model 3	−0.04 (−0.07, −0.003) *	−0.03 (−0.09, 0.04)	−0.01 (−0.07, 0.05)	−0.08 (−0.15, −0.01) *

Note: Generalized estimating equations with normal distribution and identity link were used in all models to estimate the mean change in BMI (standardized score) by change in parental warmth (standardized score, mean=0, standard deviation=1), adjusting for clustering by sibling status.

Model 1 adjusted for participant age, sex and race.

Model 2 additionally adjusted for childhood socioeconomic status (assessed by parental education).

Model 3 additionally adjusted for other childhood family environment factors (including parental abuse, whether lived with both biological parents, childhood residential area, childhood residential stability, and family religiousness).

*
 $p < .05$,

**
 $p < .01$