Published in final edited form as:

J Med Chem. 2018 October 11; 61(19): 8613–8624. doi:10.1021/acs.jmedchem.8b00498.

Structure-based Drug Design and Identification of H₂O-soluble and Low Toxic Hexacyclic Camptothecin Derivatives with Improved Efficacy in both Cancer and Lethal Inflammation Models *In Vivo*

Peichen Pan^a, Jiean Chen^c, Xijian Li^c, Miyang Li^c, Huidong Yu^d, Jean J. Zhao^{e,f}, Jing Ni^{e,f}, Xuwen Wang^a, Huiyong Sun^a, Sheng Tian^g, Feng Zhu^a, Feng Liu^g, Yong Huang^{c,*}, Tingjun Hou^{a,b,*}

^aCollege of Pharmaceutical Sciences, Zhejiang University, Hangzhou, Zhejiang 310058, China

^bState Key Lab of CAD&CG, Zhejiang University, Hangzhou, Zhejiang 310058, China

^cKey Laboratory of Chemical Genomics, School of Chemical Biology and Biotechnology, Peking University, Shenzhen Graduate School, Shenzhen, Guangdong 518055, China

^dRongene Pharma Co., Ltd., Shenzhen, Guangdong 518054, China

^eDepartment of Cancer Biology, Dana-Farber Cancer Institute, Boston, Massachusetts 02215, USA

^fDepartment of Biological Chemistry and Molecular Pharmacology, Harvard Medical School, Boston, Massachusetts 02215, USA

^gCollege of Pharmaceutical Sciences, Soochow University, Suzhou, Jiangsu 215123, China.

Abstract

Camptothecin (CPT) has been shown to block disassembly of the topoisomerase-I (Topo I)/DNA cleavable complex. However, the poor aqueous solubility, intrinsic instability and severe toxicity of CPTs have limited their clinical applications. Herein, we report the design and synthesis of H₂O-soluble and orally bioavailable hexacyclic CPT derivatives. By analysis of a virtual chemical library and cytotoxicity screening *in vitro*, **9** and **11** were identified as potential pro-drugs and chosen for further characterization *in vivo*. Both compounds exhibited remarkable anticancer and anti-inflammation efficacies in animals, and improved drug-like profiles.

*Corresponding Authors Tingjun Hou, tingjunhou@zju.edu.cn; Yong Huang, huangyong@pkusz.edu.cn.

Author Contribution

The manuscript was written with contributions from all authors. All authors have given approval to the final version of the manuscript.

Supporting Information.

Workflow of construction and screening of virtual chemical library

Solubility of analogues.

Stability of compound **9** and **11** in human liver microsomes.

Pathological anatomy of primary viscera.

NMR spectroscopic data for compounds **1** - **11**.

Molecular formula strings of the synthesized compounds (CSV).

The authors declare no competing financial interest.

Graphical Abstract

Keywords

DNA Topoisomerase I; Camptothecin; Sepsis

INTRODUCTION

Half a century has passed since the structure of Camptothecin (CPT), a plant alkaloid originally extracted from the Chinese tree, *Camptotheca acuminata*, was identified. CPT was first isolated in 1958 by Wall and Wani and has been demonstrated to be an effective agent for cancer chemotherapy.¹ However, the early clinical trials of CPT were all terminated due to its poor solubility and stability as well as unpredictable severe toxicity.²⁻⁴ In addition, the highly electrophilic lactone E ring of CPT derivatives could rapidly undergo hydrolysis to an inactive, H₂O-soluble carboxylic acid which is ionized at physiological pH (shown in Scheme 1).⁵⁻⁸

In the late 1980s, CPT was found to be capable of inhibiting the DNA topoisomerase-I (Topo I) protein, a finding which resurrected this field.⁹⁻¹⁵ Results indicated that CPT can bind to the transient Topo I/DNA cleavage complex resulting in Topo I-mediated DNA breakage, which subsequently disturbs cellular processes such as replication, transcription and DNA repair, and finally causing cell death. Mutation experiments and X-ray crystallography both confirmed that CPT interacts with three key amino residues of Topo I (Asn722, Arg364 and Asp533) as well as with DNA base pairs.^{16, 17} Non-covalent interactions between CPT and Topo I/DNA binary complex are formed while base stacking interactions and hydrogen bond interactions also play vital roles in the stabilized ternary complex (Figure 1).

Due to challenges in production and derivatization of natural products, a large number of investigations on semi-synthesis and total synthesis of CPT analogs have been reported. These have stimulated exhaustive SAR studies aiming at improving both solubility and stability. Most successes were achieved by introducing multiple substituents on the A and B rings, specifically at carbons 7, 9, 10 and 11 (Scheme 2). This exercise led to two molecules, topotecan and irinotecan, which were ultimately approved by FDA for the treatment of various cancers, including colon cancer and ovarian cancer.¹⁸⁻²³ Nevertheless, the toxicity associated with both drugs is severe, and includes, but is not limited to gastrointestinal toxicity, myelosuppression and other side effects,²⁴ which constrains the doses that can be safely administered. As a result, there has been long, but unsuccessful search for a better analogue with improved oral bioavailability, attenuated toxicity profile and a superior therapeutic window.

CPTs have long been used as anticancer agents, but a recent study indicates that blockage of Topo I by CPT suppresses the host immune response and protects mice against death from lethal inflammation *in vivo*.²⁵ Sepsis, for example, is a leading cause of death in hospitalized patients due to excessive host response to infection, and there is still a need for better treatments.²⁶ This finding expands the possibility of CPTs as a therapy against life-threatening host response caused by bacteria and/or virus infections.

In the present work, by constructing a virtual chemical library, we rationally designed and synthesized several novel hexacyclic CPT analogs. The cellular antitumor activities of these analogs were then assessed against 15 different cancer cell lines. Two potent candidates were finally identified, exhibiting improved *in vivo* efficacy in a xenograft model of human liver cancer as well as a bacterial lipopolysaccharide (LPS)-induced sepsis model.

RESULTS AND DISCUSSION

From a drug-like property perspective, the solubility of CPT is poor and its *in vivo* stability is also low. To overcome these problems, structure-activity relationship (SAR) studies have been performed in the past decades, and led to the discovery of several semisynthetic and synthetic derivatives that are currently in clinical assessment. According to the SAR information from previous studies, the aromatic rings A and B are essential for biological activity. Saturating either ring destroys the activity even at high concentrations.^{27, 28} The activity is retained in hexacyclic systems, but tetracyclic or tricyclic structures show complete loss of biological activity, indicating that at least a pentacyclic ring structure is required to preserve the activity. No activity is observed when the D ring is replaced by a benzene ring.²⁹ Hydrolysis of the lactone E ring leads to a notable decrease in anticancer activity.³⁰ Optimizations of ring A in many cases however, improve the activity of CPT, while modifications of the remainder of CPT are less acceptable.³¹

The potential of hexacyclic CPT derivatives, in which position 9 and position 10 of ring A are linked via a five- or six-membered heterocycle, was further explored in the light of the above information. In our effort to design novel hexacyclic CPTs, we first constructed a virtual chemical library in which 5- or 6-membered ring fragments were fused onto the 9,10 positions of CPT, and 682 hexacyclic CPTs were enumerated. Then all the generated

compounds were docked and ranked by the *Glide* module in Schrodinger 9.0.³² Compounds **1**, **2** and **3** (see Chart 1) were chosen from 30 top-ranked molecules based on the ease of synthesizing these compounds and potential drug-likeness properties, and were synthesized for subsequent biological assessment. In addition, further substitutions on the 1,3-oxazole ring of compound **2**, yielding compounds **4** - **7** were explored to determine the preferred sidechains. As mentioned above, the *in vivo* hydrolysis of the lactone E ring results in a marked decrease in activity. Many studies have reported esterification of CPT as an effective pro-drug strategy. Giovanella and colleagues validated that esterification of CPT at 20-OH could stabilize the lactone ring.^{33, 34} The enhanced stability and solubility by ester-CPT derivatives help to prolong circulatory retention and improve tumor accumulation, which undoubtedly lead to improved activity *in vivo*.³⁵⁻³⁷ To enhance the stability of the E ring and at the same time improve compound solubility and bioavailability, modifications on the lactone ring were carried out by introducing a similar and well-known hydrophilic chain structure to the 20-OH group of compounds **1** to **4**, producing analogues **8** to **11**.

***In Vitro* Anticancer Activity of Hexacyclic CPTs.**

The cytotoxicity of all the synthesized hexacyclic CPT derivatives was evaluated against eleven human cancer cell lines (A549, MCF-7, MDA-MB-231, HepG-2, KB, BEL-7402, SMMC-7721, MGC80-3, SK-N-SH, QGY-7703 and SGC-7901) by MTT (3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide) assay. These cell lines cover different types of human cancers, including lung cancer, breast cancer, hepatoma, oral cutaneous carcinoma, gastric cancer and neuroblastoma. The FDA-approved drug, topotecan, was used as a reference inhibitor. The results of the *in vitro* cytotoxicity of all the compounds are summarized in Table 1. Most of the designed hexacyclic CPTs exhibit anti-proliferative activity comparable or superior to that of topotecan. The cytotoxicity against two cancer cell types, HepG-2 and MGC80-3, is obviously higher compared with that of other cell lines. In particular, compound **4** can potently inhibit both HepG-2 and MGC80-3 at very low concentrations, and the inhibitory IC₅₀ values of compound **4** (0.016 and 0.018 μ M, respectively) are ~10-fold lower than those of topotecan (0.237 μ M and 0.190 μ M, respectively). The inhibitory IC₅₀ values of compound **2** against HepG-2 and MGC80-3 are 0.180 and 0.231 μ M, respectively, which are >10-fold higher than those of compound **4**, indicating that substitution in the 1,3-oxazole ring by an isopropyl group significantly increases the activity. The purpose of modifying the 20-position hydroxyl of the E ring is to improve H₂O solubility and *in vivo* bioavailability, so even though the *in vitro* activities of compounds **9** to **11** against HepG-2 are slightly decreased compared with those of the unmodified compounds, and may be caused by changes in cellular permeability and/or different sensitivity to transporter proteins (the expressions are distinct in various cells) that pump foreign substances out of cells, the *in vivo* activities of these compounds were expected to be improved.

***In Vivo* Antitumor Activity of Compounds **9** and **11**.**

Preliminary *in vitro* cell-based evaluation demonstrated that hexacyclic CPTs showed potent activity against various cancer cell lines, especially HepG-2 and MGC80-3. Before starting *in vivo* studies, we first tested the aqueous solubility of compounds **9** and **11**. Topotecan is a soluble analog of camptothecin, which is insoluble in H₂O, but its solubility is still lower

than 1 mg/mL. As shown in Figure 2a, the solubility of both compounds **9** and **11** (as sodium salts) is 5.07 and 3.40 mg/mL, respectively, much higher than that of topotecan and parental compounds **4** and **3** ($< 2 \mu\text{g/mL}$), suggesting that compounds **9** and **11** might have better therapeutic efficacy over topotecan *in vivo*, especially following oral administration. To test this hypothesis, the *in vivo* antitumor activities were assessed in Balb/C nude mice bearing an HepG-2 hepatoma xenograft. In order to determine the best doses, the maximum tolerated dose (MTD) was measured first. As can be observed in Figure 2a, both compounds **9** and **11** are tolerated better than repeated administration of topotecan. Topotecan is tolerated only up to 2 mg/kg with daily intravenous dosing for one week, while compounds **9** and **11** can be tolerated at up to 30 and 10 mg/kg dose, suggesting that the short-term toxicity of compounds **9** and **11** is lower than that of topotecan. In single intravenous administration, compounds **9** and **11** are even tolerated up to 60 and 16 mg/kg, respectively.

Based on the above results, the *in vivo* antitumor activities of topotecan, and compounds **9** and **11** in a HepG-2 xenograft model were examined using indicated dosage (Figures 2b and 2c), and the toxicities of these compounds were assessed by the observation of the body weight, general appearance and histopathologic examination. Topotecan was intravenously administrated in the highest dose (the same dose as MTD, 2 mg/kg) as a reference compound. All three compounds significantly suppress tumor growth, and both compounds **9** and **11** display dose-responsive antitumor efficacy. The final tumor growth inhibition (TGI) rates for topotecan (i.v., 2.0 mg/kg, q7d), compound **9** (i.v., 15 mg/kg, q7d) and **11** (i.v., 30 mg/kg, q7d) are 74%, 86% and 90%, respectively, suggesting that compound **9** shows superior antitumor efficacy than topotecan even at half the MTD dose (15 mg/kg). Similarly, the TGI rates of compound **11** from different doses are all higher than that of topotecan, reaching 83% (i.v., 4 mg/kg, q7d), 93% (i.v., 8 mg/kg, q7d) and 93% (p.o., 1.3 mg/kg, qd). Notably, the oral administration of compound **11** of 1.3 mg/kg gave rise to antitumor efficacy similar to that from an i.v. dose of 8 mg/kg of **11**. Though given orally daily, the total oral dose is almost identical to q7d i.v. dose of 8 mg/kg, which implies that compound **11** can be absorbed well through oral administration. The data for oral ingestion of compound **9** was not collected since most mice failed to reach the end point of this experiment. The daily oral dose of compound **9** (5 mg/kg) was lower than the MTD, but the MTD was determined via i.v. injection for up to seven days. One possible explanation is that compound **9** is metabolized into (a) more toxic compound(s) via oral administration.

In order to determine the *in vitro* activity against Topo I, both Topo I inhibitory activity assay and Topo I-mediated DNA cleavage assay were performed. Topo I and supercoiled DNA were incubated with or without the test compounds. The inhibitory activity was detected on 1% EB-free gel which could separate super-coiled DNA from relaxed and nicked DNA. As shown in the Figures 3a and 3b, CPT can inhibit Topo I activity at concentrations of both 50 μM and 100 μM . Such inhibition is represented by the increased amount of super-coiled DNA that is not relaxed or cleaved. Compound **9** exhibits weak inhibition and compound **11** is almost inactive against Topo I at the concentrations of both 50 μM and 100 μM . Similar results are observed in the Topo I-mediated DNA cleavage assay which was carried out on EB-containing gel. CPT, topotecan and compound **9** can increase the amount of nicked DNA at concentrations of 100 μM or 50 μM , but compound

11 failed to stabilize Topo I-DNA cleavage complex at a concentration of 50 μ M. Both compound **9** and **11** were hypothesized to release their parent compound **4** and **3** *in vivo*. As we can see in Figure 4, the activity of compound **3** against Topo I is high, but compound **4** is a weak inhibitor of Topo I which is similar to compound **9**. Compound **3** is much more active than compound **11** and is even slightly more active than CPT.

Pathological anatomy of viscera of all xenograft mice was also carried out and the data collected are summarized in the Supporting Information (Table S2). The dominating side effects observed are gastrointestinal flatulence and splenomegaly. Notably, oral administration of compound **11** increases the occurrence of gastrointestinal flatulence and might affect hepatic status compared with the i.v. dosage, but these side effects have little influence on mouse survival. The body weight change of mice during treatment was recorded and plotted against time, in days in Figures 5a and 5b. It was found that the body weight of the untreated mice steadily increased over 35 days, but the body weight of mice treated with topotecan actually decreased slightly. However, treatment with compounds **9** or **11** at different doses had no major influence on body weight in comparison to treatment by topotecan, which on the other hand indicates that compounds **9** and **11** are probably less toxic than much higher doses of topotecan. Compared with topotecan, compounds **9** and **11** showed improved antitumor activity without acute and short-term *in vivo* toxicity.

Compounds **9** and **11** Protect from Lethal Inflammation and Death *In Vivo*

Over the past decades, CPTs have become some of the most widely used anticancer drugs.³⁸ In 2016, CPT was reported to suppress the expression of inflammatory genes and rescue mortality caused by excessive inflammation.²⁵ This prompted us to characterize the effectiveness of compounds **9** and **11** in *in vivo* sepsis models. As shown in Figure 6a, only 20% of animals survived LPS-induced septic shock, and oral administration of 10 mg/kg compound **9** rescued 80% of animals. The protective effect of compound **11** is even better. Although it is administered in relatively lower doses, 100% of the animals orally treated with 4 mg/kg **11** were rescued (Figure 6b). Furthermore, the survival rate of the animals receiving oral administration was higher than by intravenous injection, which is in accordance with the results from *in vivo* antitumor models. This phenomenon possibly suggests that the compound may act as a pro-drug that undergoes hydrolysis and/or metabolic processes *in vivo* and the activity of the metabolites is higher, or there are other protein targets that are inhibited. In order to assess this hypothesis, the preliminary pharmacokinetic properties of compound **9** and **11** were studied. As shown in Figure 7, The concentration of both compound **9** (data not detectable) and **11** (Figure 7c) in plasma through oral administration is extremely low, and the corresponding parent compounds **4** and **3** are also not released. For i.v. administration, compound **4** is also not detected in plasma after giving 2 mg/kg compound **9**, and very low concentration of compound **3** is identified after giving 2 mg/kg compound **11** intravenously. These results indicate that both compound **9** and **11** are not hydrolyzed into their parent compound **4** and **3**. The exact metabolite(s) of compound **9** and **11** are unclear. In addition, the expressions of two inflammatory cytokines, IL-6 and TNF- α in mouse serum were suppressed when treated with compound **9** or **11** (Figure 6c and 5d). These data suggest that compounds **9** and **11** provide effective protection in LPS-induced models of lethal inflammation at the organism level.

CONCLUSIONS

CPT compounds are among the first classes of anticancer agents, which have been demonstrated to stabilize the Topo I/DNA complex.⁹ However, the poor solubility, inherent instability and unpredictable severe toxicity of CPTs are unsatisfactory^{24, 39} for therapeutic use. In the present work, a novel kind of hexacyclic CPT derivatives was first designed and synthesized. These compounds are very promising and exhibit favorable biological properties, including improved solubility, reduced toxicity and enhanced efficacy. An *in vitro* cytotoxicity assay suggests that these compounds can efficiently suppress the growth of multiple cancer cell lines, especially HepG-2 and MGC80-3. Further *in vivo* antitumor activities of pro-drugs **9** and **11** were determined using HepG-2 hepatoma xenograft models, and the results indicate that compounds **9** and **11** exhibit improved *in vivo* antitumor activity but decreased toxicity compared with topotecan. Compounds **9** and **11** can also suppress the expression of several inflammatory cytokines and rescue mortality caused by excessive inflammation in *in vivo* sepsis models. Compound **9** and **11** were designed based on the pro-drug concept, but the PK results indicate that both compounds are not hydrolyzed into their parent compound **4** and **3**, which implies that compound **9** and **11** might convert into new active compounds to work *in vivo*. These data qualify compounds **9** and **11** as not only promising anticancer agents but also potential drug candidates for treatment of lethal inflammation.

EXPERIMENTAL SECTION

General Information.

All solvents were distilled according to standard methods prior to use. All reagents and chemicals were purchased and used without further purification unless specified otherwise. Solvents for flash column chromatography were technical grade and distilled prior to use. Thin-layer chromatography (TLC) was carried out on Huanghai silica gel plates with HSGF 254. Chromatograms were visualized based on UV absorbance (254 nm) with proper stains. Flash column chromatography was conducted using Qingdao Haiyang Chemical HG/T2354-92 silica gel (200-300 mesh) with the specified solvent system in the corresponding experiment. Proton (¹H) NMR and carbon (¹³C) NMR data were recorded on Bruker nuclear resonance spectrometers (400 MHz for ¹H and 100 MHz for ¹³C). Chemical shifts (δ) in ppm are reported relative to the residual signals of chloroform (¹H 7.26 ppm and ¹³C 77.16 ppm). ¹³C NMR spectra were recorded with total proton decoupling. Multiplicities are described as: s (singlet), bs (broad singlet), d (doublet), t (triplet), q (quartet), m (multiplet); and coupling constants (J) are reported in Hertz (Hz). The purity was determined by liquid chromatography-mass spectrometry (LC-MS). The purity of all final compounds was 95% or higher.

(S)-10-Amino-4-ethyl-4,9-dihydroxy-1H-pyrano[3',4':6,7]indolizino[1,2-b]quinoline-3,14(4H,12H)-dione (S1).

65% HNO₃ was added to a solution of 10-hydroxycamptothecin (180 mg, 0.5 mmol) in AcOH (1 mL). The reaction mixture was stirred at r.t. overnight. The reaction was diluted with H₂O (2.5 mL) to give a brown suspension, which was filtered and washed with H₂O.

The resulting brown solid was suspended in MeOH (5 mL) and hydrogenated using at 50° C for 2 h using Pd/C (10% w/w, 53 mg, 0.05 mmol) and a hydrogen balloon. When TLC showed complete consumption of the starting material, the reaction mixture was filtered and washed thoroughly with MeOH. The resulting solution was concentrated to give 133 mg (70% crude yield) of compound **S1** as a pale grey solid, which was used without further purification. ¹H NMR (400 MHz, DMSO) δ 9.80 (bs, 1H), 8.73 (s, 1H), 7.73 – 7.61 (m, 1H), 7.37 (q, *J* = 9.0 Hz, 2H), 7.23 (s, 1H), 6.47 (s, 1H), 5.40 (s, 2H), 5.23 (s, 2H), 4.27 – 4.04 (m, 1H), 3.15 (d, *J* = 4.9 Hz, 1H), 1.90 – 1.78 (m, 2H), 0.87 (t, *J* = 7.3 Hz, 3H). ¹³C NMR (101 MHz, DMSO) δ 173.00 (s), 157.35 (s), 150.52 (s), 149.43 (s), 146.60 (s), 143.94 (s), 140.70 (s), 130.02 (s), 127.44 (s), 125.60 (s), 121.74 (s), 118.70 (s), 118.43 (s), 117.58 (s), 96.19 (s), 72.86 (s), 65.69 (s), 50.59 (s), 30.68 (s), 8.22 (s).

(S)-4-Ethyl-4,9-dihydroxy-3,14-dioxo-3,4,12,14-tetrahydro-1H-pyrano[3',4':6,7]indolizino[1,2-b]quinoline-10-carbaldehyde (S2).

A solution of 10-hydroxycamptothecin (100 mg, 0.27 mmol) and HMTA (80 mg, 0.55 mmol) in TFA (10 mL) was heated to reflux for 20 h under argon. The reaction mixture was concentrated. H₂O (20 mL) was added and the mixture was stirred for 1 h. Additional H₂O (20 mL) was added and the pH was adjusted to 8-9 using saturated aqueous NaHCO₃ solution. The aqueous layer was washed with EtOAc (10 mL), acidified to pH ~1.5 using 2N HCl and extracted using EtOAc (20 mL*5). The combined organic phase was washed with 1N HCl, H₂O and brine, dried over Na₂SO₄, filtered and concentrated. The residue was passed through a short plug of silica gel, thoroughly washed using a mixture of MeOH and DCM (1:50), and concentrated to give **S2** (79 mg, 73% crude yield), which was used without further purification. ¹H NMR (400 MHz, DMSO) δ 11.97 (bs, 1H), 10.61 (s, 1H), 9.50 (s, 1H), 8.13 (d, *J* = 9.3 Hz, 1H), 7.45 (d, *J* = 9.3 Hz, 1H), 7.15 (s, 1H), 6.46 (s, 1H), 5.36 (s, 2H), 5.08 (s, 2H), 1.94 – 1.69 (m, 2H), 0.87 (t, *J* = 7.3 Hz, 3H). ¹³C NMR (101 MHz, DMSO) δ 191.65 (s), 172.90 (s), 164.91 (s), 157.07 (s), 150.29 (s), 150.15 (s), 145.61 (s), 143.13 (s), 138.74 (s), 132.72 (s), 127.21 (s), 126.73 (s), 123.57 (s), 118.93 (s), 112.88 (s), 96.48 (s), 72.77 (s), 65.68 (s), 50.80 (s), 30.79 (s), 8.21 (s).

(S)-4-Ethyl-4,9-dihydroxy-3,14-dioxo-3,4,12,14-tetrahydro-1H-pyrano[3',4':6,7]indolizino[1,2-b]quinoline-10-carboxamide (S3).

S2 (880 mg, 2.2 mmol), hydroxylamine hydrochloride salt (400 mg, 5.8 mmol) and sodium formate (2 g, 29.4 mmol) were mixed in formic acid (200 mL). The reaction was refluxed overnight. The solvent was removed. The residue was passed through a short plug of silica gel, thoroughly washed using a mixture of MeOH and DCM (1:50) and concentrated. The residue was dissolved in TFA (4.8 mL). Concentrated H₂SO₄ (1.2 mL) was added slowly. The mixture was heated at 95 °C for 16 h, then cooled to r.t. and concentrated. The residue was passed through a short plug of silica gel, thoroughly washed using a mixture of MeOH and DCM (1:10), and concentrated to give **S3** (69 mg, 30% yield), which was used without further purification. ¹H NMR (400 MHz, DMSO) δ 11.66 (bs, 1H), 11.52 (bs, 1H), 9.30 (s, 1H), 8.85 (s, 1H), 8.01 (d, *J* = 9.2 Hz, 1H), 7.61 (d, *J* = 9.2 Hz, 1H), 7.23 (s, 1H), 6.51 (s, 1H), 5.39 (s, 2H), 5.20 (s, 2H), 1.86 (dd, *J* = 14.0, 6.9 Hz, 2H), 0.85 (d, *J* = 7.4 Hz, 3H). ¹³C NMR (101 MHz, DMSO) δ 172.92 (s), 157.33 (s), 157.25 (s), 150.48 (s), 149.83 (s), 146.48

(s), 146.01 (s), 143.79 (s), 132.61 (s), 131.07 (s), 127.64 (s), 127.11 (s), 123.15 (s), 118.79 (s), 110.08 (s), 96.50 (s), 72.82 (s), 65.69 (s), 50.75 (s), 30.72 (s), 8.23 (s).

(S)-8-Ethyl-8-hydroxy-8H-isoxazolo[4,5-f]pyrano[3',4':6,7]indolizino[1,2-b]quinoline-9,12(11H,14H)-dione (1).

Triethylamine (15 μ L, 0.15 mmol) was added to a solution of **S2** (50 mg, 0.13 mmol) and hydroxylamine hydrochloride (11 mg, 0.15 mmol) in EtOH (10 mL). The reaction mixture was stirred at 95 $^{\circ}$ C for 20 h. The reaction was cooled to room temperature and concentrated. The residue was dissolved in EtOAc (10 mL) and washed with H₂O (15 mL). The aqueous layer was extracted with EtOAc (10 mL*2). The combined organic layer was washed with H₂O and brine, dried over Na₂SO₄, filtered and concentrated. The residue was dissolved in THF (10 mL), and DIAD (25 mg, 0.12 mmol) was added, followed by PPh₃ (32 mg, 0.12 mmol). The mixture was stirred at r.t. for 5 h. The solvent was removed and the residue was purified by silica gel column chromatography eluting with MeOH:DCM = 1:50 to give compound **1** as a pale yellow solid (26 mg, 53% yield from **S2**). Melting point: 228.1-229.8 $^{\circ}$ C. ¹H NMR (500 MHz, DMSO-d₆) δ 9.89 (s, 1H), 9.12 (s, 1H), 8.33 (d, J = 9.3 Hz, 1H), 8.25 (d, J = 9.3 Hz, 1H), 7.33 (s, 1H), 6.53 (s, 1H), 5.41 (s, 2H), 5.34 (s, 2H), 1.87 (qd, J = 14.1, 7.3 Hz, 2H), 0.88 (t, J = 7.3 Hz, 3H). ¹³C NMR (101 MHz, DMSO-d₆) δ 172.93, 161.84, 157.20, 152.07, 150.48, 147.12, 146.48, 145.74, 133.19, 132.20, 128.07, 122.44, 119.45, 116.94, 115.01, 97.04, 72.84, 65.70, 50.87, 30.77, 8.24.

(S)-8-Ethyl-8-hydroxy-8H-oxazolo[4,5-f]pyrano[3',4':6,7]indolizino[1,2-b]quinoline-9,12(11H,14H)-dione (2).

S1 (133 mg, 0.35 mmol) was dissolved in EtOH (2 mL) and triethyl orthoformate (96 mg, 0.65 mmol) was added. The reaction mixture was heated to reflux overnight under argon. The solvent was removed and the residue was purified by silica gel chromatography eluting with MeOH:DCM = 1:50 to give compound **2** as a pale yellow solid (84 mg, 62% from **S1**). Melting point: 162.4-163.7 $^{\circ}$ C. ¹H NMR (500 MHz, DMSO-d₆) δ 9.10 (s, 1H), 9.03 (s, 1H), 8.31 (d, J = 9.2 Hz, 1H), 8.21 (d, J = 9.2 Hz, 1H), 7.35 (s, 1H), 6.52 (s, 1H), 5.43 (s, 2H), 5.33 (s, 2H), 1.87 (m, 2H), 0.88 (t, J = 7.3 Hz, 3H). ¹³C NMR (75 MHz, DMSO-d₆) δ 172.90, 156.94, 155.17, 151.74, 150.27, 147.50, 146.23, 145.42, 135.07, 130.70, 127.65, 125.75, 121.10, 119.27, 115.64, 96.84, 72.79, 65.67, 50.67, 30.81, 8.25.

(9S)-9-Ethyl-9-hydroxy-3-isopropyl-2,3-dihydro-[1,3]oxazino[5,6-f]pyrano[3',4':6,7]indolizino[1,2-b]quinoline-1,10,13(9H,12H,15H)-trione (3).

*p*TSA monohydrate (1.4 mg, 0.0075 mmol) and isobutyraldehyde (41 μ L, 0.45 mmol) were added to the solution of **S3** (60 mg, 0.15 mmol) in a mixture of toluene (0.5 mL) and DMSO (0.25). The reaction mixture was stirred at 110 $^{\circ}$ C for 16 h. Toluene was removed and the DMSO solution was diluted with DCM (5 mL) and washed with H₂O and brine. The organic layer was dried over Na₂SO₄, filtered and concentrated. The residue was purified by silica gel chromatography eluting with MeOH:DCM = 1:50 to give compound **3** as a pale yellow solid (49 mg, 71% yield from **S3**). Melting point: 256.8-258.2 $^{\circ}$ C. ¹H NMR (400 MHz, DMSO-d₆) δ 9.78 (s, 1H), 8.96 (s, 1H), 8.24 (dd, J = 9.2, 1.9 Hz, 1H), 7.56 (dd, J = 9.2, 4.3 Hz, 1H), 7.33 – 7.23 (m, 1H), 6.50 (d, J = 2.4 Hz, 1H), 5.41 (s, 2H), 5.24 (m, 3H), 2.14 (m,

1H), 1.94 – 1.78 (m, 2H), 1.09 (t, J = 6.1 Hz, 6H), 0.88 (t, J = 7.3 Hz, 3H). ^{13}C NMR (101 MHz, DMSO- d_6) δ 172.89, 163.65, 159.21, 157.17, 150.94, 150.44, 150.39, 145.72, 144.87, 136.41, 131.82, 129.33, 127.40, 122.31, 119.22, 110.53, 96.79, 88.41, 72.82, 65.69, 50.93, 30.75, 17.32, 16.35, 8.21.

(S)-8-Ethyl-8-hydroxy-2-isopropyl-8H-oxazolo[4,5-f]pyrano[3',4':6,7]indolizino[1,2-b]quinoline-9,12(11H,14H)-dione (4).

S1 (61 mg, 0.16 mmol) was dissolved in THF (1 mL) and isobutyraldehyde (17 mg, 0.24 mmol) was added. The reaction mixture was heated at 50 °C for 6 h and DDQ (55 mg, 0.24 mmol) was added in one portion. The resulting mixture was stirred at 50 °C for another 10 h. The solvent was removed and the residue was purified by silica gel chromatography eluting with MeOH:DCM = 1:50 to give compound **4** as a pale yellow solid (34 mg, 50% yield from **S1**). Melting point: 154.3-155.2°C. ^1H NMR (300 MHz, DMSO- d_6) δ 9.08 (s, 1H), 8.25 (d, J = 9.2 Hz, 1H), 8.14 (d, J = 9.2 Hz, 1H), 7.35 (s, 1H), 6.55 (s, 1H), 5.43 (s, 2H), 5.32 (s, 2H), 3.43 (d, J = 6.9 Hz, 1H), 1.87 (dd, J = 11.5, 4.5 Hz, 2H), 1.46 (d, J = 6.9 Hz, 6H), 0.88 (m, 3H). ^{13}C NMR (75 MHz, DMSO- d_6) δ 172.93, 171.85, 156.95, 151.43, 150.32, 147.93, 145.94, 145.50, 135.89, 130.32, 126.62, 125.67, 120.71, 119.13, 115.17, 96.85, 72.82, 65.63, 50.64, 30.76, 28.75, 20.52, 8.23.

(S)-2-Butyl-8-ethyl-8-hydroxy-8H-oxazolo[4,5-f]pyrano[3',4':6,7]indolizno[1,2-b]quinoline-9,12(11H,14H)-dione (5).

This compound was synthesized according to the same procedure used for compound **4** using **S1** and valeraldehyde (pale yellow solid, 58 mg, 82% yield). Melting point: 151.1-153.2°C. ^1H NMR (400 MHz, DMSO- d_6) δ 8.93 (s, 1H), 8.13 (d, J = 9.1 Hz, 1H), 8.03 (d, J = 9.1 Hz, 1H), 7.27 (s, 1H), 6.52 (s, 1H), 5.40 (s, 2H), 5.20 (s, 2H), 3.04 (t, J = 7.4 Hz, 2H), 1.85 (dd, J = 15.3, 7.7 Hz, 4H), 1.44 (dd, J = 14.7, 7.3 Hz, 2H), 0.95 (t, J = 7.3 Hz, 3H), 0.89 (t, J = 7.1 Hz, 3H). ^{13}C NMR (101 MHz, DMSO- d_6) δ 172.95, 168.26, 157.18, 151.91, 150.43, 148.20, 146.24, 145.82, 136.31, 130.86, 126.74, 126.01, 120.99, 119.31, 115.42, 96.94, 72.85, 65.71, 50.90, 30.78, 28.78, 28.06, 22.12, 14.06, 8.24.

(S)-8-Ethyl-8-hydroxy-2-isobutyl-8H-oxazolo[4,5-f]pyrano[3',4':6,7]indolizno[1,2-b]quinoline-9,12(11H,14H)-dione (6).

This compound was synthesized according to the same procedure for **4** using **S1** and isovaleraldehyde (pale yellow solid, 53 mg, 75% yield). Melting point: 150.8-152.5°C. ^1H NMR (400 MHz, DMSO- d_6) δ 8.98 (s, 1H), 8.16 (d, J = 9.1 Hz, 1H), 8.06 (d, J = 9.2 Hz, 1H), 7.29 (s, 1H), 6.52 (s, 1H), 5.41 (s, 2H), 5.24 (s, 2H), 2.94 (d, J = 7.1 Hz, 2H), 2.27 (dt, J = 13.5, 6.7 Hz, 1H), 1.92 – 1.82 (m, 2H), 1.03 (d, J = 6.6 Hz, 6H), 0.89 (t, J = 7.2 Hz, 3H). ^{13}C NMR (101 MHz, DMSO- d_6) δ 172.95, 167.51, 157.21, 152.00, 150.45, 148.25, 146.31, 145.87, 136.36, 130.97, 126.80, 126.10, 121.07, 119.35, 115.48, 96.97, 72.86, 65.71, 50.95, 37.19, 30.78, 27.51, 22.63, 8.25.

(S)-8-Ethyl-8-hydroxy-2-(pyridin-2-yl)-8H-oxazolo[4,5-f]pyrano[3',4':6,7]indolizino[1,2-b]quinoline-9,12(11H,14H)-dione (7).

This compound was synthesized according to the same procedure for **4** using **S1** and picolinaldehyde (dark red solid, 16 mg, 22% yield). Melting point > 300°C. ¹H NMR (300 MHz, DMSO-d₆) δ 9.16 (s, 1H), 8.83 (s, 1H), 8.44 – 8.32 (m, 2H), 8.23 (d, *J* = 9.3 Hz, 1H), 8.10 (s, 1H), 7.66 (s, 1H), 7.35 (s, 1H), 6.54 (s, 1H), 5.42 (s, 2H), 5.35 (s, 2H), 1.86 (dd, *J* = 12.0, 6.6 Hz, 2H), 0.89 (t, *J* = 7.0 Hz, 3H). Due to the poor solubility of this compound, ¹³C NMR could not be obtained using common NMR solvents.

(S)-4-((2-((8-Ethyl-9,12-dioxo-9,11,12,14-tetrahydro-8H-oxazolo[4,5-f]pyrano[3',4':6,7]indolizino[1,2-b]quinolin-8-yl)oxy)-2-oxoethyl)amino)-4-oxobutanoic acid (8).

A solution of compound **2** (101 mg, 0.26 mmol), Sc(OTf)₃ (79 mg, 0.16 mmol) and DMAP (94 mg, 0.78 mmol) in dry DCM (2 mL) was stirred at room temperature for 30 min then Boc-Gly-OH (136 mg, 0.78 mmol) was added in one portion. The resulting mixture was stirred at room temperature for 30 min before DCC (272 mg, 1.3 mmol) was added. The reaction was stirred at room temperature overnight, filtered through celite and concentrated. The residue was passed through a short plug of silica gel, washed using a mixture of MeOH and DCM (1:100), and concentrated. The residue was dissolved in DCM (2 mL) and TFA (1 mL) was added dropwise. The reaction mixture was stirred at rt for 30 min and concentrated. The residue was dissolved in DMF (2 mL). Succinic anhydride (97 mg, 0.97 mmol) and 4-methylpyridine (73 mg, 0.81 mmol) were added sequentially. The reaction was stirred at rt overnight. Removal of the solvent and purification using silica gel chromatography (eluent: MeOH:DCM = 1:50) afforded compound **8** as a pale yellow solid (97 mg, 68% yield). Melting point: 182.3-184.1°C. ¹H NMR (400 MHz, DMSO-d₆) δ 12.08 (s, 1H), 9.02 (s, 1H), 8.98 (s, 1H), 8.43 (t, *J* = 5.7 Hz, 1H), 8.23 (d, *J* = 9.2 Hz, 1H), 8.15 (d, *J* = 9.2 Hz, 1H), 7.13 (s, 1H), 5.50 (s, 2H), 5.35 – 5.21 (m, 2H), 4.19 (dd, *J* = 17.0, 5.8 Hz, 1H), 4.02 (dd, *J* = 17.0, 5.8 Hz, 1H), 2.49-2.37 (m, 4H), 2.22 – 2.13 (m, 2H), 0.94 (t, *J* = 7.3 Hz, 3H). ¹³C NMR (101 MHz, DMSO-d₆) δ 174.14, 174.05, 172.06, 169.58, 167.51, 156.92, 155.34, 152.02, 147.77, 146.52, 146.32, 145.58, 135.33, 131.22, 127.86, 126.11, 121.53, 119.36, 115.90, 95.59, 76.66, 66.80, 50.95, 30.92, 30.16, 29.39, 29.26, 7.98.

(S)-4-((2-((8-Ethyl-2-isopropyl-9,12-dioxo-9,11,12,14-tetrahydro-8H-oxazolo[4,5-f]pyrano[3',4':6,7]indolizino[1,2-b]quinolin-8-yl)oxy)-2-oxoethyl)amino)-4-oxobutanoic acid (9).

This compound was synthesized according to the same procedure for **8** from **4** and was obtained as a pale yellow solid (66 mg, 43% yield). Melting point: 194.8-196.4°C. ¹H NMR (400 MHz, DMSO) δ 12.07 (s, 1H), 9.06 (s, 1H), 8.44 (s, 1H), 8.21 (d, *J* = 9.0 Hz, 1H), 8.13 (d, *J* = 8.9 Hz, 1H), 7.15 (s, 1H), 5.50 (s, 2H), 5.30 (s, 2H), 4.22 – 4.14 (m, 1H), 4.03 – 3.97 (m, 1H), 3.43 – 3.38 (m, 1H), 2.43 (d, *J* = 5.5 Hz, 2H), 2.38 (s, 2H), 2.16 (d, *J* = 6.5 Hz, 2H), 1.46 (d, *J* = 6.8 Hz, 6H), 0.93 (d, *J* = 6.7 Hz, 3H). ¹³C NMR (101 MHz; DMSO-d₆) δ 7.96, 20.66, 28.84, 29.38, 30.14, 30.86, 50.97, 66.80, 76.65, 95.57, 115.65, 119.29, 121.29, 126.20, 126.83, 131.05, 136.29, 145.53, 146.35, 146.43, 148.31, 151.92, 156.97, 167.56, 169.57, 172.03, 172.22, 174.16.

(S)-4-((2-((8-Ethyl-9,12-dioxo-9,11,12,14-tetrahydro-8H-isoxazolo[4,5-f]pyrano[3',4':6,7]indolizino[1,2-b]quinolin-8-yl)oxy)-2-oxoethyl)amino)-4-oxobutanoic acid (10).

This compound was synthesized according to the same procedure for **8** from **1** (pale yellow solid, 106 mg, 75% yield). Melting point: 201.5–203.5°C. ¹H NMR (400 MHz, DMSO-d₆) δ 12.06 (s, 1H), 9.90 (s, 1H), 9.11 (s, 1H), 8.44 (t, *J* = 5.1 Hz, 1H), 8.32 (t, *J* = 10.3 Hz, 1H), 8.25 (d, *J* = 9.4 Hz, 1H), 7.15 (s, 1H), 5.48 (d, *J* = 8.6 Hz, 2H), 5.34 (s, 2H), 4.17 (dd, *J* = 17.8, 5.8 Hz, 1H), 4.00 (dd, *J* = 17.9, 5.6 Hz, 1H), 2.42 (t, *J* = 4.5 Hz, 2H), 2.40–2.35 (m, 2H), 2.16 (dd, *J* = 14.7, 7.4 Hz, 2H), 0.92 (t, *J* = 7.4 Hz, 3H). ¹³C NMR (101 MHz; DMSO-d₆) δ 7.96, 29.36, 30.12, 30.88, 50.81, 66.77, 76.63, 95.60, 115.00, 116.94, 119.35, 122.50, 128.07, 132.17, 133.08, 145.58, 146.23, 146.44, 147.11, 151.88, 156.89, 161.85, 167.50, 169.58, 172.07, 174.12.

4-((2-(((9S)-9-Ethyl-3-isopropyl-1,10,13-trioxo-1,2,3,9,10,12,13,15-octahydro-[1,3]oxazino[5,6-f]pyrano[3',4':6,7]indolizino[1,2-b]quinolin-9-yl)oxy)-2-oxoethyl)amino)-4-oxobutanoic acid (11).

This compound was synthesized according to the same procedure for **8** from **3** (pale yellow solid, 74 mg, 46% yield). Melting point: 222.4–224.8°C. ¹H NMR (400 MHz, DMSO-d₆) δ 12.05 (s, 1H), 9.81 (d, *J* = 4.3 Hz, 1H), 8.99 (s, 1H), 8.42 (t, *J* = 6.0 Hz, 1H), 8.28 (d, *J* = 9.2 Hz, 1H), 7.59 (d, *J* = 9.3 Hz, 1H), 7.11 (s, 1H), 5.48 (s, 2H), 5.33 (t, *J* = 8.9 Hz, 2H), 5.21 (s, 1H), 4.15 (dd, *J* = 18.1, 5.9 Hz, 1H), 3.99 (dd, *J* = 17.9, 5.4 Hz, 1H), 2.46–2.41 (m, 2H), 2.36 (dd, *J* = 12.1, 5.7 Hz, 2H), 2.14 (d, *J* = 5.3 Hz, 3H), 1.09 (t, *J* = 6.0 Hz, 6H), 0.93–0.86 (m, 3H). ¹³C NMR (101 MHz; DMSO-d₆) δ 7.97, 16.36, 17.34, 29.34, 30.09, 30.74, 30.83, 51.03, 66.78, 76.63, 88.44, 95.42, 110.66, 119.17, 122.50, 127.57, 129.46, 132.00, 136.43, 144.93, 145.50, 146.30, 150.94, 156.95, 159.30, 163.70, 167.55, 169.57, 172.01, 174.16.

Molecular Modeling.

Construction of a virtual chemical library of CPT derivatives and all docking simulations were performed in Schrödinger 9.0.³² The brief workflow of construction and screening of virtual library is depicted in Figure S1. The first step is to generate five-membered and six-membered ring structures from small molecule drug database which was obtained from the BindingDB database.⁴⁰ Enumeration of a virtual library was carried out based on the *Interactive Enumeration and Docking* module in Schrödinger 9.0, where CPT was set as a core-containing molecule and the ring fragment collections were incorporated into the 9- and 10-position of CPT. Preparations of the initial structure of Topo I/DNA complex (PDB codes: 1T8I) were performed using the *Protein Preparation wizard* module. All H₂O molecules were deleted from the system and the missing hydrogen atoms were added. A restrained minimization was subsequently conducted, and the receptor grid for docking simulations was produced and centered on the original CPT based on the *Receptor grid generation* module. The scaling factor for van der Waals radii and partial atomic charge cutoff value were set to 0.8 and 0.15, respectively. Preparations of CPT ligands were all performed by the *LigPrep* module with protonated states generated at pH = 7.0 ± 2.0. Finally, all the compounds were docked into the binding pocket of Topo I/DNA complex using the *Glide* module with extra precision (XP).

Cell-based Cytotoxicity Assays.

Cells were collected and re-suspended in RPMI-1640 (or DMEM) growth media + 10% fetal bovine serum (FBS). The cells were seeded on 96-well plates at the concentration of 10,000 cells/well and incubated at 37 °C in 5% CO₂ overnight. The following day, compounds with serial dilutions were added into the corresponding wells, and the plates were further incubated at 37 °C for 48 h. The number of living cells was characterized using 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) assay by adding 5 mg/ml MTT into the wells, and the plates were then cultured for 4 h. The converted dye was then dissolved in MTT buffer overnight, and the absorbance of each well was finally measured at 570 nm with the control wavelength set to 650 nm. The half-inhibitory concentration (IC₅₀) of the compound was calculated based on the fitted dose-response curve.

Topo I Inhibitory Activity and DNA Cleavage Assay.

Both the inhibitory activity and Topo I-mediated cleavage assays were carried out using the topoisomerase I drug screening kit (TopoGEN, Inc.) The reaction mixture (20 µl) contained 10 mM Tris-HCl (pH 7.9), 1 mM EDTA, pHOT1 plasmid DNA (250 ng), Topo I (7 units for inhibitory assay and 10 units for cleavage assay) and indicated drug concentrations (1% DMSO). Topo I reaction mixtures were incubated at 37°C for 30 min. Reactions were terminated by the addition of 2 µl 10% SDS, digested with 50 µg/ml proteinase K for 30 min and extracted with phenol/chloroform/isoamyl alcohol (25:24:1). After adding 2 µl 10×gel loading buffer (0.25% bromophenol blue, 50% glycerol), the samples were loaded onto the agarose gel. Two types of agarose gels (1%) were prepared. The Topo I inhibitory gel was prepared in the absence of EB (ethidium bromide), and the cleavage gel was prepared with 0.5 µg/mL EB. Gels were run at 2 V/cm for 2.5 h in 1×TAE running buffer. After electrophoresis, the EB-free gels were stained with 0.5 µg/mL EB for 25 min and de-stained in H₂O for 15 min prior to photodocumentation. All the experiments were carried out independently more than three times and the representative gels were shown.

In Vivo Xenograft Studies.

Experiments were carried out by Suzhou Xishan Zhongke Drug Research & Development (China) Co., Ltd., and all *in vivo* experiments were approved and supervised by institutional animal care and use committee (Jiangsu Province, PR China) and carried out according to the guidance for the care and use of laboratory animals. Balb/C nude mice were obtained from ShanghaiSippr-BK laboratory animal Co. Ltd. HepG-2 cells in logarithmic phase were harvested and re-suspended in PBS (5×10⁸ cells/mL). 0.2 mL cells were subcutaneously injected into the right flank of each mouse. When the tumor reached the designated size of 50 mm³, the mice were randomly assigned to treatment groups (8 mice per group). CTP compounds and topotecan were all dissolved in normal saline. Tumor volume (TV) was calculated as: $TV = (L \times W^2)/2$, where L is tumor length and W represents tumor width. Body weight change (BW) was measured using the formula below: $BW = (W/W_0) \times 100\%$, where W refers to the body weight on a specific day and W₀ is the body weight on the first day of treatment. Tumor growth inhibition (TGI) was determined as: $TGI = [1 - (T - T_0)/(C - C_0)] \times 100\%$, where T (on a specific day) and T₀ (on the first day of treatment) stand for the

mean tumor volumes of treatment groups, and C and C₀ refer to the mean tumor volumes for control groups.

Determination of Maximum Tolerated Doses (MTD).

The acute and short-term toxicity was tested following administration of drugs with increasing doses to healthy animals. Drugs were intravenously injected (single i.v. dosage or daily i.v. dosage for one week) with each group composed of three mice. Behavioral changes after each administration were monitored hourly, and survival and side effects of the mice were recorded daily for up to 28 days. MTD was determined as the dose before at least one mouse was killed in the treatment.

In Vivo Pharmacokinetics (PK).

Pharmacokinetic properties of compound **11** were studied in male C57BL/6 mice (4 mice per group) after single oral (2 mg/kg) and intravenous administration (10 mg/kg). Mice were fasted overnight with free access to H₂O and fed four h after dosing. The whole blood samples were collected via the retro-orbital sinus and plexus at indicated time points and centrifuged at 5500 rpm for 10 min to obtain plasma samples. Plasma levels of the compounds were determined by LC-MS/MS analysis.

LPS-Induced Lethal Inflammation Models.

These experiments were all carried out by CTI Biotechnology (Suzhou, China) Co., Ltd. All experiments were approved and supervised by institutional animal care and use committee (Jiangsu Province, PR China) and carried out according to the guidance for the care and use of laboratory animals. C57BL/6 mice were obtained from Shanghai Sippr-BK laboratory animal Co. Ltd. Mice were separated into eight groups (10 mice per group). For the sepsis model, mice were injected intraperitoneally with 10 mg/kg LPS (from Escherichia coli 0111, Sigma). Dexamethasone (Dex) was set as a positive control drug. For treatment groups, mice were intravenously or orally administrated assigned doses of compounds 0.5 h before LPS injection followed by the same doses of compounds 1 h, 4 h and 8 h after LPS treatment. During the experiments, mice were weighed daily and observed at least twice per day in case of death. To obtain the concentration of inflammatory cytokines, 0.5 ml blood per mouse was retro-orbitally adopted 72 h after LPS treatment, and centrifuged at 3,000 rpm for 10min to separate serum.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

ACKNOWLEDGMENT

This study was supported by the National Key R&D Program of China (2016YFA0501701; 2016YFB0201700), the National Science Foundation of China (21575128; 81302679) and Shenzhen Science and Technology Innovation Commission (JCYJ20160226105602871).

ABBREVIATIONS

MTD maximum tolerated dose

TGI tumor growth inhibition.

REFERENCES

1. Wall ME; Wani MC; Cook CE; Palmer KH; McPhail AT; Sim GA Plant antitumor agents. I. The isolation and structure of camptothecin, a novel alkaloidal leukemia and tumor inhibitor from *camptotheca acuminata*. *J. Am. Chem. Soc.* 1966, 88, 3888–3890.
2. Moertel CG; Reitemeier RJ; Schutt AJ; Hahn RG Phase ii study of streptozotocin (nsc-85998) in the treatment of advanced gastrointestinal cancer. *Cancer Chemother. Rep.* 1971, 55, 303–307. [PubMed: 5115851]
3. Schaeppi U; Fleischman RW; Cooney DA Toxicity of camptothecin (nsc-100880). *Cancer Chemother. Rep.* 3 1974, 5, 25–36. [PubMed: 4213507]
4. Gottlieb JA; Luce JK Treatment of malignant melanoma with camptothecin (nsc-100880). *Cancer Chemother. Rep.* 1972, 56, 103–105. [PubMed: 5030802]
5. Burke TG; Mi Z The structural basis of camptothecin interactions with human serum albumin: Impact on drug stability. *J. Med. Chem.* 1994, 37, 40–46. [PubMed: 8289200]
6. Hertzberg RP; Caranfa MJ; Holden KG; Jakas DR; Gallagher G; Mattern MR; Mong SM; Bartus JOL; Johnson RK; Kingsbury WD Modification of the hydroxylactone ring of camptothecin: Inhibition of mammalian topoisomerase i and biological activity. *J. Med. Chem.* 1989, 32, 715–720. [PubMed: 2537428]
7. Jaxel C; Kohn KW; Wani MC; Wall ME; Pommier Y Structure-activity study of the actions of camptothecin derivatives on mammalian topoisomerase i: Evidence for a specific receptor site and a relation to antitumor activity. *Cancer Res.* 1989, 49, 1465–1469. [PubMed: 2538227]
8. Burke TG Chemistry of the camptothecins in the bloodstream. Drug stabilization and optimization of activity. *Ann. N. Y. Acad. Sci.* 1996, 803, 29–31. [PubMed: 8993497]
9. Hsiang YH; Hertzberg R; Hecht S; Liu LF Camptothecin induces protein-linked DNA breaks via mammalian DNA topoisomerase i. *J. Biol. Chem.* 1985, 260, 14873–14878. [PubMed: 2997227]
10. Yang C; Song Z; Goto M; Liu Y i.; Hsieh KY; Morris Natschke SL; Zhao Y; Zhang J; Lee K Design, synthesis, and cytotoxic activity of novel 7-substituted camptothecin derivatives incorporating piperazinyl-sulfonylamidine moieties. *Bioorg. Med. Chem. Lett.* 2017, 27, 3959–3962. [PubMed: 28789891]
11. Zhang X; Tang K; Wang H; Liu Y; Bao B; Fang Y; Zhang X; Lu W Design, synthesis, and biological evaluation of new cathepsin b-sensitive camptothecin nanoparticles equipped with a novel multifunctional linker. *Bioconj. Chem.* 2016, 27, 1267–1275.
12. Liu Y; Chen X; Ding J; Yu L; Ma D; Ding J Improved solubility and bioactivity of camptothecin family antitumor drugs with supramolecular encapsulation by water-soluble pillar[6]arene. *ACS Omega* 2017, 2, 5283–5288. [PubMed: 30023745]
13. Zhang F; Zhu G; Jacobson O; Liu Y; Chen K; Yu G; Ni Q; Fan J; Yang Z; Xu F; Fu X; Wang Z; Ma Y; Niu G; Zhao X; Chen X Transformative nanomedicine of an amphiphilic camptothecin prodrug for long circulation and high tumor uptake in cancer therapy. *ACS Nano* 2017, 11, 8838–8848. [PubMed: 28858467]
14. Westover D; Ling X; Liu X; Lam H; Gongora C; Rio MD; Li F Abstract 829: The novel camptothecin derivative and iap inhibitor fl118 is an effective treatment for irinotecan-refractory colorectal cancer. *Cancer Res.* 2014, 74, 829. [PubMed: 24335958]
15. Capranico G; Marinello J; Chillemi G Type i DNA topoisomerases. *J. Med. Chem.* 2017, 60, 2169–2192. [PubMed: 28072526]
16. Staker BL; Feese MD; Cushman M; Pommier Y; Zembower D; Stewart L; Burgin AB Structures of three classes of anticancer agents bound to the human topoisomerase i-DNA covalent complex. *J. Med. Chem.* 2005, 48, 2336–2345. [PubMed: 15801827]
17. Staker BL; Hjerrild K; Feese MD; Behnke CA; Burgin AB; Stewart L The mechanism of topoisomerase i poisoning by a camptothecin analog. *Proc. Natl. Acad. Sci. U. S. A.* 2002, 99, 15387–15392. [PubMed: 12426403]
18. Ten BHW; Gore M; Carmichael J; Gordon A; Malfetano J; Hudson I; Broom C; Scarabelli C; Davidson N; Spaczynski M; Bolis G; Malmström H; Coleman R; Fields SC; Heron JF Topotecan

- versus paclitaxel for the treatment of recurrent epithelial ovarian cancer. *J. Clin. Oncol.* 1997, 15, 2183–2193. [PubMed: 9196130]
19. Long HJ; Bundy BN; Grendys EC; Benda JA; McMeekin DS; Sorosky J; Miller DS; Eaton LA; Fiorica JV Randomized phase iii trial of cisplatin with or without topotecan in carcinoma of the uterine cervix: A gynecologic oncology group study. *J. Clin. Oncol.* 2005, 23, 4626–4633. [PubMed: 15911865]
 20. O'Brien MER; Ciuleanu TE; Tsekov H; Shparyk Y; ũ eviá B; Juhasz G; Thatcher N; Ross GA; Dane GC; Crofts T Phase iii trial comparing supportive care alone with supportive care with oral topotecan in patients with relapsed small-cell lung cancer. *J. Clin. Oncol.* 2006, 24, 5441–5447. [PubMed: 17135646]
 21. Cunningham D; Pyrhönen S; James RD; Punt CJA; Hickish TF; Heikkilä R; Johannesen TB; Starkhammar H; Topham CA; Awad L; Jacques C; Herait P Randomised trial of irinotecan plus supportive care versus supportive care alone after fluorouracil failure for patients with metastatic colorectal cancer. *The Lancet* 1998, 352, 1413–1418.
 22. Hartmann JT; Issels RD; Nicolo KS; Grünwald V; Hertenstein B; Papesch E; Krause S; Sturm I Topotecan plus cyclophosphamide in adults with relapsed or refractory pediatric-type sarcoma: A retrospective analysis from the german sarcoma medical oncology group (aio). *Invest. New Drugs* 2015, 33, 1115–1122. [PubMed: 26163339]
 23. Kitai Y; Kawasaki T; Sueyoshi T; Kobiyama K; Ishii KJ; Zou J; Akira S; Matsuda T; Kawai T DNA-containing exosomes derived from cancer cells treated with topotecan activate a sting-dependent pathway and reinforce antitumor immunity. *The Journal of Immunology* 2017, 198, 1649–1659. [PubMed: 28069806]
 24. Creemers G; Lund B; Verweij J Topoisomerase i inhibitors: Topotecan and irinotecan. *Cancer Treat. Rev.* 1994, 20, 73–96. [PubMed: 8293429]
 25. Rialdi A; Campisi L; Zhao N; Lagda AC; Pietzsch C; Ho JSY; Martinez-Gil L; Fenouil R; Chen X; Edwards M; Metreveli G; Jordan S; Peralta Z; Munoz-Fontela C; Bouvier N; Merad M; Jin J; Weirauch M; Heinz S; Benner C; van Bakel H; Basler C; García-Sastre A; Bukreyev A; Marazzi I Topoisomerase 1 inhibition suppresses inflammatory genes and protects from death by inflammation. *Science* 2016, 352, aad7993. [PubMed: 27127234]
 26. Singer M; Deutschman CS; Seymour CW; Shankarhari M; Annane D; Bauer M; Bellomo R; Bernard GR; Chiche JD; Coopersmith CM The third international consensus definitions for sepsis and septic shock (sepsis-3). *JAMA* 2016, 315, 801–810. [PubMed: 26903338]
 27. Crow RT; Crothers DM Structural modifications of camptothecin and effects on topoisomerase i inhibition. *J. Med. Chem.* 1992, 35, 4160–4164. [PubMed: 1331458]
 28. Wani MC; Ronman PE; Lindley JT; Wall ME Plant antitumor agents. 18. Synthesis and biological activity of camptothecin analogs. *J. Med. Chem.* 1980, 23, 554–560. [PubMed: 7381856]
 29. Nicholas AW; Wani MC; Manikumar G; Wall ME; Kohn KW; Pommier Y Plant antitumor agents. 29. Synthesis and biological activity of ring d and ring e modified analogues of camptothecin. *J. Med. Chem.* 1990, 33, 972–978. [PubMed: 2155323]
 30. Wani MC; Nicholas AW; Wall ME Plant antitumor agents. 28. Resolution of a key tricyclic synthon, 5'(rs)-1,5-dioxo-5'-hydroxy-2'h,5'h,6'h-6'-oxopyrano[3',4'-f].Delta.6,8-tetrahydroindolizine: Total synthesis and antitumor activity of 20(s)- and 20(r)-camptothecin. *J. Med. Chem.* 1987, 30, 2317–2319. [PubMed: 3681902]
 31. Martino E; Della Volpe S; Terribile E; Benetti E; Sakaj M; Centamore A; Sala A; Collina S The long story of camptothecin: From traditional medicine to drugs. *Bioorg. Med. Chem. Lett.* 2017, 27, 701–707. [PubMed: 28073672]
 32. Schrödinger, M. Version 9.0. Schrödinger LLC, New York, NY 2009.
 33. Cao Z; Harris N; Kozielski A; Vardeman D; Stehlin JS; Giovannella B Alkyl esters of camptothecin and 9-nitrocamptothecin: Synthesis, in vitro pharmacokinetics, toxicity, and antitumor activity. *J. Med. Chem.* 1998, 41, 31–37. [PubMed: 9438019]
 34. Zhao H; Lee C; Sai P; Choe YH; Boro M; Pendri A; Guan S; Greenwald RB 20-o-acylcampthothecin derivatives: Evidence for lactone stabilization. *J. Org. Chem.* 2000, 65, 4601–4606. [PubMed: 10959865]

35. Greenwald RB; Pendri A; Conover C; Gilbert C; Yang R; Xia J Drug delivery systems. 2. Camptothecin 20-o-poly(ethylene glycol) ester transport forms. *J. Med. Chem.* 1996, 39, 1938–1940. [PubMed: 8642549]
36. Deshmukh M; Chao P; Kutscher HL; Gao D; Sinko PJ A series of α -amino acid ester prodrugs of camptothecin: In vitro hydrolysis and a549 human lung carcinoma cell cytotoxicity. *J. Med. Chem.* 2010, 53, 1038–1047. [PubMed: 20063889]
37. Wadkins RM; Potter PM; Vladu B; Marty J; Mangold G; Weitman S; Manikumar G; Wani MC; Wall ME; Von Hoff DD Water soluble 20 (s)-glycinate esters of 10, 11-methylenedioxycamptothecins are highly active against human breast cancer xenografts. *Cancer Res.* 1999, 59, 3424–3428. [PubMed: 10416605]
38. Takimoto CH; Wright J; Arbuck SG Clinical applications of the camptothecins. *Biochim. Biophys. Acta* 1998, 1400, 107–119. [PubMed: 9748525]
39. Mathijssen RHJ; Loos WJ; Verweij J; Sparreboom A Pharmacology of topoisomerase i inhibitors irinotecan (cpt-11) and topotecan. *Curr. Cancer Drug Targets* 2002, 2, 103–123. [PubMed: 12188913]
40. Gilson MK; Liu T; Baitaluk M; Nicola G; Hwang L; Chong J Bindingdb in 2015: A public database for medicinal chemistry, computational chemistry and systems pharmacology. *Nucleic Acids Res.* 2016, 44, D1045–D1053. [PubMed: 26481362]

Figure 1.

Binding patterns of CPT in DNA-Topo I-CPT ternary complex (PDB entry: 1T8I). TGP is a sulfhydryl modified guanosine. DA, DT and DG represent DNA adenosine, thymidine and guanosine, respectively.

Figure 2.

(a) Aqueous solubility and maximum tolerated doses (MTD) of compounds **9**, **11** and topotecan. The acute and short-term toxicity was determined after i.v. administration of drugs (single administration or repeated daily administration for seven days) with increasing doses to mice; (b) Balb/C nude mice bearing HepG-2 hepatoma xenografts were treated with compound **9** at i.v. doses of 15 and 30 mg/kg or topotecan at i.v. dose of 2.0 mg/kg once a week for 30 days followed by 2 weeks' observation; (c) Nude mice bearing HepG-2 hepatoma xenografts were treated with compound **11** at oral dose of 1.3 mg/kg once a day and i.v. doses of 4.0 and 8.0 mg/kg once a week for 30 days followed by 2 weeks' observations. Tumor volume was recorded and plotted against time, in days.

Figure 3.

Topo I inhibitory activity of CPT, compound **9** and **11** at 50 μ M (a), and at 100 μ M (b). Samples were analyzed on 1% EB-free agarose gel. Lane 1, super-coiled DNA plasmid (DNA); lane 2, DNA + Topo I; lane 3, DNA + Topo I + CPT; lane 4, DNA + Topo I + compound **9**; lane 5, DNA + Topo I + compound **11**. (c) Topo I-mediated DNA cleavage gels. Samples were analyzed on 1% EB-containing agarose gel. Lane 1, DNA; lane 2, DNA + Topo I; lane 3, DNA + Topo I + 50 μ M Topotecan; lane 4, DNA + Topo I + 50 μ M CPT; lane 5, DNA + Topo I + 100 μ M compound **9**; lane 6, DNA + Topo I + 50 μ M compound **9**; lane 7, DNA + Topo I + 100 μ M compound **11**; lane 8, DNA + Topo I + 50 μ M compound **11**.

Figure 4.

(a) Topo I inhibitory activity of CPT, compound **4** and **3** at 50 μ M. Samples were analyzed on 1% EB-free agarose gel. Lane 1, super-coiled DNA plasmid (DNA); lane 2, DNA + Topo I; lane 3, DNA + Topo I + CPT; lane 4, DNA + Topo I + compound **4**; lane 5, DNA + Topo I + compound **3**. (b) Topo I-mediated DNA cleavage gels. Samples were analyzed on 1% EB-containing agarose gel. Lane 1, DNA; lane 2, DNA + Topo I; lane 3, DNA + Topo I + CPT; lane 4, DNA + Topo I + compound **4**; lane 5, DNA + Topo I + compound **3**

Figure 5.

Change in body weight of the treated mice. (a) Balb/C nude mice bearing HepG-2 hepatoma xenografts were treated with compound **9** at i.v. doses of 15 or 30 mg/kg or topotecan at an i.v. dose of 2.0 mg/kg once a week for 30 days; (b) Nude mice bearing HepG-2 hepatoma xenografts were treated with compound **11** at an oral dose of 1.3 mg/kg once a day and i.v. doses of 4.0 and 8.0 mg/kg once a week for 30 days. Body weight percentage change was plotted against time, in days.

Figure 6.

Compounds **9** and **11** protect against LPS-induced sepsis *in vivo*. (a) Survival curves of C57BL/6C mice treated with compound **9** in response to LPS injection; (b) Survival curves of mice treated with compound **11** in response to LPS injection; (c) Expression of IL-6 in mice serum; (d) Expression of TNF-α in mice serum.

Figure 7.

Pharmacokinetic properties of compound **9** and **11**. Pharmacokinetics were studied in male C57BL/6 mice after (a) single intravenous administration of 2 mg/kg Compound **9**; (b) single intravenous administration of 2 mg/kg Compound **11**; (c) single oral administration of 10 mg/kg Compound **11**; (d) single intravenous administration of 10 mg/kg Compound **11**. Plasma levels of the compounds were determined by LC-MS/MS analysis at the indicated time points.

Scheme 1.

Equilibrium of between the lactone form and the open carboxylate form of camptothecin.

Scheme 2.

Multi-substitution at 7, 9, 10 and 11 positions.

Chart 1.
Structures and synthesis of hexacyclic CPT derivatives.

Table 1.

Anticancer activity of hexacyclic CPTs against eleven human cancer cell lines.

Cell line	Compound IC ₅₀ (μM) ^a											
	Topotecan	1	2	3	4	5	6	7	8	9	10	11
A549	1.424	1.028	2.057	2.169	1.394	1.124	3.371	1.502	0.092	0.170	1.832	6.472
MCF-7	2.135	1.157	2.314	10.846	4.640	2.135	7.865	6.438	> 10	> 10	> 10	> 10
MDA-MB-231	1.898	0.488	1.542	1.323	1.439	0.562	2.135	1.996	0.348	0.425	0.641	5.016
HepG-2	0.237	0.064	0.180	1.106	0.016	0.063	0.079	0.071	0.055	0.060	0.348	0.971
KB	2.610	3.342	2.571	9.761	2.204	2.247	2.202	2.039	0.916	0.935	7.143	7.282
BEL-7402	7.118	10.54	2.442	> 10	3.480	2.022	2.135	1.931	0.275	0.425	2.015	> 10
SMMC-7721	11.627	> 10	3.085	> 10	10.441	> 10	11.011	> 10	> 10	8.503	> 10	> 10
MGC80-3	0.190	0.514	0.231	0.390	0.018	0.085	0.090	0.054	0.064	0.062	0.275	0.971
SK-N-SH	0.093	0.463	NT ^b	0.325	NT ^b	NT ^b	NT ^b	NT ^b	0.348	0.51	0.549	0.194
QGY-7703	11.864	> 10	11.568	> 10	11.601	> 10	> 10	> 10	> 10	> 10	> 10	> 10
SGC-7901	> 10	> 10	> 10	> 10	> 10	> 10	> 10	> 10	> 10	> 10	> 10	> 10

^aValues are the mean of at least three data points.^bNot Tested.