Published in final edited form as:

Eur J Clin Nutr. 2019 March ; 73(3): 474–482. doi:10.1038/s41430-018-0285-9.

HIV and food insecurity are associated with body composition changes among pregnant and lactating Kenyan women

Elizabeth M. Widen,

Department of Nutritional Sciences, University of Texas at Austin

Irene Tsai,

School of General Studies, Columbia University

Shalean M. Collins,

Department of Anthropology, Northwestern University

Pauline Wekesa,

Kenya Medical Research Institute

Joy China,

Kenya Medical Research Institute

Natalie Krumdieck,

Weill Cornell Medical College

Joshua Miller,

Department of Anthropology, Northwestern University

Sheri D. Weiser,

School of Medicine, University of California San Francisco

Maricianah Onono,

Kenya Medical Research Institute

Sera L. Young

Department of Anthropology, Northwestern University

Institute for Policy Research, Northwestern University

Abstract

Background/Objectives: Body composition changes markedly during reproduction. In sub-Saharan Africa, impacts of HIV-infection on body composition across pregnancy and lactation in the context of Option B+ antiretroviral therapy are unknown. Therefore, we sought to evaluate the role of HIV-infection on body composition during pregnancy and lactation among Kenyan women.

CORRESPONDENCE: Sera L. Young, Department of Anthropology, Institute for Policy Research, Northwestern University, 515 Clark Street, Evanston, IL 60201, Phone: 847-491-5402, Fax: 847-467-1778, sera.young@northwestern.edu.

Supplementary information is available at *European Journal of Clinical Nutrition's* website.

Conflict of interest. Research activities and Sera Young were supported by the National Institute of Mental Health. Elizabeth Widen has received funding from the National Institutes of Health, and the Thrasher Research Fund. From August 2014-July 2015, Widen was supported by an unrestricted postdoctoral fellowship from PepsiCo Global R+D. Sheri Weiser has received funding from NIMH, NIAID, and Kaiser Community Benefits. Tsai, Collins, Wekesa, China, Krumdieck, Miller, and Onono declare no conflicts of interest.

Subjects/Methods: A cohort of pregnant women (n=333; 50.5% HIV+, receiving ART) were enrolled at 7 clinics in western Kenya. Two prenatal (mean±SD: 23.6±4.4 & 33.4±2.0 weeks gestation) and three postpartum (6, 14 and 36 weeks) measurements included: individual-level food insecurity, height, weight, fat mass (FM) and fat free mass (FFM) by bioimpedance analysis (BIA), mid-upper arm circumference (MUAC), triceps skinfold (TSF), allowing for AMA (arm muscle area) and AFA (arm fat area) derivation. Multivariable longitudinal regression models were used to relate HIV to body composition changes.

Results: In longitudinal models, HIV-infected women had lower weight ($\beta = -3.0$ kg, $p=0.003$), fat mass ($\beta = -1.5$ kg, $p=0.02$), fat-free mass ($\beta = -1.5$ kg, $p=0.01$), TSF ($\beta = -2.6$ mm, $p<0.001$), AFA ($\beta = -3.9$ cm³, $p<0.001$), and MUAC ($\beta = -1.0$ cm, $p=0.001$), but not AMA ($p=0.34$), across all observations. Food insecurity was inversely associated with AMA and MUAC postpartum (AMA β -range=-0.47 to -0.92 cm³; MUAC β -range=-0.09 to -0.15 cm, all $p<0.05$).

Conclusions: HIV-infection was associated with lower weight, fat mass, fat-free mass, TSF, AFA and MUAC values during pregnancy and lactation, while food insecurity was intermittently associated with body composition. This suggests that HIV+ and food insecure pregnant and lactating women could benefit from nutritional support.

INTRODUCTION

Body composition is an important indicator of nutritional status and health. Across life stages, body composition is influenced by both diet and infection. During pregnancy and lactation, body composition changes dynamically to support the growing fetus and infant.^{1, 2} During pregnancy, women gain fat and fat-free mass with concomitant shifts in regional body fat depots, typified by gains in arm fat area (AFA) in early and mid-pregnancy, followed by declines thereafter.^{1, 3, 4} Body composition changes are highly variable during lactation; most women experience mild and gradual weight loss, while some remain relatively weight-stable or gain weight.²

No studies have examined body composition changes across pregnancy and lactation among HIV-infected (HIV+) and HIV-uninfected (HIV-) women in a single cohort in the context of widespread adoption of Option B+ antiretroviral therapy (ART).⁵ Among HIV+ women, lower body composition and anthropometry values have been linked to adverse pregnancy outcomes,⁶⁻⁸ sub-optimal infant growth⁹, and mother-to-child transmission of HIV.¹⁰ Postnatal ART may exacerbate weight loss during six months of exclusive breastfeeding (EBF).¹¹ ART is now widely administered in accordance with World Health Organization (WHO) recommendations (e.g. initiation of lifelong ART “Option B+” at HIV diagnosis), such that understanding its role in body composition is important.⁵

Timing of HIV diagnosis relative to pregnancy could also potentially influence body composition, but to date there are no data supporting this. Body composition could be altered by the initiation and duration of ART. Further, newly diagnosed women may experience emotional distress and disengagement from care which can be deleterious for nutritional status.¹²

Food insecurity, HIV, and nutrition are linked in a multi-directional relationship, intensified during periods of heightened nutritional needs, such as reproduction or HIV infection.^{13, 14} For example, among HIV+ and HIV- Ugandan women, food insecurity was associated with adverse changes in postpartum weight and AFA.¹⁵

We examined whether HIV infection was associated with body composition changes during pregnancy and lactation among Kenyan women of mixed HIV status. We hypothesized that HIV+ women receiving ART would exhibit lower weight and greater loss of overall and regional lean mass than HIV-negative women. Relatedly, we examined how HIV diagnosis during the index pregnancy with concomitant initiation of Option B+ influenced body composition compared to HIV-diagnosis prior to the index pregnancy. Our second objective was to evaluate how food insecurity influenced body composition. We hypothesized that food insecurity would be inversely associated with body composition.

METHODS

Data are from the Pith Moromo Study, a longitudinal observational cohort designed to examine the health consequences of food insecurity in HIV+ and HIV- women in Nyanza Region, Kenya (). Study enrollment was between August 8, 2014 and February 2, 2015, and follow-up visits were conducted until April 26, 2016.

All participants received antenatal care (ANC) at Family AIDS Care and Educational Services (FACES) clinics in Kisumu, Nyahera, Rongo, Ongo, Migori, Macalder, or Nyamaraga. These clinics deliver integrated, comprehensive HIV care, treatment and prevention services. HIV+ women received comprehensive HIV services, including Option B+ ART. At the first ANC visit, women (n=371) were invited to participate if they met the following criteria: gestational age <30 weeks (assessed using last menstrual period abstracted from ANC cards), living within the study area and not planning to move, and known HIV status (assessed at ANC). Food insecurity in the past month was determined using the 9-item Individual Food Insecurity Access Scale (range: 0–27).¹⁶ Sampling was purposive by HIV and food insecurity status (Figure 1, Supplemental Table 1), such that equal numbers of HIV+ and HIV- women by tertiles of food insecurity scores (low 0–9, moderate 10–18, and severe 19–27) were enrolled. We sought to enroll equal numbers of women with a known HIV+ status (Known Positive (KP); HIV diagnosis prior to index pregnancy), and a new HIV+ diagnosis (Newly diagnosed (ND); HIV diagnosis during index pregnancy). Three hundred seventy-one women were enrolled in parent study, which was determined to be sufficient to detect differences in the primary outcomes of the study (e.g. maternal BMI) by FI status at a power of 0.8.

Visits antenatally were at enrollment (Mean±SD: 23.7±4.4 weeks gestation) and follow-up (33.4±2.0 weeks gestation), and postnatally at 6 (6.5±0.91 weeks) and 14 weeks (14.6±0.88 weeks) and 9 months (39.8±1.8 weeks), with two phone calls at 1 week and 6 months. Measurements obtained by trained research staff included: weight (Seca 874, Seca North America, Chino, CA), height (Seca 206, Seca North America, Chino, CA), mid-upper arm circumference (MUAC) with a non-stretchable retractable tape measure and triceps skinfold thickness (TSF) (Harpender calipers, Baty International, United Kingdom). Arm muscle

area (AMA) and arm fat area (AFA) were calculated.¹⁵ Bioelectrical impedance analysis (BIA) [BIA 450, Biodynamics, Seattle, Washington] was used to estimate fat-free mass and fat mass. Meeting water needs is often challenging in sub-Saharan Africa¹⁷ and many women in this cohort reported at least one experience of water insecurity in the previous month.¹⁸ To ensure hydration, which results in underestimation of fat mass by BIA¹⁹, participants consumed approximately 8oz of water and BIA was conducted approximately 90 minutes later.

At enrollment, self-report of education, marital status, and age were obtained. Household socioeconomic status was derived using principal components analysis from self-report of household assets adapted from the Kenya Integrated Household Budget Survey questionnaire²⁰, where higher scores indicate greater wealth. Measures at all visits included the Center for Epidemiologic Studies Depression (CES-D) scale,²¹ self-report of breastfeeding, morbidity (symptoms experienced at time of visit), and dietary diversity²² and time use activities (heavy/light work/sleep/leisure) within the last 24 hours. For HIV+, ART regimen and WHO clinical staging²³ were abstracted from clinical records and ANC cards.

Institutional Review Boards at Cornell University, Northwestern University and Kenya Medical Research Institute approved study procedures for Pith Moromo. All mothers provided written informed consent.

Statistical analyses

Statistical analyses were conducted using Stata 14.0 software with an α of <0.05 . Singleton participants were included in analyses if they had 2 study visits and enrollment body composition data (Figure 1). Baseline characteristics were compared between HIV+ and HIV- mothers using parametric tests for continuous, normally distributed variables, and non-parametric tests for skewed variables. Enrollment occurred at varying gestational ages, so a constructed prenatal visit timing was derived: early 2nd trimester (<21 wks); late 2nd trimester (21– <28 wk); early 3rd trimester (28– <32 wk); mid-3rd trimester (32– <36 wk), and late 3rd trimester (≥ 36 wk).

Random-effects longitudinal models were fit for each body composition outcome with indicator variables for visit and study site. Primary exposures were HIV status and time-varying individual food insecurity (continuous score). We evaluated if body composition changes varied by HIV status or food insecurity by including interaction terms between these factors and visit.

Covariates were included if the HIV β -coefficient changed by $>10\%$ in one or more of the multivariable models. Time-independent variables included: maternal height (cm), age (years), education [categorical: primary (ref), secondary, tertiary], parity (continuous), assets (continuous), and probable depression at enrollment (CES-D score ≥ 17). Time-varying variables included: fever, gastrointestinal distress, dietary diversity (score), hunger season in previous month (e.g. February to May), and proportion of heavy work to light work.

Longitudinal random-effects models evaluated associations between timing of HIV diagnosis and body composition. In HIV+ women with WHO clinical staging data (n=141, 84% of HIV+), we conducted a sensitivity analysis with adjustment for clinical stage [categorical: Stage 1 Asymptomatic (reference); Stage 2 (Mild-Severe)]. To evaluate whether breastfeeding impacted changes postpartum, we conducted a sensitivity analysis using postpartum body composition models with adjustment for EBF linked to concurrent measurements at 6 wk and 14 wk, and lagged at 9 mo (i.e. EBF at 6 mo used at 9 mo visit).

RESULTS

Body composition data were available for 333 mothers (Figure 1). Characteristics of included women were not different from those not included except that included women had lower fat mass at enrollment (18.5 ± 5.8 vs. 21.5 ± 13.8 kg, $p=0.02$).

At enrollment, most characteristics were similar between HIV+ and HIV- women (Table 1) with differences in age, parity, fever and EBF. The majority of HIV+ women were classified as WHO clinical stage I (52.4%) and II (36.7%) HIV-infection, and were receiving tenofovir/lamivudine/efavirenz (TDF/3TC/EVF) (57.7%) ART regimen (Supplemental Table 2).

Generally, HIV+ women exhibited lower mean body composition values over time, with more pronounced differences postnatally (Table 2). While some differences were not statistically significant (e.g. weight and AMA), values for TSF, AFA and fat-free mass were lower in late pregnancy and postpartum for HIV+.

In models of overall maternal body composition over time (Supplemental Table 3), we observed increases in weight during pregnancy, decreases in weight and fat-free mass postpartum, and no significant changes in fat mass. For regional body composition (Supplemental Table 4), we observed decreases in TSF and AFA across pregnancy and postpartum, and increases in AMA at visits 36 wk and MUAC at 9 mo.

The interaction between time and HIV status was not significant, thus HIV was included as a covariate. HIV was negatively associated with most body composition values (Figure 2A). Specifically, HIV-infection was associated with lower weight, fat-mass, fat-free mass, TSF, AFA and MUAC from pregnancy to 9 months postpartum, and was not associated with AMA.

Effects of food insecurity varied over time for TSF, AMA and MUAC (Figure 2B). Specifically, greater food insecurity was associated with marginally lower TSF and higher AMA at the early 2nd trimester visit, whereas during lactation, food insecurity was associated with lower AMA and MUAC, and greater TSF, suggesting muscle loss over time. Food insecurity was not associated with other outcomes.

Other factors associated with body composition included maternal age, height, probable depression, light and heavy work, hunger season in previous month, and gastrointestinal distress (Supplemental Tables 3&4).

Further, we examined the timing of HIV diagnosis as the primary exposure. Some characteristics differed depending on timing of diagnoses (Supplemental Table 5). In the models (Supplemental Table 5), newly diagnosed women did not have significant differences in body composition compared to women diagnosed prior to the index pregnancy. The effect sizes for body composition values were predominately negative among newly diagnosed women. In our sensitivity analysis evaluating the role of HIV clinical staging on body composition across pregnancy and lactation in the subset with staging data ($n=141$), we found that WHO clinical stage 2 was associated with lower AFA and TSF over time [AFA $b=-2.8 \text{ cm}^3$ (95% CI: $-5.2, -0.3$), $p=0.03$; TSF $b=-1.9 \text{ mm}$ (95% CI: $-3.5, -0.2$), $p=0.03$], but not with overall body composition (weight, fat mass, fat-free mass) or other regional measures (AMA, MUAC) (Data not shown).

Because breastfeeding may influence postpartum body composition, and we conducted additional analyses to explore this. Adjustment for breastfeeding did not appreciably alter associations between HIV and maternal body composition (data not shown). Furthermore, exclusive breastfeeding did not predict changes in body composition (data not shown).

DISCUSSION

In this first study of body composition changes across both pregnancy and lactation among women of mixed HIV status, HIV-infection was associated with lower weight, fat mass, fat-free mass, TSF, AFA and MUAC values over time. This is in support of our first hypothesis that HIV would be associated with adverse changes in body composition. Regarding timing of HIV diagnosis relative to the pregnancy, we observed that HIV diagnosis in the index pregnancy was associated with worse body composition measures over time. This suggests that newly diagnosed women may need increased nutritional support during pregnancy and lactation.

Our second hypothesis, that food insecurity would lead to adverse changes in body composition, was only supported by some of our findings. Greater food insecurity was associated intermittently with regional changes in body composition in the arm (TSF, AMA and MUAC), suggesting muscle tissue loss over time. However, food insecurity was not associated with most body composition outcomes, including overall body composition (weight, fat mass and fat-free mass) and regional arm fat depots.

Similar to other reports in HIV⁺^{4, 24, 25} and HIV⁻ populations,^{3, 26} participants exhibited changes in overall and regional body composition. For overall body composition, in longitudinal multivariate models, we observed increases in weight during pregnancy and decreases postpartum. Interestingly, we observed only significant decreases in lean mass at postpartum visits, and observed no significant differences in fat mass over time. This suggests that maternal weight changes were a proportional change in fat and lean mass during pregnancy, and that the weight changes were largely leaner mass during lactation. These decreases in overall lean mass postpartum are likely due to the large decreases in body water and shifts in other supporting tissues after delivery (e.g. uterus).^{1, 2} For regional body composition, TSF and AFA decreased across pregnancy and lactation, with the largest decrease at 9 months postpartum, suggesting arm fat was mobilized to support

breastfeeding; this is similar to a report in HIV-uninfected rural Taiwanese women.³ AMA increased from late pregnancy and across lactation, which is comparable to a previous study reporting effects in ART-naïve HIV+ Malawian women during pregnancy.⁴ At 9 months postpartum, MUAC was higher in our cohort of Kenyan women, which is similar to our findings in Ugandan HIV+ and HIV- women.¹⁵

Unadjusted overall body composition values were not markedly different across pregnancy and lactation by maternal HIV status, although some differences by HIV were observed for regional measures of arm fat (TSF & AFA). When adjusting for maternal height and covariates, maternal HIV-infection was associated with lower overall body composition values (weight, fat-mass, and lean-mass), and also regional body composition values (TSF and AFA) across pregnancy and postpartum. This was not so for regional arm muscle depots (AMA). This suggests that HIV+ women may be more prone to adverse changes in body composition during pregnancy and lactation. It is difficult to compare these findings to other studies, because prior work has only examined body composition changes during lactation by HIV status. In these two previous studies, there were no observable differences by HIV status in overall^{15, 24} or regional body composition¹⁵ at 8–24 wk postpartum²⁴ and 1 wk to 12 mo postpartum.¹⁵

In addition to the main effects of HIV-infection, we also evaluated if timing of HIV diagnosis and ART initiation were associated with changes in body composition. Although most body composition effects were lower for new HIV diagnoses, suggesting a trend towards lower values in this group, there were no significant differences between newly diagnosed and known positive groups. To our knowledge, there are no data on timing of HIV diagnosis and body composition during pregnancy and lactation to which we can compare these observations.

Similar to our previous work, in which food insecurity was inversely associated with weight and AFA among HIV+ and HIV- postpartum Ugandan women receiving ART¹⁵, we observed some adverse effects of higher food insecurity scores on body composition during lactation. However, in contrast to our previous study in Uganda,¹⁵ food insecurity was not associated with overall body composition in our Kenyan cohort. Instead, we found that food insecurity was intermittently associated with regional body composition values. Specifically, food insecurity was associated with lower AMA and MUAC postpartum, suggesting muscle tissue loss. These differential findings could be attributable to different periods of follow up (9 vs. 12 mo), hunger season exposure, or more pronounced food insecurity in Nyanza, Kenya.

In our sensitivity analysis with adjustment for EBF, associations between HIV and postpartum body composition were not altered. This is similar to our previous findings in HIV+ and HIV- Ugandan women,¹⁵ and among well-nourished HIV+ women receiving ART from Kenya, Burkina Faso and South Africa, where breastfeeding was not associated with weight loss.²⁷

We identified several other predictors of body composition changes (Supplemental Tables 3&4). Maternal age was positively associated with several measures, suggesting that

younger women may be at higher risk for poor nutritional outcomes. Probable maternal depression during pregnancy was associated with lower weight across pregnancy and lactation and a trend toward lower fat and lean mass over time. Previously, in HIV+ and HIV- Ugandan women, maternal depression was associated with weight and lean mass loss postpartum.¹⁵ Together, these findings suggest that women with impaired mental health may be more vulnerable to adverse nutrition outcomes.

Finally, we observed that exposure to the hunger season (e.g. February to May), when food stores run low, was associated with lower AMA and MUAC over time. This suggests that this period may be particularly important for nutritional interventions. This relationship is supported by findings from Uganda, where exposure to hunger season was associated with lower AMA during breastfeeding.¹⁵ While exposure to hunger season was associated with lower AMA during pregnancy in the HIV+ Malawian cohort⁴, enrollment and follow-up timing (2003–2009) may have allowed for more robust examinations of the effects of food shortages on body composition, compared to our study, where women were enrolled during a relatively short period of time (~6mo: August 2014–February 2015).

The strengths of this study include evaluation of many indicators of body composition along with a variety of environmental and demographic characteristics. Limitations include the differential enrollment timing across study participants during pregnancy, which precluded adjustment for most initial body composition values in the longitudinal models. The rurality of some of the clinics and distance between study sites necessitated the use of a portable BIA, which precluded us from using more robust body composition assessment methods. There are concerns with the validity of BIA during pregnancy;¹ however, BIA has been used in many other studies of body composition during pregnancy^{28–31} and it has been shown to predict total body water during pregnancy,³² which is the primary factor used in deriving fat-free mass. The Biodynamics BIA 450 does not have race specific proprietary equations for predicting body fat, and developed equations from a range of body types, weights, percentage fat, and ethnic heritage (Personal communications with Biodynamics Corporation, July 19, 2018). It is unknown whether race-specific equations would improve our body composition estimates.

There may also be unaccounted confounders that could explain differences between HIV+ and HIV- women. Although we had WHO clinical staging, viral load and HIV-specific comorbidities would have provided a more comprehensive picture. Due to the breadth of ART regimens, we were unable to evaluate how specific ART regimens related to body composition. Although we measured dietary diversity, we were unable to measure dietary and energy intakes. Furthermore, we assessed time use activities such as heavy or light work, but were unable to robustly measure physical activity with a gold-standard method such as accelerometry. Finally, because HIV status was determined in ANC clinics, we may have had imprecise dates of diagnoses.

In conclusion, HIV-infection was associated with adverse changes in overall and regional body composition across pregnancy and lactation. Food insecurity was associated with greater loss of arm muscle and MUAC during lactation. Adverse changes in body composition were associated with HIV, lower maternal age, physical exertion, and

depression. Taken together, this suggests that women with these characteristics may need nutritional support to ensure adequate reserves to support breastmilk production, and maintain maternal health during pregnancy and lactation.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements:

We would like to thank the Kenya Medical Research Institute (KEMRI) for providing space within the FACES clinics for the work to be conducted, ethical and logistic support to conduct this research, and oversight in Kenya. We would also like to warmly thank our study nurses: Joy China, Joyce Boke, and Tobias Odwar, and our study trackers: Benter Ogwana, Teresa Owade, and Sarah Obaje, as well as the mothers and their infants who participated in this study.

SUPPORT: Research activities and SLY were supported by the National Institute of Mental Health (NIH/NIMH K01MH098902). EW was supported by the Eunice Kennedy Shriver National Institute of Child Health & Human Development (NIH/NICHD K99HD086304); the National Institute of Diabetes and Digestive and Kidney Diseases (NIH/NIDDK T32DK091227; T32 DK007559) and by an unrestricted grant to support research in maternal and child health from PepsiCo Global R+D. The content is solely the responsibility of the authors and does not necessarily represent the views of the National Institutes of Health.

This trial was registered at <http://clinicaltrials.gov> as .

ABBREVIATIONS:

AMA	arm muscle area
AFA	arm fat area
FM	fat mass
FFM	fat-free mass
MUAC	mid-upper arm circumference
TSF	triceps skinfold thickness
WHO	World Health Organization

REFERENCES

1. Widen EM, Gallagher D. Body composition changes in pregnancy: measurement, predictors and outcomes. *Eur J Clin Nutr* 2014; 68(6): 643–652. e-pub ahead of print 2014/03/29; doi: 10.1038/ejcn.2014.40 [PubMed: 24667754]
2. Butte NF, Hopkinson JM. Body composition changes during lactation are highly variable among women. *J Nutr* 1998; 128(2 Suppl): 381s–385s. e-pub ahead of print 1998/03/21; [PubMed: 9478031]
3. Adair LS, Pollitt E, Mueller WH. Maternal anthropometric changes during pregnancy and lactation in a rural Taiwanese population. *Hum Biol* 1983; 55(4): 771–787. e-pub ahead of print 1983/12/01; [PubMed: 6674101]
4. Ramlal RT, Tembo M, Soko A, Chigwenembe M, Tohill BC, Kayira D et al. Patterns of body composition among HIV-infected, pregnant Malawians and the effects of famine season. *Matern Child Health J* 2013; 17(2): 265–273. e-pub ahead of print 2012/03/08; doi: 10.1007/s10995-012-0970-6 [PubMed: 22395817]

5. World Health Organization. Guideline on when to start antiretroviral therapy and on pre-exposure prophylaxis for HIV. WHO Press: Geneva, Switzerland, 2015.
6. Ramlal RT, Tembo M, Soko A, Chigwenembe M, Ellington S, Kayira D et al. Maternal mid-upper arm circumference is associated with birth weight among HIV-infected Malawians. *Nutr Clin Pract* 2012; 27(3): 416–421. e-pub ahead of print 2012/04/19; doi: 10.1177/0884533611435991 [PubMed: 22511656]
7. Villamor E, Dreyfuss ML, Baylin A, Msamanga G, Fawzi WW. Weight loss during pregnancy is associated with adverse pregnancy outcomes among HIV-1 infected women. *J Nutr* 2004; 134(6): 1424–1431. e-pub ahead of print 2004/06/03; [PubMed: 15173407]
8. Young S, Murray K, Mwesigwa J, Natureeba P, Osterbauer B, Achan J et al. Maternal nutritional status predicts adverse birth outcomes among HIV-infected rural Ugandan women receiving combination antiretroviral therapy. *PLoS One* 2012; 7(8): e41934 e-pub ahead of print 2012/08/11; doi: 10.1371/journal.pone.0041934 [PubMed: 22879899]
9. Widen EM, Bentley ME, Kayira D, Chasela CS, Jamieson DJ, Tembo M et al. Maternal weight loss during exclusive breastfeeding is associated with reduced weight and length gain in daughters of HIV-infected Malawian women. *J Nutr* 2013; 143(7): 1168–1175. e-pub ahead of print 2013/05/24; doi: 10.3945/jn.112.171751 [PubMed: 23700341]
10. Villamor E, Saathoff E, Msamanga G, O'Brien ME, Manji K, Fawzi WW. Wasting during pregnancy increases the risk of mother-to-child HIV-1 transmission. *J Acquir Immune Defic Syndr* 2005; 38(5): 622–626. e-pub ahead of print 2005/03/29; [PubMed: 15793376]
11. Kayira D, Bentley ME, Wiener J, Mkhomawanthu C, King CC, Chitsulo P et al. A lipid-based nutrient supplement mitigates weight loss among HIV-infected women in a factorial randomized trial to prevent mother-to-child transmission during exclusive breastfeeding. *The American journal of clinical nutrition* 2012; 95(3): 759–765. doi: 10.3945/ajcn.111.018812 [PubMed: 22258269]
12. Phillips T, Thebus E, Bekker LG, McIntyre J, Abrams EJ, Myer L. Disengagement of HIV-positive pregnant and postpartum women from antiretroviral therapy services: a cohort study. *J Int AIDS Soc* 2014; 17: 19242 e-pub ahead of print 2014/10/11; doi: 10.7448/ias.17.1.19242 [PubMed: 25301494]
13. Young SL, Plenty AH, Luwedde FA, Natamba BK, Natureeba P, Achan J et al. Household food insecurity, maternal nutritional status, and infant feeding practices among HIV-infected Ugandan women receiving combination antiretroviral therapy. *Matern Child Health J* 2014; 18(9): 2044–2053. e-pub ahead of print 2014/03/04; doi: 10.1007/s10995-014-1450-y [PubMed: 24585398]
14. Weiser SD, Young SL, Cohen CR, Kushel MB, Tsai AC, Tien PC et al. Conceptual framework for understanding the bidirectional links between food insecurity and HIV/AIDS. *The American journal of clinical nutrition* 2011; 94(6): 1729s–1739s. e-pub ahead of print 2011/11/18; doi: 10.3945/ajcn.111.012070 [PubMed: 22089434]
15. Widen EM, Collins SM, Khan H, Biribawa C, Acidri D, Achoko W et al. Food insecurity, but not HIV-infection status, is associated with adverse changes in body composition during lactation in Ugandan women of mixed HIV status. *The American journal of clinical nutrition* 2017; 105(2): 361–368. e-pub ahead of print 2017/01/06; doi: 10.3945/ajcn.116.142513 [PubMed: 28052888]
16. Natamba BK, Kilama H, Arbach A, Achan J, Griffiths JK, Young SL. Reliability and validity of an individually focused food insecurity access scale for assessing inadequate access to food among pregnant Ugandan women of mixed HIV status. *Public Health Nutr* 2015; 18(16): 2895–2905. e-pub ahead of print 2014/08/30; doi: 10.1017/s1368980014001669 [PubMed: 25171462]
17. Keino S, van den Borne B, Plasqui G. Body composition, water turnover and physical activity among women in Narok County, Kenya. *BMC public health* 2014; 14: 1212 e-pub ahead of print 2014/11/26; doi: 10.1186/1471-2458-14-1212 [PubMed: 25420458]
18. Krumdieck NR, Collins SM, Wekesa P, Mbullo P, Boateng GO, Onono M et al. Household water insecurity is associated with a range of negative consequences among pregnant Kenyan women of mixed HIV status. *Journal of water and health* 2016; 14(6): 1028–1031. e-pub ahead of print 2016/12/14; doi: 10.2166/wh.2016.079 [PubMed: 27959881]
19. Thompson DL, Thompson WR, Prestridge TJ, Bailey JG, Bean MH, Brown SP et al. Effects of hydration and dehydration on body composition analysis: a comparative study of bioelectric impedance analysis and hydrodensitometry. *The Journal of sports medicine and physical fitness* 1991; 31(4): 565–570. e-pub ahead of print 1991/12/01; [PubMed: 1806735]

20. Kenya Ministry of Planning and National Development CBoS. Kenya Integrated Household Budget Survey (KIHBS), 2004/05, Household Questionnaire. In, 2004.
21. Natamba BK, Achan J, Arbach A, Oyok TO, Ghosh S, Mehta S et al. Reliability and validity of the center for epidemiologic studies-depression scale in screening for depression among HIV-infected and -uninfected pregnant women attending antenatal services in northern Uganda: a cross-sectional study. *BMC Psychiatry* 2014; 14: 303 e-pub ahead of print 2014/11/25; doi: 10.1186/s12888-014-0303-y [PubMed: 25416286]
22. Food and Agriculture Organization. Guidelines for measuring household and individual dietary diversity. Food and Agriculture Organization: Rome, Italy, 2010.
23. World Health Organization. WHO case definitions of HIV for surveillance and revised clinical staging and immunological classification of HIV-related disease in adults and children. In: WHO Press, 2007.
24. Papathakis PC, Van Loan MD, Rollins NC, Chantry CJ, Bennish ML, Brown KH. Body composition changes during lactation in HIV-infected and HIV-uninfected South African women. *J Acquir Immune Defic Syndr* 2006; 43(4): 467–474. e-pub ahead of print 2006/09/19; doi: 10.1097/01.qai.0000243094.42276.92 [PubMed: 16980904]
25. Villamor E, Msamanga G, Spiegelman D, Peterson KE, Antelman G, Fawzi WW. Pattern and predictors of weight gain during pregnancy among HIV-1-infected women from Tanzania. *J Acquir Immune Defic Syndr* 2003; 32(5): 560–569. e-pub ahead of print 2003/04/08; [PubMed: 12679710]
26. Robinson JJ. Changes in body composition during pregnancy and lactation. *Proc Nutr Soc* 1986; 45(1): 71–80. e-pub ahead of print 1986/02/01; [PubMed: 3703866]
27. Cames C, Cournil A, de Vincenzi I, Gaillard P, Meda N, Luchters S et al. Postpartum weight change among HIV-infected mothers by antiretroviral prophylaxis and infant feeding modality in a research setting. *AIDS (London, England)* 2014; 28(1): 85–94. e-pub ahead of print 2014/01/15; doi: 10.1097/01.aids.0000433243.24481.c3
28. Toro-Ramos T, Sichieri R, Hoffman DJ. Maternal fat mass at mid-pregnancy and birth weight in Brazilian women. *Annals of human biology* 2016; 43(3): 212–218. e-pub ahead of print 2015/09/24; doi: 10.3109/03014460.2015.1032348 [PubMed: 26392036]
29. Fattah C, Farah N, Barry SC, O'Connor N, Stuart B, Turner MJ. Maternal weight and body composition in the first trimester of pregnancy. *Acta obstetrica et gynecologica Scandinavica* 2010; 89(7): 952–955. e-pub ahead of print 2010/04/13; doi: 10.3109/00016341003801706 [PubMed: 20380598]
30. Kent E, O'Dwyer V, Fattah C, Farah N, O'Connor C, Turner MJ. Correlation between birth weight and maternal body composition. *Obstetrics and gynecology* 2013; 121(1): 46–50. e-pub ahead of print 2012/12/13; doi: <http://10.1097/AOG.0b013e31827a0052> [PubMed: 23232753]
31. Larciprete G, Valensise H, Vasapollo B, Altomare F, Sorge R, Casalino B et al. Body composition during normal pregnancy: reference ranges. *Acta diabetologica* 2003; 40 Suppl 1: S225–232. e-pub ahead of print 2003/11/18; doi: 10.1007/s00592-003-0072-4 [PubMed: 14618479]
32. Lukaski HC, Siders WA, Nielsen EJ, Hall CB. Total body water in pregnancy: assessment by using bioelectrical impedance. *The American journal of clinical nutrition* 1994; 59(3): 578–585. e-pub ahead of print 1994/03/01; [PubMed: 8116533]

Figure 1.

Participant flow diagram of HIV+ and HIV- Kenyan Women. HIV-, HIV-uninfected; HIV+, HIV-infected.

Figure 2.

Predicted β (95% CI) effects of HIV (A) and food insecurity (B) on maternal body composition over time across pregnancy and lactation. Results shown are estimated from multivariable random-effects longitudinal models that adjust for maternal height and covariates. Results for food insecurity include an interaction term between study visit timing and food insecurity. (A) Predicted effects comparing women with HIV+ infection to HIV- women on body composition across pregnancy and lactation. (B) Predicted effects of a 1-

unit increase in individual food insecurity access score on TSF, AMA and MUAC at each observation.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 1.

Characteristics of HIV+ and HIV- Kenyan women (*n* = 333)

	All (n=333)	HIV+ (n=168)	HIV- (n=165)	p-value ¹
Age, y	24.9±4.9	26.2±5.0	23.6±4.4	<0.001
Married, %	91.0	88.7	93.3	0.14
Education, %				
<=Primary	59.2	64.5	53.9	0.09
Secondary	31.5	29.0	34.0	
>=Tertiary	9.3	6.5	12.2	
Asset index ²	6.8±3.1	6.8±3.2	6.9±2.9	0.61
Gestational age at enrollment, weeks	23.6±4.4	22.9±4.7	24.3±4.0	0.001
Gestational age at enrollment, %				
Early 2 nd trimester (<21)	26.4	32.1	20.6	0.05
Late 2 nd trimester (>=21, <28)	47.5	44.6	50.3	
Early 3 rd trimester (>=28, <32)	26.1	23.2	29.1	
Parity, n	2.0±1.7	2.3±1.7	1.6±1.7	<0.001
Maternal BMI at enrollment, kg/m ²	24.0±3.5	23.8±3.5	24.2±3.5	0.38
Maternal height at enrollment, cm	162.3±6.2	161.8±6.1	162.8±6.3	0.15
Gastrointestinal ailments at enrollment, %	23.7	25.6	21.8	0.42
Fever-related illness at enrollment, %	15.9	11.3	20.6	0.02
Probable depression at enrollment, % ³	59.8	61.9	57.6	0.42
Dietary diversity (0–8) at enrollment, score	4.2±1.2	4.1±1.3	4.3±1.2	0.29
Food insecurity (0–27) at enrollment, score	13.0±5.1	12.9±5.0	13.1±5.2	0.75
Time allocation at enrollment (hr)				
Heavy Work	1.9±2.4	1.9±2.4	1.9±2.4	0.82
Light Work	10.6±3.2	10.6±3.0	10.7±3.3	0.86
Leisure	2.5±2.4	2.6±2.5	2.5±2.4	0.55
Sleep	8.9±1.3	8.8±0.9	9.0±1.6	0.24
Breastfeeding				
Exclusive, %				
6 weeks	59.5	64.9	53.9	0.04
14 weeks	56.46	67.9	44.9	<0.001
6 months	21.3	31.6	10.9	<0.001

	All (n=333)	HIV+ (n=168)	HIV- (n=165)	t	p-value
Breastfeeding at 9 mo, ⁴ %	95.9	91.8	100		0.002
Clinic site, %					
Kisumu	19.5	22.0	17.0		0.08
Makder	10.8	10.7	10.9		
Migori	11.1	11.3	10.9		
Nyahera	14.4	12.5	16.4		
Nyamaraga	11.7	11.9	11.5		
Ongo	15.0	13.1	17.0		
Rongo	17.4	18.5	16.4		

Mean±SD, all such values.

¹ Comparison between HIV+ and HIV- women;

² Asset index derived from principal components analysis of possession of twenty different household assets, where households with higher values represent greater household wealth relative to others in the sample.

³ CES-D score 17.

⁴ Breastfeeding at 9 mo available for 221 women.

CES-D, Center for Epidemiologic Studies Depression Scale; BMI, body mass index; KEDH, Kisumu East District Hospital.

Table 2.

Maternal body composition values among HIV+ and HIV- Kenyan mothers across pregnancy and lactation¹

		HIV- (n)		HIV+ (n)		Weight, kg		BMI, kg/m ²		FM, kg		FFM, kg		AEA, cm ³		AMA, cm ³		TSF, mm		MUAC, cm	
Pregnancy	Early 2 nd Tri.	34	54	66.1 (12.8)	61.7 (8.4)	24.5 (4.4)	23.5 (3.0)	19.9 (6.5)	18.8 (5.2)	18.3 (6.6)	15.6 (6.0) ²	23.7 (10.2)	19.8 (9.0)	41.2 (12.6)	40.6 (7.6)	18.3 (6.6)	15.6 (6.0) ¹	28.3 (3.9)	27.4 (3.1)		
	Late 2 nd Tri.	83	75	62.7 (9.6)	61.7 (10.0)	23.8 (3.2)	24.0 (4.0)	18.1 (5.7)	17.8 (6.0)	16.2 (6.6)	14.7 (6.3)	20.5 (9.7)	18.2 (9.6)	39.5 (5.6)	39.8 (7.9)	16.2 (6.6)	14.7 (6.3)	27.3 (2.8)	26.8 (3.0)		
	Early 3 rd Tri.	66	53	65.1 (10.2)	64.1 (10.9)	24.6 (3.4)	24.1 (3.2)	19.5 (5.5)	18.9 (6.8)	15.6 (6.8)	14.1 (4.3)	19.4 (10.1)	17.4 (6.4)	38.2 (8.3)	39.6 (9.5)	15.6 (6.8)	14.1 (4.3)	26.7 (3.4)	26.6 (3.3)		
	Mid 3 rd Tri.	111	123	67.2 (10.6)	64.7 (8.9)	25.5 (3.4)	24.8 (3.0)	20.8 (5.9)	19.7 (4.9)	14.8 (5.5)	13.3 (4.7) ²	18.6 (8.4)	16.5 (6.8) ²	40.1 (9.0)	40.3 (8.3)	14.8 (5.5)	13.3 (4.7) ²	27.0 (3.2)	26.5 (3.1)		
	Late 3 rd Tri.	16	19	66.0 (7.9)	64.4 (6.9)	25.0 (3.0)	24.7 (2.0)	18.9 (5.0)	18.3 (3.9)	15.2 (5.6)	11.1 (4.7) ²	18.7 (7.2)	14.0 (6.5)	40.0 (9.2)	44.2 (17.2)	15.2 (5.6)	11.1 (4.7) ²	27.1 (2.5)	26.7 (4.3)		
Postpartum																					
	6 wk	134	130	62.2 (11.1)	60.0 (9.9)	23.8 (3.6)	22.9 (3.5) ²	19.9 (10.0)	18.2 (6.3)	14.5 (5.7)	13.0 (5.0) ²	18.5 (8.6)	16.1 (7.2) ³	42.1 (9.7)	40.4 (8.9)	14.5 (5.7)	13.0 (5.0) ²	27.4 (3.3)	26.4 (3.4) ²		
	14 wk	136	124	62.7 (14.1)	60.0 (9.0)	24.0 (4.7)	22.9 (3.2) ²	20.3 (10.8)	17.7 (5.1) ³	15.7 (7.2)	13.2 (6.0) ³	19.9 (10.2)	16.6 (8.7) ³	40.7 (9.2)	41.9 (9.4)	15.7 (7.2)	13.2 (6.0) ³	27.4 (3.2)	26.9 (3.2)		
	36 wk	116	120	62.4 (14.8)	59.8 (11.0)	23.9 (5.3)	22.7 (3.9)	18.1 (7.0)	18.9 (14.7)	12.9 (8.1)	10.5 (7.2) ²	16.9 (11.4)	13.4 (9.9) ³	46.3 (15.9)	46.1 (12.9)	12.9 (8.1)	10.5 (7.2) ²	27.9 (4.0)	27.1 (3.4)		

¹ Mean(SD). Comparisons by HIV Status

² <0.05;

³ p 0.01.

BMI, body mass index; FM, fat mass; FFM, fat-free mass; AFA, arm fat area; AMA, arm muscle area; TSF, triceps skinfold; Tri., trimester; MUAC, mid-upper arm circumference.