Published in final edited form as:

Psychiatry Res Neuroimaging. 2019 February 28; 284: 21–28. doi:10.1016/j.psychres.2018.12.016.

Aberrations of anterior insular cortex functional connectivity in nontreatment-seeking alcoholics

Meredith E Halcomb^{a,b,e}, Evgeny J Chumin^{a,b,c,e}, Joaquín Goñi^{d,e}, Mario Dzemidzic^{a,b,e}, Karmen K Yoder^{*,a,b,c,e,f}

^aDepartment of Radiology and Imaging Sciences, Indiana University School of Medicine, Indianapolis, IN, USA

^bIndiana University Center for Neuroimaging, Indiana University School of Medicine, Indianapolis, IN, USA

^cStark Neurosciences Research Institution, Indiana University School of Medicine, Indianapolis, IN, USA

^dPurdue Institute for Integrative Neuroscience, Purdue University, West-Lafayette, IN, USA

^eDepartment of Neurology, Indiana University School of Medicine, Indianapolis, IN USA

^fDepartment of Psychology, Indiana University Purdue University Indianapolis, Indianapolis, IN, USA

Abstract

An emergent literature suggests that resting state functional magnetic resonance imaging (rsfMRI) functional connectivity (FC) patterns are aberrant in alcohol use disorder (AUD) populations. The salience network (SAL) is an established set of brain regions prominent in salience attribution and valuation, and includes the anterior insular cortex (AIC). The SAL is thought to play a role in AUD through directing increased attention to interoceptive cues of intoxication. There is very little information on the salience network (SAL) in AUD, and, in particular, there are no data on SAL FC in currently drinking, nontreatment seeking individuals with AUD (NTS). rsfMRI data from 16 NTS and 21 social drinkers (SD) were compared using FC correlation maps from ten seed regions

*Corresponding Author: Karmen K. Yoder, Ph.D., Ph: 317.963.7507, Fx: 317.963.7547, kkyoder@iupui.edu.

Contributors

KKY designed the study; secured all regulatory approvals; obtained funding for the study; supervised all aspects of the study, including but not limited to, subject recruitment, data acquisition and processing, and statistical analysis; contributed to the intellectual content of the manuscript; and approved the final version of the manuscript.

MEH assisted with subject recruitment and study day execution; conducted all image processing and quality assurance procedures; conducted the statistical analyses; and provided the initial drafts of the manuscript.

EJC helped build, test, and validate the in-house imaging pipeline; supervised the data processing and QA and contributed to the intellectual content of the manuscript.

JG initiated the in-house pipeline components; supervised the data processing and QA; and contributed to the intellectual content of the manuscript.

MD designed the MRI sequences; supervised all data acquisition, processing, and QA; assisted with the design of the statistical analysis; and contributed intellectual content to the manuscript.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Conflicts of interest: None of the authors have any conflicts of interest to declare

of interest in the bilateral AIC. As anticipated, SD subjects demonstrated greater insular FC with frontal and parietal regions. We also found that, compared to SD, NTS had higher insular FC with hippocampal and medial orbitofrontal regions. The apparent overactivity in brain networks involved in salience, learning, and behavioral control in NTS suggests possible mechanisms in the development and maintenance of AUD.

Keywords

alcohol use disorder; resting state functional magnetic resonance imaging; salience network; interoception; alcoholism; hippocampus; orbitofrontal cortex

1. Introduction

Heavy alcohol consumption can result in maladaptive behaviors and physiological damage and is a prevalent, debilitating disorder. Recent research estimates that, within a twelve-month period, approximately 13.9% of the United States population met the Diagnostic and Statistical Manual version V (DSM-5) criteria for diagnosis of alcohol use disorder (AUD) (Grant et al., 2015; Stahre et al., 2014). Decades of preclinical investigation have pointed to numerous neural mechanisms and pathways implicated in the development and maintenance of AUD. With the advent of advanced neuroimaging techniques, we can now examine how the disorder manifests *in vivo*. Although evidence from neuroimaging indicates aberrant structural integrity and functional activity in AUD populations (Chumin et al., 2018; Courtney et al., 2013; Fortier et al., 2014; Rogers et al., 2012; Schacht et al., 2013; Schulte et al., 2010; Wang et al., 1993), there is still much to learn about how specific circuitry is altered in AUD.

Recent advances in functional magnetic resonance imaging (fMRI) acquisition and analysis have greatly enhanced our ability to study critical brain systems involved in AUD. In particular, functional connectivity (FC) analysis of fMRI data can quantify the coherence and strength of functional connections between distinct brain regions (Beckmann et al., 2005; Fox et al., 2005; Greicius et al., 2003). FC is often evaluated during a “resting state”, while the subject is lying still in the scanner, without any task being performed. These task-free fMRI data can be used to extract resting state networks (RSNs), which consist of brain regions that are temporally – and presumably functionally -- coupled (Greicius, 2008; Greicius et al., 2003; Lowe et al., 2000). There is now an emergent literature on alterations in FC within the AUD population, although most of the work has been done in abstinent individuals (Camchong et al., 2013; Chanraud et al., 2011; Jansen et al., 2015; Müller-Oehring et al., 2015; Sjoerds et al., 2017; Zhu et al., 2015). There are only a few studies that report findings in either at-risk drinkers (Vergara et al., 2017) or currently drinking alcohol-dependent subjects (Weiland et al., 2014; Zhu et al., 2015), albeit with equivocal results. Weiland et al. (2014) found that currently-drinking AUD individuals had widespread lower FC compared to controls, whereas Zhu et al. (2015) detected higher FC in multiple networks. Although the Vergara (2017) study did not focus on AUD per se, one of their findings was lower FC in the insula in at-risk alcohol drinkers, supportive of our laboratory’s interest in the potential role of the salience network in alcoholism.

The salience network (SAL) is defined as a functional circuit comprised of the dorsal anterior cingulate cortex (ACC), portions of the anterior insular cortex (AIC), and medial prefrontal cortex (Seeley et al., 2007). The SAL mediates interoceptive awareness and attentional shifts toward relevant (salient) internal and external stimuli. As part of this perceptual role, it is responsible for assigning valence and relative value strength to emotionally relevant sensory information. Ultimately, the SAL interacts with other brain networks to direct attention and modulate behavioral output, including motivated behaviors (Goulden et al., 2014; Menon and Uddin, 2010; Zhou et al., 2018). Given that alcohol dependence involves seeking out and obtaining alcohol (i.e., motivated behaviors) despite adverse consequences, we hypothesized that the SAL may have altered function in currently-drinking individuals with AUD.

There are also numerous studies that support a role for the insula in addictive behaviors. For example, drug-abstinent populations (including addictions to nicotine, cocaine and alcohol) have higher cue-induced insular activation (Kilts et al., 2004; McBride et al., 2006; Myrick et al., 2003; Tapert et al., 2004). Greater insular activation was also noted in a group of alcohol-dependent women during a task involving alcohol cues and the likelihood to drink despite negative outcomes (Arcurio et al., 2015). Human studies and preclinical work both show that lesions of the insula mitigate drug seeking behaviors (Contreras et al., 2007; Naqvi et al., 2007), suggesting that the insula is integral to the development and maintenance of alcohol and substance use disorders.

At present, there is no information regarding the FC of the insula to other brain regions in currently drinking, non-treatment-seeking (NTS) alcoholics. Given the prominent role of the AIC in the salience network and its potential influence in drug-seeking behaviors, detecting aberrations of insular FC in an NTS sample would greatly enhance our understanding of how insular mechanisms may mediate drinking behavior in AUD. Thus, our primary goal was to perform a whole-brain resting state FC study in NTS subjects and social drinkers (SD), using seeds placed in the AIC. Because insular function has been documented to be lateralized (Cauda et al., 2011; Duerden et al., 2013; Sander and Scheich, 2005), we evaluated functional connectivity of the left and right anterior insula separately. We hypothesized that connectivity between the AIC and regions associated with reward processing, salience attribution, and behavioral control would be altered in NTS. The potential impact of this work lies in the possibility that, if the functional connectivity of the salience network is altered in AUD, it may be feasible to develop neuroimaging protocols that will utilize SAL function as a marker to predict treatment response and risk for relapse. Additionally, such protocols could be used to track longitudinal changes in the SAL, which could lead to testing existing and future therapies for alcoholism for the ability to modulate SAL function.

2. Methods

2.1 Subjects

Study procedures were approved by the Indiana University Institutional Review Board. Informed consent from each subject was obtained upon determination that breath alcohol concentration was 0 mg%. Seventeen non-treatment-seeking alcoholics (NTS) and 22 social

drinkers (SD) completed the study protocol. NTS subjects met DSM-IV criteria for alcohol dependence as determined by the Semi-structured Assessment for the Genetics of Alcoholism (Bucholz et al., 1994). NTS subjects were not actively seeking treatment and had not sought treatment within the last year. Exclusionary criteria included: contraindications for MRI, positive urine pregnancy screen, positive urine toxicology screen for illicit substances (except marijuana), current use of any psychotropic medication, history or presence of any neurological, psychiatric disorders (including substance abuse or dependence), or major medical disorders. Demographics and drinking characteristics are listed in Table 1. Supplemental Materials Table 1 lists the percentage of subjects who reported any prior substance use during the course of their lifetime.

2.2 General Study Procedures

The MRI data reported herein were acquired as part of two separate PET protocols. MRI acquisition sequences were identical for all subjects. MRIs were completed in the morning prior to any subsequent experimental procedures. Both PET studies included NTS and SD subjects according to the criteria described above. Upon subjects' arrival, sobriety was confirmed by measurements of breath alcohol concentration (BrAC) with an Alcotest 6510 (Dräger Manufacturing, Irving, TX). Urine samples were collected for drug and pregnancy testing. The Clinical Withdrawal Assessment for Alcohol, Revised (CIWA-AR) was administered to NTS subjects periodically to monitor for alcohol withdrawal. (Sullivan et al., 1989). Subjects underwent a 90 minute magnetic resonance imaging session, which started with a high resolution anatomical scan followed by a 10 minute resting state functional MRI scan as detailed below. Subjects were instructed to lay still with their eyes closed and to keep their mind clear. All subjects reported after the scan that they had remained awake and complied with all instructions.

2.3 Image Acquisition

Imaging was performed on a Siemens 3T Prisma (Erlangen, Germany) using a 64-channel head coil array. A high-resolution anatomic volume (3D magnetization prepared rapid gradient echo (MP-RAGE); $1.05 \times 1.05 \times 1.2 \text{ mm}^3$ or $1.00 \times 1.00 \times 1.00 \text{ mm}^3$ voxels) was used for positioning the blood oxygenation level dependent (BOLD) resting state data acquisition. Five hundred whole-brain BOLD volumes were acquired using a multiband (MB) echo-planar imaging (EPI) sequence (Center for Magnetic Resonance Research at the University of Minnesota, gradient echo, repetition/echo time TR/TE=1200/29 ms, flip angle 65° , field-of-view $220 \times 220 \text{ mm}$, matrix 88×88 , 54 2.5 mm slices, $2.5 \times 2.5 \times 2.5 \text{ mm}^3$ voxel, slice acceleration factor = 3) (Smith et al., 2013). The first 25 seconds of each resting state scan were excluded to ensure that only volumes with steady state magnetization would be considered for statistical inferences. BOLD acquisition was preceded by a pair of MB=3 spin echo field mapping scans (3 A-P and 3 P-A phase direction volumes, TR=1560 ms, TE = 49.80 ms) with imaging and voxel sizes identical to the BOLD acquisition.

2.4 Image Preprocessing

Preprocessing was performed with a Matlab-based processing pipeline using FMRIB Software Library (FSL version 5.0.9) (Smith et al., 2004) as detailed in (Amico et al., 2017; Contreras et al., 2017). Resting state functional connectivity data were processed in native

BOLD EPI space. T1-weighted MP-RAGE volumes were denoised prior to brain masking and extraction (FSL “bet”) and then nonlinearly transformed (FSL’s FLIRT and FNIRT) to the Montreal Neurological Institute (MNI) brain template space. Image processing included BOLD volume unwarping using FSL’s topup/applytopup (using spin echo field mapping scans), slice acquisition timing correction, head motion realignment, registration to T1, demeaning and detrending, band pass filtering (0.009-0.08 Hz), and normalization to mode 1000.

Based upon Power et al.’s approach, motion regressors from the realignment and their derivatives were regressed out (Power et al., 2012; Power et al., 2014). We used 3 image quality control metrics (frame displacement, “DVARs” [D refers to the temporal derivative of BOLD time courses, VARs refers to RMS variance over voxels], and whole-brain standard deviation of BOLD signal) to identify and exclude (“scrub”) outlier BOLD volumes. Outlier cutoffs were set at 0.3 mm for frame displacement (characterized as medium head motion) and for 1.7 standardized DVARs (Nichols, 2017; Parker et al., 2017; Power et al., 2015). Any BOLD volume defined as an outlier was “scrubbed” from statistical inferences by excluding it from the calculation of correlation coefficients. There were no significant differences between groups in frame displacement, DVARs, or percentage volumes scrubbed (Supplementary Figure 1). In addition, two subjects whose scans had >50% outlier volumes were dropped from the analyses (one SD subject and one NTS subject); thus, the final dataset was comprised of 21 SD and 16 NTS subjects. Principal component analysis (PCA) regression was performed with the Matlab *pca* function after band-pass filtering so that all components were within the frequency range of interest (Chai et al., 2012; Power et al., 2015). The first five signals obtained by PCA from each of the three tissue compartments were regressed out to address confounding effects of physiologic noise and residual head motion within eroded masks of the whole-brain gray matter, white matter and cerebrospinal fluid of the lateral ventricles. This included global signal regression (Power et al., 2012; Power et al., 2015).

2.5 Regions of Interest and Functional Connectivity

A priori insular seed regions of interest (ROIs) were created using the MarsBar toolbox in SPM12 (Brett et al., 2002). The $5 \times 5 \times 5$ mm³ cubical seeds were based upon previous characterization of insular FC (Cauda et al., 2012; Cauda et al., 2011). We selected a subset of the Cauda seeds (“2”, “5”, “6”, “8”, and “9”, according to their nomenclature; Table 2) within the anterior insular cortex (AIC), which is a node of the salience network (Figure 1). Seed regions were created in the 1 mm isotopic voxel MNI space and were nonlinearly transformed to native space of each subject. Mean BOLD time series from each seed region was extracted and correlated with time series of all voxels to create whole-brain Pearson’s correlation coefficient maps. Then, a variance-stabilizing Fisher z-transformation was performed to create a Z-statistic map. Native space Z-statistic maps of each subject were inversely transformed to MNI space using FSL’s registration and smoothed by a 6 mm full width at a half maximum Gaussian kernel.

2.6 Statistics

Independent *t*-tests were performed to test for group differences in age, education, and recent drinking metrics (drinks consumed per drinking day and drinks consumed per week). Chi-squared tests were used to assess group differences in the ratios of sex and smoking status.

Ten *Z*-statistic images of each subject (one for each seed) were entered into an SPM12 group random effects factorial model, with Group(2; NTS, SD) and Seed(10; 5 seeds $\times$ 2 hemispheres) treated as fixed effects. Group was modeled as an independent factor, while seed was a dependent factor. Contrasts were specified to yield “average” connectivity of left seeds, right seeds, or all seeds (main effect). The design matrix presented in Supplementary Figure 2 illustrates the covariance structure of the model, which reflects the repeated-measures nature of the data (with 10 statistical maps contributed by each subject). We tested for group differences in the whole-brain connectivity maps using the SD > NTS and NTS > SD contrasts. Statistical significance was inferred at the cluster level ($p_{FWE} < 0.05$) using family wise error (FWE) correction for multiple comparisons across the whole brain, with a cluster-forming height threshold set to $p = 0.001$, uncorrected (Eklund et al., 2016). To simplify the interpretation and avoid “the double subtraction pitfall,” we reported significant group differences in FC only in areas that showed positive FC with the seed region. This allowed us to exclude areas that were not associated or were anticorrelated with the seed regions. Specifically, SD>NTS findings were constrained by an inclusive mask from the positive SD contrast (SD subjects only; voxel-wise $p < 0.05$, uncorrected). Similarly, the NTS positive contrast was used as an inclusive mask when reporting NTS>SD results. Analyses were conducted both with and without sex, age, and education as covariates.

We also examined an SPM multiple regression model across all subjects that included recent drinking as indexed by number of drinks consumed per week. This measure of alcohol intake was chosen as it better describes overall drinking patterns relative to drinks consumed per drinking day. The dependent variable in these regression models were each subject’s *Z*-statistic connectivity map (entered into the model as described above; left and right seeds were averaged for each side and were and tested independently within the model).

3. Results

3.1 Demographics

There were no significant group differences in sex, age, smoking status or education. As anticipated, groups differed on drinks per drinking day (Dr/DD) and drinks per week (Dr/Wk), with the NTS consuming significantly greater amounts of alcohol, $ps < 0.001$.

3.2 Functional Connectivity Patterns Across All Subjects

For both the left and right AIC, the main effects of insular FC across all subjects were similar to previously reported findings (Cauda et al., 2011). Conjunction analyses confirmed significant overlap in FC in both SD and NTS subjects (see Supplementary Material, Figure 3). There was significant connectivity between the AIC and other nodes of the SAL, the cerebellum, subcortical regions, and the prefrontal cortex (see Supplementary Material

Figures 3 and 5; Tables 2 and 3). Below, we present the group differences from the right and left AIC seed analyses. All reported group differences remained significant when age, sex, and education were included as covariates.

3.3 Between Group Differences

Significant differences in the patterns of AIC FC were present between NTS and SD subjects. Overall, SD subjects demonstrated greater FC than NTS, mostly with frontal and parietal regions from both left and right hemisphere AIC seed regions (Table 3). However, NTS subjects had greater FC than SD in temporal, limbic, and orbitofrontal regions (Table 4).

3.4 SD > NTS Contrast

Subjects in the SD group displayed greater bilateral FC with frontoparietal regions, including the right superior orbital gyrus (SOG) and the right inferior frontal gyrus (IFG) (Figures 3 and 4A). The left AIC seeds showed an enhanced FC with the right superior parietal lobule (SPL), left calcarine gyrus (CalG), and the left inferior frontal gyrus (Figure 3). The right AIC seeds showed greater FC with the posterior cerebellum, the left superior parietal lobule (SPL), and the right putamen (PUT) (Figures 3 and 4B).

3.5 NTS > SD Contrast

Compared to SD, NTS subjects had greater insular connectivity with several regions (Figure 3; Table 4). The left AIC seeds had greater FC with the anterior cerebellum, the right fusiform gyrus (FFG), and the left posterior insular cortex (PIC). Seeds located in the right hemisphere displayed greater FC with bilateral parahippocampal gyri (PHG; Figure 4A), medial orbitofrontal cortex (mOFC; Figure 4B), the left hippocampus (HIP; Figure 4A), and the left supramarginal gyrus (SMG).

3.6 Alcohol intake correlations

When multiple regression models were corrected for multiple comparisons across all subjects, there were no associations between insular FC and recent alcohol intake. This finding, however, was not unexpected. First, drinking rates were self-reported, which can be inherently unreliable. There was also high variability in the metric of drinks per week in both groups. Both factors could have contributed to a lack of a relationship between FC and drinking behavior.

4. Discussion

This study was the first to detect significant differences in FC related to the salience network (SAL) in an NTS sample. Overall, there were disparate patterns of insular connectivity between NTS and SD subjects, which occurred in regions associated with memory formation (Miendlarzewska et al., 2016; Zola-Morgan et al., 1989), stimulus valuation, reinforcement, and decision making (Kringelbach and Rolls, 2004). The SAL integrates information about interoceptive sensations and perceptions about external stimuli, and assigns valence to these experiences (Craig, 2010; Naqvi and Bechara, 2010; Seeley et al., 2007). Additional work suggests that the insula provides representations of emotional states

linked to stimuli, as well as any cues associated with the stimuli (Craig and Craig, 2009; Gasquoin, 2014; Wager and Barrett, 2017). The present findings suggest that an insular mechanism may be involved in the maintenance of hazardous drinking behavior in currently-drinking AUD individuals, perhaps through exaggerated attribution of positive valence to intoxication and/or cues associated with alcohol consumption.

4.1 Greater Insular FC in NTS

In the NTS sample, there was greater insular FC with bilateral hippocampal regions, which are closely associated with memory formation and encoding. There is abundant evidence for alterations in the hippocampal regions of subjects with long-term alcohol exposure (Livy et al., 2003; Riley and Walker, 1978; Zhou et al., 2011). As with other types of learning, the hippocampus is involved with the short- and long-term memory formation necessary for drug-associated learning (Jones and Bonci, 2005). Therefore, it is possible that the higher insular-hippocampal connectivity observed in our NTS sample could have real-world consequences. For example, greater FC between the AIC and bilateral hippocampal regions in the NTS group could indicate a tendency for formation of abnormally strong associations with alcohol-related interoceptive states (e.g., intoxication) and/or the cues associated with drinking. If the memory of intoxication is represented more intensely or with more positive valence, then this could exert greater influence over behaviors such as alcohol seeking and consumption, relative to representations such as the negative consequences of alcohol consumption. Thus, a disproportionate assignment of positive value to alcohol-related interoception and associated environmental cues could contribute to the sustained alcohol use in NTS populations.

NTS subjects also displayed greater FC between the right AIC and the medial OFC (mOFC), a region involved with reward valuation and encoding (Elliott et al., 2000; Kringelbach and Rolls, 2004; O'Doherty et al., 2001; Sescousse et al., 2010), as well as decision-making based on expected outcomes of actions (Schoenbaum et al., 2006). If the AIC and mOFC are working in synchrony, it is possible that the mOFC may be compromised in its ability to modulate behavioral decisions based on the skewed reward value of alcohol and associated environmental stimuli. This could also help explain why individuals with AUD continue to drink despite negative outcomes.

The present results may be counter-intuitive in the face of other neuroimaging data that suggests that addicts have lower OFC function (Bolla et al., 2003; London et al., 2000; Schoenbaum and Roesch, 2005), and preclinical data that indicate that drugs of abuse compromise the ability of the OFC to modulate behavior and learning (for review, Schoenbaum et al., 2006). However, it is consistent with overactivation of the OFC observed in drug addicts during exposure to drug-related cues (Goldstein et al., 2007; London et al., 2000; Tomasi et al., 2015). It should be noted that this is the first study in a non-abstinent AUD population.

4.2 Lower FC in NTS

Relative to SD, NTS subjects had lower insular connectivity with prefrontal nodes of the frontoparietal attention network (FPN). The FPN is considered a “flexible hub” of cognitive

control (Cole et al., 2013), and is essential in driving goal-directed behavior (Cole et al., 2014). Given the extensive influence of the FPN on controlling behavior, degradation of this system could lead to an inability to regulate behavior. Thus, lower FPN–AIC connectivity in NTS may inhibit cognitive and behavioral control over alcohol consumption despite recurrent negative consequences.

Connectivity between the insula and putamen has been confirmed in healthy populations (Cauda et al., 2011; Di Martino et al., 2008; Postuma and Dagher, 2006). NTS subjects displayed lower right insular FC with the right anterior caudate and anterior putamen as compared to SD subjects. This is consistent with previous research in cocaine addicts (McHugh et al., 2013). The caudate and putamen are integral components of the basal ganglia, and contribute to both motor and cognitive processes (DeLong et al., 1984; Marchand et al., 2008; Packard and Knowlton, 2002; Seger and Cincotta, 2005; White, 2009). The putamen in particular is heavily implicated in mediating habit formation, a prominent aspect of AUD (Tricomi et al., 2009; Yin and Knowlton, 2006). The diminished insular-anterior striatal FC observed here may suggest potential alterations in anterior putamen function in NTS subjects that could contribute to habitual alcohol consumption.

The primary limitation of this study is a modest sample size. Although we found significant results, a larger sample could permit us to detect more discrete regions that may have compromised insular connectivity in AUD. The sample size also did not permit us to test for effects of sex, age, or smoking status. While we utilized an *a priori* seed-based approach, whole-brain connectomics approaches may provide more comprehensive information about functional abnormalities in AUD. Although no subjects endorsed falling asleep during the scan, it is possible they could have briefly fallen asleep without being aware this occurred. Future studies could implement the use of the Stanford Sleepiness Scale to evaluate states of consciousness immediately prior to and after resting state scans to gain insight into the subjects' level of consciousness and its changes during imaging (Hoddes et al., 1973; Vaghi et al., 2017). We also cannot make any assumptions about the neurochemistry of the neuronal populations that are contributing to the BOLD signal. Finally, there are conceptual limitations with our explanation of the data. It could be the case that the observed differences in the SAL are independent of drinking behaviors, and may instead reflect a broader, non-specific effect of years of chronic drinking on insular circuitry in AUD. Alternatively, such differences could reflect premorbid differences in FC that could be related to risk factors for development of AUD. Carefully planned longitudinal studies would be required to disentangle these possible interpretations.

In summary, a mounting body of evidence strongly indicates alterations in overall FC in AUD individuals. The results extend previous findings to include information regarding insular connectivity patterns in currently-drinking AUD subjects. Together, This research illustrates a potential impact of the insula (and by extension, the salience network) on addictive behaviors. Future work with task-based fMRI will be needed to test specific hypotheses about how neural circuits that mediate salience attribution, behavioral control, and habit formation are altered in AUD.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

The authors would like to thank the research MRI technologists of the Indiana Institute of Biomedical Imaging Sciences. In addition, we would like to acknowledge the assistance of study coordinators Karen Hile and James Walters. This work was supported by the National Institute of Alcohol Abuse and Alcoholism grants R01 AA01835402 (KKY), R21 AA02265002 (KKY), and T32 AA00746532 (MEH).

References

- Amico E, Marinazzo D, Di Perri C, Heine L, Annen J, Martial C, Dziedzic M, Kirsch M, Bonhomme V, Laureys S, Goni J, 2017 Mapping the functional connectome traits of levels of consciousness. *Neuroimage* 148, 201–211. [PubMed: 28093358]
- Arcurio LR, Finn PR, James TW, 2015 Neural mechanisms of high-risk decisions-to-drink in alcohol-dependent women. *Addict Biol* 20(2), 390–406. [PubMed: 24373127]
- Beckmann CF, DeLuca M, Devlin JT, Smith SM, 2005 Investigations into Resting-State Connectivity Using Independent Component Analysis. *Philosophical Transactions: Biological Sciences* 360(1457), 1001–1013. [PubMed: 16087444]
- Bolla K, Eldreth D, London E, Kiehl K, Mouratidis M, Contoreggi C.e., et al, Matochik J, Kurian V, Cadet J, Kimes A., 2003 Orbitofrontal cortex dysfunction in abstinent cocaine abusers performing a decision-making task. *Neuroimage* 19(3), 1085–1094. [PubMed: 12880834]
- Brett M, Anton J-L, Valabregue R, Poline J-B, 2002 Region of interest analysis using the MarsBar toolbox for SPM 99. *Neuroimage* 16(2), S497.
- Bucholz KK, Cadoret R, Cloninger CR, Dinwiddie SH, Hesselbrock V, Nurnberger J Jr, Reich T, Schmidt I, Schuckit MA, 1994 A new, semi-structured psychiatric interview for use in genetic linkage studies: a report on the reliability of the SSAGA. *Journal of studies on alcohol* 55(2), 149–158. [PubMed: 8189735]
- Camchong J, Stenger VA, Fein G, 2013 Resting-State Synchrony in Short-Term Versus Long-Term Abstinent Alcoholics. *Alcoholism: Clinical and Experimental Research* 37(5), 794–803.
- Cauda F, Costa T, Torta DME, Sacco K, D'Agata F, Duca S, Geminiani G, Fox PT, Vercelli A, 2012 Meta-analytic clustering of the insular cortex: Characterizing the meta-analytic connectivity of the insula when involved in active tasks. *Neuroimage* 62(1), 343–355. [PubMed: 22521480]
- Cauda F, D'Agata F, Sacco K, Duca S, Geminiani G, Vercelli A, 2011 Functional connectivity of the insula in the resting brain. *Neuroimage* 55(1), 8–23. [PubMed: 21111053]
- Chai XJ, Castanon AN, Ongir D, Whitfield-Gabrieli S, 2012 Anticorrelations in resting state networks without global signal regression. *Neuroimage* 59(2), 1420–1428. [PubMed: 21889994]
- Chanraud S, Pitel A-L, Pfefferbaum A, Sullivan EV, 2011 Disruption of Functional Connectivity of the Default-Mode Network in Alcoholism. *Cerebral cortex* 21(10), 2272–2281. [PubMed: 21368086]
- Chumin EJ, Goni J, Halcomb ME, Durazzo TC, Dziedzic M, Yoder KK, 2018 Differences in White Matter Microstructure and Connectivity in Nontreatment-Seeking Individuals with Alcohol Use Disorder. *Alcoholism: Clinical and Experimental Research* 42(5), 889–896.
- Cole MW, Repovs G, Anticevic A, 2014 The frontoparietal control system: A central role in mental health. *The Neuroscientist: a review journal bringing neurobiology, neurology and psychiatry* 20(6), 652–664.
- Cole MW, Reynolds JR, Power JD, Repovs G, Anticevic A, Braver TS, 2013 Multi-task connectivity reveals flexible hubs for adaptive task control. *Nature Neuroscience* 16,1348. [PubMed: 23892552]
- Contreras JA, Goni J, Risacher SL, Amico E, Yoder K, Dziedzic M, West JD, McDonald BC, Farlow MR, Sporns O, Saykin AJ, 2017 Cognitive complaints in older adults at risk for Alzheimer's disease are associated with altered resting-state networks. *Alzheimer's & Dementia : Diagnosis, Assessment & Disease Monitoring* 6, 40–49.

- Contreras M, Ceric F, Torrealba F, 2007 Inactivation of the Interoceptive Insula Disrupts Drug Craving and Malaise Induced by Lithium. *Science* 318(5850), 655–658. [PubMed: 17962567]
- Courtney KE, Ghahremani DG, Ray LA, 2013 Fronto-striatal functional connectivity during response inhibition in alcohol dependence. *Addiction Biology* 18(3), 593–604. [PubMed: 23240858]
- Craig AD, Craig A, 2009 How do you feel--now? The anterior insula and human awareness. *Nature reviews neuroscience* 10(1).
- Craig ADB, 2010 The sentient self. *Brain Structure & Function* 214(5-6), 563–577. [PubMed: 20512381]
- DeLong M, Alexander G, Georgopoulos A, Crutcher M, Mitchell S, Richardson R, 1984 Role of basal ganglia in limb movements. *Human neurobiology* 2(4), 235–244. [PubMed: 6715208]
- Di Martino A, Scheres A, Margulies DS, Kelly AMC, Uddin LQ, Shehzad Z, Biswal B, Walters JR, Castellanos FX, Milham MP, 2008 Functional Connectivity of Human Striatum: A Resting State fMRI Study. *Cerebral Cortex* 18(12), 2735–2747. [PubMed: 18400794]
- Duerden EG, Arsalidou M, Lee M, Taylor MJ, 2013 Lateralization of affective processing in the insula. *Neuroimage* 78, 159–175. [PubMed: 23587690]
- Eklund A, Nichols TE, Knutsson H, 2016 Cluster failure: Why fMRI inferences for spatial extent have inflated false-positive rates. *Proceedings of the National Academy of Sciences* 113(28), 7900–7905.
- Elliott R, Dolan RJ, Frith CD, 2000 Dissociable Functions in the Medial and Lateral Orbitofrontal Cortex: Evidence from Human Neuroimaging Studies. *Cerebral Cortex* 10(3), 308–317. [PubMed: 10731225]
- Fortier CB, Leritz EC, Salat DH, Lindemer E, Maksimovskiy AL, Shepel J, Williams V, Venne JR, Milberg WP, McGlinchey RE, 2014 Widespread Effects of Alcohol on White Matter Microstructure. *Alcoholism: Clinical and Experimental Research* 38(12), 2925–2933.
- Fox MD, Snyder AZ, Vincent JL, Corbetta M, Van Essen DC, Raichle ME, 2005 The Human Brain Is Intrinsically Organized into Dynamic, Anticorrelated Functional Networks. *Proceedings of the National Academy of Sciences of the United States of America* 102(27), 9673–9678. [PubMed: 15976020]
- Gasquoine PG, 2014 Contributions of the insula to cognition and emotion. *Neuropsychology review* 24(2), 77–87. [PubMed: 24442602]
- Goldstein RZ, Tomasi D, Rajaram S, Cottone LA, Zhang L, Maloney T, Telang F, Alia-Klein N, Volkow ND, 2007 Role of the anterior cingulate and medial orbitofrontal cortex in processing drug cues in cocaine addiction. *Neuroscience* 144(4), 1153–1159. [PubMed: 17197102]
- Goulden N, Khusnulina A, Davis NJ, Bracewell RM, Bokde AL, McNulty JP, Mullins PG, 2014 The salience network is responsible for switching between the default mode network and the central executive network: Replication from DCM. *Neuroimage* 99, 180–190. [PubMed: 24862074]
- Grant BF, Goldstein RB, Saha TD, et al., 2015 Epidemiology of dsm-5 alcohol use disorder: Results from the national epidemiologic survey on alcohol and related conditions iii. *JAMA Psychiatry* 72(8), 757–766. [PubMed: 26039070]
- Greicius M, 2008 Resting-state functional connectivity in neuropsychiatric disorders. *Current Opinion in Neurology* 21(4), 424–430. [PubMed: 18607202]
- Greicius MD, Krasnow B, Reiss AL, Menon V, 2003 Functional Connectivity in the Resting Brain: A Network Analysis of the Default Mode Hypothesis. *Proceedings of the National Academy of Sciences of the United States of America* 100(1), 253–258. [PubMed: 12506194]
- Hoddes E, Zarcone V, Smythe H, Phillips R, Dement WC, 1973 Quantification of sleepiness: a new approach. *Psychophysiology* 10(4), 431–436. [PubMed: 4719486]
- Jansen JM, Holst RJ, Brink W, Veltman DJ, Caan MWA, Goudriaan AE, 2015 Brain function during cognitive flexibility and white matter integrity in alcohol-dependent patients, problematic drinkers and healthy controls. *Addiction Biology* 20(5), 979–989. [PubMed: 25477246]
- Jones S, Bonci A, 2005 Synaptic plasticity and drug addiction. *Current Opinion in Pharmacology* 5(1), 20–25. [PubMed: 15661621]
- Kilts CD, Gross RE, Ely TD, Drexler KP, 2004 The neural correlates of cue-induced craving in cocaine-dependent women. *American Journal of Psychiatry* 161(2), 233–241. [PubMed: 14754771]

- Kringelbach ML, Rolls ET, 2004 The functional neuroanatomy of the human orbitofrontal cortex: evidence from neuroimaging and neuropsychology. *Progress in Neurobiology* 72(5), 341–372. [PubMed: 15157726]
- Livy DJ, Miller EK, Maier SE, West JR, 2003 Fetal alcohol exposure and temporal vulnerability: effects of binge-like alcohol exposure on the developing rat hippocampus. *Neurotoxicology and Teratology* 25(4), 447–458. [PubMed: 12798962]
- London ED, Ernst M, Grant S, Bonson K, Weinstein A, 2000 Orbitofrontal cortex and human drug abuse: functional imaging. *Cerebral cortex* 10(3), 334–342. [PubMed: 10731228]
- Lowe MJ, Dzemidzic M, Lurito JT, Mathews VP, Phillips MD, 2000 Correlations in Low-Frequency BOLD Fluctuations Reflect Cortico-Cortical Connections. *NeuroImage* 12(5), 582–587. [PubMed: 11034865]
- Marchand WR, Lee JN, Thatcher JW, Hsu EW, Rashkin E, Suchy Y, Chelune G, Starr J, Barbera SS, 2008 Putamen coactivation during motor task execution. *Neuroreport* 19(9), 957–960. [PubMed: 18521000]
- McBride D, Barrett SP, Kelly JT, Aw A, Dagher A, 2006 Effects of expectancy and abstinence on the neural response to smoking cues in cigarette smokers: an fMRI study. *Neuropsychopharmacology* 31(12), 2728. [PubMed: 16598192]
- McHugh MJ, Demers CH, Braud J, Briggs R, Adinoff B, Stein EA, 2013 Striatal-insula circuits in cocaine addiction: implications for impulsivity and relapse risk. *The American Journal of Drug and Alcohol Abuse* 39(6), 424–432. [PubMed: 24200212]
- Menon V, Uddin LQ, 2010 Saliency, switching, attention and control: a network model of insula function. *Brain Struct Funct* 214(5-6), 655–667. [PubMed: 20512370]
- Miendlarzewska EA, Bavelier D, Schwartz S, 2016 Influence of reward motivation on human declarative memory. *Neuroscience & Biobehavioral Reviews* 61, 156–176. [PubMed: 26657967]
- Müller-Oehring EM, Jung Y-C, Pfefferbaum A, Sullivan EV, Schulte T, 2015 The Resting Brain of Alcoholics. *Cerebral Cortex* 25(11), 4155–4168. [PubMed: 24935777]
- Myrick H, Anton RF, Li X, Henderson S, Drobos D, Voronin K, George MS, 2003 Differential Brain Activity in Alcoholics and Social Drinkers to Alcohol Cues: Relationship to Craving. *Neuropsychopharmacology* 29, 393.
- Naqvi NH, Bechara A, 2010 The insula and drug addiction: an interoceptive view of pleasure, urges, and decision-making. *Brain Structure and Function* 214(5-6), 435–450. [PubMed: 20512364]
- Naqvi NH, Rudrauf D, Damasio H, Bechara A, 2007 Damage to the Insula Disrupts Addiction to Cigarette Smoking. *Science* 315(5811), 531–534. [PubMed: 17255515]
- Nichols TE, 2017 Notes on creating a standardized version of DVARS. arXiv preprint arXiv: 1704.01469.
- O'Doherty J, Kringelbach ML, Rolls ET, Hornak J, Andrews C, 2001 Abstract reward and punishment representations in the human orbitofrontal cortex. *Nature Neuroscience* 4, 95. [PubMed: 11135651]
- Packard MG, Knowlton BJ, 2002 Learning and memory functions of the basal ganglia. *Annual review of neuroscience* 25(1), 563–593.
- Parker D, Liu X, Razlighi QR, 2017 Optimal slice timing correction and its interaction with fMRI parameters and artifacts. *Medical image analysis* 35, 434–445. [PubMed: 27589578]
- Postuma RB, Dagher A, 2006 Basal Ganglia Functional Connectivity Based on a Meta-Analysis of 126 Positron Emission Tomography and Functional Magnetic Resonance Imaging Publications. *Cerebral Cortex* 16(10), 1508–1521. [PubMed: 16373457]
- Power JD, Barnes KA, Snyder AZ, Schlaggar BL, Petersen SE, 2012 Spurious but systematic correlations in functional connectivity MRI networks arise from subject motion. *NeuroImage* 59(3), 2142–2154. [PubMed: 22019881]
- Power JD, Mitra A, Laumann TO, Snyder AZ, Schlaggar BL, Petersen SE, 2014 Methods to detect, characterize, and remove motion artifact in resting state fMRI. *NeuroImage* 84, 320–341. [PubMed: 23994314]
- Power JD, Schlaggar BL, Petersen SE, 2015 Recent progress and outstanding issues in motion correction in resting state fMRI. *NeuroImage* 105, 536–551. [PubMed: 25462692]

- Riley J, Walker D, 1978 Morphological alterations in hippocampus after long-term alcohol consumption in mice. *Science* 201(4356), 646–648. [PubMed: 566953]
- Rogers BP, Parks MH, Nickel MK, Katwal SB, Martin PR, 2012 Reduced Fronto-Cerebellar Functional Connectivity in Chronic Alcoholic Patients. *Alcoholism: Clinical and Experimental Research* 36(2), 294–301.
- Sander K, Scheich H, 2005 Left auditory cortex and amygdala, but right insula dominance for human laughing and crying. *Journal of cognitive Neuroscience* 17(10), 1519–1531. [PubMed: 16269094]
- Schacht JP, Anton RF, Myrick H, 2013 Functional neuroimaging studies of alcohol cue reactivity: a quantitative meta-analysis and systematic review. *Addiction Biology* 18(1), 121–133. [PubMed: 22574861]
- Schoenbaum G, Roesch M, 2005 Orbitofrontal cortex, associative learning, and expectancies. *Neuron* 47(5), 633–636. [PubMed: 16129393]
- Schoenbaum G, Roesch MR, Stalnaker TA, 2006 Orbitofrontal cortex, decision-making and drug addiction. *Trends in Neurosciences* 29(2), 116–124. [PubMed: 16406092]
- Schulte T, Müller-Oehring EM, Pfefferbaum A, Sullivan EV, 2010 Neurocircuitry of emotion and cognition in alcoholism: contributions from white matter fiber tractography. *Dialogues in Clinical Neuroscience* 12(4), 554–560. [PubMed: 21319499]
- Seeley WW, Menon V, Schatzberg AF, Keller J, Glover GH, Kenna H, Reiss AL, Greicius MD, 2007 Dissociable intrinsic connectivity networks for salience processing and executive control. *J Neurosci* 27(9), 2349–2356. [PubMed: 17329432]
- Seger CA, Cincotta CM, 2005 The Roles of the Caudate Nucleus in Human Classification Learning. *The Journal of Neuroscience* 25(11), 2941–2951. [PubMed: 15772354]
- Sescousse G, Redoute J, Dreher J-C, 2010 The Architecture of Reward Value Coding in the Human Orbitofrontal Cortex. *The Journal of Neuroscience* 30(39), 13095–13104. [PubMed: 20881127]
- Sjoerds Z, Stufflebeam SM, Veltman DJ, Brink W.V.d., Penninx BWJH, Douw L, 2017 Loss of brain graph network efficiency in alcohol dependence. *Addiction Biology* 22(2), 523–534. [PubMed: 26692359]
- Smith SM, Beckmann CF, Andersson J, Auerbach EJ, Bijsterbosch J, Douaud G, Duff E, Feinberg DA, Griffanti L, Harms MP, Kelly M, Laumann T, Miller KL, Moeller S, Petersen S, Power J, Salimi-Khorshidi G, Snyder AZ, Vu AT, Woolrich MW, Xu J, Yacoub E, Ugurbil K, Van Essen DC, Glasser MF, Consortium, W.U.-M.H., 2013 Resting-state fMRI in the Human Connectome Project. *Neuroimage* 80, 144–168. [PubMed: 23702415]
- Smith SM, Jenkinson M, Woolrich MW, Beckmann CF, Behrens TEJ, Johansen-Berg H, Bannister PR, De Luca M, Drobnjak I, Flitney DE, Niazy RK, Saunders J, Vickers J, Zhang Y, De Stefano N, Brady JM, Matthews PM, 2004 Advances in functional and structural MR image analysis and implementation as FSL. *Neuroimage* 23(Supplement 1), S208–S219. [PubMed: 15501092]
- Stahre M, Roeber J, Kanny D, Brewer RD, Zhang X, 2014 Contribution of excessive alcohol consumption to deaths and years of potential life lost in the United States. *Prev Chronic Dis* 11, E109. [PubMed: 24967831]
- Sullivan JT, Sykora K, Schneiderman J, Naranjo CA, Sellers EM, 1989 Assessment of alcohol withdrawal: the revised clinical institute withdrawal assessment for alcohol scale (CIWA-Ar). *Addiction* 84(11), 1353–1357.
- Tapert SF, Brown GG, Baratta MV, Brown SA, 2004 fMRI BOLD response to alcohol stimuli in alcohol dependent young women. *Addictive Behaviors* 29(1), 33–50. [PubMed: 14667419]
- Tomasi D, Wang GJ, Wang R, Caparelli EC, Logan J, Volkow ND, 2015 Overlapping patterns of brain activation to food and cocaine cues in cocaine abusers. *Human brain mapping* 36(1), 120–136. [PubMed: 25142207]
- Tricomi E, Balleine Bernard W, O'Doherty John P, 2009 A specific role for posterior dorsolateral striatum in human habit learning. *European Journal of Neuroscience* 29(11), 2225–2232. [PubMed: 19490086]
- Vaghi MM, Vertes PE, Kitzbichler MG, Apergis-Schoute AM, van der Flier FE, Fineberg NA, Sule A, Zaman R, Voon V, Kundu P.J.B.p., 2017 Specific frontostriatal circuits for impaired cognitive flexibility and goal-directed planning in obsessive-compulsive disorder: evidence from resting-state functional connectivity. *81(8)*, 708–717.

- Vergara VM, Liu J, Claus ED, Hutchison K, Calhoun V, 2017 Alterations of resting state functional network connectivity in the brain of nicotine and alcohol users. *Neuroimage* 151, 45–54. [PubMed: 27864080]
- Wager TD, Barrett LF, 2017 From affect to control: Functional specialization of the insula in motivation and regulation. *bioRxiv*, 102368.
- Wang GJ, Volkow ND, Roque CT, Cestaro VL, Hitzemann RJ, Cantos EL, Levy AV, Dhawan AP, 1993 Functional importance of ventricular enlargement and cortical atrophy in healthy subjects and alcoholics as assessed with PET, MR imaging, and neuropsychologic testing. *Radiology* 186(1), 59–65. [PubMed: 8416587]
- Weiland BJ, Sabbineni A, Calhoun VD, Welsh RC, Bryan AD, Jung RE, Mayer AR, Hutchison KE, 2014 Reduced Left Executive Control Network Functional Connectivity Is Associated with Alcohol Use Disorders. *Alcoholism: Clinical and Experimental Research* 38(9), 2445–2453.
- White NM, 2009 Some highlights of research on the effects of caudate nucleus lesions over the past 200 years. *Behavioural brain research* 199(1), 3–23. [PubMed: 19111791]
- Yin HH, Knowlton BJ, 2006 The role of the basal ganglia in habit formation. *Nature Reviews Neuroscience* 7(6), 464. [PubMed: 16715055]
- Zhou Y, Friston KJ, Zeidman P, Chen J, Li S, Razi A, 2018 The Hierarchical Organization of the Default, Dorsal Attention and Salience Networks in Adolescents and Young Adults. *Cerebral Cortex* 28(2), 726–737. [PubMed: 29161362]
- Zhou Z, Yuan Q, Mash DC, Goldman D, 2011 Substance-specific and shared transcription and epigenetic changes in the human hippocampus chronically exposed to cocaine and alcohol. *Proceedings of the National Academy of Sciences* 108(16), 6626–6631.
- Zhu X, Dutta N, Helton SG, Schwandt M, Yan J, Hodgkinson CA, Cortes CR, Kerich M, Hall S, Sun H, Phillips M, Momenan R, Lohoff FW, 2015 Resting-state functional connectivity and presynaptic monoamine signaling in Alcohol Dependence. *Human Brain Mapping* 36(12), 4808–4818. [PubMed: 26368063]
- Zola-Morgan S, Squire L, Amaral D, Suzuki W, 1989 Lesions of perirhinal and parahippocampal cortex that spare the amygdala and hippocampal formation produce severe memory impairment. *The Journal of Neuroscience* 9(12), 4355–4370. [PubMed: 2593004]

Highlights

- Little is known about resting state functional connectivity (FC) of the salience network in nontreatment-seeking alcoholics.
- We analyzed resting state functional magnetic resonance imaging data from 16 Non-treatment seeking alcoholics (NTS) and 21 social drinkers, using seed regions in the anterior insular cortex.
- Novel findings showed that NTS has higher insular FC with hippocampal regions and medial orbitofrontal cortex.
- Hyperactive interactions between networks involved with interoception, salience attribution, learning, and behavioral control may underlie the development and maintenance of alcohol use disorders.

Figure 1.

Anterior insula seed regions. 5mm^3 cubical seed regions are shown in an axial view of the right hemisphere (R). Coordinates denote the centroid of each seed, and are in MNI space. Numerical designations for each seed are from Cauda et al. 2011, 2012.

Figure 2.

Functional connectivity patterns of the left and right anterior insula seeds (panels A and B, respectively). Regions in which SD have greater FC compared to NTS are shown in magenta, while regions where NTS have greater FC than SD are indicated in cyan. Statistical inferences were made with a cluster-level significance $p_{FWE} < 0.05$, FWE-corrected for the whole-brain multiple comparisons with the cluster-forming threshold, $p = 0.001$, uncorrected.

Figure 3.

Additional regions in which insular FC is greater in SD than in NTS. The left AIC seeds had enhanced FC with the right superior orbital gyrus (SOG) (Panel A; sagittal view). The right AIC seeds had higher FC with the right anterior putamen (Panel B; coronal view). Statistical inferences were made with a cluster-level significance $p_{FWE} < 0.05$ (FWE-corrected for the whole-brain multiple comparisons with the cluster-forming threshold, $p = 0.001$, uncorrected).

Figure 4.

Additional regions with greater FC in NTS than SD. Data in both panels are from the right AIC seed analysis. NTS have higher insular connectivity with bilateral hippocampal regions (Panel A; coronal view) as well as the caudal portion of medial OFC (Panel B, midline sagittal view). Statistical inferences were made with a cluster-level significance $p_{FWE} < 0.05$ (FWE-corrected for the whole-brain multiple comparisons with the cluster-forming threshold, $p = 0.001$, uncorrected).

Table 1.

Subject Demographics and Characteristics

	SD (<i>n</i> = 21)	NTS (<i>n</i> = 16)	Significance
Sex	10 M; 11 F	11 M; 5 F	$\chi^2(1) = 1.52, p = 0.32$
Age	35.3 (± 11.4)	37.3 (± 11.6)	$t(35) = -0.32, p = 0.60$
Edu	15.6 (± 2.1)	14.1 (± 2.5)	$t(35) = 1.97, p = 0.06$
Race	17 W; 3 AA; 1 As	12 W; 3 AA; 1 AI	
Drinks/day	2.5 (± 0.8)	5.7 (± 3.5)	$t(35) = -4.00, p < 0.001$
Drinks/week	8.7 (± 5.8)	30.0 (± 22.0)	$t(35) = -4.078, p < 0.001$
Smokers	4	6	$\chi^2(1) = 1.57, p = 0.274$

M: Male; F: Female; *n.s.*: non-significant; W: White; AA: African American; As: Asian; AI: American Indian.

Table 2.

MNI Coordinates for Anterior Insular Cortex Seed Regions

ROI	x	y	z	mm ³
2 L	-37	8	-7	125
2 R	40	9	-8	125
5 L	-33	12	4	125
5 R	38	10	0	125
6 L	-39	2	1	125
6 R	40	2	0	125
8 L	-31	12	9	125
8 R	34	11	8	125
9 L	-36	1	8	125
9 R	36	0	7	125

Table 3.

Regions with greater FC in SD than NTS (SD > NTS Contrasts)

LEFT AIC SEEDS					RIGHT AIC SEEDS				
Region	Cluster		Peak		Region	Cluster		Peak	
	<i>k</i>	<i>p</i> (FWE)	<i>Z</i>	MNI Coordinate		<i>k</i>	<i>p</i> (FWE)	<i>Z</i>	MNI Coordinate
<i>R Superior orbital gyrus</i>	5169	<0.001	6.26	31 65 -3	<i>L Cerebellum</i>	3349	<0.001	4.62	-8 -75 -16
<i>R Superior parietal lobule</i>	2578	<0.001	5.69	43 -50 70	<i>L Superior parietal lobule</i>	1918	0.003	4.97	-45 -50 66
<i>L Calcarine gyrus</i>	1798	0.005	4.26	-11 -67 6	<i>R Inferior frontal gyrus</i>	1368	0.019	4.69	36 33 28
<i>L Inferior frontal gyrus</i>	1411	0.017	4.64	-57 32 11	<i>R Putamen</i>	1343	0.021	4.46	21 7 11
<i>R Inferior frontal gyrus</i>	1251	0.029	4.14	48 16 14	<i>R Superior Orbital gyrus</i>	1148	0.042	4.14	29 65 -4

Statistical inferences made using cluster-level significance $p_{FWE} < 0.05$, FWE-corrected for the whole-brain multiple comparisons with the cluster-forming threshold, $p = 0.001$, uncorrected.

Table 4.

Regions with greater FC in NTS than SD (NTS > SD Contrasts)

LEFT AIC SEEDS					RIGHT AIC SEEDS				
Region	Cluster		Peak		Region	Cluster		Peak	
	<i>k</i>	<i>p</i> (FWE)	<i>Z</i>	MNI Coordinates		<i>k</i>	<i>p</i> (FWE)	<i>Z</i>	MNI Coordinates
<i>L Cerebellum</i>	2958	<0.001	4.68	−21 −36 −28	<i>R medial OFC</i>	2437	0.001	5.12	2 32 −25
<i>R Fusiform Gyrus</i>	2096	0.002	5.34	33 −56 −10	<i>L Hippocampus</i>	1807	0.004	4.70	−34 −16 −19
<i>R Posterior insular cortex</i>	1187	0.037	4.52	34 −25 20	<i>L Supramarginal gyrus</i>	1439	0.015	4.73	−70 −21 15
					<i>R Parahippocampal gyrus</i>	1334	0.022	5.72	19 −7 −26

Statistical inferences made using cluster-level significance $p_{FWE} < 0.05$, FWE-corrected for the whole-brain multiple comparisons with the cluster-forming threshold, $p = 0.001$, uncorrected.