Published in final edited form as:

Public Health Nutr. 2019 September ; 22(13): 2472–2478. doi:10.1017/S1368980019000843.

Associations between frequency of food shopping at different food sources and fruit and vegetable intake among rural residents in upstate New York

Brian K. Lo¹, Sarah Megiel¹, Evelyn Liu¹, Sara C. Foltz², Meredith L. Graham¹, Rebecca A. Seguin¹

¹Division of Nutritional Sciences, Cornell University, Ithaca, NY, USA 14853

²Friedman School of Nutrition Science and Policy, Tufts University, Boston, MA, USA 02111

Abstract

Objective—To examine the frequency of shopping at different food sources, and the associations between shopping at different food sources and fruit and vegetable (FV) intake among upstate New York rural residents.

Design—Cross-sectional study. Descriptive statistics and linear mixed models were used.

Setting—Eighty-two rural communities in upstate New York, USA.

Participants—465 adults (82.3% female, mean age = 51.5, mean BMI = 31.7).

Results—Within one's community, the majority of participants reported often going to supermarkets (73.1%). Many participants sometimes or occasionally shopped at superstores (48.0%), convenience stores (57.9%), small grocery stores or local markets (57.2%), farmers' markets or FV stores (66.6%), dollar stores (51.5%), pharmacies (46.0%), or farm stands or community supported agricultures (56.8%). Most participants had never utilized food banks or food pantries (94.0%), community gardens (92.7%), or home food delivery (91.9%). While frequent visits to farmers' markets or farm stands were associated with higher fruit intake ($P < 0.001$), frequent visits to food co-ops or food hubs were associated with lower fruit intake ($P = 0.004$). Frequent visits to convenience stores ($P = 0.002$) and dollar stores ($P = 0.004$) were associated with lower vegetable intake. When FV intakes were combined, frequent visits to farmers' markets or farm stands ($P < 0.001$) were associated with higher FV intake, and frequent visits to convenience stores ($P = 0.005$) were associated with lower FV intake.

Corresponding author: Rebecca A. Seguin, PhD, rs946@cornell.edu, Tel: (607) 255-8250, Fax: (607) 254-1681.

Authorship:

R.A.S., S.C.F., and M.L.G. designed the study and wrote the protocol. M.L.G. and B.L. implemented the study. B.L. conducted the statistical analysis. B.L., S.M., and E.L. wrote the first draft of the manuscript. R.A.S., S.C.F., and M.L.G. provided substantial comments and feedback on preliminary drafts of this paper. All authors contributed to and have approved the final manuscript.

Conflict of interest:

None.

Ethical standards disclosure:

This study was conducted according to the guidelines laid down in the Declaration of Helsinki and all procedures involving human subjects/patients were approved by the Cornell University Institutional Review Board (Protocol# 1402004505). Written informed consent was obtained from all subjects/patients.

Conclusions—Findings from the present study provide important insight for informing future food environment interventions related to helping rural residents to consume adequate FV.

Keywords

Rural; shopping behavior; fruit and vegetable; food environment

Introduction

Although idyllic images of farm life are often perceived as facilitating rural residents' access to fresh produce, rural populations are actually less likely to consume adequate fruit and vegetables (FV) to maintain physical health in comparison to their non-rural counterparts ⁽¹⁾. Only 25% of rural residents consume five or more daily servings of FV, compared to 40% of the general population in the United States ⁽²⁾. This difference in FV consumption, along with other dietary disparities, may contribute to higher rates of obesity and chronic disease among rural populations ^(3–5).

It is important to understand the factors that contribute to lower intakes of FV. The small but growing literature on rural food environments has found that one such factor may be rural residents' limited access to affordable, healthy foods ^(6–9). There is some evidence that in comparison to urban and suburban areas, rural communities struggle to maintain and attract large food retailers and grocery stores due to their declining customer base, aging ownership, lack of distribution infrastructure, and lack of small business capital ⁽¹⁰⁾. Therefore, rural residents often rely on a variety of both traditional and non-traditional sources, such as retail food outlets, mobile food vendors, farm stands, dollar stores, pharmacies, convenience stores within a gas station, informal ordering groups, gardening, and hunting ^(11–17). The shopping behaviors of rural residents among these sources is unknown, however, as is how shopping at these sources affects their diet quality.

Given that the rural foodscape is unique and complex, information on where rural residents shop for food and how that relates to their diet quality is vital for informing the development of relevant strategies to improve rural healthy food access and for encouraging healthy eating among rural residents. The present study aims to examine 1) the frequency of shopping at different food sources and 2) the associations between shopping at different food sources and FV intake among rural residents in upstate New York.

Methods

Context

The present study was conducted within the context of a randomized, community-based intervention trial aimed at cardiovascular disease risk reduction among rural midlife and older women: *Strong Hearts, Healthy Communities* (SHHC) ⁽¹⁸⁾. A total of 179 participants were enrolled in the community-based trial and completed a detailed baseline online survey prior to randomization in 2017. Those participants were asked to identify one to five of their closest family members and/or friends, who were then invited to complete a similar online questionnaire (n=497).

The present study used baseline data gathered from SHHC participants and their family members and friends. We excluded individuals who: did not fill out the questionnaire (n=6), were < 18 years old (n=14), did not provide a household address (n=8), did not live in a rural area defined by the US Census Bureau (n=122) ⁽¹⁹⁾, or did not live in New York State (n=61). In the present study, a rural area is defined as “all population, housing, and territory not included within an urbanized area or urban cluster;” urbanized areas are areas with a population of 50,000 or more, and urban clusters are areas with populations of between 2,500 and 50,000 ⁽¹⁹⁾. This definition captures “a wide variety of settlements, from densely settled small towns and ‘large-lot’ housing subdivisions on the fringes of urban areas, to more sparsely populated and remote areas,” which better reflects the changing settlement patterns of the US ⁽¹⁹⁾. The US Census Bureau further divides counties into three rurality categories: (1) completely rural: counties that have a population that is 100% rural; (2) mostly rural: counties that have a population that is 50-99.9% rural; and (3) mostly urban: counties that have a population that is less than 50% rural ⁽¹⁹⁾. Participants who lived in mostly urban counties were excluded from analysis. The final sample for the present study consisted of a total of 465 individuals (from this point onwards referred to as “participants”) with 160 SHHC study participants and 305 of their family members and friends.

Measures

Frequency of visiting different types of food sources—To examine participants’ shopping patterns, we adapted a previously used question to ask participants ‘When you go shopping for food in your community/town, how often do you go to each of the following locations?’ ⁽²⁰⁾ Options included the following: supermarket, superstore, convenience store, small grocery or local market, farmers’ market or FV store, dollar store, pharmacy, farm stand or community supported agriculture (CSA), food bank or food pantry, food co-op or food hub, community garden, and home food delivery. Response options for each question ranged from 1 to 4: 1=Never, 2=Occasionally, 3=Sometimes, and 4=Often, as well as the option ‘This food resource is not present in my community/town.’ To ensure reporting consistency, local examples of different food sources and definitions of “local community/town” were provided within the survey. We defined community/town as the community or town that is closest to participants’ home, including where they go to work, school, or shop. Participants were also asked to indicate whether they did the majority of their food shopping within their community (Yes/No). The survey instrument used in the present study is provided as Supplementary Material 1.

Fruit and vegetable intake—Participants self-reported their FV intake by answering the following questions from the My Life Check - Life’s Simple 7 developed by the American Heart Association: ‘How much fruit (in cups) do you eat in an average day?’ and ‘How many vegetables (in cups) do you eat in an average day?’ ⁽²¹⁾ Examples of one cup equivalents were provided to assist participants with recall of portion sizes.

Sociodemographic variables—Participants reported their age, sex, education, employment status, marital status, car access, weight, height, and home address.

Statistical analyses

Descriptive statistics were used to characterize participants' demographics, frequency of visiting different food sources, and FV intake. We found that the distributions of fruit, vegetable, and FV combined intakes were right-skewed; therefore, we applied square root transformations to these variables for analysis. Similarly, since the distributions of shopping frequency at food bank or food pantry, food co-op or food hub, community garden, and home food delivery were skewed to the lower end (see Table 2), these food sources' shopping frequency variables were dichotomized into "Ever" versus "Never" by collapsing "Occasionally," "Sometimes," and "Often" into "Ever" for analysis.

Linear mixed models were used to examine participants' FV intake as a function of their frequency of shopping at different food sources. Models controlled for age, sex, body mass index (BMI), education, employment status, marital status, and whether participants did the majority of their food shopping in their community. Participants' town of residence and SHHC participants' family and friend referrals were treated as random effects. Results from all models are reported as estimated marginal means of the squared root transformations of FV intakes. To reduce the chance of observing false positive findings, type 1 error was set at 0.01 to account for the multiple tests that were conducted, and a Bonferroni correction was also applied to account for multiple between-group comparisons within each model. All analyses were conducted using SPSS version 25 (SPSS Inc., Chicago, USA).

Results

Table 1 summarizes participant characteristics. The majority of participants were female (82.3%) with a mean age of 51.5 (SD 14.4) and a mean BMI of 31.7 (SD 7.8) (29.0% overweight and 52.5% obese). The majority of participants had completed education beyond high school (71.0%), were employed (73.1%), were either partnered or married (71.1%), and had access to a car for food shopping (95.3%). Two-thirds of participants reported doing the majority of their food shopping in their community (66.7%). Median daily fruit intake was 1.0 cup (IQR=1.5) and median daily vegetable intake was 1.5 cups (IQR=1.0). Median total daily FV intake was 3.0 cups (IQR=2.0).

Frequency of shopping at different food sources

Table 2 outlines participants' frequency of shopping at different food sources. When shopping for food within one's community, the majority of participants reported often going to supermarkets (73.1%). Many participants sometimes or occasionally shopped at superstores (48.0%), convenience stores (57.9%), small grocery stores or local markets (57.2%), farmers' markets or FV stores (66.6%), dollar stores (51.5%), pharmacies (46.0%), or farm stands or CSAs (56.8%). Almost all participants had never utilized food banks or food pantries (94.0%), food co-ops or food hubs (87.8%), community gardens (92.7%), or home food delivery (91.9%). None of the participants chose "This food resource is not present in my community/town" as a response to any of the shopping locations. However, this response may not have been visible to viewers online. Therefore, missing values in this question likely reflect that these food sources are not available in participants' communities.

Associations between frequency of shopping at different food sources and FV intake

While frequent visits to farmers' markets or farm stands were associated with higher mean squared root fruit intake ($P<0.001$), frequent visits to food co-ops or food hubs were associated with lower fruit intake ($P=0.004$). On the other hand, frequent visits to convenience stores ($P=0.002$) and dollar stores ($P=0.004$) were associated with lower mean squared root vegetable intake. Interestingly, people who never visited a community garden consumed more vegetables than those who did ($P=0.008$). When FV intakes were analyzed together, frequent visits to farmers' markets or farm stands remained associated with higher mean squared root FV intake ($P<0.001$), and frequent visits to convenience stores also remained associated with lower mean squared root FV intake ($P=0.005$). All significant pairwise comparisons are indicated in Table 3.

Discussion

This study is among the first to examine rural residents' shopping behaviors and their associations with FV intake^(22,23). We found that rural residents who shop at farmers' markets or farm stands more frequently consume more FV. Similar relationships have been supported in other studies^(22–24). Previous studies have suggested that increasing visitation to farmers' markets could be an effective strategy to enhance FV intake because such markets generally carry produce of higher quality, allow consumers to interact with growers, and expose consumers to nutrition information, cooking instructions, and recipe ideas^(25–28). Due to the cross-sectional nature of our data, it could be that people with a high intake of FV simply tend to shop at farmers' markets or farm stands more often. In contrast, we found that frequent visits to convenience stores are associated with lower FV intake, and frequent visits to dollar stores are associated with lower vegetable intake. Such findings are similar to a study conducted in southwestern Ontario, Canada⁽²⁰⁾. Other studies also found that the lower availability of produce at convenience stores and dollar stores was associated with fewer produce purchases⁽²⁹⁾. However, no association was found between visits to convenience stores and dollar stores, and fruit intake. Unexpectedly, we found that visits to food co-ops or food hubs were associated with lower fruit intake, and visits to community gardens were associated with lower vegetable intake. This could be biased by the fact that only 12.2% of participants reported utilizing food co-ops or food hubs, and only 7.3% of them reported visiting community gardens.

We found no associations between FV intake and shopping frequency at other food sources. The lack of associations between FV intake and utilization of supermarkets, food banks, and home food delivery could be due to the lack of variation among participants' responses (73.1% reported often going to supermarkets, 94.0% never going to food banks, and 91.9% never using home food delivery). Our results were in contradiction with three urban studies which found associations between higher frequency shopping at supermarkets, food co-ops, and community gardens, and increased produce intake^(20,30,31). The discrepancies between our study and others may have been the result of differences in locational and rural/urban characteristics as well as in differences between the methods used in assessing shopping behaviors and FV intake.

Our data on rural residents' frequency of shopping at different food sources provides insight for informing future interventions. More than half of the participants reported going to convenience stores and dollar stores for food. Given that these food outlets are common in rural communities and our study showed that visits to these stores may be associated with lower vegetable intake, these food retail settings might serve as important venues for public health interventions to improve rural residents' diet quality. Likewise, home food delivery was rarely used by rural residents, and future studies could examine the feasibility of home food delivery services to enhance rural residents' access to FV, particularly among older adults and the disabled, who may lack the ability to drive or travel independently for food.

This study has some limitations. First, the cross-sectional nature of the present study prevented the examination of the causality of participants' FV intake. Second, FV intake in cups was self-reported by participants, and may have been subject to measurement error. Multiple methods of FV intake assessment should be used, such as the National Cancer Institute's Fruit & Vegetable Intake Screener or the Automated Self-Administered 24-Hour (ASA24) Dietary Assessment Tool, and, ideally, other objective measures of FV intake including dermal carotenoid scans or blood carotenoids ^(32, 33). Third, our study only examined participants' shopping behavior within their community, and 33.3% of participants reported doing the majority of their food shopping outside of their community. However, our analyses controlled for whether participants shopped for the majority of their foods within their community. Future studies should also examine rural residents' food shopping behaviors outside of their community. Fourth, our results may not be generalized to other rural populations. The majority of participants in this study were midlife to older, employed females who were overweight or obese, and who had received education beyond high school. Fifth, we did not assess participants' income in the present study. Given the associations found in our study and previous research associating income with food shopping at different sources ^(34, 35), including income is important for further understanding. Finally, our survey instrument may not have captured all the available rural food resources, particularly those promoted through word-of-mouth. Some of the food sources examined in our study, such as food hubs or co-ops, may not have been familiar or applicable to rural residents. Future studies should better characterize the unique foodscape in rural communities.

Conclusions

We found that visits to different food sources contribute differently to rural residents' FV intake. Findings from the present study provide important insight for informing future food environment interventions related to helping rural residents to consume adequate FV.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements:

We are grateful to the rural residents who participated in our study and Judy Ward (Cornell) for her work and contributions to this research.

Financial support:

This study was supported by grant R01 HL120702 from the National Institutes of Health and the National Heart, Lung, and Blood Institute (NHLBI). The funders/sponsors had no role in the design, analysis or writing of this article.

References

1. Dean WR & Sharkey JR (2011) Rural and urban differences in the associations between characteristics of the community food environment and fruit and vegetable intake. *J Nutr Educ Behav* 43, 426–433. [PubMed: 21616721]
2. Lutfiyya MN, Chang LF & Lipsky MS (2012) A cross-sectional study of US rural adults' consumption of fruits and vegetables: do they consume at least five servings daily? *BMC Public Health* 12, 280. [PubMed: 22490063]
3. Jackson JE, Doescher MP, Jerant AF et al. (2005) A national study of obesity prevalence and trends by type of rural county. *J Rural Health* 21, 140–148. [PubMed: 15859051]
4. Befort CA, Klemp JR, Austin HL et al. (2012) Outcomes of a weight loss intervention among rural breast cancer survivors. *Breast Cancer Res Treat* 132, 631–639. [PubMed: 22198470]
5. Matthews KA, Croft JB, Liu Y et al. (2017) Health-related behaviors by urban-rural county classification — United States, 2013. *MMWR Surveill Summ.* 66, 1–8.
6. Sharkey JR, Johnson CM & Dean WR (2010) Food access and perceptions of the community and household food environment as correlates of fruit and vegetable intake among rural seniors. *BMC Geriatr* 10, 32. [PubMed: 20525208]
7. Campbell EA, Shapiro MJ, Welsh C et al. (2017) Healthy food availability among food sources in rural Maryland counties. *J Hunger Environ Nutr* 12, 328–341. [PubMed: 29242739]
8. Bailey JM (2010) Rural Grocery Stores: Importance and Challenges. Retrieved from Center for Rural Affairs website: <http://files.cfra.org/pdf/rural-grocery-stores.pdf> (accessed August 2018)
9. Dean WR, Sharkey JR & Johnson CM (2011) Food insecurity is associated with social capital, perceived personal disparity, and partnership status among older and senior adults in a largely rural area of central Texas. *J Nutr Gerontol Geriatr* 30, 169–186. [PubMed: 21598165]
10. Pinard CA, Fricke HE, Smith TM et al. (2016) The future of the small rural grocery store: a qualitative exploration. *Am J Health Behav* 40, 749–760. [PubMed: 27779943]
11. Morton L & Blanchard T (2007) Starved for access: life in rural America's food deserts. *Rural Realities* 1, 1–10.
12. Blanchard T & Lyson T (2010) Retail concentration, food deserts, and food disadvantaged communities. *Food Cult Soc* 13, 460.
13. Sharkey JR, Dean WR & Johnson CM (2012) Use of vendedores (mobile food vendors), pulgas (flea markets), and vecinos o amigos (neighbors or friends) as alternative sources of food for purchase among Mexican-origin households in Texas border colonias. *J Acad Nutr Diet* 112, 705–710. [PubMed: 22709775]
14. Valdez Z, Dean WR & Sharkey JR (2012) Mobile and home-based vendors' contributions to the retail food environment in rural South Texas Mexican-origin settlements. *Appetite* 59, 212–227. [PubMed: 22531289]
15. Van Hoesen J, Bunkley B & Currier C (2013) A GIS-based methodology toward refining the concept of rural food deserts: a case study from Rutland County, Vermont. *J Agric Food Syst Community Dev* 3, 61–76.
16. Wegener J & Hanning RM (2010) Concepts and measures of 'alternative' retail food outlets: considerations for facilitating access to healthy, local food. *J Hunger Environ Nutr* 5, 158–173.
17. Yousefian A, Leighton A, Fox K et al. (2011) Understanding the rural food environment—perspectives of low-income parents. *Rural Remote Health* 11, 1631. [PubMed: 21513422]
18. Seguin RA, Eldridge G, Graham ML et al. (2016) Strong Hearts, healthy communities: a rural community-based cardiovascular disease prevention program. *BMC Public Health* 16, 86. [PubMed: 26822982]

19. Ratcliffe M, Burd C, Holder K et al. (2016) Defining Rural at the U.S. Census Bureau (ACSGEO-1). Washington DC: U.S. Census Bureau.
20. Minaker LM, Olstad DL, Thompson ME et al. (2016) Associations between frequency of food shopping at different store types and diet and weight outcomes: findings from the NEWPATH study. *Public Health Nutr* 19, 2268–2277. [PubMed: 26956712]
21. American Heart Association (2017) My Life Check - Life's Simple 7 http://www.heart.org/HEARTORG/Conditions/My-Life-Check---Lifes-Simple-7_UCM_471453_Article.jsp#.W0kVL9hKjR1 (accessed July 2018).
22. Pitts SBJ, Gustafson A, Wu Q et al. (2014) Farmers' market use is associated with fruit and vegetable consumption in diverse southern rural communities. *Nutr J* 13, 1. [PubMed: 24405527]
23. Jilcott Pitts SB, Wu Q, McGuirt JT et al. (2013) Associations between access to farmers' markets and supermarkets, shopping patterns, fruit and vegetable consumption and health indicators among women of reproductive age in eastern North Carolina, USA. *Public Health Nutr* 16, 1944–1952. [PubMed: 23701901]
24. Freedman DA, Choi SK, Hurley T et al. (2013) A farmers' market at a federally qualified health center improves fruit and vegetable intake among low-income diabetics. *Prev Med*. 56, 288–292. [PubMed: 23384473]
25. Feagan R (2008) Direct marketing: towards sustainable local food systems? *Local Environ* 13, 161–167.
26. Brown A (2002) Farmers' market research 1940-2000: an inventory and review. *Am J Altern Agric* 17, 167–176.
27. DeWitt E, McGladrey M, Liu E et al. (2017) Social marketing campaign at farmers' markets to encourage fruit and vegetable purchases in rural obese counties. *J Nutr Educ Behav* 49, S109.
28. Stewart H (2018) Shopping at Farmers' Markets and Roadside Stands Increases Fruit and Vegetable Demand. USDA Economic Research Service. <https://www.ers.usda.gov/amber-waves/2018/march/shopping-at-farmers-markets-and-roadside-stands-increases-fruit-and-vegetable-demand/> (accessed July 2018).
29. Ruff RR, Akhund A & Adjoian T (2016) Small convenience stores and the local food environment: an analysis of resident shopping behavior using multilevel modeling. *Am J Health Promot* 30, 172–180. [PubMed: 25806566]
30. Zenk SN, Schulz AJ, Hollis-Neely T, et al. (2005) Fruit and vegetable intake in African Americans: income and store characteristics. *Am J Prev Med* 29, 1–9. [PubMed: 15958245]
31. Alaimo K, Packnett E, Miles RA, et al. (2008) Fruit and vegetable intake among urban community gardeners. *J Nutr Educ Behav* 40, 94–101. [PubMed: 18314085]
32. Shim JS, Oh K, & Kim HC (2014) Dietary assessment methods in epidemiologic studies. *Epidemiol Health* 36.
33. Subar AF, Laurence SF, Janet A et al. (2015) Addressing current criticism regarding the value of self-report dietary data, 2. *J Nutr* 145, 2639–2645. [PubMed: 26468491]
34. Ledoux TF & Vojnovic I (2013) Going outside the neighborhood: the shopping patterns and adaptations of disadvantaged consumers living in the lower eastside neighborhoods of Detroit, Michigan. *Health Place* 19, 1–14. [PubMed: 23142639]
35. Stern D, Robinson WR, Ng SW et al. (2015) US household food shopping patterns: dynamic shifts since 2000 and socioeconomic predictors. *Health Aff* 34, 1840–1848.

Table 1.

Participant characteristics of SHHC trial participants and their family members and friends in rural upstate New York (2017/2018)

	Mean	SD
Age (years) (n=457)	51.5	14.4
BMI (kg/m ²) (n=458)	31.7	7.8
	n	%
Sex (n=464)		
<i>Female</i>	382	82.3
<i>Male</i>	82	17.7
Education (n=465)		
<i>High school or below</i>	135	29.0
<i>Vocational school or some college</i>	107	23.0
<i>College graduate</i>	118	25.4
<i>Post-graduate</i>	105	22.6
Employment (n=464)		
<i>Yes</i>	339	73.1
<i>No</i>	125	26.9
Marital status (n=464)		
<i>Single</i>	134	28.9
<i>Partnered or married</i>	330	71.1
Car access for food shopping (n=464)		
<i>Yes</i>	442	95.3
<i>No</i>	22	4.7
Do majority of food shopping in own community (n=463)		
<i>Yes</i>	309	66.7
<i>No</i>	154	33.3
	Median	Interquartile range
Daily fruit intake (cups) (n=464)	1.0	1.5
Daily vegetable intake (cups) (n=462)	1.5	1.0
Total daily fruit and vegetable intake (cups) (n=462)	3.0	2.0

Table 2.

Frequency of shopping at different food sources among SHHC trial participants and their family members and friends in rural upstate New York (2017/2018)

	Never		Occasionally		Sometimes		Often	
	n	%	n	%	n	%	n	%
Supermarket (n=431, 7.3% missing)	20	4.6	37	8.6	59	13.7	315	73.1
Superstore (n=391, 15.9% missing)	53	13.6	94	24.0	94	24.0	150	38.4
Convenience store (n=453, 2.6% missing)	163	36.0	187	41.3	75	16.6	28	6.2
Small grocery store or local market (n=415, 10.8% missing)	125	30.1	143	34.5	94	22.7	53	12.8
Farmers' market or FV store (n=441, 5.2% missing)	60	13.6	166	37.6	128	29.0	87	19.7
Dollar store (n=451, 3.0% missing)	194	43.0	168	37.3	64	14.2	25	5.5
Pharmacy (n=440, 5.4% missing)	197	44.8	130	29.6	72	16.4	41	9.3
Farm stand or CSA (n=428, 8.0% missing)	137	32.0	150	35.1	93	21.7	48	11.2
Food bank or food pantry (n=434, 6.7% missing)	408	94.0	12	2.8	8	1.8	6	1.4
Food co-op or food hub (n=400, 14.0% missing)	351	87.8	31	7.8	11	2.8	7	1.8
Community garden (n=357, 23.2% missing)	331	92.7	14	3.9	10	2.8	2	0.6
Home food delivery (n=405, 12.9% missing)	372	91.9	21	5.2	8	2.0	4	1.0

Associations between frequency of shopping at different food sources and FV intake among SHHC trial participants and their family members and friends in rural upstate New York (2017/2018)

Table 3.

Food Source	Fruit intake						Vegetable intake						FV intake					
	Never		Occasionally		Sometimes		Often		Never		Occasionally		Sometimes		Often		μ	SE
Supermarket	* μ	SE	μ	SE	μ	SE	μ	SE	μ	SE	μ	SE	μ	SE	μ	SE	μ	SE
	1.28	0.13	1.12	0.09	1.02	0.08	1.08	0.05	1.11	0.11	1.17	0.08	1.27	0.07	1.22	0.04	1.69	0.13
Superstore			1.08	0.07	1.10	0.07	1.06	0.06	1.27	0.07	1.23	0.06	1.26	0.06	1.12	0.05	1.79	0.09
	1.22	0.08																
Convenience store			1.05	0.05	0.99	0.07	1.18	0.10	1.30 ^{††}	0.05	1.24	0.04	1.08 ^{††}	0.06	1.12	0.09	1.78 ^{†††}	0.06
	1.16	0.05																
Small grocery store or local market			1.15	0.06	1.10	0.07	1.04	0.08	1.22	0.05	1.28	0.05	1.11	0.05	1.21	0.07	1.67	0.06
	1.08	0.06																
Farmers' market or FV store			1.14 [§]	0.05	1.04 [‡]	0.06	1.27 ^{†‡‡}	0.07	1.13	0.06	1.22	0.05	1.22	0.05	1.34	0.06	1.50 ^{§§}	0.08
	0.86 ^{†‡§}	0.07																
Dollar store			1.08	0.05	1.12	0.08	1.11	0.11	1.25 ^{***}	0.04	1.23	0.04	1.03 ^{***}	0.06	1.11	0.09	1.69	0.05
	1.07	0.05																
Pharmacy			1.06	0.06	1.10	0.07	1.21	0.09	1.26	0.04	1.19	0.05	1.16	0.06	1.21	0.08	1.71	0.06
	1.09	0.05																
Farm stand or CSA			1.11	0.06	1.17	0.07	1.19	0.08	1.22	0.05	1.20	0.05	1.24	0.06	1.37	0.07	1.66	0.06
	1.04	0.05																
Food Source	Never		Occasionally		Sometimes		Often		Never		Occasionally		Sometimes		Often		Ever	
	μ	SE	μ	SE	μ	SE	μ	SE	μ	SE	μ	SE	μ	SE	μ	SE	μ	SE
Food bank or food pantry	1.08	0.04	1.13	0.10	1.13	0.08	1.13	0.09	1.22	0.04	1.12	0.09	1.12	0.09	1.59	0.04	1.68	0.11
Food co-op or food hub	1.12 ^{//}	0.04	0.89 ^{//}	0.08	0.89 ^{//}	0.08	0.89 ^{//}	0.08	1.23	0.04	1.16	0.07	1.16	0.07	1.51	0.05	1.71	0.09
Community garden	1.11	0.04	1.02	0.10	1.02	0.10	1.02	0.10	1.25 ^{††‡}	0.04	1.02 ^{††‡}	0.09	1.02 ^{††‡}	0.09	1.53	0.04	1.71	0.10
Home food delivery	1.09	0.05	1.00	0.09	1.00	0.09	1.00	0.09	1.23	0.04	1.22	0.08	1.22	0.08	1.60	0.05	1.86	0.09

* All μ represent the estimated marginal squared root mean intakes; all models controlled for age, sex, BMI, education, employment status, marital status, and whether participants did the majority of their food shopping in their community; participants' town of residence and SHHC participants' family and friend referrals were treated as random effects

[†] Often vs Never: =0.40, SE=0.09, P<0.001

[‡] Often vs Sometimes: =0.23, SE=0.07, P=0.006

[§] Occasionally vs Never: =0.28, SE=0.08, P=0.002

^{//} Ever vs Never: =-0.23, SE=0.08, P=0.004

^{††} Sometimes vs Never: =-0.22, SE=0.06, P=0.002

^{***} Sometimes vs Never: =-0.22, SE=0.06, P=0.005

^{††‡} Ever vs Never: =-0.23, SE=0.08, P=0.008

Sometimes vs Never: =-0.27, SE=0.08, P=0.003
\$\$ Often vs Never: =0.38, SE=0.09, P<0.001
/// Often vs Sometimes: =0.24, SE=0.07, P=0.006

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript