Published in final edited form as:

J Solution Chem. 2019 July ; 48(7): 983–991. doi:10.1007/s10953-018-0836-7.

Binary Mixtures of Ionic Liquids in Aqueous Solution: Towards an Understanding of their Salting-In/Salting-Out Phenomena

Catarina M. S. S. Neves¹, Teresa B. V. Dinis¹, Pedro J. Carvalho¹, Bernd Schröder¹, Luís M. N. B. F. Santos², Mara G. Freire¹, João A. P. Coutinho^{1,*}

¹CICECO - Aveiro Institute of Materials, Department of Chemistry, University of Aveiro, 3810-193 Aveiro, Portugal

²CIQUP-Centro de Investigação em Química da Universidade do Porto, Departamento de Química e Bioquímica, Faculdade de Ciências, Universidade do Porto, Rua do Campo Alegre, 687, 4169-007 Porto, Portugal

Abstract

The order of the salting-in or salting-out inducing ability of ions on the aqueous solubility of macromolecules in aqueous solutions is known as the Hofmeister series. Taking into account that ionic liquids (ILs) are constituted by ions, they can exert similar effects on the solubility of other ILs in aqueous media. In order to expand the knowledge on the salting-in/-out ability of ILs, experimental studies on the solubility of 1-butyl-3-methylimidazolium bis(trifluoromethylsulfonylimide) in water in presence of other IL/salts were conducted at 298.15 K at atmospheric pressure. Both the impact of the anion and cation of the IL were evaluated with the following ILs/salts: 1-butyl-3-methylimidazolium chloride, 1-butyl-3-methylimidazolium hydrogensulfate, cholinium bis(trifluoromethylsulfonyl)imide, 1-butyl-1-methylpyrrolidinium bis(trifluoromethylsulfonyl)imide and lithium bis(trifluoromethylsulfonyl)imide, in a wide composition range. As happens with common salts, both salting-in and salting-out effects exerted by ILs were observed, with a higher impact exerted by the IL anion on the salting-out phenomenon. These data allow to better understand the ILs impact when designing liquid-liquid separation processes.

Keywords

Ionic Liquids; Solubility; Water; Salting-in effect; Salting-out effect; Hofmeister series

*Corresponding author: jcoutinho@ua.pt Tel: +351 234 401 507, Fax: +351 234 370 084.

CMSSN 0000-0002-8322-6210

TBVD 0000-0002-7645-8929

PJC 0000-0002-1943-0006

BS 0000-0002-3760-900X

LMNBFS 0000-0003-3040-0358

MGF 0000-0001-8895-0614

JAPC 0000-0002-3841-743X

1 Introduction

Ionic liquids (ILs) belong to the molten salts group and are usually formed by a large organic cation and a smaller organic or inorganic anion, allowing them to remain liquid below 373 K at atmospheric pressure [1]. One of the most interesting feature of ILs is their design ability for a specific task, by a particular combination of cations and anions [2].

The use of ILs as alternative solvents for a range of applications has attracted large interest [3–5]. Hydrophobic ILs have been used in the creation of liquid-liquid extraction systems with aqueous media [6], whereas hydrophilic ones have been used in the creation of aqueous biphasic systems (ABS) by the addition of a salting-out species [7]. In both types of systems, the manipulation of the ILs solubility in water by adding specific compounds could lead to the development of efficient extraction and purification systems to be used in separation processes [8–13]. Over the years, the ILs mutual solubilities with water have been studied and different IL structural effects have been addressed, namely the cation core [14–16], the alkyl chain length [17–20], the cation isomerism [18, 21, 22] and the anion nature [14]. However, when dealing with real aqueous matrices that contain, for instance, metabolites and salts, significant changes in the ILs solubility can occur. Freire *et al.* [23, 24] showed that the presence of salts in aqueous media has a significant influence on the mutual solubilities of water and ILs, following the salting-in/salting-out trends of the Hofmeister series [25]. The authors found that salting-out inducing ions (high charge density) and salting-in inducing ions (low charge density) act by different mechanisms [23, 24]. While the former act mainly through an entropic effect resulting from the formation of hydration complexes, the salting-in results from a direct ion binding of the low charge density ions to the hydrophobic moieties of the IL [23]. Additionally, Shahriari *et al.* showed the effect of salts on promoting the phase separation in ABS, *i.e.* in phase separation of mixtures comprising an hydrophilic IL and water [26]. The results gathered showed that the salts ability to form two phases follows the Hofmeister series, and that the salt ions molar entropy of hydration closely correlates with the salting-out effect [26].

Although the Hofmeister series is a well-known qualitative approach ranking the ions influence on the solubility of charged molecules, further research in order to rank IL ions in this series is still scarce [27–29]. Yang [28] and Zhao *et al.* [27] showed that the ILs ions, which follow the Hofmeister series, in the stability and activity of enzymes, and showed that ILs composed of chaotropic cations and kosmotropic anions can be valuable for the application of ILs in biotransformations. On the other hand, Constantinescu and Weingärtner [29] ranked the cations and anions of ILs in the Hofmeister series by evaluating the stability of ribonuclease A in different ILs aqueous solutions. Since ILs are constituted by ions, they may exert similar (salting-in/salting-out) effects of traditional salts on the solubility of other ILs in aqueous media. The aim of this work is to expand the knowledge of the salting-in/-out ability of ILs by evaluating their effect on the aqueous solubility of a given hydrophobic IL. For that purpose, experimental data on the solubility of 1-butyl-3-methylimidazolium bis(trifluoromethylsulfonyl)imide, $[\text{C}_4\text{C}_1\text{im}][\text{NTf}_2]$, in aqueous solutions containing other ILs were determined at 298.15 K at atmospheric pressure. Both the influence of the cation core and anion of the IL added to the system on the solubility of $[\text{C}_4\text{C}_1\text{im}][\text{NTf}_2]$ in water were evaluated.

2 Experimental Section

2.1 Chemicals used

The study of solubility of 1-butyl-3-methylimidazolium bis(trifluoromethanesulfonyl)imide, $[\text{C}_4\text{C}_1\text{im}][\text{NTf}_2]$ (99 % purity, halides < 100 ppm), in water was carried out in presence of the following ILs and salt: 1-butyl-3-methylimidazolium hydrogensulphate, $[\text{C}_4\text{C}_1\text{im}][\text{HSO}_4]$ (> 98 % purity, halides < 100 ppm); 1-butyl-3-methylimidazolium chloride, $[\text{C}_4\text{C}_1\text{im}]\text{Cl}$ (99 % purity); 1-butyl-1-methylpyrrolidinium bis(trifluoromethanesulfonyl)imide, $[\text{C}_4\text{C}_1\text{pyr}][\text{NTf}_2]$ (99 % purity, halides < 100 ppm); cholinium bis(trifluoromethanesulfonyl)imide, $[\text{N}_{1112}\text{OH}][\text{NTf}_2]$ (99 % purity, halides < 100 ppm); and lithium bis(trifluoromethanesulfonyl)imide, $\text{Li}[\text{NTf}_2]$ (99 % purity, halides < 100 ppm). With the exception of $\text{Li}[\text{NTf}_2]$, which has a melting temperature between (507-511) K [30], all remaining compounds fall in ILs class according to their general characterization of melting temperatures below 373 K. Figure 1 depicts the chemical structure of the ILs/salts investigated. To avoid any ambiguity, all these compounds will be hereafter referred as salts. All of these salts were supplied by Iolitec. Before their use, they were purified (to reduce the content of water and other volatile components) by drying individual samples at circa 353 K for at least 48 h under constant stirring and moderate vacuum conditions (≈ 0.1 Pa). After this procedure, ^1H , ^{13}C , and ^{19}F NMR analyses confirmed the purity of the samples used - respective spectra given in the ESI. Double-distilled water, passed by a reverse osmosis system and further treated with a MilliQ plus 185 water purification apparatus, was used in the preparation of all ($[\text{C}_4\text{C}_1\text{im}][\text{NTf}_2]$ + salt + water) ternary mixtures. Purity analyses of water revealed resistivity values of $18.2 \text{ M}\Omega\cdot\text{cm}$ and a TOC (Total Organic Carbon content) lower than $5 \mu\text{g}\cdot\text{dm}^{-3}$.

2.2 Experimental Procedure

The solubility of $[\text{C}_4\text{C}_1\text{im}][\text{NTf}_2]$ in aqueous solutions of salts was determined at 298.15 K at atmospheric pressure. The solubility of pure $[\text{C}_4\text{C}_1\text{im}][\text{NTf}_2]$ in water at 298 K was taken from the literature, corresponding to a value of $1.72 \times 10^{-2} \text{ mol}\cdot\text{kg}^{-1}$ [17]. Mixtures of $[\text{C}_4\text{C}_1\text{im}][\text{NTf}_2]$ + salt were gravimetrically prepared within $\pm 10^{-4}$ g, with the salt present at a concentration ranging between 500 and 100,000 ppm, followed by the addition of 15 mL of double-distilled water (the mass weighted of each component is given in Table S1 in the ESI). The ternary mixture was vigorously agitated and allowed to reach complete phase separation and equilibrium for at least 48 h. This period proved to be the time required to guarantee a complete separation of the two phases and that no further variations in mole fraction solubilities occurred [18]. The phases in equilibrium, kept in the glass vials, were stabilized for 24 h in an aluminum block designed for the purpose, which is further placed in an isolated air bath capable of maintaining the temperature within ± 0.01 K. The temperature control was achieved with a PID temperature controller driven by a calibrated Pt100 (class 1/10) temperature sensor inserted in the aluminum block. The solubility of $[\text{C}_4\text{C}_1\text{im}][\text{NTf}_2]$ in water was determined by UV-spectroscopy, using a SHIMADZU UV-1700 Pharma-Spec Spectrometer, at a wavelength of 211 nm, and using a calibration curve established before measuring the IL solubility (given in Fig. S6 in the ESI). For that, and according to the detection limits of the equipment (given in Table S1 in the ESI), the dilution factor was circa 1:500 (w/w) times. The solubility results are an average of at least five independent

measurements. The detailed values and respective associated uncertainty at 95% of confidence level are given in Table S1 in the ESI. A scheme of this procedure is depicted in Fig. 2. Since the mixtures with $[\text{C}_4\text{C}_1\text{im}]\text{Cl}$ and $[\text{C}_4\text{C}_1\text{im}][\text{HSO}_4]$ have a common cation, ionic chromatography (IC) was used to quantify $[\text{HSO}_4]^-$ and Cl^- at the water-rich phase. The quantitative analyses of $[\text{HSO}_4]^-$ and Cl^- water-rich phase was measure in a Dionex ionic chromatography system Model: Series 2000/SP using the anion IC column Model IonPac AS12A with background conductivity suppression technology. The aqueous phase was diluted immediately after the sampling in order to avoid further phase separation and finally diluted to the ionic chromatography concentration level of the calibration range (10^{-3} to 10^{-5} mol·dm $^{-3}$) of the IC analytical system. The molality of each IL in water phase is given in Table S1 in the Supporting Information.

3 Results and discussion

The impact of the addition of a second IL or salt in the $[\text{C}_4\text{C}_1\text{mim}][\text{NTf}_2]$ solubility in water was here investigated. This IL was chosen for being one of the “standard” hydrophobic ILs, with a high temperature and chemical stability, and because its solubility was previously widely studied in presence of a large diversity of salts [23, 24, 31]. In this work, each salt was added to $[\text{C}_4\text{C}_1\text{im}][\text{NTf}_2]$ in the concentration range of (500-100,000) ppm for the IL-salt binary mixture, and the solubility of $[\text{C}_4\text{C}_1\text{mim}][\text{NTf}_2]$ in water was measured at 298.15 K and atmospheric pressure. All systems here studied comprise always a similar cation or anion, thus avoiding ion exchange. The experimental results obtained are depicted in Fig. 3 and presented in Table S1 in the Supporting Information.

The influence of ILs/salts, *e.g.* at $\approx 10,000$ ppm, in the $[\text{C}_4\text{C}_1\text{im}][\text{NTf}_2]$ solubility in water follows the order $[\text{C}_4\text{C}_1\text{im}][\text{HSO}_4] > [\text{C}_4\text{C}_1\text{im}]\text{Cl} > \text{Li}[\text{NTf}_2] > [\text{N}_{1112}\text{OH}][\text{NTf}_2] > [\text{C}_4\text{C}_1\text{pyrr}][\text{NTf}_2]$. These results clearly establish that the change of the salt anion has a major impact on decreasing the solubility of $[\text{C}_4\text{C}_1\text{im}][\text{NTf}_2]$, or on inducing the salting-out effect, in good agreement with previous works [23, 24, 31]. Although all salts lead to the $[\text{C}_4\text{C}_1\text{im}][\text{NTf}_2]$ salting-out at higher concentrations, at lower concentrations of salt, mainly below 1,000 ppm, an increase in the $[\text{C}_4\text{C}_1\text{im}][\text{NTf}_2]$ solubility is observed for some salts. This inversion in behavior, *i.e.* a salting-in effect at low concentrations of salt followed by a salting-out at higher concentrations was also observed by Freire *et al.* [23], when evaluating the effect of traditional salts in the $[\text{C}_4\text{C}_1\text{im}][\text{NTf}_2]$ solubility [32]. This effect was also reported when ILs were added to a poly(ethylene glycol)/water mixture [33]. The authors explained that ILs with shorter alkyl chain can produce some water-structuring (salting-out) effect through the polar parts of the cation and anion, while the non-polar parts are responsible for the water-structure-breaking (salting-in) effect [33].

At the salting-out regions, the higher the ability of each salt ion to be hydrated, the higher its impact on decreasing the $[\text{C}_4\text{C}_1\text{im}][\text{NTf}_2]$ solubility in water, as addressed by the differentiation of their hydrogen-bond basicity's ($[\text{C}_4\text{C}_1\text{im}][\text{HSO}_4]$, 0.95 [34] > $[\text{C}_4\text{C}_1\text{im}]\text{Cl}$, 0.70 [34]) and hydrogen-bond acidity's ($[\text{N}_{1112}\text{OH}][\text{NTf}_2]$, 1.491 [35] > $[\text{C}_4\text{C}_1\text{pyrr}][\text{NTf}_2]$, 0.433 [35]). The solubility of $[\text{C}_4\text{C}_1\text{im}][\text{NTf}_2]$ in water decreases in the order $[\text{C}_4\text{C}_1\text{pyr}]^+ > [\text{N}_{1112}\text{OH}]^+ > \text{Li}^+$, being the lithium salt the one that promotes a strong salting-out effect, while the pyrrolidinium almost does not affect the solubility of the IL, even up to

concentrations of 100,000 ppm. The cholinium-based salt has a similar behavior to the lithium salt at lower concentration; however, at the highest concentration studied its salting-out effect is less pronounced. This behavior follows the Hofmeister series [25, 36] generally applied to common salts, because ammonium-based cations are weaker salting-out species than Li^+ . In general, organic cations have a lower effect on inducing salting-out than inorganic ones, mainly because of their low-charge density and lower ability to be hydrated. In what concerns the anion effect, ILs containing the common cation $[\text{C}_4\text{C}_1\text{im}]^+$ were used combined with the anions $[\text{HSO}_4]^-$ and Cl^- . These salts ability to salt-out $[\text{C}_4\text{C}_1\text{im}][\text{NTf}_2]$ follows the order: $[\text{HSO}_4]^- > \text{Cl}^-$. The anion $[\text{HSO}_4]^-$ is a stronger salting-out agent than Cl^- , although at the highest concentration studied both compounds are able to completely exclude $[\text{C}_4\text{C}_1\text{im}][\text{NTf}_2]$ from the aqueous medium. The IL with the halogen anion induces the $[\text{C}_4\text{C}_1\text{im}][\text{NTf}_2]$ salting-in at concentrations up to 7,000 ppm, having thus a behavior similar to that previously observed and discussed with NaCl [23].

According to the results of Fig. 3, the IL/salt effects on the $[\text{C}_4\text{C}_1\text{im}][\text{NTf}_2]$ solubility in water closely follows the Hofmeister series [37]. Furthermore, IL/salt anions induce a more significant salting-out effect than the corresponding cations, in agreement with previous studies on the effect of traditional salts over the ILs solubility in water [23], which is due to their higher hydration capability and polarization [38]. Like with common salts, ILs affect the solubility of charged molecules, such as other ILs. The experimental data obtained in this work reveal that IL ions seem to follow the Hofmeister series towards their ability to promote salting-in/-out effects through other ILs, and with IL anions presenting a more significant contribution in the salting-out effect. However, a more complex framework, particularly at the molecular-level, is still required in order to better understand IL ions specific effects.

This study contributes to the understanding of the solubility of a hydrophobic IL in water in presence of additional ILs/salts, allowing to design ILs for a given application and with a predicted solubility in water, as well as in the design of the solubility of a panoply of charged compounds in water by the addition of ILs.

4 Conclusions

The influence of ILs/salts ions in the aqueous solubility of other IL was here determined and discussed in terms of salting-in/salting-out phenomena. The results obtained show that the solubility of $[\text{C}_4\text{C}_1\text{im}][\text{NTf}_2]$ in water at 298.15 K displays a salting-in region at low concentrations of the salt added, followed by a salting-out region at higher concentrations. These effects depend on the IL/salt nature and its concentration, although the anion plays the main role on defining the salting-out regions due to its higher hydration capability and polarizability. The strength of the salting-in/-out phenomena allow classifying IL ions according to the Hofmeister series. The understanding of these phenomena helps in the design of solvents with a predicted solubility in water, and therefore on the tailoring of liquid-liquid processes involving ILs.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This work was developed in the scope of the projects CICECO-Aveiro Institute of Materials (Ref. FCT UID/CTM/50011/2013), financed by national funds through the FCT/MEC and when applicable co-financed by FEDER under the PT2020 Partnership Agreement. Catarina M. S. S. Neves and Teresa B. V. Dinis also acknowledge FCT for their grants (SFRH/BPD/109057/2015 and SFRH/BD/130958/2017, respectively). P. J. Carvalho acknowledges FCT for a contract under the Investigador FCT 2015, Contract No. IF/00758/2015. The research leading to reported results has received funding from the European Research Council under the European Union's Seventh Framework Programme (FP7/2007-2013)/ERC grant agreement no. 337753.

References

1. Rogers RD, Seddon KR. Chemistry. Ionic Liquids - Solvents of the Future? Science. 2003; 302:792–793. [PubMed: 14593156]
2. Freemantle M. Designer Solvents - Ionic Liquids may Boost Clean Technology Development. Chem Eng News. 1998; 76:32–37.
3. Visak ZP. Some Aspects of Ionic Liquids as Diverse and Versatile Sustainable Solvents. J Solution Chem. 2012; 41:1673–1695.
4. Plechkova NV, Seddon KR. Applications of Ionic Liquids in the Chemical Industry. Chem Soc Rev. 2008; 37:123–150. [PubMed: 18197338]
5. Durga, G, Mishra, A. Ionic Liquids: Industrial Applications Encyclopedia of Inorganic and Bioinorganic Chemistry. John Wiley & Sons, Ltd; Chichester, UK: 2016. 1–13.
6. Abraham MH, Zissimos AM, Huddleston JG, Willauer HD, Rogers RD, Acree WE. Some Novel Liquid Partitioning Systems: Water–Ionic Liquids and Aqueous Biphasic Systems. Ind Eng Chem Res. 2003; 42:413–418.
7. Freire MG, Cláudio AFM, Araújo JMM, Coutinho JAP, Marrucho IM, Canongia Lopes JN, Rebelo LPN. Aqueous Biphasic Systems: a Boost Brought About by Using Ionic Liquids. Chem Soc Rev. 2012; 41:4966–4995. [PubMed: 22692625]
8. Gutowski KE, Broker GA, Willauer HD, Huddleston JG, Swatoski RP, Holbrey JD, Rogers RD. Controlling the Aqueous Miscibility of Ionic Liquids: Aqueous Biphasic Systems of Water-Miscible Ionic Liquids and Water-Structuring Salts for Recycle, Metathesis, and Separations. J Am Chem Soc. 2003; 125:6632–6633. [PubMed: 12769563]
9. Pei Y, Wang J, Wu K, Xuan X, Lu X. Ionic Liquid-Based Aqueous Two-Phase Extraction of Selected Proteins. Sep Purif Technol. 2009; 64:288–295.
10. Wellens S, Thijs B, Möller C, Binnemans K. Separation of Cobalt and Nickel by Solvent Extraction with Two Mutually Immiscible Ionic Liquids. Phys Chem Chem Phys. 2013; 15:9663–9669. [PubMed: 23665857]
11. Usui H, Yamamoto Y, Yoshiyama K, Itoh T, Sakaguchi H. Application of Electrolyte Using Novel Ionic Liquid to Si Thick Film Anode of Li-Ion Battery. J Power Sources. 2011; 196:3911–3915.
12. Vander Hoogerstraete T, Binnemans K. Highly Efficient Separation of Rare Earths from Nickel and Cobalt by Solvent Extraction with the Ionic Liquid Trihexyl(tetradecyl)phosphonium Nitrate: a Process Relevant to the Recycling of Rare Earths from Permanent Magnets and Nickel Metal Hydride Batterie. Green Chem. 2014; 16:1594–1606.
13. Wellens S, Goovaerts R, Möller C, Luyten J, Thijs B, Binnemans K. A Continuous Ionic Liquid Extraction Process for the Separation of Cobalt from Nickel. Green Chem. 2013; 15:3160.
14. Freire MG, Neves CMSS, Carvalho PJ, Gardas RL, Fernandes AM, Marrucho IM, Santos LMNBF, Coutinho JAP. Mutual Solubilities of Water and Hydrophobic Ionic Liquids. J Phys Chem B. 2007; 111:13082–13089. [PubMed: 17958353]
15. Freire MG, Neves CMSS, Ventura SPM, Pratas MJ, Marrucho IM, Oliveira J, Coutinho JAP, Fernandes AM. Solubility of Non-Aromatic Ionic Liquids in Water and Correlation Using a QSPR Approach. Fluid Phase Equilib. 2010; 294:234–240.

16. Neves CMSS, Rodrigues AR, Kurnia KA, Esperança JMSS, Freire MG, Coutinho JAP. Solubility of Non-Aromatic Hexafluorophosphate-Based Salts and Ionic Liquids in Water Determined by Electrical Conductivity. *Fluid Phase Equilib.* 2013; 358:50–55.
17. Freire MG, Carvalho PJ, Gardas RL, Marrucho IM, Santos LMNBF, Coutinho JAP. Mutual Solubilities of Water and the $[C_n\text{mim}][\text{Tf}_2\text{N}]$ Hydrophobic Ionic Liquids. *J Phys Chem B.* 2008; 112:1604–1610. [PubMed: 18201080]
18. Freire MG, Neves CMSS, Shimizu K, Bernardes CES, Marrucho IM, Coutinho JAP, Lopes JNC, Rebelo LPN. Mutual Solubility of Water and Structural/Positional Isomers of N-Alkylpyridinium-Based Ionic Liquids. *J Phys Chem B.* 2010; 114:15925–15934. [PubMed: 21077599]
19. Martins MAR, Neves CMSS, Kurnia KA, Luís A, Santos LMNBF, Freire MG, Pinho SP, Coutinho JAP. Impact of the Cation Symmetry on the Mutual Solubilities Between Water and Imidazolium-Based Ionic Liquids. *Fluid Phase Equilib.* 2014; 375:161–167.
20. Freire MG, Santos LMNBF, Fernandes AM, Coutinho JAP, Marrucho IM. An Overview of the Mutual Solubilities of Water-Imidazolium-Based Ionic Liquids Systems. *Fluid Phase Equilib.* 2007; 261:449–454.
21. Martins MAR, Neves CMSS, Kurnia KA, Santos LMNBF, Freire MG, Pinho SP, Coutinho JAP. Analysis of the Isomerism Effect on the Mutual Solubilities of bis(trifluoromethylsulfonyl)imide-Based Ionic Liquids with Water. *Fluid Phase Equilib.* 2014; 381:28–35.
22. Rotrekl J, Storch J, Velišek P, Schröer W, Jacquemin J, Wagner Z, Husson P, Bendová M. Liquid Phase Behavior in Systems of 1-Butyl-3-alkylimidazolium bis{(trifluoromethyl)sulfonyl}imide Ionic Liquids with Water: Influence of the Structure of the C5 Alkyl Substituent. *J Solution Chem.* 2017; 46:1456–1474.
23. Freire MG, Carvalho PJ, Silva AMS, Santos LMNBF, Rebelo LPN, Marrucho IM, Coutinho JAP. Ion Specific Effects on the Mutual Solubilities of Water and Hydrophobic Ionic Liquids. *J Phys Chem B.* 2009; 113:202–211. [PubMed: 19072549]
24. Freire MG, Neves CMSS, Silva AM, Santos LMNBF, Marrucho IM, Rebelo LPN, Shah JK, Maginn EJ, Coutinho JAP. ^1H NMR and Molecular Dynamics Evidence for an Unexpected Interaction on the Origin of Salting-in/Salting-out Phenomena. *J Phys Chem B.* 2010; 114:2004–2014. [PubMed: 20088550]
25. Hofmeister F. “Zur Lehre von der Wirkung der Salze” [Title translation: About the Science of the Effect of Salts]. *Arch Exp Pathol Pharmacol.* 1888; 24:247–260.
26. Shahriari S, Neves CMSS, Freire MG, Coutinho JAP. Role of the Hofmeister Series in the Formation of Ionic-Liquid-Based Aqueous Biphasic Systems. *J Phys Chem B.* 2012; 116:7252–7258. [PubMed: 22594382]
27. Zhao H, Campbell SM, Jackson L, Song Z, Olubajo O. Hofmeister Series of Ionic Liquids: Kosmotropic Effect of Ionic Liquids on the Enzymatic Hydrolysis of Enantiomeric Phenylalanine Methyl Ester. *Tetrahedron: Asymmetry.* 2006; 17:377–383.
28. Yang Z. Hofmeister Effects: an Explanation for the Impact of Ionic Liquids on Biocatalysis. *J Biotechnol.* 2009; 144:12–22. [PubMed: 19409939]
29. Constantinescu D, Weingärtner H, Herrmann C. Protein Denaturation by Ionic Liquids and the Hofmeister Series: A Case Study of Aqueous Solutions of Ribonuclease A. *Angew Chemie Int Ed.* 2007; 46:8887–8889.
30. Andriola A, Singh K, Lewis J, Yu L. Conductivity, Viscosity, and Dissolution Enthalpy of LiNTf_2 in Ionic Liquid BMINTF₂. *J Phys Chem B.* 2010; 114:11709–14. [PubMed: 20735031]
31. Neves CMSS, Held C, Mohammad S, Schleinitz M, Coutinho JAP, Freire MG. Effect of Salts on the Solubility of Ionic Liquids in Water: Experimental and Electrolyte Perturbed-Chain Statistical Associating Fluid Theory. *Phys Chem Chem Phys.* 2015; 17
32. Voet D, Voet JG, Pratt CW. *Fundamentals of Biochemistry.* John Wiley & Sons Inc; New York: 1999.
33. Visak ZP, Canongia Lopes JN, Rebelo LPN. Ionic Liquids in Polyethylene Glycol Aqueous Solutions: Salting-in and Salting-out Effects. *Monatshefte für Chemie - Chem Mon.* 2007; 138:1153–1157.

34. Cláudio AFM, Swift L, Hallett JP, Welton T, Coutinho JAP, Freire MG. Extended Scale for the Hydrogen-Bond Basicity of Ionic Liquids. *Phys Chem Chem Phys*. 2014; 16:6593–6601. [PubMed: 24569531]
35. Kurnia KA, Lima F, Cláudio AFM, Coutinho JAP, Freire MG. Hydrogen-Bond Acidity of Ionic Liquids: an Extended Scale. *Phys Chem Chem Phys*. 2015; 17:18980–18990. [PubMed: 26129926]
36. López-León T, Jódar-Reyes AB, Ortega-Vinuesa JL, Bastos-González D. Hofmeister Effects on the Colloidal Stability of an IgG-Coated Polystyrene Latex. *J Colloid Interface Sci*. 2005; 284:139–148. [PubMed: 15752795]
37. Tomé LIN, Varanda FR, Freire MG, Marrucho IM, Coutinho JAP. Towards an Understanding of the Mutual Solubilities of Water and Hydrophobic Ionic Liquids in the Presence of Salts: The Anion Effect. *J Phys Chem B*. 2009; 113:2815–2825. [PubMed: 19708113]
38. Kunz W, Henle J, Ninham BW. Zur Lehre von der Wirkung der Salze (about the science of the effect of salts): Franz Hofmeister's historical papers. *Curr Opin Colloid Interface Sci*. 2004; 9:19–37.

Fig. 1.
Chemical structures of the ILs and salt used in this work.

Fig. 2.
Scheme of the experimental procedure for sample preparation and quantification used in this work.

Fig. 3.

Solubility, and respective associated uncertainties at 95% of confidence level, of $[C_4C_1im][NTf_2]$ in aqueous solutions at different concentrations of: $\square$, $Li[NTf_2]$; $\triangle$, $[N_{1112}OH][NTf_2]$; $\circ$, $[C_4C_1pyr][NTf_2]$; $\diamond$, $[C_4C_1im][HSO_4]$; $*$, $[C_4C_1im]Cl$, at 298.15 K and atmospheric pressure; $u(T) = 0.1$ K and $u(p) = 0.05$. The lines have no physical meaning are only guides for the eye.