Published in final edited form as:

J Immigr Minor Health. 2019 December ; 21(6): 1200–1207. doi:10.1007/s10903-019-00868-8.

Prevalence and Correlates of Injecting with Visitors from the United States Among People Who Inject Drugs in Tijuana, Mexico

A. Bórquez¹, R. S. Garfein¹, D. Abramovitz¹, L. Liu², L. Beletsky^{1,3}, D. Werb^{1,4}, S. R. Mehta¹, G. Rangel^{5,6}, C. Magis-Rodríguez⁷, P. González-Zúñiga¹, S. A. Strathdee¹

¹Department of Medicine, School of Medicine, University of California San Diego, 9500 Gilman Drive, MC-0725, La Jolla, CA 92093-0725, USA

²Department of Family Medicine and Public Health, University of California San Diego, La Jolla, CA, USA

³School of Law and Bouvé College of Health Sciences, Northeastern University, Boston, MA, USA

⁴Li Ka Shing Knowledge Institute, St. Michael's Hospital, Toronto, Canada

⁵Comisión de Salud Fronteriza México-Estados Unidos, Tijuana, Mexico

⁶El Colegio de la Frontera Norte, Tijuana, Mexico

⁷CENSIDA, Mexico City, Mexico

Abstract

Cross-border infectious disease transmission is a concern related to drug tourism from the U.S. to Mexico. We assessed this risk among people who inject drugs (PWID) in Tijuana, Mexico. We measured the prevalence and identified correlates of injecting with PWID visiting from the U.S. among PWID in Tijuana using univariable and multivariable logistic regression. Of 727 participants, 18.5% injected during the past 6 months in Mexico with U.S. PWID described mostly as friends (63%) or acquaintances (26%). Injecting with U.S. PWID was independently associated with higher education [adjusted odds ratio (aOR) = 1.13/year], deportation from the U.S. (aOR = 1.70), younger age at first injection (aOR = 0.96/year), more lifetime overdoses (aOR = 1.08), and, in the past 6 months, backloading (aOR = 4.00), syringe confiscation by the police (aOR = 3.02) and paying for sex (aOR = 2.98; all p-values < 0.05). Nearly one-fifth of PWID in Tijuana recently injected with U.S. PWID, and their reported risk behaviors could facilitate cross-border disease transmission.

Keywords

HIV; HCV; Injection drug use; Deportation; Drug tourism

*R. S. Garfein rgarfein@ucsd.edu.

A. Bórquez and R.S. Garfein are co-lead authors who both contributed equally to the conceptualization, data analysis, interpretation of results, manuscript drafting and final approval of this manuscript.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Introduction

Whether in the context of labor migration [1], travel [2–5] or unstable housing, [6–8] population mobility is a recognized risk factor for infectious disease transmission through bridging populations, with mobile individuals being exposed to higher infection prevalence or engaging in higher risk behaviors at their destination vs. place of origin (or the converse). In the context of drug use, differences in drug availability, price, quality or legality between settings has been shown to promote travel among people who inject drugs (PWID) [9]. This is particularly relevant on the U.S.-Mexico border, where spillover drugs from trafficking into the U.S. can increase availability and lower prices of drugs in Mexican border cities. San Diego, CA and Tijuana, Mexico share one of the world's busiest international border crossings [10]. Given high prevalence of HIV and hepatitis C virus (HCV) infection among PWID in Tijuana (3.5% [11] and 96% [12]) and in San Diego (3.2% [13] and 66% [14]), this study investigated infection risk and demographic and behavioral patterns among PWID in Tijuana who have crossborder injecting partners.

Among PWID in Tijuana enrolled in a 2006–2007 study [15], 56% reported ever injecting with PWID from the U.S. These participants had higher education, were more likely to speak English, and to have ever spent time in the U.S., been deported from the U.S. or been incarcerated. Exposure to incarceration and law enforcement can exacerbate HIV associated risk behaviors among PWID in Tijuana [16–18]. In this study, injecting risk behaviors and prevalence of HIV, syphilis and tuberculosis infection were not associated with injecting with PWID from the U.S.

Drug use epidemics are shaped by changes in the structural environment. Rhodes's risk environment framework describes how morbidity and mortality associated with drug use are influenced by determinants exogenous to the individual. We assume that changes in drug and immigration policies affecting Mexico and the U.S. influence cross-border injecting patterns and associated risks. Deportation rates through removals (as opposed to voluntary returns) were high during the 2008–2016 U.S. presidential administration [19], which likely increased PWID deportations to Tijuana, potentially facilitating cross-border injecting because U.S. PWID may visit Mexico to inject with deported peers. Additionally, in 2009, Mexico enacted drug law reforms allowing the possession of most drugs in small quantities for personal consumption [20, 21]. Lower legal penalties might have motivated some U.S. PWID to seek drugs in Mexico and affected the nature of cross-border interactions. A cohort study of PWID recruited in San Diego in 2012–2014 found that 18% had injected drugs while in Mexico in the past 6 months; this group shared syringes less often while injecting in Mexico than in the U.S [14]. We hypothesized that injecting in Tijuana with PWID visiting from the U.S. would be associated with deportation, that it would not be associated with HIV risk behaviors and that risk behaviors would decrease when injecting with U.S. PWID.

We used baseline data (2011–2012) from a cohort study of PWID in Tijuana to describe the prevalence of injecting with PWID visiting from the U.S. and identify sociodemographic factors, injecting and sexual behaviors associated with this practice. In addition, we assessed injecting practices and locations specific to times participants injected with PWID from the

U.S. while in Mexico and described their relationships. This information is key to estimating the potential for cross-border outbreaks of infectious diseases and inform the design of interventions.

Methods

Participants

We analyzed baseline data from a cohort study of PWID living in Tijuana, described elsewhere [22]. Eligibility criteria included injecting drugs within the past month, age at least 18 years, able to speak Spanish or English, and current Tijuana residency with no plans to move away in the following 2 years.

Data Collection

Participants were recruited in 2011–2012 using convenience sampling. They completed the baseline assessment in Spanish or English administered by trained interviewers using computer assisted personal interviewing technology in a private setting. Upon completing the interview, participants were tested for HIV infection and received pre- and post-test counseling accompanied by referrals for care as needed. Participants received the equivalent of \$20 USD for completing the baseline assessment. Written informed consent was obtained prior to completing any study procedures. The University of California San Diego Human Research Protection Program and the Institutional Review Board for Xochicalco University approved the study protocol.

Measures

Study interviews captured information on drug use and sexual behaviors, including information on injecting with PWID visiting from the U.S. Specifically, the outcome variable (injected with a PWID from the U.S. in Mexico in the past 6 months—yes vs. no) was constructed based on the participants' responses to the following question: "In the last 6 months, how many different people who live in the U.S. did you inject drugs with in Mexico?"

Potential correlates were explored based on theoretical subject matter considerations and included socio-demographic characteristics, encounters with law enforcement and injecting and sexual risk behaviors. Variables referred to the past 6 months or lifetime as indicated in Table 1.

Socio-demographic variables included age, sex, educational attainment, income level and main source of income, as well as exposures to the U.S. and Mexico, assessed as country of birth (Mexico vs. other), number of years lived in Tijuana, languages spoken, and history of deportation from the U.S. Law enforcement variables included history of arrest and incarceration, syringe confiscation and harassment by the police. Injection risk included age at first injection, duration of injecting, frequency of injecting, types of drugs injected, borrowing (i.e. receptive sharing) and lending (i.e. distributive sharing) used syringes for injecting, and sharing other injection paraphernalia. We also assessed access to sterile syringes, drug overdose history, and treatment for alcohol or drug use. Sexual risks included

having a regular partner and condom use with this partner, number of casual partners, engagement in selling or buying sex, having had sex with an HIV-positive partner and having had sex in the U.S. We conducted descriptive analyses of variables characterizing injecting behaviors specific to times they injected with U.S. PWID in Mexico and defining their relationships with those U.S. PWID based on the following questions: “The next questions refer to the people who live in the U.S. that you injected drugs with in the past 6 months in Mexico. What was your relationship to these people?” and “Where did you meet these people?” (see Table 3 for response options).

Statistical Analyses

Among 734 cohort participants, 727 (99%) had responses for the dependent variable and were included in this analysis. We calculated baseline descriptive statistics for socio-demographic and behavioral factors among participants in our two outcome groups (i.e., those who did and did not inject in Tijuana with PWID visiting from the U.S. within the past 6 months). We presented frequencies and proportions for binary factors and means and standard deviations for continuous factors (Table 1).

Univariable and multivariable logistic regression models with empirical standard error estimation via generalized estimating equations (GEE) were used to identify factors associated with injecting drugs with PWID visiting from the U.S. in Mexico [23, 24]. GEE is commonly used to estimate parameters for a wide variety of generalized linear models (including logistic regression) when correlation between observations might be present. Since our data was collected via convenience sampling (as opposed to random or other type of probability sampling), some observations may be correlated and use of GEE to obtain robust standard errors is warranted. Population averaged estimates for odds ratios along with corresponding 95% confidence intervals and p-values yielded by the univariable GEE analyses are presented in Table 1. Variables that yielded $p \leq 0.10$ in univariable models were considered for inclusion in multivariable models, based on the acceptable cut-off range [25, 26]. To create the final multivariable model, we used a “purposeful selection of variables” approach [27, 28] guided by subject matter significance, relationships among potential correlates (e.g., correlations, confounding, and interactions) and statistical significance. Multi-collinearity was assessed and was ruled out by appropriate values of the variance inflation factors and condition indexes. All two-way interactions between the main correlates were also assessed. A liberal $p \leq 0.10$ significance level was used and the interaction significant at this level was evaluated by calculating the corresponding simple main effects. In the multivariable model, with the exception of interaction terms, $p \leq 0.05$ was interpreted as being statistically significant and $0.05 < p \leq 0.10$ was interpreted as being borderline significant. All statistical analyses were conducted using SAS, version 9.4.

Results

Among 727 PWID in Tijuana, mean age was 37.4 (range 18–63) years; 62% were men; 93% were born in Mexico; and 50% could speak English (Table 1). Overall, 132 (18.5%) had injected in Mexico with a PWID visiting from the U.S. during the past 6 months.

Univariable Analysis of Factors by Injecting with PWID Visiting from the U.S.

Odds of injecting with U.S. PWID were lower among those born in Mexico or having lived their entire lives in Tijuana (Table 1). Odds were higher among those who could speak English, were deported from the U.S., had moved to Tijuana because of U.S. deportation and had higher education and annual income. HIV prevalence was 3.6% overall and did not differ between the groups. All those who tested HIV-positive were previously unaware of their positive sero-status, and thus, were not receiving treatment.

Regarding injection risk, odds of injecting with U.S. PWID were lower among those who initiated injecting in Mexico and higher among those who injected heroin and either cocaine or methamphetamine (polydrug use), engaged in receptive and distributive syringe sharing, divided drugs with a used syringe, shared other injection paraphernalia, reported difficulty obtaining unused syringes for injecting in the past 6 months and those who experienced more lifetime overdoses.

Regarding sexual behavior, odds of injecting with U.S. PWID were higher among those having a regular partner and reporting never using condoms with that partner in the past 6 months, among those having bought sex and those having had sex in the U.S. in the past 6 months; but lower among those identifying as heterosexual. Having a higher number of casual partners and sex with an HIV-positive partner had a borderline significant association with injecting with U.S. PWID.

In terms of law enforcement exposure, participants with more arrests and times spent in jail in the past 6 months had higher odds of injecting with U.S. PWID, as well as those who had syringes confiscated by the police without being arrested and those harassed by the police.

Multivariable Analysis of Factors by Injecting with PWID Visiting from the U.S.

Injecting with PWID from the U.S. was independently associated with higher education [adjusted odds ratio (aOR) = 1.13 per year, $p = 0.001$], having moved to Tijuana because of deportation from the U.S. (aOR = 1.70, $p = 0.02$), younger age at first injection (aOR = 0.96 per year, $p = 0.03$), dividing drugs with someone else with a used syringe in the past 6 months (aOR = 4.00, $p < 0.001$), more lifetime overdose events (aOR = 1.08 per event, $p = 0.05$), having syringes confiscated by the police without being arrested in the past 6 months (aOR = 3.02, $p < 0.001$) and giving money or other commodities in exchange for sex in the past 6 months (aOR = 2.98, $p < 0.001$; Table 2).

A borderline significant ($p = 0.09$) two-way interaction showed that men, but not women, had higher odds of injecting with PWID from the U.S. if they had given money or other commodities in exchange for sex. All variables in Table 2 remained significant in a model that included the interaction term.

Relationship and Behaviors with PWID from the U.S.

PWID from the U.S. were characterized by participants as friends (63%), acquaintances (26%), strangers (7%) or sex trade clients (5%). They mostly reported meeting U.S. PWID in outside locations (47%) or at their or someone else's home (23%) (Table 3). During injecting events with PWID from the U.S., most reported distributive (67%) or receptive

(62%) syringe sharing and sharing other injection paraphernalia (70%) in the past 6 months (Table 3); however, prevalence of any distributive and receptive syringe sharing among PWID who injected with U.S. PWID was higher overall (88% and 90%, respectively, in the past 6 months) than while injecting with U.S. PWID (Table 1).

Discussion

Nearly one-fifth of PWID who reside in Tijuana reported injecting with PWID visiting from the U.S. in the past 6 months. While previous studies investigated injecting drug use with U.S. PWID among Tijuana PWID, this is the first to characterize these interactions in a defined time period (i.e. past 6 months) and location (i.e. in Mexico). As found by Wagner et al., compared to PWID who did not inject with U.S. PWID, those who did were more likely to be male, more educated and to have been deported from the U.S. However, contrary to previous findings, they were also more likely to report HIV-associated risk factors including backloading drugs for injection, higher numbers of overdoses, paying for sex, and having syringes confiscated by the police. This study also contributes new information regarding the nature of relationships between PWID from Tijuana and San Diego and their injecting risk behaviors during these interactions. U.S. visitors with whom they injected were mainly considered friends and acquaintances, but some were strangers or sex trade clients and rates of distributive and receptive syringe sharing with U.S. PWID were high.

Being deported from the U.S. and higher education were independently associated with injecting with U.S. PWID. In bivariate analyses, speaking English (which was correlated with education) and having started injecting in the U.S. were also associated with injecting with U.S. PWID. These are all potential indicators of having stronger links to the U.S. facilitating relationships with PWID visiting from the U.S. Quantitative and qualitative studies among PWID in Tijuana showed associations between deportation and high risk drug using behaviors, including trying new drugs [29] and greater syringe sharing [30].

The high prevalence of syringe sharing during encounters with U.S. PWID is worrying as it suggests potential for cross-border disease transmission. Reported prevalence of syringe sharing was higher than those reported in the parallel study in San Diego [14] among U.S. PWID who injected in Mexico (62% vs 35% for receptive syringe sharing), although differences in study design could account for some variation. Nonetheless, as was found among PWID in San Diego, fewer injecting risk behaviors were reported when injecting with someone visiting from the U.S. compared to their behaviors overall, suggesting safer injecting practices during these encounters. It is possible that PWID from the U.S. perceive higher risk when injecting in Mexico and practice safer injecting, leading to lower risk behaviors among PWID in Tijuana. If true, this finding suggests that PWID can reduce their risk in specific circumstances and supports the potential for change through peer-based network-based interventions. Indeed, the majority of U.S. PWID were considered friends, although one-third were acquaintances, strangers or sex trade clients suggesting that the HIV/HCV infection status of many injection partners from the U.S. was unknown. Education and prevention messages among PWID in the U.S. about the increased risk of exposure to HIV and HCV in Tijuana resulting from the lower access to HIV [31, 32] and harm reduction [33, 34] services could motivate them to further protect themselves and

PWID in Tijuana. Qualitative research regarding these relationships and the motivations and circumstances of these encounters would inform the design of interventions.

PWID who paid for sex in the past 6 months were more likely to have injected with PWID visiting from the U.S. in the same period. Female sex workers in Tijuana have elevated prevalence of HIV (4.8%), active syphilis (11.8%) and chlamydia infection (17%) [35]. In bivariate analyses, unprotected sex with a regular partner, sex with someone while in the U.S. and homosexual orientation were significantly associated with injecting with U.S. PWID. These findings suggest that PWID in Tijuana who inject with U.S. PWID could bridge multiple groups at increased risk for sexual and bloodborne infections.

Having syringes confiscated by the police without being arrested in the past 6 months was also independently associated with injecting with U.S. PWID. In bivariate analyses, this group also had a higher arrest frequency and higher exposure to police harassment and incarceration, which are associated with greater injecting and sexual behaviors among this population [16–18, 36]. Higher law enforcement exposure might be due to injecting in public spaces more often, which prior studies have shown to be associated with high injecting risk profiles [37, 38]. Indeed, PWID reported mostly injecting with U.S. PWID in outside locations, including in the streets and in the Tijuana river canal, where many deported PWID congregate. This suggests mobile, street-based interventions may be most effective at preventing risk during cross-border encounters.

Importantly, overdose was prevalent among PWID who injected with U.S. PWID, although the association was borderline significant. This is consistent with the higher risk profile exhibited by this group, including engagement in polydrug use, and emphasizes the need to prioritize this population with harm reduction services, including naloxone.

Limitations

This study relied on self-reported behaviors that are subject to problems with recall, lack of precision and socially desirable responding. To minimize these problems, the survey was administered by highly trained interviewers skilled at eliciting accurate responses by establishing respectful, non-judgemental rapport. Some participants might not have known that their injection partners were from the U.S.; therefore, our findings could underestimate the prevalence of injecting with PWID visiting from the U.S. This study involved a cross-sectional analysis of baseline data, which precludes drawing causal inferences. Further, this sample of PWID might not be representative of all PWID in Tijuana given the hidden nature of this population.

Conclusions

Our study supports prior research showing that ‘drug tourism’ among U.S. PWID in the border region is common. A history of deportation was independently associated with injecting with PWID from the U.S.; thus, our findings challenge U.S. federal government proposals to increase deportation, because such policies could paradoxically contribute to HIV and HCV transmission. Indeed, deportation and crossborder injecting were associated with a range of high risk behaviors and exposures that could facilitate cross-border disease

transmission. However, our study also found that PWID in Tijuana reduced their HIV risk behaviors during encounters with U.S. PWID, suggesting that unilateral and/ or binational interventions could reduce risk on both sides of the border.

Acknowledgements

The authors wish to thank the study participants for sharing their detailed histories, without which this research would not have been possible. We also wish to thank the Tijuana and San Diego research staff members whose compassion for the study population and dedication to the research objectives helped to acquire high quality data for this analysis.

Funding This study was funded by the National Institutes of Health (R37 DA019829). RSG also received funding from the National Institutes for Health (R01 DA031074).

References

1. Weine SM, Kashuba AB. Labor migration and HIV risk: a systematic review of the literature. *AIDS Behav.* 2012;16 (6):1605–21. [PubMed: 22481273]
2. Delany-Moretlwe S, Bello B, Kinross P, Oliff M, Chersich M, Kleinschmidt I, et al. HIV prevalence and risk in long-distance truck drivers in South Africa: a national cross-sectional survey. *Int J Std AIDS.* 2014;25 (6):428–38. [PubMed: 24352131]
3. Garcia PJ, Fazio B, Bayer AM, Lizarraga AG, Chiappe M, La Rosa S, et al. Sexual health knowledge and practices and STI/ HIV prevalence among long-distance truck drivers in Peru. *SAGE Open Med.* 2017;5:2050312117746308.
4. Lacerda R, Gravato N, McFarland W, Rutherford G, Iskrant K, Stall R, et al. Truck drivers in Brazil: prevalence of HIV and other sexually transmitted diseases, risk behavior and potential for spread of infection. *AIDS.* 1997;11 (Suppl 1):15–9.
5. Lee D, Bell DC, Hinojosa M. Drug use, travel and HIV risk. *AIDS Care.* 2002;14 (4):443–53. [PubMed: 12204148]
6. Marshall BD, Kerr T, Shoveller JA, Patterson TL, Buxton JA, Wood E. Homelessness and unstable housing associated with an increased risk of HIV and STI transmission among street-involved youth. *Health Place.* 2009;15 (3):753–60. [PubMed: 19201642]
7. Elifson KW, Sterk CE, Theall KP. Safe living: the impact of unstable housing conditions on HIV risk reduction among female drug users. *AIDS Behav.* 2007;11 (6 Suppl):45–55. [PubMed: 17828451]
8. Corneil TA, Kuyper LM, Shoveller J, Hogg RS, Li K, Spittal PM, et al. Unstable housing, associated risk behaviour, and increased risk for HIV infection among injection drug users. *Health Place.* 2006;12 (1):79–85. [PubMed: 16243682]
9. Rachlis B, Brouwer KC, Mills EJ, Hayes M, Kerr T, Hogg RS. Migration and transmission of blood-borne infections among injection drug users: understanding the epidemiologic bridge. *Drug Alcohol Depend.* 2007;90 (2–3):107–19. [PubMed: 17485179]
10. U.S. General Services Administration. The San Ysidro Land Port of Entry Project Facts. 2018 <https://www.gsa.gov/about-us/regions/welcome-to-the-pacific-rim-region-9/land-ports-of-entry/san-ysidro-land-port-of-entry/san-ysidro-lpoe-project-facts>. Accessed 24 Sept 2018.
11. Steffanie Strathdee PI. Proyecto el Cuete phase IV. NIDA grant R01 DA0 19829; 2010–2020.
12. White EF, Garfein RS, Brouwer KC, Lozada R, Ramos R, Firestone-Cruz M, et al. Prevalence of hepatitis C virus and HIV infection among injection drug users in two Mexican cities bordering the U.S. *Salud Publica Mex.* 2007;49 (3):165–72. [PubMed: 17589770]
13. Centers for Disease Control and Prevention. Infection HIV, Prevention R and testing behaviors among persons who inject drugs—National HIV Behavioral Surveillance: Injection Drug Use, 20 U.S. Cities. 2015 <https://www.cdc.gov/hiv/pdf/library/reports/surveillance/cdc-hiv-hssr-nhbs-pwid-2015.pdf>. Accessed 19 Sept 2018.

14. Horyniak D, Wagner KD, Armenta RF, Cuevas-Mota J, Hendrickson E, Garfein RS. Cross-border injection drug use and HIV and hepatitis C virus seropositivity among people who inject drugs in San Diego, California. *Int J Drug Policy*. 2017;47:9–17. [PubMed: 28683432]
15. Wagner KD, Pollini RA, Patterson TL, Lozada R, Ojeda VD, Brouwer KC, et al. Cross-border drug injection relationships among injection drug users in Tijuana, Mexico. *Drug Alcohol Depend*. 2011;113 (2–3):236–41. [PubMed: 20889270]
16. Pinedo M, Burgos JL, Zuniga ML, Perez R, Macera CA, Ojeda VD. Police victimization among persons who inject drugs along the U.S.–Mexico border. *J Stud Alcohol Drugs*. 2015;76 (5):758–63. [PubMed: 26402356]
17. Pollini RA, Brouwer KC, Lozada RM, Ramos R, Cruz MF, Magis-Rodriguez C, et al. Syringe possession arrests are associated with receptive syringe sharing in two Mexico-US border cities. *Addiction*. 2008;103 (1):101–8. [PubMed: 18028520]
18. Beletsky L, Lozada R, Gaines T, Abramovitz D, Staines H, Vera A, et al. Syringe confiscation as an HIV risk factor: the public health implications of arbitrary policing in Tijuana and Ciudad Juarez. *Mexico J Urban Health*. 2013;90 (2):284–98. [PubMed: 22806453]
19. U.S. Department of Homeland Security. Yearbook of immigration statistics, aliens removed or returned: fiscal years 1892 to 2016. 2017. <https://www.dhs.gov/immigration-statistics/yearbook/2016/table39>. Accessed 9 Sept 2018.
20. DECRETO por el que se reforman. Adicionan y derogan diversas disposiciones de la Ley General de Salud, del Código Penal Federal y del Código Federal de Procedimientos Penales. Diario oficial de la federación. 2009.
21. Mackey TK, Werb D, Beletsky L, Rangel G, Arredondo J, Strathdee SA. Mexico's "ley de narcomenudeo" drug policy reform and the international drug control regime. *Harm Reduct J*. 2014;11 (1):31. [PubMed: 25395346]
22. Robertson AM, Garfein RS, Wagner KD, Mehta SR, Magis-Rodriguez C, Cuevas-Mota J, et al. Evaluating the impact of Mexico's drug policy reforms on people who inject drugs in Tijuana, B.C., Mexico, and San Diego, CA, United States: a binational mixed methods research agenda. *Harm Reduct J*. 2014;11:4. [PubMed: 24520885]
23. Liang KY, Zeger S. Longitudinal data analysis using generalized linear models. *Biometrika*. 1986;73 (1):13–22.
24. Hardin J, Hilbe J. Generalized estimating equations. London: Chapman and Hall/CRC; 2003.
25. Bendal RB, Afifi AA. Comparison of stopping rules in forward regression. *J Am Stat Assoc*. 1977;72:46–53.
26. Mickey RM, Greenland S. The impact of confounder selection criteria on effect estimation. *Am J Epidemiol*. 1989;129 (1):125–37. [PubMed: 2910056]
27. Hosmer DW, Lemeshow S. Applied logistic regression. New York: Wiley; 2000.
28. Hosmer DW, Lemeshow S. Applied survival analysis: regression modeling of time to event data. New York: Wiley; 1999.
29. Robertson AM, Rangel MG, Lozada R, Vera A, Ojeda VD. Male injection drug users try new drugs following U.S. deportation to Tijuana, Mexico. *Drug Alcohol Depend*. 2012;120 (1–3):142–8. [PubMed: 21835559]
30. Ojeda VD, Robertson AM, Hiller SP, Lozada R, Cornelius W, Palinkas LA, et al. A qualitative view of drug use behaviors of Mexican male injection drug users deported from the United States. *J Urban Health*. 2011;88 (1):104–17. [PubMed: 21246301]
31. Moyer LB, Brouwer KC, Brodine SK, Ramos R, Lozada R, Cruz MF, et al. Barriers and missed opportunities to HIV testing among injection drug users in two Mexico–US border cities. *Drug Alcohol Rev*. 2008;27 (1):39–45. [PubMed: 18034380]
32. Smith LR, Patterson TL, Magis-Rodriguez C, Ojeda VD, Burgos JL, Rojas SA, et al. Engagement in the HIV care continuum among key populations in Tijuana, Mexico. *AIDS Behav*. 2016;20 (5):1017–25. [PubMed: 26354518]
33. Werb D, Wagner KD, Beletsky L, Gonzalez-Zuniga P, Rangel G, Strathdee SA. Police bribery and access to methadone maintenance therapy within the context of drug policy reform in Tijuana, Mexico. *Drug Alcohol Depend*. 2015;148:221–5. [PubMed: 25655577]

34. Smith DM, Werb D, Abramovitz D, Magis-Rodriguez C, Vera A, Patterson TL, et al. Predictors of needle exchange program utilization during its implementation and expansion in Tijuana, Mexico. *Am J Addict.* 2016;25 (2):118–24. [PubMed: 26765073]
35. Patterson TL, Semple SJ, Fraga M, Bucardo J, de la Torre A, Salazar J, et al. Comparison of sexual and drug use behaviors between female sex workers in Tijuana and Ciudad Juarez, Mexico. *Subst Use Misuse.* 2006;41 (10–12):1535–49. [PubMed: 17002992]
36. Pollini RA, Alvelais J, Gallardo M, Vera A, Lozada R, Magis Rodriguez C, et al. The harm inside: injection during incarceration among male injection drug users in Tijuana, Mexico. *Drug Alcohol Depend.* 2009;103 (1–2):52–8. [PubMed: 19386448]
37. Mazhnaya A, Tobin KE, Owczarzak J. Association between injection in public places and HIV/HCV risk behavior among people who use drugs in Ukraine. *Drug Alcohol Depend.* 2018;189:125–30. [PubMed: 29913323]
38. Ickowicz S, Wood E, Dong H, Nguyen P, Small W, Kerr T, et al. Association between public injecting and drug-related harm among HIV-positive people who use injection drugs in a Canadian setting: a longitudinal analysis. *Drug Alcohol Depend.* 2017;180:33–8. [PubMed: 28865390]

Table 1

Descriptive statistics and univariable associations between sociodemographic, behavioral and law enforcement factors and injection drug use in the past 6 months in Mexico with people visiting from the United States

Variables	<u>Injected in Mexico with U.S. PWID</u>		Odds ratio	95% confidence interval	p-value
	Yes (n = 132)	No (n = 595)			
<i>Socio-demographic characteristics</i>					
Age, mean (SD)	38.3 (10.0)	37.2 (8.7)	1.01	(0.99, 1.04)	0.280
Male gender (reference: female), N (%)	89 (67.4)	364 (61.2)	1.31	(0.88, 1.96)	0.181
Country of birth: Mexico vs. other, N (%)	118 (89.4)	558 (93.8)	0.56	(0.29, 1.07)	0.078
Lived in Tijuana entire life ^a , N (%)	26 (19.7)	168 (28.2)	0.62	(0.39, 0.99)	0.046
Number of years lived in Tijuana, mean (SD)	19.5 (14.2)	23.3 (14.0)	0.98	(0.97, 0.99)	0.008
Ever lived in the U.S., N (%)	103 (24.0)	29 (9.8)	2.91	(1.87, 4.53)	< 0.001
Able to speak English, N (%) ^a	97 (73.5)	250 (42.0)	3.82	(2.51, 5.82)	< 0.001
Ever deported from the U.S., N (%) ^a	70 (53.0)	227 (38.2)	1.83	(1.25, 2.68)	0.002
Moved to Tijuana due to deportation from the U.S., N (%) ^a	45 (34.4)	119 (20.0)	2.09	(1.38, 3.16)	< 0.001
Number of years of education, mean (SD)	8.6 (3.3)	7.6 (3.2)	1.11	(1.05, 1.19)	< 0.001
Average monthly income: 2500 pesos vs. < 2500 pesos, N (%)	90 (68.2)	268 (45.3)	2.58	(1.73, 3.85)	< 0.001
Principal source of income: selling drugs, N (%)	3 (2.3)	3 (0.5)	4.59	(0.92, 23.00)	0.064
Principal source of income: prostitution/sex work, N (%)	27 (20.5)	112 (18.8)	1.11	(0.69, 1.77)	0.666
Tested positive for HIV (at baseline) ^a , N (%)	5 (3.8)	21 (3.5)	1.08	(0.40, 2.91)	0.885
<i>Injecting risk</i>					
Age when first injected, mean (SD)	19.5 (5.8)	21.3 (6.8)	0.95	(0.92, 0.99)	0.006
Duration (years) of injection, mean (SD)	18.7 (10.6)	15.9 (9.4)	1.03	(1.01, 1.05)	0.004
First injected in Mexico ^a , N (%)	81 (61.4)	448 (75.3)	0.52	(0.35, 0.77)	0.001
Injected once per day, N (%) ^a	125 (94.7)	564 (95.1)	0.92	(0.39, 2.14)	0.844
Injected heroin, N (%) ^a	126 (95.5)	567 (95.3)	1.04	(0.42, 2.56)	0.937
Injected cocaine, N (%) ^a	18 (13.6)	37 (6.2)	2.38	(1.31, 4.32)	0.005
Injected methamphetamine, N (%) ^a	56 (42.4)	150 (25.3)	2.18	(1.47, 3.23)	< 0.001
Any distributive needle sharing, N (%) ^a	116 (87.9)	410 (68.9)	3.27	(1.89, 5.67)	< 0.001
Any receptive needle sharing, N (%) ^a	119 (90.2)	400 (67.2)	4.46	(2.46, 8.11)	< 0.001
Divided drugs with someone else by using a syringe (backloading), N (%) ^a	111 (84.7)	328 (55.2)	4.50	(2.72, 7.44)	< 0.001
Used a cooker/cotton/water with or after someone else, N (%) ^a	112 (85.5)	373 (62.8)	3.49	(2.09, 5.84)	< 0.001
Got syringes to inject drugs most often from pharmacist, N (%) ^a	44 (33.3)	257 (43.2)	0.66	(0.44, 0.98)	0.038

Variables	<u>Injected in Mexico with U.S. PWID</u>		Odds ratio	95% confidence interval	p-value
	Yes (n = 132)	No (n = 595)			
Found it hard to get new, unused syringes for injecting drugs, N (%) ^a	36 (27.3)	98 (16.5)	1.90	(1.22, 2.95)	0.004
Got syringes from a syringe exchange program, N (%) ^a	18 (13.6)	40 (6.7)	2.19	(1.21, 3.96)	0.009
Number of times overdosed ever, mean (SD)	2.3 (2.7)	1.6 (2.5)	1.10	(1.03, 1.17)	0.003
Ever received professional help for drug or alcohol use ^a , N (%)	83 (62.9)	330 (55.5)	1.36	(0.92, 2.01)	0.120
<i>Sexual risk</i>					
Sexual orientation: heterosexual vs. homosexual/bisexual/other, N (%)	117 (88.6)	557 (93.6)	0.53	(0.28, 1.00)	0.050
Had a regular sex partner, N (%) ^a	85 (64.4)	305 (51.3)	1.71	(1.16, 2.53)	0.007
Never used a condom with regular partner, N (%) ^a	72 (54.5)	240 (40.5)	1.76	(1.21, 2.58)	0.003
Number of casual sex partners, mean (SD) [*]	8.4 (45.5)	3.9 (16.6)	1.01	(1.00, 1.01)	0.055
Received money/goods/etc in exchange for sex, N (%) ^a	49 (37.1)	173 (29.4)	1.42	(0.96, 2.11)	0.082
Gave money/goods/etc in exchange for sex, N (%) ^a	24 (18.2)	37 (6.2)	3.35	(1.92, 5.82)	< 0.001
Ever had sex with HIV infected person, N (%) ^a	7 (5.3)	13 (2.2)	2.50	(0.98, 6.40)	0.055
Had sex in the US (past year), N (%) ^a	18 (13.6)	19 (3.2)	4.77	(2.43, 9.37)	< 0.001
<i>Encounters with law enforcement</i>					
Arrested, N (%) ^a	71 (54.2)	299 (50.3)	1.17	(0.80, 1.71)	0.414
Number of times arrested, mean (SD) [*]	6.9 (20.4)	3.8 (8.6)	1.02	(1.01, 1.03)	0.005
Spent time in jail, N (%) ^a	65 (49.2)	214 (36.0)	1.73	(1.18, 2.53)	0.005
Had syringes confiscated by police without arrest, N (%) ^a	28 (21.2)	54 (9.1)	2.70	(1.63, 4.46)	< 0.001
Harassed by police, N (%) ^a	105 (79.5)	383 (64.4)	2.15	(1.37, 3.39)	0.001

^{*} Reference period was the past 6 months, bold: significant at p = 0.05 level, odds ratios are from univariable logistic models fitted via GEE

^a Yes vs. no

Table 2

Factors independently associated with injection drug use in the past 6 months in Mexico with people visiting from the United States

Variable	Adjusted odds ratio	95% confidence interval		p-value
Age at first injection (per year)	0.96	0.93	0.99	0.027
Male gender (reference: female)	1.21	0.77	1.90	0.414
Number of years of education (per year)	1.13	1.05	1.21	0.001
Moved to Tijuana due to deportation from the U.S.	1.70	1.08	2.68	0.022
Divided drugs with someone else by using a syringe (backloading) *	4.00	2.33	6.88	< 0.001
Had syringes confiscated by police without being arrested *	3.02	1.73	5.28	< 0.001
Number of times overdosed in lifetime (per event)	1.08	1.00	1.18	0.049
Gave money/goods/etc. in exchange for sex *	2.98	1.69	5.24	< 0.001

* Reference period was the past 6 months. Adjusted odds ratios are from multivariable logistic models fitted via GEE

Table 3

Characteristics of injecting events with PWID from the United States among participants who reported injecting in the past 6 months in Mexico with people visiting from the United States

Variable	Percent
<i>Injecting risk past 6 months (yes vs. no) (N= 132)</i>	
Any distributive needle sharing	67
Any receptive needle sharing	62
Ever used a cooker/cotton/water with or after someone else	70
<i>Participant's relationship with PWID from the U.S.: (N= 132)*</i>	
Family member	3
Friend	63
Sex partner (not from sex trade)	3
Client (paying partner)	5
Acquaintance	26
Stranger	7
Other	2
<i>Place (s) participant met PWID from the U.S.: (N= 132)*</i>	
Home	11
Someone else's home	12
Shooting gallery	11
Outside location (construction site, alleyway, on the street, vacant lot, park, freeway overpass/bridge, canal)	47
Bar/club	12
Brothel/strip club	1
Public restroom	0
Other	19

PWID people who inject drugs

* Participants could check multiple responses