Published in final edited form as:

Obes Res Clin Pract. 2019 ; 13(4): 395–397. doi:10.1016/j.orcp.2019.05.001.

Do the Holidays Impact Weight and Self-Weighing Behavior Among Adults Engaged in a Behavioral Weight Loss Intervention?

Margaret C. Fahey, MA^a, Robert C. Klesges, PhD^{b,c}, Mehmet Kocak, PhD^b, Jiajing Wang^b, Gerald W. Talcott, PhD^{b,c}, Rebecca A. Krukowski, PhD^b

^aThe University of Memphis, Department of Psychology, 400 Innovation Drive Memphis, TN, USA, 38111

^bDepartment of Preventive Medicine, University of Tennessee Health Science Center, 66 N Pauline Street Memphis, TN, USA, 38105

^cUniversity of Virginia, Department of Public Health Sciences, School of Medicine, 1215 Lee Street, Charlottesville, VA, USA, 22908

Adults in the United States (U.S.) typically gain weight [4–9 kilograms (kg)], during the holiday period (i.e., mid-November–January) [1–5]. This weight gain is often maintained throughout the year [3, 5], perhaps, contributing to the rising prevalence of overweight [body mass index (BMI) > 25.0 kg/m²] and obesity (> 30.0 kg/m²) [6]. Weight gain is attributed to changes in weight-related behaviors (e.g., self-weighing) [3,7], which may be due to less physical activity in colder temperatures and holiday-based eating.

Previous weight maintenance interventions (i.e., conjugated linoleic acid supplementation, group support, cognitive-behavioral treatment focused on self-monitoring) were successful in preventing weight gain during the holiday period [1,7, 8], although one self-monitoring-focused intervention found that only participants with the greatest adherence did not gain weight during this time [9]. However, it is unknown how the holidays influence weight change in a weight loss seeking population, self-weighing frequency, and as well as how timing of this period within an intervention impacts long-term success [1, 7–9].

To our knowledge, only one study used electronic scales (e-scales) to facilitate more frequent measurements [10,11] and only in the general population [10,11], not in a weight loss seeking population. The current study aims to use e-scale measurements to examine

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Ethical approval.

All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards.

Conflict of Interest:

Margaret C. Fahey declared no conflicts of interest. Robert C. Klesges declared no conflicts of interest. Mehmet Kocak declared no conflict of interests. Jiajing Wang declared no conflicts of interest. Gerald W. Talcott declared no conflicts of interest. Rebecca A. Krukowski declared no conflicts of interest.

changes in weight and self-weighing frequency over the holiday period among military personnel enrolled in a weight loss intervention. This study will also examine the influence of enrollment and completion during the holiday period on treatment outcomes. This sample may have increased motivation to manage their weight due to military fitness and appearance standards [12].

Methods

Sample

Active duty military personnel [>18 years of age, BMI > 25.9 kg/m²] who expected to be stationed at least one year at Joint Base San Antonio, TX were enrolled. There was an equal distribution of gender and BMI category, and diversity with race (19.9% African American, 14.0% Other races) and ethnicity (18.7% Hispanic/Latino). Participant characteristics and main outcomes are reported elsewhere [13].

Design

Protocol was approved by the IRB of the 59th Medical Wing in San Antonio, TX and acknowledged by IRB at the University of Tennessee Health Science Center. After obtaining informed consent, one-week of self-monitoring (i.e., diet, physical activity), and healthcare provider approval, participants were randomized to a counselor-initiated (CI) or self-paced (SP) weight loss intervention (based on the Look AHEAD intensive lifestyle intervention) [14]. Personalized weight loss, calorie, and exercise goals were the same between conditions. At baseline visit, participants were given BodyTrace™ e-scale and encouraged to self-weigh daily. Weight was uploaded to personalized webpages using automatic cellular transmission, accessible by participants and counselors. Conditions differed in amount of initiation needed for receiving intervention components. In the CI condition, counselors provided 16 weekly, then 8 every other week, and then 4 monthly email feedback messages focused on self-monitoring and weight loss progress. The SP condition could request the same number of feedback messages [15].

Measures

Participants reported demographics and height was collected using a wall-mounted stadiometer at baseline by study staff. At baseline and 12-months, participants were weighed in kg on a calibrated scale (Tanita BWB-800S). BMI was calculated from baseline height and weight. Body weights were collected from e-scale measurements. For the 12-month follow-up, if in-clinic weight was missing, e-scale weight closest ($\pm$ 30 days) to 12-months was used [16,17]. Missing weights ($n = 42$) were then imputed as baseline weight carried forward.

Analysis

Mean (%) weight change, rate of weekly weight change, and mean number of times participants self-weighed, were compared from pre-holiday (October 1st - November 14th), to the holiday (November 15th - January 1st), and post-holiday (January 2nd - February 14th) periods overall and by participant demographics, condition, and days from randomization (< 90, 91–180, 181–270, and > 270 days) using Wilcoxon-Mann Whitney and Kruskal-Wallis

tests. Dates were consistent with previous studies [3, 5]. Annual quarterly time periods were used to compare time from randomization. To be included in rate of weight change analyses, individuals need to self-weigh more than two times at least 14 days apart in each period.

Logistic regressions examined the relationship between enrollment during holiday (versus other times) and 12-month weight change (i.e., 2.3% and 5%). Logistic regressions examined difference between completing final study visit during holiday (versus other times) in relationship to 12-month weight change. Criteria were based on standard definitions (i.e., < 2.3% maintenance, 5% loss) [18].

Results

Most self-weighed at least weekly during the pre-holiday, holiday, and post-holiday periods (i.e., 89.57%, 81.82%, 83.63%, respectively). From the pre-holiday to the holiday period, there was not significant weight change overall (i.e., $M = + 0.18 (+ 2.16) \text{ kg}$, $p = 0.30$; $+ 0.22\%$, $p = .30$) (Table 1), but this differed by days from randomization ($p < 0.0001$). Those enrolled within 90 days of the holidays (i.e., mid-August - mid-November) lost on average $- 0.98\%$ ($- 0.86 \text{ kg}$). Those enrolled > 90 days before holiday (i.e., before mid-August) gained weight ($M = + 0.25\%$ to $+ 0.93\%$). Rate of weekly weight change differed from the pre-holiday to the holiday period overall [$M = + 0.17 (0.45) \text{ kg}$; $p = 0.001$] and differed by race ($p = 0.04$) and days from randomization ($p = 0.026$) (Table 1).

Participants self-weighed less frequently ($M = -1.09$, $p = 0.01$) during the holiday than the pre-holiday period (Table 1), which differed by days from randomization ($p = 0.006$). Those enrolled within 90 days of the holidays self-weighed, on average, 5.18 times *more* frequently during the holidays than the pre-holiday period. Participants enrolled > 90 days before the holidays (i.e., before mid-August) self-weighed *less* ($M = -1.72$ to -5.00 times).

From the holiday to the post-holiday periods, there was not significant weight change overall ($M = + 0.10 (1.95) \text{ kg}$, $p = 0.41$; $+ 0.12\%$; $p = 0.37$), which differed by days from randomization ($p < 0.0001$) (Table 2). Those enrolled within 90 days of the post-holiday period lost $- 0.85\%$ ($- 0.68 \text{ kg}$) of weight. Those enrolled > 90 days before the post-holiday period gained weight ($M = +0.35$ to $+ 0.60\%$). Rate of weekly weight change differed from the holiday to post-holiday periods overall [$M = - 0.11 (0.41) \text{ kg}$, $p = 0.007$] and differed by race ($p = 0.01$) and gender ($p = 0.04$) (Table 2).

Controlling for condition, those enrolled during the holiday period were more likely to lose 2.3% of baseline weight at 12-months compared to those enrolled during other times (odds ratio: 2.49, 95% confidence interval: 1.03 – 6.27; $p = 0.04$). There was not a difference in enrollment periods for 5% weight loss at 12-months ($p = 0.55$). Adjusting for condition, there was no difference in achieving 2.3% or 5% weight change between those who completed their final 12-month visit within holiday period versus other times.

Discussion

Participants, on average, did not experience significant percent or total weight change between the pre-holiday, holiday, and post-holiday periods. However, they gained weight

more rapidly during the holidays compared to other times. Participants, perhaps due to increased weight loss motivation in this population, gained less weight overall (+ 0.18 kg) [1], and gained weight *less* rapidly compared to previous samples [7–9]. Despite differences in samples and protocols, results extend previous findings that intervention engagement prevents gains during the holidays [1, 8]. Importantly, the longer participants had engaged in the intervention, the more weight they gained and the less frequently they self-weighed during this time. As noted previously [7,9], interventionists might encourage more frequent self-monitoring behavior during holidays, particularly for those who enroll in August or earlier in the year.

Importantly for future trials, holiday period enrollment did not negatively impact weight loss success and perhaps facilitated improved outcomes. Because early self-monitoring adherence and weight loss is associated with long-term weight maintenance [19], “braving” the holiday period at the beginning of treatment, when participants are the most motivated, may increase self-efficacy. Despite potential time constraints (e.g., scheduling), enrollment during the holidays may increase long-term self-monitoring, or perhaps, participants who enroll during the holidays are most motivated to make behavioral changes. Surprisingly, when self-monitoring adherence is typically lowest [20], completing the intervention within the holiday period did not impact outcomes. Further, researchers should examine both percent weight change and rate of weight changes when observing the holiday impact on weight, to facilitate comparability between studies. Qualitative research also would help to understand how holidays influence weight and self-weighing.

These participants were active duty military personnel, perhaps limiting generalizability due to differences in holiday practices (e.g., stationed away from extended family) or in body composition. However, prevalence of overweight and obesity in the U.S. military is high [21], and this sample was more diverse in gender, race, and ethnicity [1,7–9]. Enrollment period comparisons had small samples and were collected from one site; thus, findings should be replicated.

Conclusion

Using e-scale measurements, this study captured changes in weight and self-weighing behavior among weight-loss-seeking adults throughout the U.S. holiday period. More recent intervention enrollment promoted more self-weighing behavior and resulted in better weight management during this time. Results suggest weight loss intervention enrollment prior to and during the holiday period does not negatively impact, and might improve, treatment success.

References

1. Andersson I, Rossner S The Christmas factor in obesity research. *International J Obes* 1991; 16 (12): 1013–1015.
2. Cook CM, Subar AF, Troiano RP, Schoeller DA. (2012). Relation between holiday weight gain and total energy expenditure among 40- to 69-y-old men and women (OPEN study). *Am J of Clin Nutr* 2012; 95(3): 726–731. [PubMed: 22301936]
3. Schoeller DA. The effect of holiday weight gain on body weight. *Physiol Behav* 2014; 134:66–69. [PubMed: 24662697]

4. Stevenson JL, Krishnan S, Stoner MA, Goktas Z, & Cooper JA. (2013). Effects of exercise during the holiday season on changes in body weight, body composition and blood pressure. *Eur J of Clin Nutr* 2013; 67(9): 944–949. [PubMed: 23695203]
5. Yanovski JA, Yanovski SZ, Nguyen TT, O'Neil PM, Sebring NG. A prospective study of holiday weight gain. *N Eng J Med* 2000; 23 (12): 861–867.
6. Ogden CL, Fryar CD, Carroll MD, Flegal KM. Mean body weight, height, and body mass index, United States 1960–2002. Advance data from vital and health statistics; no 347 Hyattsville, Maryland: National Center for Health Statistics 2004.
7. Boutelle KN, Kirschenbaum DS, Baker RC, Mitchell ME. How can obese weight controllers minimize weight gain during the high-risk holiday season? By self-monitoring very consistently. *Health Psych* 1999; 18 (4): 364–368.
8. Watras AC, Buchholz AC, Close RN, Zhang Z, Schoeller DA. The role of conjugated linoleic acid in reducing body fat and preventing holiday weight gain. *Int J Obes* 2007; 31: 481–487.
9. Baker RC, Kirschenbaum DS. Weight control during the holidays: Highly consistent self-monitoring as a potentially useful coping mechanism. *Heath Psych* 1998; 17 (4): 367–370.
10. Mehrang S, Helander E, Chieh A, Korhonen I. Seasonal weight variation patterns in seven countries located in northern and southern hemispheres. Conference Proceedings of the Annual International Conference of the IEEE Engineering in Medicine and Biology Society, 2016, 2475–2478.
11. Helander EE. (2016). Weight gain over the holidays in three countries. *N Eng J Med* 2016; 375 (12): 1200–1202.
12. Dall TM, Zhang Y, Chen YJ, Askarinam Wagner RC, Hogan PF, Fagan NK, et al. Cost associated with being overweight and with obesity, high alcohol consumption, and tobacco use within the military health system's TRICARE prime-enrolled population. *Am J Health Promot* 2007; 22: 120–139. [PubMed: 18019889]
13. Krukowski RA, Hare ME, Talcott GW, Gladney LA, Johnson KC, Richey PA, Kocak M, Keller PL, Hryshko-Mullen A, Klesges RC. Dissemination of the Look AHEAD Intensive Lifestyle Intervention in the United States Military: A Randomized Controlled Trial. *Obesity* (Silver Spring) 2018; 26 (10): 1558–1565. [PubMed: 30277030]
14. Look AHEAD Research Group. (2006). The Look AHEAD study: a description of the lifestyle intervention and the evidence supporting it. *Obesity*, 14(5), 737–752. [PubMed: 16855180]
15. Krukowski RA, Hare ME, Talcott GW, Johnson KC, Richey PA, Kocak M, et al. Dissemination of the Look AHEAD intensive lifestyle intervention in the United States Air Force: Study rationale, design, and methods. *Contemp Clin Trials* 2015; 40: 232–239 [PubMed: 25545025]
16. Ross KM, Wing RR. Concordance of in-home 'smart' scale measurement with body weight measured in-person. *Obes Sci & Pract* 2016; 2 (2): 224–2248. [PubMed: 27499884]
17. Pebley K, Klesges RC, Talcott GW, Kocak M, & Krukowski RA. E-scale measurements in comparison to clinic weight measurements. *Obesity* In press.
18. Jensen MD, Ryan DH, Donato KA, et al. Guidelines (2013) for managing overweight and obesity in adults. *Obesity* 2014; 22: S1–S410.
19. Nackers LM, Ross KM, Perri MG. The association between rate of initial weight loss and long-term success in obesity treatment: Does slow and steady win the race? *Int J Behav Med* 2010; 17: 161–167. [PubMed: 20443094]
20. Burke LE, Wang J, Sevick MA. Self-monitoring in weight loss: A systematic review. *J Am Diet Assoc* 2011; 111(1): 92–102. [PubMed: 21185970]
21. Tanofsky-Kraff M, Sbrocco T, Theim KR, Cohen LA, Mackey ER, Stice E, Henderson JL, McCreight SJ, Bryant EJ, Stephens MB. Obesity and the US military family. *Obesity* 2013; 21: 2205–2220. [PubMed: 23836452]

Table 1.

Pre-Holiday to Holiday Period Comparisons by Participant Characteristics

	Weight Change (%)	
	N	M (SD)
Overall	163	.18 (2.16)
Days from randomization		
< 90 (<i>mid-August to mid-November</i>)	44	−0.98 (1.75) ***
91–180 (<i>mid-May to mid-August</i>)	43	0.25 (1.89) ***
181–270 (<i>mid-February to mid-May</i>)	54	0.90 (3.27) ***
> 270 (<i>Before mid-February</i>)	22	0.93 (1.65) ***
	Rate of Weekly Weight Change	
Overall	71	0.17 (.45) **
Race		
African American	16	0.38 (0.47) *
Other	13	0.28 (0.39) *
Caucasian	42	0.06 (0.43) *
Days from randomization		
< 90 (<i>mid-August to mid-November</i>)	14	0.45 (0.33) *
91–180 (<i>mid-May to mid-August</i>)	21	0.18 (0.52) *
181–270 (<i>mid-February to mid-May</i>)	24	0.03 (0.06) *
> 270 (<i>Before mid-February</i>)	12	0.07 (0.05) *
	Self-Weighing Frequency	
Overall	163	−1.09 (11.31) **
Days from randomization		
< 90 (<i>mid-August to mid-November</i>)	44	5.18 (14.70) **
91–180 (<i>mid-May to mid-August</i>)	43	−4.70 (9.55) **
181–270 (<i>mid-February to mid-May</i>)	54	−1.72 (7.85) **
> 270 (<i>Before mid-February</i>)	22	−5.00 (8.95) **

Note.

 $p < .001$ **
 $p < .01$ *
 $p < .05$

Table 2.

Holiday to Post-Holiday Period Comparisons by Participant Characteristics

	Weight Change (%)	
	N	M (SD)
Overall	171	.12 (2.21)
Days from randomization		
< 90 (<i>Mid-November to Mid-February</i>)	43	−0.85 (2.10) ***
91–180 (<i>Mid-August to Mid-November</i>)	48	0.60 (2.10) ***
181–270 (<i>Mid-May to Mid-August</i>)	39	0.35 (2.59) ***
> 270 (<i>Before Mid-May</i>)	41	0.36 (1.80) ***
	Rate of Weekly Weight Change	
Overall	152	−0.11 (0.41) **
Race		
African American	31	−0.32 (0.47) **
Other	22	−0.09 (0.29) **
Caucasian	99	−0.04 (0.40) **
Gender		
Male	73	−0.17 (0.39) *
Female	79	−0.05 (0.43) *
	Self-Weighing Frequency	
Overall	171	−0.46 (10.66)
Days from randomization		
< 90 (<i>Mid-November to Mid-February</i>)	43	3.74 (13.62)
91–180 (<i>Mid-August to Mid-November</i>)	48	−2.48 (8.07)
181–270 (<i>Mid-May to Mid-August</i>)	39	−1.74 (10.82)
> 270 (<i>Before Mid-May</i>)	41	−1.29 (8.59)

Note.

 $p < .001$ **
 $p < .01$ *
 $p < .05$