Published in final edited form as:

Curr Opin Neurobiol. 2007 October ; 17(5): 587–592. doi:10.1016/j.conb.2007.11.003.

Integration of light-controlled neuronal firing and fast circuit imaging

Raag D. Airan, Elbert Hu, Ragu Vijaykumar, Madhuri Roy, Leslie A. Meltzer, Karl Deisseroth*

Department of Bioengineering, Department of Psychiatry and Behavioral Sciences, Stanford University, Stanford, CA

Summary

For understanding normal and pathological circuit function, capitalizing on the full potential of recent advances in fast optical neural circuit control will depend crucially on fast, intact-circuit readout technology. First, millisecond-scale optical control will be best leveraged with simultaneous millisecond-scale optical imaging. Second, both fast circuit control and imaging should be adaptable to intact-circuit preparations from normal and diseased subjects. Here we illustrate integration of fast optical circuit control and fast circuit imaging, review recent work demonstrating utility of applying fast imaging to quantifying activity flow in disease models, and discuss integration of diverse optogenetic and chemical genetic tools that have been developed to precisely control the activity of genetically-specified neural populations. Together these neuroengineering advances raise the exciting prospect of determining the role specific cell types play in modulating neural activity flow in neuropsychiatric disease.

Genetically-targeted circuit control technologies

Several technologies recently have been developed to control the activity of genetically-specified neural populations. Chemically-triggered genetic silencing technologies include MISTs [1••], AlstR [2••], GluClαβ [3••], and modified GABA-A receptors [4•]; all of these allow for inducible and reversible “knock out” of genetically-specified cell populations from neural circuits after application or washout of appropriate small molecule drugs. This chemical-genetic strategy (reviewed in this issue by Tervo and Karpova) already has shown feasibility in awake, behaving animals performing amphetamine-induced rotational behavior, a classic behavioral assay of Parkinson’s Disease models [3••].

Complementing these elegant chemical-genetic neural control methods are recently developed optogenetic tools that allow for fast stimulation and inhibition of genetically-targeted neural populations on the millisecond timescale. Transgenic expression of the light-

*Correspondence should be addressed to K.D.: Karl Deisseroth, M.D., Ph.D., Department of Bioengineering, W083 Clark Center, 318 Campus Drive West, Stanford, University, Phone: (650) 736-4325, Fax: (650) 725-7799, deissero@stanford.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

activated cation channel channelrhodopsin-2 from *C. reinhardtii* (ChR2) confers ability to depolarize neurons with blue (~470nm) light [5•], while expression of the light-activated chloride pump halorhodopsin from *N. pharaonis* (NpHR) permits hyperpolarization of cells with yellow (~580nm) light [6•]. Both of these tools have been demonstrated to permit precise stimulation and inhibition of individual spikes in neurons with millisecond resolution [5•, 6•, 7], together could modulate neuronal synchronization [8], and have been shown to operate in intact tissue without addition of chemical cofactors [6•, 7]. Other relevant technological advances include light-activated ionotropic glutamate receptors (LiGluR) [9•] modified by an exogenous chemical photoswitch; rhodopsin itself, which can confer neuronal silencing under optical control [10•]; transmembrane receptors that can activate neurons in response to flash uncaging of exogenous chemical ligands [11•]; and technology for precise stimulation in prescribed spatial patterns [12•], as in glutamate uncaging methods [13•], that will allow fast circuit mapping with optogenetic technologies.

Integrating fast optical control with fast optical imaging of activity flow

The chemical and optogenetic tools described above in principle allow characterization of the contribution of genetically-defined cell types to neural activity flow through intact circuits. Complementing these control strategies, recent advances in high-speed optical imaging now allow for quantification of activity flow through intact neural circuits with high spatiotemporal resolution. Activity imaging presents clear advantages over alternative techniques in quantifying whole-circuit activity flow. For example, traditional electrode-based techniques provide relatively limited information on the spatial distribution of evoked activity or the type of active cells. Conversely, fMRI and PET can determine the spatial distribution of activity across the entire brain, albeit with coarse spatial and temporal resolution. However, circuit imaging with fast optical activity reporters could in principle permit quantification of activity of intact neural circuits on the millisecond timescale and with near-micron spatial resolution [14,15•]; and when combined with chemical or optogenetic control tools could illuminate how specific cell types control neural activity flow.

Imaging of neural activity typically involves either calcium or voltage-sensitive reporters. Calcium dyes (e.g. Fura-2) report activity through binding to calcium and have higher signal to noise ratios (SNR) than voltage-sensitive dyes, but also much slower overall kinetics, limiting their use in quantifying millisecond-scale kinetics of activity propagation [16]. Voltage-sensitive dyes (e.g. di-4-ANEPPS, RH1691) report activity changes on much faster timescales (<1ms), although correspondingly reduced SNRs have limited their application relative to calcium dyes [17,18•]. Both modalities have been applied to imaging neural activity in culture, slice, and *in vivo* preparations [16,19] and choice of reporter is heavily dependent on the experiment in question (see [16] for a review of these considerations). Emerging genetically-encodeable reporters of both calcium (e.g. GCaMP2, Camgaroo) and membrane voltage (e.g. hVOS, VSFP1) will allow a further level of specificity in quantifying activity flow of genetically-defined neural populations [16].

The acute slice preparation provides an excellent experimental system for determining the changes in neural activity flow that underlie behavior, with both optical and behavioral

assays conducted in the same animal. While Ca^{2+} -Fura-2 imaging (excitation wavelength 340nm) can be integrated with ChR2/NpHR optical control in acute slices [6••], fast VSDI technology also can be integrated with fast optical control (Figure 1,2). We have described experimental and analytical methods for quantitative high-speed voltage-sensitive dye imaging in acute hippocampal slices [20••], and circuit activity controlled by ChR2 (excited at ~470nm) can be imaged with the voltage-sensitive dye RH-155, which has a voltage-sensitive absorption band at ~700nm that is sufficiently spectrally separated from the excitation peaks of ChR2 and NpHR to permit smoothly integrated imaging of activity (Figure 1) [21•,22•]. Indeed, integrated millisecond-scale optical control and imaging can be achieved, here using acute brain slices from Thy1::ChR2-YFP transgenic mice [12••,23••]. These mice express ChR2 in excitatory projection neurons, and in the hippocampus ChR2 is chiefly expressed in the CA1 subfield (Figure 2a). Imaging optically-evoked neural activity flow on the millisecond-timescale in this preparation can be conducted, and data can be collected and quantified from the entire imaged circuit (Figure 2), in principle allowing determination of the real-time impact of genetically defined cells on intact neural circuit dynamics.

Preclinical disease models

Could this kind of method also be applied to quantifying altered circuit dynamics in neuropsychiatric disease models? For the slice preparation method to be effective, stable changes in the circuit of interest would have to be involved in the pathophysiology of the neuropsychiatric phenotype. A strategy for quantitative delineation of the changes to neural circuitry underlying neuropsychiatric disease would entail (Figure 3): 1) identification of a validated animal model and target brain regions; 2) measurement of the disease-relevant behavior using a validated behavioral assay; 3) preparation of acute slices from the hypothesized brain regions labeled with the activity reporter; 4) fast imaging of the spatiotemporal dynamics of optically, chemically, or electrically evoked activity within the brain area of interest; and finally 5) analysis of the imaging data to identify features of evoked activity linked to the behavioral state of the individual animal. This experimental paradigm has already been applied to determining the role of key hippocampal pathways in an animal model of depression [20••]. In this study, animal models of depression and its treatment were induced using combinations of chronic mild stress (CMS), an ethologically-relevant well-validated model of depression [24], and widely-used antidepressants (fluoxetine and imipramine). The modified Porsolt forced swim test (FST) served as a behavioral endpoint, given its predictive validity in assaying models of depression and antidepressant activity [25]; and high-speed voltage sensitive dye imaging (VSDI) was employed to quantify alterations in hippocampal activity flow associated with this model of neuropsychiatric disease.

VSDI of acute hippocampal formation slices demonstrated that activity percolation was decreased in the dentate gyrus and increased in CA1 following CMS; a pattern reversed by antidepressant treatment [20••], defining a neurophysiological endophenotype of depression-relevant behavior. This metric correlated strongly with FST scores on an individual animal basis ($r^2 > 0.55$), across all treatment groups. Interestingly, amplitude of neural circuit activity at any one point was not consistently altered; instead, affective state

modulation controlled the areal spread of activity, a result that would have been very difficult to identify without an imaging approach. This identification of a neurophysiological endophenotype may be combined with chemical and optogenetic control tools to allow for cell-type based dissection. For example, hippocampal neurogenesis has been suggested to have a pivotal role in depression pathophysiology and treatment [26]. By selectively expressing ChR2 and NpHR in targeted cell populations, it may be possible to quantitatively determine the causal influence of newborn neurons or other cell types in mediating the antidepressant-induced changes in circuit dynamics that have been observed [20• •].

Many other animal models of disease symptomatology could be studied with these methods. While no animal model completely simulates all the signs, symptoms, and etiologies of any human disease (e.g. environmental and genetic influences on diabetes, suicidality in depression), neuropsychiatric disease models do simulate core symptoms (e.g. anhedonia in depression). These models are validated based on several sources: face validity – the animal behavior models the human symptoms well; construct validity – the model construction parallels known etiology of the disease; and predictive validity – the reaction of the model to a given treatment predicts reaction of the human disease to analogous treatment (e.g., drug) [27]. By understanding the contribution of target cell types and brain regions to the physiology underlying the behavioral phenotype of a particular animal model, we will gain understanding of the physiology that underlies the corresponding human neuropsychiatric symptoms.

Neurological diseases may also be addressed with combined high-speed optical imaging and control. For instance, while DBS to inhibit the subthalamic nucleus (STN) is effective in treating some Parkinson's Disease symptoms, it is unclear which cell populations (e.g. excitatory or inhibitory neurons) mediate the relevant changes in activity within this structure [28,29]. To determine the role of these populations in mediating successful DBS, inhibitory optogenetic probes (e.g. NpHR) or chemical genetic probes (VAMP/Syb MIST) [1• •] can be expressed in excitatory neurons in the STN (e.g. by driving expression with the selective promoter CaMKII α). Furthermore, while it is known that dopamine depletion in the basal ganglia is important in Parkinson's disease, there is still controversy regarding the exact quantitative role of different dopamine-receptor expressing populations in movement-related deficits associated with the disorder [29]. By imaging basal ganglia activity flow following millisecond-scale optogenetic control of D1- or D2-expressing neural populations, a more quantitative model of this circuit may result that could guide future pharmacological or electrical stimulation-based treatments.

As a final perspective, it is important to note that imaging whole-circuit activity in the awake, behaving animal is close to realization. Activity imaging of cerebellar and cortical neural networks has already been achieved *in vivo* using both calcium and voltage sensitive reporters [19,22•,30], and similar studies may be completed in deep-brain structures with the Schnitzer system for endoscopic imaging of neural circuitry in freely moving animals [31• •]. Loading techniques such as *in vivo* electroporation allow for precise and efficient activity-dye labeling of neurons [32•] and genetically-encodeable reporters of both membrane potential and calcium described above can be delivered with injection of viral vectors or using transgenic animals [16].

Conclusions

By driving expression of optogenetic probes (or their chemical counterparts) in specific cell types, circuit elements can be controlled to determine their role in behavior. With high-speed imaging, activity propagation can be quantified both in normal circuits and in animal models of neuropsychiatric disease. When these two classes of fast optical technology are combined (Figure 1,2), a high-speed experimental platform results that may further our quantitative understanding of neuropsychiatric diseases (Figure 3), extend our ability to design and screen for novel treatments, and deepen our understanding of neural circuit dynamics arising from properties of individual cellular components.

Acknowledgments

We thank the entire Deisseroth lab for helpful discussions. This work was supported by NIDA, NIMH, and the NIH Director's Pioneer Award, as well as by NARSAD, APIRE, and the Snyder, Culpeper, Coulter, Klingenstein, Whitehall, McKnight, and Albert Yu and Mary Bechmann Foundations (KD) and the Stanford MSTP (RDA).

References and Recommended Reading

Papers of particular interest have been highlighted as:

- of special interest
 - of outstanding interest
- 1••. Karpova AY, Tervo DGR, Gray NW, Svoboda K. Rapid and Reversible Chemical Inactivation of Synaptic Transmission in Genetically Targeted Neurons. *Neuron*. 2005; 48:727–735. [PubMed: 16337911]
 - 2••. Tan EM, Yamaguchi Y, Horwitz GD, Gosgnach S, Lein ES, Goulding M, Albright TD, Callaway EM. Selective and Quickly Reversible Inactivation of Mammalian Neurons In Vivo Using the *Drosophila* Allatostatin Receptor. *Neuron*. 2006; 51:157–170. [PubMed: 16846851]
 - 3••. Lerchner W, Xiao C, Nashmi R, Slimko EM, van Trigt L, Lester HA, Anderson DJ. Reversible Silencing of Neuronal Excitability in Behaving Mice by a Genetically Targeted, Ivermectin-Gated Cl-Channel. *Neuron*. 2007; 54:35–49. [PubMed: 17408576]
 - 4•. Wulff P, Goetz T, Leppa E, Linden A-M, Renzi M, Swinny JD, Vekovischeva OY, Sieghart W, Somogyi P, Korpi ER, et al. From synapse to behavior: rapid modulation of defined neuronal types with engineered GABAA receptors. *Nat Neurosci*. 2007; 10:923–929. [PubMed: 17572671]
 - 5••. Boyden ES, Zhang F, Bamberg E, Nagel G, Deisseroth K. Millisecond-timescale, genetically targeted optical control of neural activity. *Nat Neurosci*. 2005; 8:1263. [PubMed: 16116447]
 - 6••. Zhang F, Wang L-P, Brauner M, Liewald JF, Kay K, Watzke N, Wood PG, Bamberg E, Nagel G, Gottschalk A, et al. Multimodal fast optical interrogation of neural circuitry. *Nature*. 2007; 446:633–639. [PubMed: 17410168]
 7. Zhang F, Wang L-P, Boyden ES, Deisseroth K. Channelrhodopsin-2 and optical control of excitable cells. *Nat Meth*. 2006; 3:785–792.
 8. Han X, Boyden ES. Multiple-color optical activation, silencing, and desynchronization of neural activity, with single-spike temporal resolution. *PLoS ONE*. 2007;2.
 - 9•. Szobota S, Gorostiza P, Del Bene F, Wyart C, Fortin DL, Kolstad KD, Tulyathan O, Volgraf M, Numano R, Aaron HL, et al. Remote Control of Neuronal Activity with a Light-Gated Glutamate Receptor. *Neuron*. 2007; 54:535–545. [PubMed: 17521567]
 - 10•. Li X, Gutierrez DV, Hanson MG, Han J, Mark MD, Chiel H, Hegemann P, Landmesser LT, Herlitze S. Fast noninvasive activation and inhibition of neural and network activity by vertebrate

- rhodopsin and green algae channelrhodopsin. *PNAS*. 2005; 102:17816–17821. [PubMed: 16306259]
- 11•. Miesenbock G, Kevrekidis IG. OPTICAL IMAGING AND CONTROL OF GENETICALLY DESIGNATED NEURONS IN FUNCTIONING CIRCUITS. *Annual Review of Neuroscience*. 2005; 28:533–563.
 - 12•. Wang H, Peca J, Matsuzaki M, Matsuzaki K, Noguchi J, Qiu L, Wang D, Zhang F, Boyden E, Deisseroth K, et al. High-speed mapping of synaptic connectivity using photostimulation in Channelrhodopsin-2 transgenic mice. *PNAS*. 2007; 104:8143–8148. [PubMed: 17483470]
 - 13•. Shoham S, O'Connor DH, Sarkisov DV, Wang SSH. Rapid neurotransmitter uncaging in spatially defined patterns. *Nat Meth*. 2005; 2:837–843.
 14. Ang CW, Carlson GC, Coulter DA. Hippocampal CA1 Circuitry Dynamically Gates Direct Cortical Inputs Preferentially at Theta Frequencies. *J Neurosci*. 2005; 25:9567–9580. [PubMed: 16237162]
 - 15•. Mann EO, Suckling JM, Hajos N, Greenfield SA, Paulsen O. Perisomatic Feedback Inhibition Underlies Cholinergically Induced Fast Network Oscillations in the Rat Hippocampus In Vitro. *Neuron*. 2005; 45:105. [PubMed: 15629706]
 16. Knopfel T, Diez-Garcia J, Akemann W. Optical probing of neuronal circuit dynamics: genetically encoded versus classical fluorescent sensors. *Trends in Neurosciences*. 2006; 29:160–166. [PubMed: 16443289]
 17. Tominaga T, Tominaga Y, Yamada H, Matsumoto G, Ichikawa M. Quantification of optical signals with electrophysiological signals in neural activities of Di-4-ANEPPS stained rat hippocampal slices. *Journal of Neuroscience Methods*. 2000; 102:11. [PubMed: 11000407]
 - 18•. Petersen CCH, Grinvald A, Sakmann B. Spatiotemporal Dynamics of Sensory Responses in Layer 2/3 of Rat Barrel Cortex Measured In Vivo by Voltage-Sensitive Dye Imaging Combined with Whole-Cell Voltage Recordings and Neuron Reconstructions. *J Neurosci*. 2003; 23:1298–1309. [PubMed: 12598618]
 19. Grinvald A, Hildesheim R. VSDI: A NEW ERA IN FUNCTIONAL IMAGING OF CORTICAL DYNAMICS. *Nature Reviews Neuroscience*. 2004; 5:874. [PubMed: 15496865]
 - 20•. Airan R, Meltzer LA, Roy M, Gong Y, Chen H, Deisseroth K. High-speed imaging reveals neurophysiological links to behavior in an animal model of depression. *Science*. 2007 Aug 10; 317(5839):819–23. [PubMed: 17615305]
 - 21•. London JA, Zecevic D, Cohen LB. Simultaneous optical recording of activity from many neurons during feeding in Navanax. *J Neurosci*. 1987; 7:649–661. [PubMed: 2435862]
 - 22•. Petersen CCH, Sakmann B. Functionally Independent Columns of Rat Somatosensory Barrel Cortex Revealed with Voltage-Sensitive Dye Imaging. *J Neurosci*. 2001; 21:8435–8446. [PubMed: 11606632]
 - 23•. Arenkiel BR, Peca J, Davison IG, Feliciano C, Deisseroth K, Augustine GJ, Ehlers MD, Feng G. In Vivo Light-Induced Activation of Neural Circuitry in Transgenic Mice Expressing Channelrhodopsin-2. *Neuron*. 2007; 54:205–218. [PubMed: 17442243]
 24. Willner P. Chronic Mild Stress (CMS) Revisited: Consistency and Behavioural-Neurobiological Concordance in the Effects of CMS. *Neuropsychobiology*. 2005; 52:90–110. [PubMed: 16037678]
 25. Cryan JF, Valentino RJ, Lucki I. Assessing substrates underlying the behavioral effects of antidepressants using the modified rat forced swimming test. *Neuroscience & Biobehavioral Reviews*. 2005; 29:547. [PubMed: 15893822]
 26. Warner-Schmidt JL, Duman RS. Hippocampal neurogenesis: Opposing effects of stress and antidepressant treatment. *Hippocampus*. 2006; 16:239–249. [PubMed: 16425236]
 27. Willner P, Mitchell P. The validity of animal models of predisposition to depression. *Behavioural Pharmacology*. 2002; 13:169–188. [PubMed: 12122308]
 28. Benabid AL. Deep brain stimulation for Parkinson's disease. *Current Opinion in Neurobiology*. 2003; 13:696–706. [PubMed: 14662371]
 29. Bevan MD, Atherton JF, Baufreton J. Cellular principles underlying normal and pathological activity in the subthalamic nucleus. *Current Opinion in Neurobiology*. 2006; 16:621–628. [PubMed: 17084618]

30. Diez-Garcia J, Akemann W, Knopfel T. In vivo calcium imaging from genetically specified target cells in mouse cerebellum. *NeuroImage*. 2007; 34 :859–869. [PubMed: 17161628]
- 31•. Flusberg BA, Cocker ED, Piyawattanametha W, Jung JC, Cheung ELM, Schnitzer MJ. Fiber-optic fluorescence imaging. *Nat Meth*. 2005; 2:941–950.
- 32•. Nagayama S, Zeng S, Xiong W, Fletcher ML, Masurkar AV, Davis DJ, Pieribone VA, Chen WR. In Vivo Simultaneous Tracing and Ca²⁺ Imaging of Local Neuronal Circuits. *Neuron*. 2007; 53:789–803. [PubMed: 17359915]

Figure 1. Design for integrating fast optical control and recording technologies

- a) Voltage-sensitive absorption band of RH-155 overlaid on action spectra of ChR2, NpHR; spectral separation suggests that high-speed all-optical control and recording of neural activity flow can be conducted with this combination of optical tools.
- b) Schematic of microscopy apparatus for high-speed all-optical control and recording of circuit activity.

Figure 2. Experimental integration of high-speed all-optical circuit control and imaging

a) *Top:* (Left to right) *Brightfield:* image showing a 350μm horizontal hippocampal slice prepared from a Thy1::ChR2-YFP transgenic mouse (6wks) stained with RH-155 (0.1mg/mL for 3hrs). White boxes indicate sources of VSDI traces in CA1 and cortex (ctx); field recording electrode can be seen in upper middle region of image. Scale bar 350μm. *YFP:* image showing labeling of neurons with ChR2-YFP primarily in CA1 of hippocampus; same field as brightfield image. *5Hz, 20Hz:* sample VSDI images of neural activity flow in CA1 and cortex following ChR2-mediated optical stimulation (10ms pulses of 470nm light delivered by the Sutter DG-4 high-speed optical switch); same field as brightfield and YFP images. Note the lack of optical signal in non-YFP expressing CA3; the optical signal seen in adjacent cortex is synaptically induced as shown in (b). *Bottom:* Single-pixel VSDI traces corresponding to white boxes for 5Hz (left) and 20Hz (right) optical stimulation. Blue dashes indicate timing of 10ms blue light pulses used to stimulate ChR2. Scale bars: 0.1% dF/F and 200ms. *Insets:* sample field recording traces for one optically stimulated event delivered in trains of 10 pulses at 5Hz (left) and 20Hz (right). Scale bars: 0.5mV and 20ms.

b) Optical recording of optically-evoked neural activity flow following pharmacological modulation. Blue light directly stimulates CA1 neurons and triggers stimulation of adjacent cortex via excitatory synapses as CNQX (10μM) and D-APV (25μM) block cortical responses to optical stimulation determined by VSDI. Image scale bar: 350μm. VSDI trace scale bars: 0.1% dF/F and 200ms.

Figure 3. Experimental design for using high-speed all-optical control and recording to probe neuropsychiatric disease.