Cochrane
Library

Cochrane Database of Systematic Reviews

Human albumin infusion for treating oedema in people with nephrotic syndrome (Review)

Ho JJ, Adnan AS, Kueh YC, Ambak NJ, Van Rostenberghe H, Jummaat F

Ho JJ, Adnan AS, Kueh YC, Ambak NJ, Van Rostenberghe H, Jummaat F.
Human albumin infusion for treating oedema in people with nephrotic syndrome.
Cochrane Database of Systematic Reviews 2019, Issue 7. Art. No.: CD009692.
DOI: [10.1002/14651858.CD009692.pub2](https://doi.org/10.1002/14651858.CD009692.pub2).

www.cochranelibrary.com

TABLE OF CONTENTS

HEADER	1
ABSTRACT	1
PLAIN LANGUAGE SUMMARY	2
SUMMARY OF FINDINGS	3
BACKGROUND	5
OBJECTIVES	5
METHODS	5
RESULTS	8
Figure 1.	8
Figure 2.	9
DISCUSSION	10
AUTHORS' CONCLUSIONS	11
ACKNOWLEDGEMENTS	11
REFERENCES	12
CHARACTERISTICS OF STUDIES	14
ADDITIONAL TABLES	18
APPENDICES	19
CONTRIBUTIONS OF AUTHORS	22
DECLARATIONS OF INTEREST	22
SOURCES OF SUPPORT	23
DIFFERENCES BETWEEN PROTOCOL AND REVIEW	23

[Intervention Review]

Human albumin infusion for treating oedema in people with nephrotic syndrome

Jacqueline J Ho¹, Azreen S Adnan², Yee Cheng Kueh³, Nurul Jannah Ambak⁴, Hans Van Rostenberghe⁵, Fauziah Jummaat⁶

¹Department of Paediatrics, RCSI & UCD Malaysia Campus (formerly Penang Medical College), George Town, Malaysia. ²Management Science University (MSU) Medical Centre, Shah Alam, Malaysia. ³Unit Biostatistics and Research Methodology, School of Medical Sciences, Universiti Sains Malaysia, Kubang Kerian, Malaysia. ⁴Centre for Research on Women and Gender (KANITA), Universiti Sains Malaysia, Penang, Malaysia. ⁵Department of Paediatrics, School of Medical Sciences, Universiti Sains Malaysia, Kubang Kerian, Malaysia. ⁶Department of Obstetrics and Gynecology, School of Medical Sciences, University Science Malaysia, Kota Bharu, Malaysia

Contact address: Yee Cheng Kueh, Unit Biostatistics and Research Methodology, School of Medical Sciences, Universiti Sains Malaysia, Kubang Kerian, Kelantan, 16150, Malaysia. yckueh@usm.my.

Editorial group: Cochrane Kidney and Transplant Group.

Publication status and date: New, published in Issue 7, 2019.

Citation: Ho JJ, Adnan AS, Kueh YC, Ambak NJ, Van Rostenberghe H, Jummaat F. Human albumin infusion for treating oedema in people with nephrotic syndrome. *Cochrane Database of Systematic Reviews* 2019, Issue 7. Art. No.: CD009692. DOI: [10.1002/14651858.CD009692.pub2](https://doi.org/10.1002/14651858.CD009692.pub2).

Copyright © 2019 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

ABSTRACT

Background

Oedema is a common clinical symptom in people with nephrotic syndrome and human albumin has been widely used in the treatment of oedema by increasing vascular volume and this inducing diuresis. It may be used with or without diuretics such as furosemide. However, the quantitative contribution of human albumin in treating oedema is not fully understood. If human albumin were found to be effective and safe in the treatment of oedema, it could help clinicians to develop therapeutic strategies to improve the management of diuretic resistance associated with nephrotic syndrome.

Objectives

This review aimed to examine the benefits and harms of human albumin infusion for treating oedema associated with nephrotic syndrome.

Search methods

We searched the Cochrane Kidney and Transplant Register of Studies up to 23 June 2019 through contact with the Information Specialists using search terms relevant to this review. Studies in the Specialised Register are identified through searches of CENTRAL, MEDLINE, and EMBASE, conference proceedings, the International Clinical Trials Register (ICTRP) Search Portal and ClinicalTrials.gov.

Selection criteria

We included randomised controlled trials (RCTs) and quasi-RCTs evaluating the effect of human albumin infusion compared with placebo or no intervention, human albumin with diuretics compared with diuretic alone, human albumin compared with diuretics and other treatments, clinical outcomes, death, quality of life, kidney function and adverse effects in people with nephrotic syndrome. We excluded cross-over studies but data for the first period was to be included if available.

Data collection and analysis

Standard methods of the Cochrane Collaboration were used. Two authors independently assessed eligibility, risk of bias, study quality and extracted data. We calculated mean difference (MD) for continuous data with 95% confidence intervals (CI). We assessed the certainty of the evidence using GRADE.

Main results

One study met our inclusion criteria (26 children with minimal change nephrotic syndrome) and 11 were excluded (nine cross-over studies, one where albumin was not used for nephrotic syndrome and one where authors did not state whether the children had oedema). Risk of bias for the included study was unclear for selection bias, high for performance and detection bias, low for attrition bias, and high for selective reporting. The included study compared albumin plus furosemide with an equal volume of dextrose. Of our prespecified outcomes, the authors reported clinical improvement as weight change, serum sodium and adverse outcomes (blood pressure). The authors reported a greater weight loss in the albumin treated group initially but no difference overall at 10 days. However, the data in the text and the figures were inconsistent so we could not confirm the authors statements (very low certainty evidence). It is uncertain whether albumin infusion improves serum sodium when compared with an equal volume of dextrose (MD 2.00 mEq/L, 95% CI -0.09 to 4.09), systolic blood pressure (MD 2.00 mmHg, 95% CI -3.52 to 7.52) or diastolic blood pressure (MD 2.00 mmHg, 95% CI -4.29 to 8.29). Death, quality of life, and kidney function were not reported.

Authors' conclusions

We identified only one small study that was relevant to our review, therefore we are unable to draw any conclusions regarding the use of human albumin with or without diuretics in nephrotic syndrome. More RCTs are needed.

PLAIN LANGUAGE SUMMARY

Human albumin infusion for treating oedema in people with nephrotic syndrome

What is the issue?

Nephrotic syndrome is a kidney disorder that causes the body to excrete high quantities of protein in urine. The loss of protein causes the body to retain fluid that would normally be excreted in the urine and swelling (oedema) typically starting at the ankles and sometimes involving the hands and face.

Oedema is often treated by a diuretic (a medication that stimulates the kidney to increase the excretion of salt and water). However, in nephrotic syndrome diuretics alone often don't work. Human albumin has been used to replace the protein lost. This might result in the oedema fluid being drawn back into the blood circulation so it can be excreted by the kidneys. It might be given alone or in combination with a diuretic. We wanted to find out whether albumin could treat oedema in people with nephrotic syndrome.

What did we do?

We searched for randomised controlled trials (RCTs) or quasi-RCTs comparing (1) albumin with a placebo or no treatment, (2) albumin with a diuretic, (3) albumin combined with a diuretic compared with a diuretic alone, or (4) albumin with other treatments. We excluded cross-over studies but would have included data for the first period before the cross-over if available. The last search was done in June 2019.

What did we find?

The review found one small RCT (26 patients) comparing human albumin plus the diuretic furosemide with placebo suitable for inclusion. We found nine studies on people with nephrotic syndrome that tested these comparisons but these were 'cross-over' studies which we judged not suitable. To find out whether there was any improvement after albumin, the study measured weight loss and serum sodium. The adverse effect measured was blood pressure. Although the authors reported increased weight loss we were not able to confirm this due to inconsistency between the data reported in the table and the text. There was no change in serum sodium or blood pressure. We judged these outcomes all to be of very low certainty. Death, quality of life, and kidney function were not reported.

Conclusions

Because there was only one small study found we cannot tell whether albumin is effective in people with nephrotic syndrome and we do not know from the studies we looked at whether it is safe. There is no evidence in adults. We judged the evidence to be very low certainty. Therefore RCTs are needed.

SUMMARY OF FINDINGS

Summary of findings for the main comparison. Human albumin infusion plus furosemide compared to glucose for treating people with nephrotic syndrome

Human albumin infusion compared to placebo, diuretics or other treatment for treating oedema in people with nephrotic syndrome

Patient or population: patients of all ages with oedema due to nephrotic syndrome

Setting: health care

Intervention: human albumin infusion plus furosemide

Comparison: placebo, diuretics or other treatment

Outcomes	Anticipated absolute effects* (95% CI)		Relative effect (95% CI)	No. of participants (studies)	Quality of the evidence (GRADE)	Comments
	Risk with other intervention or placebo	Risk with human albumin infusion				
Weight loss rate up to day 10 after treatment	Unable to report data due to lack of clarity		-	26 (1)	⊕⊕⊕⊕ VERY LOW ^{1 2 3 4}	Weight loss rate was reported in the text at day 1, day 4, day 7 and day 10. However the values stated in text are inconsistent with the values depicted in figure (Fig 1; Lee 2000a). The authors reported no difference up to day 10
Others clinical improvement parameters: Serum sodium	At day 5, the mean serum sodium was 2 mEq/L higher (0.09 lower to 4.09 higher) than the control group		-	26 (1)	⊕⊕⊕⊕ VERY LOW ^{1 2 3 4}	Other clinical improvement parameters such as increased urine volume; reductions in ascites; pleural effusion; penile or scrotal oedema were not measured or reported in the included study The serum sodium was reported in the text at day 1, day 3 and day 5. The authors reported no difference up to day 5
Death	No deaths reported over the 12-month study period					Death due to kidney failure was not measured or reported in the included study
Kidney function	There was a higher reported mean spot urine protein/creatinine ratio in the albumin treated group at Day 2 and 4 after treatment but not on Day 6, 8 and 10					Serum creatinine, glomerular filtration rate were not measured or reported in the included study
Quality of life	See comments					Quality of life scale score was not measured or reported in the included study

Adverse effects:	At day 5, the systolic blood pressure was 2 mmHg higher (3.52 lower to 7.52 higher) than the control group	-	26 (1)	⊕⊕⊕⊕ VERY LOW ^{1 2 3 4 5}	Mean blood pressure was reported but there was no data on hypertension or hypotension. Plasma volume overload, pulmonary oedema, need for dialysis or ventilation to control plasma volume overload were not measured or reported in the only included study The systolic blood pressure was reported at day 1, day 3 and day 5. The authors reported no difference up to day 5
------------------	--	---	--------	---------------------------------------	--

***The risk in the intervention group** (and its 95% CI) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

CI: Confidence interval; **MD:** Mean difference

GRADE Working Group grades of evidence

High certainty: We are very confident that the true effect lies close to that of the estimate of the effect

Moderate certainty: We are moderately confident in the effect estimate: The true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different

Low certainty: Our confidence in the effect estimate is limited: The true effect may be substantially different from the estimate of the effect

Very low certainty: We have very little confidence in the effect estimate: The true effect is likely to be substantially different from the estimate of effect

¹ Downgraded one level for study limitations because the majority of the components of risk of bias assessment for the one included study were judged as 'unclear' risk

² Downgraded one level because there was only one study included and inconsistency with other studies could not be assessed

³ Downgraded one level for imprecision because of small sample size and wide confidence interval including potential for a decrease and increase in the outcome

⁴ There was only one study included, thus publication bias could not be assessed

⁵ Downgraded one level for indirectness

BACKGROUND

Description of the condition

Nephrotic syndrome is a condition characterised by generalised oedema (swelling caused by fluid retention), massive proteinuria (protein in the urine), hypoalbuminaemia (abnormally low levels of albumin in blood serum), and hypercholesterolaemia (high levels of cholesterol in the blood). The severity of oedema varies among patients, and severe oedema, sometimes called anasarca, may result in prolonged immobilisation, thus predisposing patients to thromboembolic events (blood clots), osteoporosis (loss of bone density) and infection. Traditionally, diuretics (drugs to promote urine production) have been the mainstay of treatment for oedema but even large doses of potent diuretics or combinations of diuretics sometimes have no effect (Brater 1985).

Description of the intervention

Intravenous infusion of albumin has been advocated as an effective method of treating oedema resulting from nephrotic syndrome. Albumin has been used for people whose oedema is resistant to diuretics and for low blood volume. Several doses in series are usually given by infusion over a few consecutive days. Albumin may be given concurrently with, or instead of, diuretics. Several potential mechanisms for diuretic resistance have been identified. They include decreased delivery of diuretic to the urinary site of action due to renal insufficiency or hypoalbuminaemia (Beermann 1977; Rose 1977); binding of diuretic to urinary albumin (Kirchner 1990; Voelker 1989); or decreased sensitivity of the nephron to diuretic (Kirchner 1991; Villeneuve 1986). The quantitative contribution of each potential mechanism has not been delineated. Thus, clinicians have been frustrated by the lack of sufficient information to develop therapeutic strategies targeted toward the pathophysiology of diuretic resistance.

How the intervention might work

The pathophysiology of oedema in nephrotic syndrome is explained by two theories, popularly described as the 'underfill' and 'overfill' theories. The classical hypothesis relates renal sodium and water retention to hypovolaemia (decreased blood plasma volume; underfill theory). Loss of protein by the kidneys decreases plasma albumin concentration. The resulting reduction of plasma oncotic pressure increases fluid movement out of the vessels into the interstitium decreasing plasma volume (Humphreys 1992). Reduction in plasma volume activates the sympathetic nervous system (Herman 1989) and the renin-angiotensin-aldosterone axis (Meltzer 1979; Tulassay 1989), promoting vasopressin secretion (Usberti 1984), and suppressing atrial natriuretic peptide (ANP) (Perico 1993). Interaction of these systems stimulates renal water and sodium retention (Tulassay 1986; Tulassay 1987; Usberti 1995). In children with nephrotic syndrome, clinical symptoms of hypovolaemia are occasionally observed but most studies failed to document decreased plasma volumes. One study that did document and report decreased plasma volumes (Saatci 1996) suggested albumin infusion might increase osmotic pressure and draw extracellular fluid back into the vascular compartment, reducing oedema. An increasing number of studies in adult patients and in experimental animal models of nephrotic syndrome indicate that intrarenal mechanisms are responsible for sodium and water retention (Brown 1982; Brown

1984; Brown 1985; Dorhout Mees 1984; Geers 1984; Ichikawa 1983) (overfill theory). In this situation, albumin infusion may have a minimal effect on oedema. Albumin infusion has potentially harmful effects including fluid overload, pulmonary oedema (Vande Walle 2004) and may increase death in critically ill patients (Roberts 2011).

Why it is important to do this review

Human albumin is expensive to administer, potentially harmful, and any benefits may be short-lived. It is therefore important that human albumin use be evaluated to better understand its efficacy and safety for treating oedema associated with nephrotic syndrome.

OBJECTIVES

This review aimed to examine the benefits and harms of human albumin infusion for treating oedema associated with nephrotic syndrome.

METHODS

Criteria for considering studies for this review

Types of studies

All randomised controlled trials (RCTs) and quasi-RCTs (RCTs in which allocation to treatment was obtained by alternation, use of alternate medical records, date of birth or other predictable methods) were included. We intended to include cross-over studies where they reported outcomes that we considered to not be subject to a period effect. If the period effect existed, we would obtain the first period data from the studies' corresponding authors.

Types of participants

Patients of all ages with oedema due to nephrotic syndrome from any cause were included.

Types of interventions

1. Human albumin infusion compared with placebo or no intervention. If other treatments are used, such as salt restriction, both groups should receive these interventions.
2. Human albumin with diuretics compared with diuretics alone.
3. Human albumin compared with diuretics.
4. Human albumin compared with other treatments.

Types of outcome measures

Primary outcomes

- Clinical improvement of oedema: weight loss; increased urine volume; reductions in ascites (measured by reduction in abdominal girth); pleural effusion (on chest X-ray); penile or scrotal oedema; serum sodium.

Secondary outcomes

- Death: death (all causes); death due to kidney failure
- Acute kidney injury (AKI) and need for acute dialysis treatment
- Kidney function measures: serum creatinine (SCr); glomerular filtration rate (GFR) (measured as creatinine clearance (CrCl) or estimated GFR)
- Quality of life (QoL)

- Adverse events: plasma volume overload; hypertension; pulmonary oedema; need for dialysis or ventilation to control plasma volume overload.

Search methods for identification of studies

Electronic searches

We searched the [Cochrane Kidney and Transplant Register of Studies](#) up to 23 June 2019 through contact with the Information Specialist using search terms relevant to this review. The Register contains studies identified from the following sources:

1. Monthly searches of the Cochrane Central Register of Controlled Trials (CENTRAL)
2. Weekly searches of MEDLINE OVID SP
3. Handsearching of kidney-related journals and the proceedings of major kidney conferences
4. Searching of the current year of EMBASE OVID SP
5. Weekly current awareness alerts for selected kidney and transplant journals
6. Searches of the International Clinical Trials Register (ICTRP) Search Portal and ClinicalTrials.gov.

Studies contained in the Register are identified through searches of CENTRAL, MEDLINE, and EMBASE based on the scope of Cochrane Kidney and Transplant. Details of search strategies, as well as a list of handsearched journals, conference proceedings and current awareness alerts, are available on the [Cochrane Kidney and Transplant website](#).

See [Appendix 1](#) for search terms used in strategies for this review.

Searching other resources

1. Reference lists of review articles, relevant studies and clinical practice guidelines.
2. Letters seeking information about unpublished or incomplete studies to investigators known to be involved in previous studies.
3. Grey literature sources (e.g. abstracts, dissertations and theses), in addition to those already included in the Cochrane Kidney and Transplant Register of Studies, was not searched.

Data collection and analysis

Selection of studies

The search strategy ([Appendix 1](#)) was used to obtain titles and abstracts of studies that may be relevant to the review. The titles and abstracts were screened independently by two authors (ASA and FJ), who discarded studies that were not applicable; however, studies and reviews that might include relevant data or information on potential studies for inclusion were retained initially. Two authors independently assessed retrieved abstracts, and where necessary the full text of these studies to determine whether they satisfy the inclusion criteria.

Data extraction and management

Data extraction was carried out independently by two authors (NJA and KYC) using standard data extraction forms. If more than one publication of one study was found, reports would have been grouped together and the publication with the most complete data would have been used in the analyses. Where

relevant outcomes were only published in earlier versions, this data would have been used. Any discrepancy between published versions is highlighted in the [Characteristics of included studies](#). Any further information required from the original author was requested by written correspondence and any relevant information obtained in this manner would have been included in the review. Disagreements were resolved in consultation with JJH and ASA.

Assessment of risk of bias in included studies

Two review authors (NJA and KYC) independently assessed risk of bias of the included study using the risk of bias assessment tool in the *Cochrane Handbook for Systematic Reviews of Interventions* ([Higgins 2011](#)). We resolved any disagreement by discussion and consulting a third person (ASA) and sometimes a fourth person (JJH).

The following items were independently assessed by two authors (NJA and KYC) using the risk of bias assessment tool ([Higgins 2011](#)) ([Appendix 2](#)).

- Was there adequate sequence generation (selection bias)?
- Was allocation adequately concealed (selection bias)?
- Was knowledge of the allocated interventions adequately prevented during the study?
 - * Participants and personnel (performance bias)
 - * Outcome assessors (detection bias)
- Were incomplete outcome data adequately addressed (attrition bias)?
- Are reports of the study free of suggestion of selective outcome reporting (reporting bias)?
- Was the study apparently free of other problems that could put it at a risk of bias?

Measures of treatment effect

For dichotomous outcomes (e.g. acute kidney injury and frequency of relapse), we would have reported the pooled risk ratio (RR) with 95% Confidence intervals (95% CI). If studies reported death data, we would have reported hazard ratios (HR) with its 95% CI.

For continuous scales of measurements (e.g. weight loss), we recorded the mean, standard deviation (SD) and total number of participants in both the treatment and control groups and calculated the mean difference (MD) with the 95% CI. If studies reported the MD and standard error (SE), we would have calculated the standard deviation (SD). If studies had reported outcomes on different scales (e.g. QoL) and we judged them to have reported the same thing, we would have used standardised mean difference (SMD)

Unit of analysis issues

For some of our pre-specified outcomes the intervention was not suitable for cross-over study design (death and quality of life). Although nephrotic syndrome is a relatively chronic stable condition the course of oedema is not, occurring intermittently and sometimes resolving spontaneously. The half-life of albumin is 17 to 19 days and clinical effects have been reported to last for 27 days ([Nicholson 2000](#); [Peters 1996](#)). It is likely that the duration of these clinical effects would be shorter in a dose-dependent fashion in people with albuminuria but there is no data available on what a suitable washout period might be in the presence of

albuminuria. Since there was no literature to guide us we judged a carry-over effect might be expected for any study with a washout period shorter than the half-life of albumin. Secondly, because of the fluctuating nature of the disease, particularly in adults, if the washout period is long it is possible that during this period oedema might spontaneously resolve, improve or fluctuate or alternatively improvement or deterioration could be due to the treatment administered during the first period. If data were available for the first period of a cross-over study, we would have included this in the meta-analysis and analyse the data as a parallel study. We attempted to contact all authors for data from the first period data but none of the authors responded to our request.

We do not anticipate that cluster RCTs would be used. If we had encountered any cluster RCTs we would have included these along with any individually randomised studies, if there was little heterogeneity, and analysed them at the level of the individual by making adjustments for cluster size. If the inter-cluster correlation coefficient (ICC) was not described, we would have estimated this based on other sources in the literature. We would then have conducted sensitivity analyses based on the source of the ICC. If we had encountered any studies with multiple treatment groups, we planned to analyse only two of the groups.

Dealing with missing data

We intended to note levels of attrition and would have attempted to contact authors for missing data. If data were not available we would have described the level of attrition, attempted to explore whether there was any pattern associated with missing data and analyse the available data. However, we did attempt to contact the authors of all the excluded cross-over studies to determine whether we could get data for the first period but did not receive any responses. We did not intend to impute missing data. If SDs were missing from continuous data, and we were unable to obtain these from the original investigators, we intended to calculate these if sufficient data are available.

As far as possible, we would have conducted all analyses on an intention-to-treat basis.

Assessment of heterogeneity

If we had included more than one study heterogeneity would have been assessed using a χ^2 test on $N-1$ degrees of freedom, with an alpha of 0.05 used for statistical significance and with the I^2 test (Higgins 2003). I^2 values of 25%, 50% and 75% correspond to little or none, moderate and considerable levels of heterogeneity.

Assessment of reporting biases

We intended to construct funnel plots for analyses if there were more than 10 or more studies and if funnel plot asymmetry was found, we would have attempted to explore if this effect was due to publication bias or due to other reasons. Where reports were available, we would have checked for selective outcomes reporting by comparing study protocols or methods sections with results presented in final study reports.

Data synthesis

If we had included more than one study, we planned that data would be pooled using the random effects model but we would have performed a sensitivity analysis using fixed effect model

to ensure robustness of the model chosen and susceptibility to outliers.

Subgroup analysis and investigation of heterogeneity

If we had sufficient data, we intended to conduct subgroup analyses to explore possible sources of heterogeneity. We planned to carry out subgroup analyses for the following subgroups.

1. Participant age or disease type (predominantly children and predominantly adult) or (predominantly minimal change nephrotic syndrome and other causes)
2. Dose of albumin (low dose, high dose, low total dose, high total dose)
3. Timing of use of diuretics (early, such as before or at the beginning of the albumin infusion; or late, such as after midway through infusion or at completion).

We intended to look at the primary outcome and adverse effects for each of these subgroups, and where possible, the risk difference (RD) with 95% CI would be calculated.

Sensitivity analysis

We intended to do a sensitivity analysis testing the inclusion of cross-over studies with relatively long washout periods which might have been long enough to take into account the half-life of albumin in people with albuminuria. We would have explored the effects of study quality by conducting sensitivity analyses. We would have also looked at the effect of using fixed-effect versus random-effects models for meta-analysis.

'Summary of findings' tables

We presented the main results of the review in a 'Summary of findings' table. These tables present key information concerning the certainty (quality) of the evidence, the magnitude of the effects of the interventions examined, and the sum of the available data for the main outcomes (Schunemann 2011a). The 'Summary of findings' table also include an overall grading of the evidence related to each of the main outcomes using the GRADE (Grades of Recommendation, Assessment, Development and Evaluation) approach (GRADE 2008; GRADE 2011). The GRADE approach defines the quality of a body of evidence as the extent to which one can be confident that an estimate of effect or association is close to the true quantity of specific interest. The quality of a body of evidence involves consideration of within-trial risk of bias (methodological quality), directness of evidence, heterogeneity, precision of effect estimates and risk of publication bias (Schunemann 2011b). We presented the following outcomes in the 'Summary of findings' tables.

1. Weight loss rate
2. Other clinical improvement parameters
3. Death
4. Kidney function
5. Quality of life
6. Adverse effects

RESULTS

Description of studies

Results of the search

Through database searching, we identified a total 294 of records, from CENTRAL (195), EMBASE (88) and the Cochrane Kidney and

Transplant Specialised Register (11). Two hundred fifty-five records were left after duplicates were removed; 238 records were excluded after title and abstract screening and the remaining 17 records (12 studies) were assessed for eligibility. After full-text review, we identified one study that met our inclusion criteria ([Lee 2000a](#)) and 11 studies were excluded ([Figure 1](#)). We did not find any ongoing studies.

Figure 1. Study flow diagram.

Included studies

We included one study ([Lee 2000a](#)). The study recruited all children with nephrotic syndrome with clinical or biopsy evidence of minimal change nephrotic syndrome. The report included 26 children with nephrotic syndrome who went into remission after 8 weeks of treatment. Remission was defined as an absence of albumin in the urine for three consecutive days.

The treatment group received 1 g/kg of albumin and 2 mg/kg of furosemide for 1 to 3 days and the control group received an equivalent amount of 5% dextrose solution. Both groups received steroids and diuretics as part of the treatment regime. Clinical outcomes were reported and patients were followed up for one year.

Outcomes measured were weight change, blood pressure, serum sodium, urine protein/creatinine ratio, mean time to complete remission, recurrence rate at 12 months and number of relapses at 12 months were reported. Percent weight loss was reported at days 1, 4, 7 and 10. However we were not able to report the quantitative data for this outcome due to lack of clarity of the data. Where outcomes were reported at several time points we have reported those time points we judged to be the most clinically relevant. Serum sodium was not included in our prespecified measures of clinical improvement but we considered this to be a

valid measure and made a post hoc decision to include it. We had prespecified hypertension as an adverse effect. We judged included blood pressure as a measure that might give some information on hypertension.

Excluded studies

We excluded 11 studies that did not meet our inclusion criteria. In one study the participants had oedema but not due to nephrotic syndrome ([Chalasani 2001](#)) and nine met our inclusion criteria for participants and interventions but were excluded because they were cross-over studies ([Akciek 1995a](#); [Dharmaraj 2009](#); [Fliser 1999](#); [Ghafari 2005](#); [Ghafari 2011](#); [Kanitkar 2010](#); [Na 2001](#); [Phakdeekitcharoen 2012](#); [Sjostrom 1989](#)). Raw data on the study variables of patients on the first period (before cross-over) could not be obtained within the papers or from the authors. The remaining study compared the effect of albumin plus an oral fish protein extract with albumin alone on serum albumin levels in children with NS and hypoalbuminaemia. The authors did not state whether the children had oedema. Indeed children with 'anasarca oedema' were excluded ([Herumuryawan 2017](#)). Further details are found in the table [Characteristics of excluded studies](#).

The full text publications for two abstracts published in Paediatric Nephrology ([Kanitkar 2010](#)) and Nephrology ([Ghafari 2005](#)) could not be located.

Risk of bias in included studies

The assessment of risk of bias of the included study is summarised in [Figure 2](#)

Figure 2. Risk of bias graph: review authors' judgements about each risk of bias item presented as percentages across all included studies.

The authors reported [Lee 2000a](#) as a RCT. They included all patients with a diagnosis of minimal change nephrotic syndrome. However, this report includes those who went into complete remission by 8 weeks of treatment. Therefore, the actual number recruited and randomised, the number who did not go into remission, if any, and when in the course of the study these children were excluded is unclear.

Allocation

The authors did not describe the method of generation of the random sequence and the method of allocation concealment. We considered the risk of bias for both these two domains be unclear.

Blinding

The control group received a volume of dextrose equivalent to the volume of albumin in the intervention group. However, it is unclear whether this was intended as a placebo and if so how they blinded the furosemide. Since the two solutions look different we judged this to be unlikely. No other attempt at blinding was described.

For reported outcomes that were laboratory-test results we judged that the lack of blinding would be unlikely to have an influence on these outcomes and judged these to be at low risk of bias whereas clinical outcomes were judged at high risk.

Incomplete outcome data

Although it is unclear whether some patients were excluded either pre- or post-intervention we considered this to be an unclear risk to selection bias. All those reported had complete follow-up data to 12 months. We judged attrition bias to be at low risk.

Selective reporting

The study was not registered on a clinical trials register and the protocol was not available and not all expected outcomes were reported. Therefore we judged this domain to be at high risk of bias

Other potential sources of bias

None detected.

Effects of interventions

See: [Summary of findings for the main comparison Human albumin infusion plus furosemide compared to glucose for treating people with nephrotic syndrome](#)

Human albumin plus furosemide versus placebo or no treatment

See: [Summary of findings for the main comparison](#) for the main findings of the comparison of human albumin infusion plus furosemide compared to glucose.

Clinical Improvement of oedema

Weight loss

[Lee 2000a](#) reported a greater weight loss in the albumin group in the first few days but stated that there no overall significant difference in weight loss up to Day 10. However, we were unable to report the data due to lack of clarity. Data reported in the text of the report was not consistent with data shown in the figure, thus we could not confirm the authors' statements. We judged the evidence to be of very low certainty.

Other measures of clinical improvement

[Lee 2000a](#) did not report any of our other prespecified measures of clinical improvement. However the authors reported serum

sodium. It is uncertain whether albumin improves serum sodium when compared with an equal volume of dextrose (1 study, 26 participants: MD 2.00 mEq/L, 95% CI -0.09 to 4.09; very low certainty evidence).

Death

Lee 2000a reported no deaths over the 12 months of follow-up.

Acute kidney injury

Acute kidney injury or treatment with acute dialysis was not reported.

Kidney function measures

Lee 2000a reported spot urine protein/creatinine ratio as a measure of kidney function. Although this was not included in our prespecified measures of kidney function measures we considered this a valid measure. There was a higher mean spot urine protein/creatinine ratio in the albumin treated group at Day 2 and 4 after treatment but not on Day 6, 8 and 10,

Quality of life

Lee 2000a did not report quality of life.

Adverse events

The only prespecified adverse effect reported in Lee 2000a was blood pressure. This was reported as mean systolic and diastolic pressure in each group. The authors defined hypertension but did not give data on its incidence. It is uncertain whether albumin compared with dextrose results in any change in systolic blood pressure (Day 1: MD 3.00 mmHg, 95% CI -2.54 to 8.54; Day 3: MD 1.00 mmHg, 95% CI -3.85 to 5.85; Day 5: MD 2.00 mmHg, 95% CI -3.52 to 7.52; very low certainty evidence).

Other non-prespecified outcomes reported

Lee 2000a reported that remission duration to first relapse was shorter in the albumin-treated group than the control group (94 (63.5) vs 190 (106.4) days, (MD -96.00, 95% CI -163.36 to -28.64, one study 26 patients). There was no difference between the two groups in the frequency of relapse within one year of remission (77% versus 46%, RR 1.67, 95% CI 0.86 to 3.22). We judged the evidence to be of low certainty. Other reported outcomes in Lee 2000a can be found in Table 1.

Other pre-specified outcomes

Lee 2000a did not report any of the following outcomes: increased urine volume; reductions in ascites (measured by reduction in abdominal girth); pleural effusion (on chest X-ray); penile or scrotal oedema; serum creatinine (SCr); glomerular filtration rate (GFR) (measured as creatinine clearance (CrCl) or estimated GFR); plasma volume overload; pulmonary oedema; need for dialysis or ventilation to control plasma volume overload.

Other pre-specified comparisons

The following comparisons have not been investigated:

1. Human albumin compared to placebo or no treatment
2. Human albumin with diuretics compared with diuretics alone
3. Human albumin compared with diuretics.

DISCUSSION

Summary of main results

We found only one small study that reported very few of our pre-specified outcomes (Lee 2000a). The limited data available does not show any clinical improvement with albumin and furosemide in children with minimal change nephrotic syndrome. They also showed no difference in blood pressure or serum sodium concentrations. There is uncertainty about all these outcomes.

We excluded nine small studies involving a total of 117 participants evaluating the role of human albumin in treating nephrotic syndrome that were cross-over studies. The washout period for seven studies was between 24 and 72 hours. Ghafari 2005) did not report the washout period and Phakdeekitcharoen 2012 recorded a washout period of 2 weeks. These excluded studies all reported only one of our prespecified primary outcomes (urine volume) and one reported a secondary outcome (GFR). We were unable to obtain data for the first period for any of the cross-over studies.

Overall completeness and applicability of evidence

There were no studies on the use of human albumin in adults with nephrotic syndrome. Therefore, there is almost no evidence for or against this commonly used intervention in the management of nephrotic syndrome and what little evidence that does exist does not support or oppose its use. The one included study did not report most of our secondary outcomes. We are unable to explain why the treatment group had a shorter duration of remission except that in this small study a chance finding is possible. We found nine cross-over studies, all of which were excluded because we could not obtain data for the first period. In addition the washout period was judged to be too short. In addition, only one of the nine studies, which provided individual patient data, did not have statistical information available to appropriately incorporate these into a meta-analysis.

Quality of the evidence

Of the five pre-specified outcomes we chose to assess using the GRADE approach only one was reported, weight loss, a measure of our primary outcome, clinical improvement. We made a post hoc decision that serum sodium was also a valid measure of clinical improvement and have included this in the summary of findings table. We judged both the reported measures of clinical improvement to be very low certainty evidence. We downgraded for study limitations, imprecision and single study results since inconsistency with other studies could not be assessed. We judged that blood pressure might also provide information on whether hypertension was an adverse effect and incorporated systolic blood pressure into the summary of findings table. We judged this to be very low certainty evidence. We downgraded for study limitations, imprecision, indirectness and inconsistency (since this could not be assessed as there was only one study).

Potential biases in the review process

All studies were independently assessed by two review authors as recommended in Cochrane Handbook for Systematic Review of Intervention (Higgins 2011). Although we made an effort to identify relevant studies through different databases and resources, it is possible that relevant but unpublished data may have been missed. We also attempted to contact all included and excluded

study authors for further information. We believe this intervention would be unlikely to be subject to publication bias due to industry involvement.

One potential bias in this review is that we excluded cross-over studies because we judged the washout period to be too short and we were not able to obtain data for the first period in order to analyse these as parallel studies. We found that the nine studies we excluded because they were cross-over design all met our inclusion criteria for patients and intervention and they all reported our primary outcome. We judged that a cross-over design would be possible for this intervention for short-term outcomes only if the washout period were sufficiently long to exclude any possible carry over effect of the albumin infusion. A potential weakness is that there is insufficient data to support our choice of washout period in people with albuminuria due to nephrotic syndrome and our choice is a judgement rather than based on evidence. If the washout period is considerable, fluctuations in the albumin levels and degree of oedema that occur with this condition could be shorter than a period effect. Therefore, in the course of excluding these studies we have excluded a large body of currently available research. Even if applicable, analysis of all but one of these cross-over studies would not have been possible because of insufficient statistical information to incorporate them appropriately into a meta-analysis.

We reported outcomes that were not part of our pre-specified outcomes and thus possibly could introduce bias to the review process. However, we judged that these outcomes might reveal important adverse effects of the intervention that might warrant further research.

Agreements and disagreements with other studies or reviews

Our search did not identify any other review evaluating clinical improvement of oedema due to nephrotic syndrome. However we found a recent systematic review and meta-analysis undertaken to look at the efficacy and safety of human albumin solutions as part of fluid volume expansion and resuscitation in critically unwell adults with sepsis of any severity (Patel 2014). The findings of this review are not relevant to our patient group. However, of note, the authors reported that human albumin did not result in increased death. Epidemiological studies examining the long term safety of human albumin including serious adverse events have not reported any deaths thought to be related to the use of albumin and non-fatal adverse events appear to be rare (Vincent 2003; von Hoegen

2001). Two Cochrane reviews on the effect of human albumin in other populations (Roberts 2011; Fortin 2007) do not report non-fatal adverse events and there was uncertainty about the effect of albumin on death.

AUTHORS' CONCLUSIONS

Implications for practice

Since we found only one small study we are unable to draw any conclusions regarding the use of human albumin with or without diuretics in nephrotic syndrome.

Implications for research

Due to wide current usage there is a pressing need to establish the role of human albumin in nephrotic syndrome. The role of human albumin alone needs to be established against no treatment or other supportive care. The role of low sodium albumin preparations also needs to be established. Large well designed RCTS are required to compare human albumin with other interventions currently used in the management of oedema in nephrotic syndrome, such as furosemide and other diuretics, either in combination with human albumin or alone. It is important that participants be adequately described in terms of their albumin status and this includes quantification of the degree of the oedema and albuminuria at study entry, preferably by using 24-hour urine protein-creatinine index. The presence of any pre-existing CKD, dialysis or pre-dialysis measure, and coexisting liver cirrhosis also should be documented. We recommend that the primary outcome be reported in a variety of ways and that urine volume be reported in such a way that it can be incorporated into a meta-analysis. Adverse effects of albumin infusions are important since albumin infusions can be associated with morbidity (pulmonary oedema) and death. Therefore, we recommend that adverse effects need to be carefully assessed in any studies of albumin infusions for any indication.

ACKNOWLEDGEMENTS

We wish to thank Elaine Beller for helpful statistical advice. We also thank the referees for the comments and feedback during the preparation of this review. We thank Hyeong Sik Ahn and team from Cochrane Korea for with help in translation of an included study written in Korean language. Zainal Darus, Nephrology Unit, and Nik Zainal Nik Ismail formerly of Dept of Paediatrics, Universiti Sains Malaysia were involved in the development of the concept and had input into the protocol.

REFERENCES

References to studies included in this review

Lee 2000a {published data only}

Lee SJ, Jeong JA, Hwang SJ, Lee SJ. The effect of human albumin infusion on the clinical course of steroid sensitive nephrotic syndrome. *Journal of the Korean Society of Pediatric Nephrology* 2000;**4**(2):102-10. [CENTRAL: CN-01045918]

References to studies excluded from this review

Akcicek 1995a {published data only}

Akcicek F, Basci A, Ozbasli C, Yalniz MT, Dorhout Mees EJ. Diuretic effects of intravenous albumin (A) furosemide (F) and the combination (F+A) in the nephrotic syndrome (NS) [abstract]. 12th International Congress of Nephrology; 1993 Jun 13-18; Jerusalem, Israel. 1993:87. [CENTRAL: CN-00550483]

Akcicek F, Yalniz T, Basci A, Ok E, Mees EJ. Diuretic effect of frusemide in patients with nephrotic syndrome: is it potentiated by intravenous albumin?. *BMJ* 1995;**310**(6973):162-3. [MEDLINE: 7833755]

Chalasani 2001 {published data only}

Chalasani N, Gorski JC, Horlander JC Sr, Craven R, Hoen H, Maya J, et al. Effects of albumin/furosemide mixtures on responses to furosemide in hypoalbuminemic patients. *Journal of the American Society of Nephrology* 2001;**12**(5):1010-6. [MEDLINE: 11316860]

Dharmaraj 2009 {published data only}

Dharmaraj R, Hari P, Bagga A. Randomized cross-over trial comparing albumin and frusemide infusions in nephrotic syndrome. *Pediatric Nephrology* 2009;**24**(4):775-82. [MEDLINE: 19142668]

Hari P, Bagga A. Does albumin infusion augment frusemide induced diuresis in edematous children with nephrotic syndrome? [abstract no: M-PO-0834]. 4th World Congress of Nephrology. 19th International Congress of the International Society of Nephrology (ISN); 2007 Apr 21-25; Rio de Janeiro, Brazil. Rio de Janeiro, Brazil, 2007:285.

Fliser 1999 {published data only}

Fliser D, Zurbruggen I, Mutschler E, Bischoff I, Nussberger J, Franek E, et al. Coadministration of albumin and furosemide in patients with the nephrotic syndrome. *Kidney International* 1999;**55**(2):629-34. [MEDLINE: 9987087]

Fliser D, Zurbruggen I, Mutschler E, Ritz E. Furosemide and albumin coadministration in patients with the nephrotic syndrome [abstract]. *Nephrology Dialysis Transplantation* 1997;**12**(9):A115. [CENTRAL: CN-00261361]

Ghafari 2005 {published data only}

Ghafari A, Makhdooni K, Ahmadpoor P, Khalifani M, Darazam IA. Does albumin potentiate the diuretic effect of furosemide in nephrotic patients? [abstract no: T-PO30043]. *Nephrology* 2005;**10**(Suppl):A170.

Ghafari 2011 {published data only}

Ghafari A, Mehdizadeh A, Alavi-Darazam I, Rahimi E, Kargar C, Sepehrvand N. Co-administration of albumin-furosemide in patients with the nephrotic syndrome. *Saudi Journal of Kidney Diseases & Transplantation* 2011;**22**(3):471-5. [MEDLINE: 21566302]

Herumuryawan 2017 {published data only}

Herumuryawan M, Hardaningsih G. Effect of striped snakehead fish (*ophiocephalusstriatus*) extract supplement pills compared to human albumin infusion on albumin serum, lipid profile, malondialdehyde and IL-8 serum level on nephrotic syndrome. *Pakistan Journal of Medical & Health Sciences* 2017; Vol. 11, issue 4:1601-6. [EMBASE: 620394056]

Kanitkar 2010 {published data only}

Kanitkar M, Garg A, Venkateshwar V. 5% vs 20% albumin in nephrotic syndrome [abstract no: 60]. *Pediatric Nephrology* 2010;**25**(9):1806-7. [EMBASE: 70438162]

Na 2001 {published data only}

Jeon US, Earm JH, Kim DH, Huh WS, Han JS, Kim S, et al. Does albumin infusion enhance the diuretic action of furosemide in nephrotic syndrome? [abstract no: A0424]. *Journal of the American Society of Nephrology* 1997;**8**(Program & Abstracts):88-9A. [CENTRAL: CN-00445917]

Na KY, Han JS, Kim YS, Ahn C, Kim S, Lee JS, et al. Does albumin preinfusion enhance the diuretic action of furosemide in patients with the nephrotic syndrome? [abstract no: A0590]. *Journal of the American Society of Nephrology* 2000;**11**(Sept):109A. [CENTRAL: CN-00644262]

Na KY, Han JS, Kim YS, Ahn C, Kim S, Lee JS, et al. Does albumin preinfusion potentiate diuretic action of furosemide in patients with nephrotic syndrome?. *Journal of Korean Medical Science* 2001;**16**(4):448-54. [MEDLINE: 11511790]

Phakdeekitcharoen 2012 {published data only}

Phakdeekitcharoen B, Boonyawat K. The added-up albumin enhances the diuretic effect of furosemide in patients with hypoalbuminemic chronic kidney disease: a randomized controlled study. *BMC Nephrology* 2012;**13**:92. [MEDLINE: 22931630]

Sjostrom 1989 {published data only}

Sjostrom PA, Odland BG, Beermann BA, Karlberg BE. Pharmacokinetics and effects of frusemide in patients with the nephrotic syndrome. *European Journal of Clinical Pharmacology* 1989;**37**(2):173-80. [MEDLINE: 2792172]

Additional references

Beermann 1977

Beermann B, Dalen E, Lindström B. Elimination of furosemide in healthy subjects and in those with renal failure. *Clinical Pharmacology & Therapeutics* 1977;**22**(1):70-8. [MEDLINE: 872498]

Brater 1985

Brater DC. Resistance to loop diuretics. Why it happens and what to do about it. *Drugs* 1985;**30**(5):427-43. [MEDLINE: 3905337]

Brown 1982

Brown EA, Markandu ND, Sagnella GA, Squires M, Jones BE, MacGregor GA. Evidence that some mechanism other than the renin system causes sodium retention in nephrotic syndrome. *Lancet* 1982;**2**(8310):1237-40. [MEDLINE: 6128546]

Brown 1984

Brown EA, Markandu ND, Sagnella GA, Jones BE, MacGregor GA. Lack of effect of captopril on the sodium retention of the nephrotic syndrome. *Nephron* 1984;**37**(1):43-8. [MEDLINE: 6371561]

Brown 1985

Brown EA, Markandu N, Sagnella GA, Jones BE, MacGregor GA. Sodium retention in nephrotic syndrome is due to an intrarenal defect: evidence from steroid-induced remission. *Nephron* 1985;**39**(4):290-5. [MEDLINE: 3982574]

Dorhout Mees 1984

Dorhout Mees EJ, Geers AB, Koomans HA. Blood volume and sodium retention in the nephrotic syndrome: a controversial pathophysiological concept. *Nephron* 1984;**36**(4):201-11. [MEDLINE: 6369161]

Fortin 2007

Fortin PM, Bassett K, Musini VM. Human albumin for intradialytic hypotension in haemodialysis patients. *Cochrane Database of Systematic Reviews* 2010, Issue 11. [DOI: [10.1002/14651858.CD006758.pub2](https://doi.org/10.1002/14651858.CD006758.pub2)]

Geers 1984

Geers AB, Koomans HA, Boer P, Dorhout Mees EJ. Plasma and blood volumes in patients with the nephrotic syndrome. *Nephron* 1984;**38**(3):170-3. [MEDLINE: 6493410]

GRADE 2008

Guyatt GH, Oxman AD, Vist GE, Kunz R, Falck-Ytter Y, Alonso-Coello P, et al. GRADE: an emerging consensus on rating quality of evidence and strength of recommendations. *BMJ* 2008;**336**(7650):924-6. [MEDLINE: 18436948]

GRADE 2011

Guyatt G, Oxman AD, Akl EA, Kunz R, Vist G, Brozek J, et al. GRADE guidelines: 1. Introduction-GRADE evidence profiles and summary of findings tables. *Journal of Clinical Epidemiology* 2011;**64**(4):383-94. [MEDLINE: 21195583]

Herman 1989

Herman PJ, Sawin LL, DiBona GF. Role of renal nerves in renal sodium retention of nephrotic syndrome. *American Journal of Physiology* 1989;**256**(5 Pt 2):F828-9. [MEDLINE: 2719117]

Higgins 2003

Higgins JP, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses. *BMJ* 2003;**327**(7414):557-60. [MEDLINE: 12958120]

Higgins 2011

Higgins JPT, Green S (editors). *Cochrane Handbook for Systematic Reviews of Interventions* Version 5.1.0 [updated March 2011]. The Cochrane Collaboration, 2011. Available from www.cochrane-handbook.org.

Humphreys 1992

Humphreys MH, Valentin J, Qui C, Ying W, Muldowney WP, Gardner DG. Underfill and overflow revisited: mechanisms of nephrotic edema. *Transactions of the American Clinical & Climatological Association* 1992;**104**:47-59. [MEDLINE: 1343447]

Ichikawa 1983

Ichikawa I, Rennke HG, Hoyer JR, Badr KF, Scholf N, Troy JL, et al. Role for intrarenal mechanisms in the impaired salt excretion of experimental nephrotic syndrome. *Journal of Clinical Investigation* 1983;**71**(1):91-103. [MEDLINE: 6848563]

Kirchner 1990

Kirchner KA, Voelker JR, Brater DC. Intratubular albumin blunts the response to furosemide—a mechanism for diuretic resistance in nephrotic syndrome. *Journal of Pharmacology & Experimental Therapeutics* 1990;**252**(3):1097-101. [MEDLINE: 2319462]

Kirchner 1991

Kirchner KA, Voelker JR, Brater DC. Binding inhibitors restore furosemide potency in tubule fluid containing albumin. *Kidney International* 1991;**40**(3):418-24. [MEDLINE: 1724020]

Meltzer 1979

Meltzer J, Keim HJ, Lyragh JH, Sealey JE, Jan KM, Chien S. Nephrotic syndrome: vasoconstriction and hypervolemic types indicated by renin-sodium profiling. *Annals of Internal Medicine* 1979;**91**(5):688-96. [MEDLINE: 496101]

Nicholson 2000

Nicholson JP, Wolmarans MR, Park GR. The role of albumin in critical illness. *British Journal of Anaesthesia* 2000;**85**(4):599-610. [MEDLINE: 11064620]

Patel 2014

Patel A, Laffan MA, Waheed U, Brett SJ. Randomised trials of human albumin for adults with sepsis: systematic review and meta-analysis with trial sequential analysis of all-cause mortality.[Erratum in: *BMJ*. 2014;349:g4850]. *BMJ* 2014;**349**:g4561. [MEDLINE: 25099709]

Perico 1993

Perico N, Remuzzi G. Edema of the nephrotic syndrome: the role of the atrial peptide system. *American Journal of Kidney Diseases* 1993;**22**(3):355-66. [MEDLINE: 8372830]

Peters 1996

Peters TJ. Metabolism: albumin in the body. All about albumin. Biochemistry, Genetics and Medical Applications. San Diego: Academic Press, 1996:188-250.

Roberts 2011

Roberts I, Blackhall K, Alderson P, Bunn F, Schierhout G. Human albumin solution for resuscitation and volume expansion in

critically ill patients. *Cochrane Database of Systematic Reviews* 2011, Issue 10. [DOI: [10.1002/14651858.CD001208.pub4](https://doi.org/10.1002/14651858.CD001208.pub4)]

Rose 1977

Rose HJ, O'Malley K, Pruitt AW. Depression of renal clearance of furosemide in man by azotemia. *Clinical Pharmacology & Therapeutics* 1977;**21**(2):141-6. [MEDLINE: 837633]

Saatci 1996

Saatci U, Ozen S, Ozdemi S, Aras T. Atrial natriuretic peptide in childhood nephrotic syndrome. *Nephrology Dialysis Transplantation* 1996;**11**(5):905. [MEDLINE: 8671924]

Schunemann 2011a

Schünemann HJ, Oxman AD, Higgins JP, Vist GE, Glasziou P, Guyatt GH. Chapter 11: Presenting results and 'Summary of findings' tables. In: Higgins JP, Green S (editors). *Cochrane Handbook for Systematic Reviews of Interventions* Version 5.1.0 [updated March 2011]. The Cochrane Collaboration, 2011. Available from www.cochrane-handbook.org.

Schunemann 2011b

Schünemann HJ, Oxman AD, Higgins JP, Deeks JJ, Glasziou P, Guyatt GH. Chapter 12: Interpreting results and drawing conclusions. In: Higgins JP, Green S (editors). *Cochrane Handbook for Systematic Reviews of Interventions* Version 5.1.0 [updated March 2011]. The Cochrane Collaboration, 2011. Available from www.cochrane-handbook.org.

Tulassay 1986

Tulassay T, Rascher W, Ganten D, Schärer K, Lang RE. Atrial natriuretic peptide and volume changes in children. *Clinical & Experimental Hypertension - Part A, Theory & Practice* 1986;**8**(4-5):695-701. [MEDLINE: 2944674]

Tulassay 1987

Tulassay T, Rascher W, Lang RE, Seyberth HW, Schärer K. Atrial natriuretic peptide and other vasoactive hormones in nephrotic syndrome. *Kidney International* 1987;**31**(6):1391-5. [MEDLINE: 2956451]

Tulassay 1989

Tulassay T, Rascher W, Shärer K. Intra- and extrarenal factors of oedema formation in the nephrotic syndrome. *Pediatric Nephrology* 1989;**3**(1):92-100. [MEDLINE: 2702097]

Usberti 1984

Usberti M, Federico S, Meccariello S, Ciancarruso B, Balletta M, Pecoraro C, et al. Role of plasma vasopressin in the impairment

of water excretion in nephrotic syndrome. *Kidney International* 1984;**25**(2):422-9. [MEDLINE: 6727137]

Usberti 1995

Usberti M, Gazotti R, Poiesi C, Avanzo L, Ghielmi S. Considerations on the sodium retention in nephrotic syndrome. *American Journal of Nephrology* 1995;**15**(1):38-47. [MEDLINE: 7872363]

Vande Walle 2004

Vande Walle J, Mael R, Raes A, Vandekerckhove K, Donckerwolcke R. ARF in children with minimal change nephrotic syndrome may be related to functional changes of the glomerular basal membrane. *American Journal of Kidney Diseases* 2004;**43**(3):399-404. [MEDLINE: 14981597]

Villeneuve 1986

Villeneuve JP, Verbeeck RK, Wilkinson GR, Branch RA. Furosemide kinetics and dynamics in patients with cirrhosis. *Clinical Pharmacology & Therapeutics* 1986;**40**(1):14-20. [MEDLINE: 3720175]

Vincent 2003

Vincent JL, Wilkes MM, Navickis RJ. Safety of human albumin--serious adverse events reported worldwide in 1998-2000. *British Journal of Anaesthesia* 2003;**91**(5):625-30. [MEDLINE: 14570782]

Voelker 1989

Voelker JR, Jameson DM, Brater DC. In vitro evidence that urine composition affects the fraction of active furosemide in the nephrotic syndrome. *Journal of Pharmacology & Experimental Therapeutics* 1989;**250**(3):772-8. [MEDLINE: 2778714]

von Hoegen 2001

von Hoegen I, Waller C. Safety of human albumin based on spontaneously reported serious adverse events. *Critical Care Medicine* 2001;**29**(5):994-6. [MEDLINE: 11378610]

References to other published versions of this review

Ho 2012

Ho JJ, Adnan AS, Darus Z, Nik Ismail NZ, Van Rostenberghe H. Human albumin infusion for treating oedema in people with nephrotic syndrome. *Cochrane Database of Systematic Reviews* 2012, Issue 3. [DOI: [10.1002/14651858.CD009692](https://doi.org/10.1002/14651858.CD009692)]

CHARACTERISTICS OF STUDIES

Characteristics of included studies [ordered by year of study]

Lee 2000a

Methods

- Study design: parallel RCT
- Study duration: 31 months (March 1997 to September 1999)
- Study follow-up period: at least one year (authors did not specified the duration of follow-up period but they had reported the relapse rate within 1 year after complete remission)

Lee 2000a (Continued)

Participants	<ul style="list-style-type: none"> Country: Korea Setting: single centre (Department of Pediatrics at the Ewha University Mokdong Hospital) Relevant health status: patients with nephrotic syndrome (biopsy proven or clinically compatible to minimal change nephrotic syndrome) Number: treatment group (13); control group (13) Mean age $\pm$ SD (years): treatment group (5.3 ± 2.9); control group (7.1 ± 4.2) Sex (M/F): treatment group (9/4); control group (11/2) Other relevant information (e.g. comorbidities): not reported Exclusion criteria: not reported
Interventions	<p>Treatment group</p> <ul style="list-style-type: none"> 20% human albumin 1g/kg and furosemide 2 mg/kg was administered IV for 1 to 3 days The timing of the furosemide in relation to the albumin infusion was not reported <p>Control group</p> <ul style="list-style-type: none"> Equivalent amount of 5% glucose solution was administered IV for 1 to 3 days <p>Co-interventions</p> <ul style="list-style-type: none"> All patients in both groups received corticosteroids and other diuretics (spironolactone and hydrochlorothiazide)
Outcomes	<ul style="list-style-type: none"> Daily percent weight loss rate at days 1 to 10 SBP and DBP at admission to day 5 Serum sodium concentration at admission, days 3 and 5 Serum potassium concentration at admission, days 3 and 5 Urine PCR at admission, days 2, 4, 6, 8, 10, 12, and 14 Time to complete remission in days: defined as negative urine protein for 3 consecutive days Recurrence of nephrotic syndrome after complete remission
Notes	<ul style="list-style-type: none"> The time to remission was reported as means. Although not clearly state but since the patients all had minimal change nephrotic syndrome and complete remission is expected, we judged that all patients went into remission Some results were imputed from the figures and graphs in the published paper Funding source for this study is not reported We attempted to contact the study authors for additional information via email. but have not received any reply The article is in Korean language and was translated by Cochrane Korea

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	<p>Quote: "We prospectively randomised the patients with nephrotic syndrome..."</p> <p>Comment: method of randomisation was not reported</p>
Allocation concealment (selection bias)	Unclear risk	There is no description on how allocation was implemented in the study although authors reported that patients were randomised to intervention and control groups
Blinding of participants and personnel (performance bias)	High risk	Control group received an equivalent dose of dextrose

Lee 2000a (Continued)

All outcomes		Comment: It is not clear whether this was intended to act as a placebo. However since the two solutions look different we judged there was no blinding of participants and personnel. No blinding of furosemide was described
Blinding of outcome assessment (detection bias) Clinical outcomes	Unclear risk	Insufficient information to permit judgement. Comment: Unclear for clinical outcomes.
Blinding of outcome assessment (detection bias) Laboratory outcomes	Low risk	We judged that outcomes reporting laboratory data would be at low risk of detection bias
Incomplete outcome data (attrition bias) All outcomes	Low risk	All patient outcome data reported Quote: "These 26 steroid-sensitive patients with nephrotic syndrome (13 in the albumin-treated group, 13 in the control group) that exhibited complete remission ... within 8 weeks of the steroid therapy were followed-up to observe the clinical course"
Selective reporting (reporting bias)	High risk	No protocol available. Not all the expected outcomes were reported
Other bias	Low risk	The study appears to be free of other sources of bias

DBP - diastolic blood pressure; IV - intravenously; M/F - male/female; PCR - protein-creatinine ratio; RCT - randomised controlled trial; SBP - systolic blood pressure; SD - standard deviation

Characteristics of excluded studies [ordered by study ID]

Study	Reason for exclusion
Akciek 1995a	A total of 12 participants with nephrotic syndrome with minimal lesions or a well preserved glomerular histology on kidney biopsy. Four participants were excluded from the study (one became temporarily anuric, two lost their oedema before completion, and one failed to comply). The study was conducted in Turkey and compared human albumin, furosemide, and a combination of these two interventions in three separate periods in random order and each separated by 48 hours. The statistical methods used in the analysis are unclear and it is unlikely that the study could be appropriately incorporated into a meta-analysis. This study was excluded because cross-over design was applied. In addition the washout period was judged too short. The raw data for the first period (before the cross-over) could not be obtained for meta-analysis.
Chalasani 2001	The study was a RCT that included 13 patients with cirrhosis of varying causes. Four intervention groups were involved: (1) albumin alone; (2) furosemide alone; (3) premixed furosemide and albumin; (4) separately infused furosemide and albumin
Dharmaraj 2009	The researchers enrolled 16 children with nephrotic syndrome and significant oedema (age range: 3 to 18 years). They received either a combination of human albumin and furosemide infusion or furosemide infusion alone using a cross-over design. However the order of receiving treatments was randomised as a cluster. Two participants were withdrawn from the furosemide infusion group (one for metabolic alkalosis, one with peritonitis) and three from human albumin and furosemide infusion group (two because they were oedema free after the first intervention, and one for hypokalaemia). This study reported the primary outcome (urine volume) as a median and interquartile range and could therefore not be included in the quantitative analysis. However, this study was excluded because cross-over design with washout period of 48 hours was applied and all patients received the same treatment during the first period. This study includes one abstract report which we judged to be a preliminary report of 10 of the 16 patients reported in the full paper.

Study	Reason for exclusion
Fliser 1999	Nine nephrotic syndrome patients with oedema (mean age 48 years) were enrolled in the study. Six patients had biopsy proven membranous glomerulonephritis and three minimal change glomerulonephritis/focal segmental glomerulosclerosis). They were each given three treatments in a random order on the fifth, seventh and ninth study days: (1) human albumin, (2) furosemide, (3) human albumin and a furosemide infusion. The statistical data was insufficient for incorporating appropriately into a meta-analysis. However, this study was excluded because cross-over design was applied. The washout periods were reported as one day and raw data for the first period could not be obtained for meta-analysis. We judged the washout period to be too short.
Ghafari 2005	There were 10 patients with nephrotic syndrome on standardized sodium chloride intake, in random order on three separate days, administered IV for 60 minutes either, (1) 1mg/kg furosemide plus a sham infusion, (2) 1mg/kg furosemide plus 200 mL of a 20% solution of human albumin, or (3) sham infusion plus 200 mL of a 20% solution of human albumin. The appropriate paired statistic to incorporate this study into the meta-analysis was not reported. The washout period between periods was not reported. The outcomes were compared for three groups based on urinary volume, sodium excretion and GFR. However, this study was excluded because raw data for the first period before cross-over could not be obtained for meta-analysis. This is a conference abstract and the full article could not be obtained
Ghafari 2011	Included 10 nephrotic syndrome patients with proteinuria of more than 3.5 g in 24 hours, hypoalbuminaemia (plasma albumin level < 3.5 g/dL), oedema and hyperlipidaemia. There were three study arms each with two infusions, (1) 2 mg/kg furosemide plus sham (100 mL normal saline), (2) human albumin (0.5 g/kg) plus sham (100 mL normal saline); and (3) human albumin (0.5 g/kg) plus furosemide 2 mg/kg. The methods used for statistical analysis were inappropriate for incorporation of this study into a meta-analysis. However, this study was excluded because of its cross-over design with a washout period of 48 hours and raw data for the first period (before participant cross-over) could not be obtained for meta-analysis
Herumuryawan 2017	This parallel RCT involved 35 children with nephrotic syndrome aged 2 to 14 years with a normal GFR and serum albumin level of 1 to 2.5 g/dL. Patients with anasarca oedema were excluded. It is not clear whether any of the patients had any form of oedema. The participants were randomised into two groups: (1) human albumin and striped snakehead fish extract protein supplement pills, and (2) human albumin alone. The outcomes reported were serum albumin, total cholesterol and weight. We judged this study to not meet our objective of assessing the effectiveness of albumin in people with nephrotic syndrome with oedema and the comparison did not meet any of our 5 pre-specified comparisons
Kanitkar 2010	There were 24 people with nephrotic syndrome for 15 to 144 months with anasarca were randomised into two groups: 20% of albumin and 5% of albumin. The outcomes were compared for both groups based on weight loss and urinary volume. There is insufficient clarity reported to incorporate the data appropriately into a meta-analysis. However, this study was excluded because a cross-over design with washout period of 48 hours was applied and raw data for the first period (before participants were crossed over) could not be obtained for meta-analysis. We judged the washout period to be too short. This is an conference abstract and the full article could not be obtained
Na 2001	A total of seven patients with nephrotic syndrome aged from 17 to 74 years were enrolled in the study. The outcomes were compared for the effects of furosemide alone with furosemide immediately following an albumin infusion. The appropriate paired statistic to incorporate this study into the meta-analysis was not reported. However, this study was excluded because a cross-over design with wash out period of two days was applied and raw data for the first period (before participant cross-over) could not be obtained for meta-analysis
Phakdeekitcharoen 2012	The researchers included 24 patients with stable CKD (no fluctuation of GFR of more than 10 per cent in two months) who presented with oedema. There were two study periods with two weeks apart. Participants received a single dose of IV furosemide alone for one period of 24 hours and frusemide in combination with human albumin for the other period. The study analysis did not take into account the cross-over design. However this study was excluded because a cross-over de-

Study	Reason for exclusion
	sign was applied and we judged that two weeks might be insufficient washout. Raw data for the first period (before participants were crossed over) could not be obtained for meta-analysis
Sjostrom 1989	Five patients were enrolled in this cross-over study. The patients were diagnosed with nephrotic syndrome and allocated over three periods (at days 1, 2 and 5). All patients received the furosemide only for period one and were randomly assigned to either albumin and furosemide or dextran and furosemide. The primary outcome was the change in plasma and urinary albumin concentration. Individual patient data was reported. However, this study was excluded because although individual period data was available for the first period all patients received the same allocated treatment in this period and it is not possible to tell which treatment they received for the second period. This could not be obtained from the authors. We judged the washout period to be insufficient

CKD - chronic kidney disease; GFR - glomerular filtration rate; IV - intravenous; RCT - randomised controlled trial

ADDITIONAL TABLES

Table 1. Reported outcomes

	Albumin group (n = 13) Mean (SD)	Control group (n = 13) Mean (SD)	Mean difference (95% CI)	P-value
Clinical improvement				
Weight loss (in %) ^a	Reported as no difference; clarity of quantitative data uncertain			
Increased urine volume	Not reported	Not reported	Not reported	--
Reduction in ascites				
Pleural effusion				
Penile or scrotal oedema				
Serum sodium^a				
Day 3	135 (3.1)	137 (2.4)	-2.00 (-4.13 to 0.13)	0.078
Day 5	137 (2.4)	135 (3.0)	2.00 (-0.09 to 4.09)	0.073
Secondary outcomes				
Death	Not reported	Not reported	Not reported	--
Acute kidney injury and need for dialysis				
Kidney function measures				
Serum creatinine				
Glomerular filtration rate				

Table 1. Reported outcomes (Continued)

Spot urine protein-creatinine ratio ^a				
Day 2	31.9 (14.07)	8.0 (3.22)	23.90 (16.05 to 31.75)	< 0.001
Day 4	13.7 (3.78)	4.9 (5.12)	5.12 (5.34 to 12.26)	< 0.001
Day 6	4.4 (3.6)	2.1 (2.32)	2.30 (-0.03 to 4.63)	0.065
Day 8	3.6 (4.28)	1.5 (2.11)	2.10 (-0.49 to 4.69)	0.126
Day 10	1.8 (2.24)	0.8 (0.42)	1.00 (-0.24 to 2.24)	0.127
Quality of life	Not reported			
Adverse events				
Plasma volume overload	Not reported			
Hypertension ^b				
Mean systolic blood pressure				
Day 1	103 (3.7)	100 (9.5)	3.00 (-2.54 to 8.54)	0.299
Day 3	100 (6.0)	99 (6.6)	1.00 (-3.85 to 5.85)	0.690
Day 5	100 (6.0)	98 (8.2)	2.00 (-3.52 to 7.52)	0.485
Mean diastolic blood pressure				
Day 1	63 (9.4)	60.2 (9.8)	2.80 (-4.58 to 10.18)	0.434
Day 3	59 (5.5)	62 (10.4)	-3.00 (-9.40 to 3.40)	0.367
Day 5	62 (7.5)	60 (8.8)	2.00 (-4.29 to 8.29)	0.539
Pulmonary oedema	Not reported			
Need for ventilation to control plasma volume overload				

Data based on one study (26 participants) ([Lee 2000a](#)); CI = Confidence interval

^a No difference at baseline

^b Incidence of hypertension not reported

APPENDICES

Appendix 1. Electronic search strategies

Database	Search terms
CENTRAL	1. "nephrotic syndrome":ti,ab,kw

(Continued)

2. nephrosis:ti,ab,kw
3. "minimal change disease":ti,ab,kw
4. (minimal next change next glomerulo*):ti,ab,kw
5. (#1 OR #2 OR #3 OR #4)
6. (edema OR oedema OR edematous OR oedematous OR edemic OR oedemic OR anasarca):ti,ab,kw
7. #5 OR #6
8. MeSH descriptor Albumins, this term only
9. "albumin":ti,ab,kw
- 10.(#8 OR #9)
- 11.(#6 AND #10)

MEDLINE

1. exp Nephrosis/
2. nephrotic syndrome.tw.
3. nephrosis.tw.
4. minimal change disease.tw.
5. minimal change glomerulo\$.tw.
6. or/1-5
7. Edema/
8. (edema or oedema or edematous or oedematous or edemic or oedemic).tw.
9. anasarca.tw.
- 10.or/7-9
- 11.or/6,10
- 12.Albumins/
- 13.Serum Albumin/
- 14.albumin.tw.
- 15.or/12-14
- 16.and/11,15

EMBASE

1. Nephrotic Syndrome/
2. Lipoid Nephrosis/
3. Nephrosis/
4. Minimal Change Disease/
5. Minimal Change Glomerulonephritis/
6. nephrotic syndrome.tw.
7. nephrosis.tw.
8. minimal change disease.tw.
9. minimal change glomerulo\$.tw.
- 10.or/1-9
- 11.Edema/
- 12.(edema or oedema or edematous or oedematous or edemic or oedemic).tw.
- 13.Ansarca/
- 14.anasarca.tw.
- 15.or/11-14
- 16.or/10,15
- 17.Albumin/
- 18.Human Albumin/
- 19.Human Serum Albumin/
- 20.Recombinant Human Serum Albumin/
- 21.Serum Albumin/
- 22.albumin.tw.
- 23.or/17-22

(Continued)

24.16 and 23

Appendix 2. Risk of bias assessment tool

Potential source of bias	Assessment criteria
Random sequence generation Selection bias (biased allocation to interventions) due to inadequate generation of a randomised sequence	<p><i>Low risk of bias:</i> Random number table; computer random number generator; coin tossing; shuffling cards or envelopes; throwing dice; drawing of lots; minimization (minimization may be implemented without a random element, and this is considered to be equivalent to being random).</p> <p><i>High risk of bias:</i> Sequence generated by odd or even date of birth; date (or day) of admission; sequence generated by hospital or clinic record number; allocation by judgement of the clinician; by preference of the participant; based on the results of a laboratory test or a series of tests; by availability of the intervention.</p> <p><i>Unclear:</i> Insufficient information about the sequence generation process to permit judgement.</p>
Allocation concealment Selection bias (biased allocation to interventions) due to inadequate concealment of allocations prior to assignment	<p><i>Low risk of bias:</i> Randomisation method described that would not allow investigator/participant to know or influence intervention group before eligible participant entered in the study (e.g. central allocation, including telephone, web-based, and pharmacy-controlled, randomisation; sequentially numbered drug containers of identical appearance; sequentially numbered, opaque, sealed envelopes).</p> <p><i>High risk of bias:</i> Using an open random allocation schedule (e.g. a list of random numbers); assignment envelopes were used without appropriate safeguards (e.g. if envelopes were unsealed or non-opaque or not sequentially numbered); alternation or rotation; date of birth; case record number; any other explicitly unconcealed procedure.</p> <p><i>Unclear:</i> Randomisation stated but no information on method used is available.</p>
Blinding of participants and personnel Performance bias due to knowledge of the allocated interventions by participants and personnel during the study	<p><i>Low risk of bias:</i> No blinding or incomplete blinding, but the review authors judge that the outcome is not likely to be influenced by lack of blinding; blinding of participants and key study personnel ensured, and unlikely that the blinding could have been broken.</p> <p><i>High risk of bias:</i> No blinding or incomplete blinding, and the outcome is likely to be influenced by lack of blinding; blinding of key study participants and personnel attempted, but likely that the blinding could have been broken, and the outcome is likely to be influenced by lack of blinding.</p> <p><i>Unclear:</i> Insufficient information to permit judgement</p>
Blinding of outcome assessment Detection bias due to knowledge of the allocated interventions by outcome assessors.	<p><i>Low risk of bias:</i> No blinding of outcome assessment, but the review authors judge that the outcome measurement is not likely to be influenced by lack of blinding; blinding of outcome assessment ensured, and unlikely that the blinding could have been broken.</p> <p><i>High risk of bias:</i> No blinding of outcome assessment, and the outcome measurement is likely to be influenced by lack of blinding; blinding of outcome assessment, but likely that the blinding could have been broken, and the outcome measurement is likely to be influenced by lack of blinding.</p> <p><i>Unclear:</i> Insufficient information to permit judgement</p>
Incomplete outcome data Attrition bias due to amount, nature or handling of incomplete outcome data.	<p><i>Low risk of bias:</i> No missing outcome data; reasons for missing outcome data unlikely to be related to true outcome (for survival data, censoring unlikely to be introducing bias); missing outcome data balanced in numbers across intervention groups, with similar reasons for missing data across groups; for dichotomous outcome data, the proportion of missing outcomes compared with observed event risk not enough to have a clinically relevant impact on the intervention effect esti-</p>

(Continued)

mate; for continuous outcome data, plausible effect size (difference in means or standardized difference in means) among missing outcomes not enough to have a clinically relevant impact on observed effect size; missing data have been imputed using appropriate methods.

High risk of bias: Reason for missing outcome data likely to be related to true outcome, with either imbalance in numbers or reasons for missing data across intervention groups; for dichotomous outcome data, the proportion of missing outcomes compared with observed event risk enough to induce clinically relevant bias in intervention effect estimate; for continuous outcome data, plausible effect size (difference in means or standardized difference in means) among missing outcomes enough to induce clinically relevant bias in observed effect size; 'as-treated' analysis done with substantial departure of the intervention received from that assigned at randomisation; potentially inappropriate application of simple imputation.

Unclear: Insufficient information to permit judgement

Selective reporting

Reporting bias due to selective outcome reporting

Low risk of bias: The study protocol is available and all of the study's pre-specified (primary and secondary) outcomes that are of interest in the review have been reported in the pre-specified way; the study protocol is not available but it is clear that the published reports include all expected outcomes, including those that were pre-specified (convincing text of this nature may be uncommon).

High risk of bias: Not all of the study's pre-specified primary outcomes have been reported; one or more primary outcomes is reported using measurements, analysis methods or subsets of the data (e.g. subscales) that were not pre-specified; one or more reported primary outcomes were not pre-specified (unless clear justification for their reporting is provided, such as an unexpected adverse effect); one or more outcomes of interest in the review are reported incompletely so that they cannot be entered in a meta-analysis; the study report fails to include results for a key outcome that would be expected to have been reported for such a study.

Unclear: Insufficient information to permit judgement

Other bias

Bias due to problems not covered elsewhere in the table

Low risk of bias: The study appears to be free of other sources of bias.

High risk of bias: Had a potential source of bias related to the specific study design used; stopped early due to some data-dependent process (including a formal-stopping rule); had extreme baseline imbalance; has been claimed to have been fraudulent; had some other problem.

Unclear: Insufficient information to assess whether an important risk of bias exists; insufficient rationale or evidence that an identified problem will introduce bias.

CONTRIBUTIONS OF AUTHORS

1. Developed the concept of this study: ASA
2. Draft the protocol: ASA and JJH with input from other authors
3. Study selection: NJA and KYC with input from ASA and JJH
4. Extract data from studies: ASA and HVR
5. Enter data into RevMan: NJA, KYC
6. Carry out the analysis: NJA, KYC, ASA with input from JJH
7. Interpret the analysis: JJH, NJA, ASA and KYC with input from other authors
8. Draft the final review: JJH and KYC with input from all authors
9. Disagreement resolution: JJH
10. Update the review: JJH and KYC

DECLARATIONS OF INTEREST

- Jacqueline Ho: none known
- Azreen Adnan: none known

- Yee Cheng Kueh: none known
- Nurul Jannah Ambak: none known
- Hans Van Rostenberghe: none known
- Fauziah Jummaat: none known

SOURCES OF SUPPORT

Internal sources

- Penang Medical College, Malaysia.
- Universiti Sains Malaysia, Kota Bharu, Malaysia.

External sources

- No sources of support supplied

DIFFERENCES BETWEEN PROTOCOL AND REVIEW

In the protocol we stated that we would not include cross-over studies. Since we found studies using the cross-over design we have elaborated in the methods under the section on 'Unit of analysis issues', why cross-over studies might not be suitable. We have also mentioned this in the sensitivity analysis. We have also added a section to the methods after the section on assessment of risk of bias, assessing the quality of evidence using the GRADE approach and selected seven outcomes to include in a summary of findings table. We have also added serum sodium as new measure of the primary outcome, clinical improvement. In the protocol we did not specify that we would include it in a summary of findings table.