Published in final edited form as:

Org Lett. 2019 July 05; 21(13): 5172–5177. doi:10.1021/acs.orglett.9b01769.

Enantio- and Diastereoselective Synthesis of Functionalized Carbocycles by Cu-Catalyzed Borylative Cyclization of Alkynes with Ketones

Joseph M. Zanghi, Shuang Liu, Simon J. Meek

Department of Chemistry, The University of North Carolina at Chapel Hill, Chapel Hill, North Carolina 27599-3290.

Abstract

A single-pot Cu-catalyzed enantio- and diastereoselective tandem hydroboration/borylative cyclization of alkynes with ketones for the synthesis of carbocycles is reported. The reaction proceeds via a desymmetrization and generates four contiguous stereocenters, including an all-carbon quaternary center. The method provides rapid access to [6,5]-, [5,5]-bicycles, and cyclopentane products. Catalyst controlled diastereoselectivity by selection of bisphosphine ligand is noted. Utility of the products is demonstrated by site- and chemoselective transformations that afford valuable alkenyl and allyl organoborons.

Graphical Abstract

Multi-component tandem reactions, wherein subsequent reactions occur as a result of functionality generated in a previous step, allow for the formation of several bonds in a single operation, without the need of additional reagents.¹ Such reactions minimize the economic, environmental, and labor costs associated with multi-step syntheses.¹ Of particular interest, are tandem catalytic reactions that stereoselectively form all-carbon quaternary stereocenters, structural motifs that remain a significant synthetic challenge in organic chemistry.² One strategy to generate quaternary stereocenters is through

Corresponding Author sjmeek@unc.edu.

Supporting Information.

Experimental procedures and spectral and analytical data for all products. This material is available free of charge via the Internet at <http://pubs.acs.org>.

Notes

Authors declare no competing financial interests.

enantioselective desymmetrization processes.³ Accordingly, incorporating an enantioselective desymmetrization step into a tandem reaction would rapidly generate stereochemical complexity in a highly efficient manner.⁴ In addition to the growing importance of constructing all-carbon quaternary centers, alkyl organoborons have become important intermediates in the synthesis of pharmaceuticals and agrochemicals due to the range of stereospecific functional group transformations they undergo.^{5,6,7}

Recently we reported a Cu-catalyzed enantio- and diastereoselective tandem borylation/1,2-addition of alkenyl boronic esters to ketones (Scheme 1B).⁸ To construct more complex structures in a simpler protocol, we sought to develop a process to efficiently construct carbocycles bearing four contiguous stereocenters.

In order to achieve these objectives, we decided to examine two changes (Scheme 1C): (1) Catalytic desymmetrization of 1,3-diketones substrates that access stereodefined, highly substituted hydrindanes, diquinanes, and cyclopentanes bearing two boronic esters. Ideally, the two C–B bonds could be sequentially functionalized, offering access to a range of stereodefined products. (2) Change from alkenyl borons to alkynes due to their ease of accessibility. Notably, alkynes pose an inherent site-selectivity challenge as hydroboration can result in α - and β -borylation isomers.⁹

A number of studies related to Cu-catalyzed bis-borylation of alkynes have been reported; however, trapping of the resulting copper–C(sp³) with an electrophile other than proton has not been reported to the best of our knowledge.¹⁰ Enantioselective Cu-catalyzed bis-hydroboration of terminal alkynes and SiMe₃-protected alkynes have been reported.¹¹ Related enantioselective Cu–B(pin) initiated carboboration reactions of olefins have also been disclosed.^{12,13} In this regard, Cu-catalyzed conjugate borylation of Michael acceptors, and subsequent aldol addition of the Cu-enolate with a diketone has been shown intramolecularly (Scheme 1A).^{4b}

We initiated our studies by investigating the catalytic desymmetrization of cyclohexanedione **1a** to afford bis-boryl hydrindane **3a** (Table 1). Treatment of alkyne **1a** with 5 mol % CuCl, 6 mol % (*R*)-BINAP (**L1**), 6 mol % KO*t*-Bu, 2.2 equivalents of B₂(pin)₂, and *t*-BuOH (2.0 equiv) in THF at 22 °C for 18 h afforded **3a** in 65% yield, 5:1 dr, and 99:1 er. Examination of a number of other chiral phosphine ligands (**L2–L6**) generated either significantly lower conversion or diastereoselectivity. In contrast, it was found that employing furyl-MeO-Biphep (**L7**) greatly improved dr while maintaining good yield, and excellent enantioselectivity providing **3a** in 64% yield, 12:1 dr, and 99:1 er. Furthermore, it was found that increasing the catalyst loading to 10 mol %, and switching from *t*-BuOH to MeOH, led to an improved 78% yield of **3a** while maintaining high stereoselectivity (9:1 dr, and >99:1 er).

With conditions in hand for the alkyne desymmetrization, we next examined the substrate scope of the tandem reaction (Table 1). In general, the yields are moderate to good, and enantio- and diastereoselectivity are generally excellent. Product **3a** was isolated in 70% yield, 13:1 dr, and >99:1 er. Additional substitution on the 6-membered ring was tolerated (**3b**) (X-ray structure of **3b** was obtained (Scheme 5A) (*vide infra*). Prenyl containing

substrate **1c** was found to cyclize to **3c** with lower diastereoselectivity (5:1 dr) but in excellent enantioselectivity (98:2 er, and >99:1 er). Incorporation of benzyl (**1d**) and thiophene (**1e**) substituents led to **3d** and **3e** in 60% (>20:1 dr) and 65% (3:2 dr), respectively. Notably, upon isolation, the minor diastereomer of **3e** decomposes affording a single diastereomer (>20:1 dr, and 94:6 er). Terminal alkenes are also tolerated, and bis-boron **3f** is isolated in 46% yield, 17:1 dr, and in 88:12 er; the lower er is possibly due to coordination of the unhindered alkene to the catalyst.

For [5,5]-fused carbocycles (entries 7–9), however, **L1** was found to be a slightly better ligand compared to **L7**. Methyl-substituted diquinane **5a** is formed in 69% yield, 2:1 dr, and 90:10 er.¹⁴ Increasing the size of the group on the quaternary center to an allyl (**1h**) or n-propyl group (**1i**) provided products **5b** and **5c** in 57% yield (>20:1 dr, 89:11 er) and 64% yield (>20:1 dr, 94:6 er), respectively. Diquinane **5c** could be recrystallized once to 97.5:2.5 er. For cyclopentane products **6a–c**, yields again were moderate, but reactions proceed with generally excellent enantio- and diastereoselectivities (Table 1). For example, product **6a**, formed in 56% NMR yield, was isolated as the tris-boronate (>20:1 dr, >99:1 er).¹⁵ Simply switching from Me to Et provided tertiary alcohol bis-boronate **6b** in 42% yield as a single stereoisomer (>20:1 dr, >99:1 er). Prenyl-substituted diketone **1l** was found to undergo smooth cyclization to afford **6c** in 17:1 dr (>99:1 er), when compared to the [6,5]-prenyl **3c** formed in 5:1. The alkyne cyclization strategy was also tested for the generation of 6-membered rings to form decalins (entry 13, Table 1). Unfortunately, large amounts of double hydroboration are observed (<2% conversion to **7**); starting from the vinyl boronic ester failed to improve the reaction (see the SI for details).

Next, in order to determine the efficiency of the alkyne hydroboration and carboboration steps, we undertook a brief comparison of the catalytic reaction starting from the pre-formed alkenyl boronic esters versus alkyne (Scheme 3). The reactions of the three vinyl boronic esters **2a**, **2g**, and **2j** were examined and the results illustrated in Scheme 3. With 1.1 equivalent of B₂(pin)₂ and 1.0 equivalent of *t*-BuOH, cyclization to cyclopentane products **3a**, **5a**, and **6a** resulted in only marginal improvement in yield compared to starting from the corresponding alkynes; indicating efficient alkyne hydroboration.

A proposed mechanism for the Cu-catalyzed tandem alkyne cyclization method is shown in Scheme 4, which depicts dual catalytic cycles that proceed contemporaneously.^{9a,10} Accordingly, a Cu-catalyzed alkyne boryl–protonation cycle occurs followed by a carboboration cycle. In the hydroboration cycle, transmetalation of (L)Cu–OR **8** with B₂(pin)₂ generates (L)Cu–B(pin) complex **9**. Regioselective *syn* migratory insertion of **9** across alkyne **10**, is followed by protonation of alkenyl copper **11** by MeOH to liberate *trans*-β-vinyl boronic ester **12**. In the carboboration cycle, migratory insertion of **9** across vinyl boronic ester **12** sets the absolute stereochemistry, which is followed by diastereoselective cyclization (**13**→**14**). Protonation of Cu-bound product **14** releases the product and regenerates catalyst **8**.

As noted prior, the X-ray structure of tertiary alcohol **3b** was obtained (Scheme 5A). The structure shows a *cis* [6,5] ring juncture, with the B(pin) *anti* to the hydroxyl group.¹⁶ Importantly, borylation of the *E*-alkenyl boron with (*R*)-furyl-biphep derived catalyst results

in *R* stereochemistry (e.g., in **13**). The *syn* relationship between the two B(pin) moieties in **3b** can be explained by a stereoretentive mode of addition by the (L)Cu-alkyl to the ketone. A result rationalized by use of the small furylphosphine ligand in combination with a methyl group on the substrate, which results in a sterically accessible (L)Cu-alkyl (e.g., **13**).

Late on in our study, surprisingly, we found that when alkyne **1a** is cyclized with BDPP (**L8**), *syn, anti, syn*-diastereomer **15** is formed instead in 97% yield and 11:1 dr (Scheme 5B). With other [6,5] products we found **L8** afforded the *same* diastereomers (e.g., **3f**) as **L7** and **L1** in >20:1 dr. These unexpected results encouraged us to further investigate the stereochemistry. The X-ray structure of **3f** was obtained (Scheme 5A). Remarkably, the stereochemistry of three of the stereocenters were different compared to **3b**. As the stereochemistry of the boronic ester γ to the hydroxyl is set in initial hydroboration, we assigned this stereocenter as *R* for all substrates based on the absolute configuration of **3b**. The *anti* boronic esters of **3f** indicate a stereoinvertive cyclization, observed in previous work by our group and others.^{8,17} To account for this, we propose that the cyclization occurs invertively for R substituents larger than Me (e.g., **3f**) due to increased steric repulsion between the R group and the PAr₂ phosphine ligand, which prevents coordination of Cu to the ketone (Scheme 5C). The change in stereochemistry at the ring junction is due to cyclization on the other diastereotopic ketone due to unfavorable B(pin) $\leftrightarrow$ R steric interaction. The stereochemistry of the [5,5] products were confirmed as *syn, anti, anti* based on the X-ray structure of **5b** (Scheme 5A). Since reactions with **L1**, **L7**, and **L8** result in increasing dr for methyl and with high dr for larger substituents, all [5,5] products predominantly arise from a stereoinvertive cyclization pathway.¹⁸ Cyclopentane products **6a–c**, have been assigned by analogy; **6a** as stereoretentive *syn, anti, syn*, and **6b–c** stereoinvertive *syn, anti, anti*. This is also supported by the decrease in dr for **6a** going from **L7** $\rightarrow$ **L1**.¹⁹

The synthetic utility of organodiboron carbocycles accessible through the tandem Cu-catalyzed bis-borylation/cyclization protocol is highlighted through several site- and chemoselective transformations depicted in Scheme 6. (1) Oxidation of **3c** with H₂O₂/NaOH affords triol **18** in 73% yield (Scheme 6A). (2) Exposure of the tertiary alcohol products to Martin's sulfurane,²⁰ results in site- and chemoselective eliminations that are dependent on the hydrocarbon bicycle. For example, treatment of [6,5]-product **3a** with Martin's sulfurane effects a site-selective dehydration to afford alkenyl boron **19** in 59% yield (Scheme 6B). Subjecting benzyl-derivative **3d** to the same reaction conditions also provides dehydrated alkenyl boron product **20** in 78% yield, despite the change in relative stereochemistry. This is in direct contrast to [5,5]-bicycles, which when treated with Martin's sulfurane result in an *anti* boron-hydroxyl elimination (Scheme 6C). For example, allyl borons **21** and **22** are generated in 57% and 68% isolated yield from **5b** and **5c** respectively. We believe the divergence in reactivity arises from the σ_{C-B} aligning better with the carbocation p-orbital in the more planar [5,5]-bicycle versus the σ_{C-H} . To the best of our knowledge, this reactivity is the first time an *anti* boron-hydroxyl elimination has been reported for a tertiary alcohol, and is the first example of Martin's sulfurane used in transformation of an organoboron.²¹ Further transformation of allyl boronic ester **22** to alcohol **23** was accomplished by a

stereospecific allylation with formaldehyde, generating vicinal all-carbon quaternary stereocenters in 73% yield and 18:1 dr (Scheme 6D).

In conclusion, we present a catalytic tandem enantio- and distereoselective method for the preparation of borylated carbocycles. We demonstrate the formation of four contiguous stereocenters, including an all-carbon quaternary center, in good yield and selectivity (up to >98:2 dr, and >99:1 er). Stereoinvertive and stereoretentive cyclization pathways were found to be operative, and dependent on ligand and substrate. Site- and chemoselective eliminations with Martin's sulfuran highlight valuable alkenyl and allyl organoborons accessible through the method. Development of additional catalytic carboboration methods continue to be investigated in these laboratories.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

ACKNOWLEDGMENT

Financial support was provided by the United States National Institutes of Health, Institute of General Medical Sciences (R01GM116987) and the University of North Carolina at Chapel Hill. We thank Quentin Bruch, Tia Cervarich, and Dr. Chun-Hsing Chen of UNC for X-ray structure elucidation of **3b**, **3f**, and **5b**. AllyChem is acknowledged for donations of B₂(pin)₂.

REFERENCES

- (1). (a)For selected reviews, see:Tietze LF Domino Reactions in Organic Synthesis. Chem. Rev 1996, 96, 115–136. [PubMed: 11848746] (b)Pellissier H Stereocontrolled Domino Reactions. Chem. Rev 2013, 113, 442–524. [PubMed: 23157479] (c)Volla CMR; Atodiresei I; Rueping M Catalytic C–C Bond-Forming Multi-Component Cascade or Domino Reactions: Pushing the Boundaries of Complexity in Asymmetric Organocatalysis. Chem. Rev 2014, 114, 2390–2431. [PubMed: 24304297] (d)Epe G; Didier D; Marek I Stereocontrolled Formation of Several Carbon–Carbon Bonds in Acyclic Systems. Chem. Rev 2015, 115, 9175–9206. [PubMed: 25768205]
- (2). (a)For selected reviews, see:Corey EJ; Guzman-Perez. The Catalytic Enantioselective Construction of Molecules with Quaternary Carbon Stereocenters. A. Angew. Chem. Int. Ed 1998, 37, 388–401.(b)Peterson EA, Overman LE. Contiguous Stereogenic Quaternary Carbons: A Daunting Challenge in Natural Products Synthesis. Proc. Natl. Acad. Sci 2004, 101, 11943–11948. [PubMed: 15232003] (c)Christoffers J; Baro A Stereoselective Construction of Quaternary Stereocenters. Adv. Synth. Catal 2005, 347, 1473–1482.(d)Marek I; Minko Y; Pasco M; Mejuch T; Gilboa N; Chechik H; Das JP All-Carbon Quaternary Stereogenic Centers in Acyclic Systems through the Creation of Several C–C Bonds per Chemical Step. J. Am. Chem. Soc 2014, 136, 2682–2694. [PubMed: 24512113] (e)Quasdorf KW; Overman LE Catalytic Enantioselective Synthesis of Quaternary Carbon Stereocenters. Nature. 2014, 516, 181–191. [PubMed: 25503231] (f)Long R; Huang J; Gong J; Yang Z Direct Construction of Vicinal All-Carbon Quaternary Stereocenters in Natural Product Synthesis. Nat. Prod. Rep 2015, 32, 1584–1601. [PubMed: 26334685] (g)Liu Y; Han S-J; Liu W-B; Stoltz BM Catalytic Enantioselective Construction of Quaternary Stereocenters: Assembly of Key Building Blocks for the Synthesis of Biologically Active Molecules. Acc. Chem. Res 2015, 48, 740–751. [PubMed: 25715056] (h)Alam R; Vollgraff T Eriksson L. Szabó. Synthesis of Adjacent Quaternary Stereocenters by Catalytic Asymmetric Allylboration. J. Am. Chem. Soc 2015, 137, 11262–11265. [PubMed: 26316158]
- (3). For a recent review of the area, see: Zeng X-P; Cao ZY; Wang Y-H; Zhou F; Zhou J Catalytic Enantioselective Desymmetrization Reactions to All-Carbon Quaternary Stereocenters. Chem. Rev 2016, 116, 7330–7396. [PubMed: 27251100]

- (4). (a) For representative examples of tandem conjugate addition/aldol desymmetrization of 1,3-diketones, see: Bocknack BM; Wang L-C; Krische MJ Desymmetrization of Enone-Diones via Rhodium-Catalyzed Diastereo- and Enantioselective Tandem Conjugate Addition-Aldol Cyclization. *Proc. Natl. Acad. Sci. U. S. A* 2004, 101, 5421–5424. [PubMed: 15024093] (b) Burns AR; Gonzalez JS; Lam HW Enantioselective Copper(I)-Catalyzed Borylative Aldol Cyclizations of Enone Diones. *Angew. Chem. Int. Ed* 2012, 51, 10827–10831.
- (5). Hall DG In *Boronic Acids: Preparation and Applications in Organic Synthesis and Medicine*; Wiley-VCH: Weinheim, 2011.
- (6). (a) For reviews of Suzuki-Miyaura cross-couplings, see: Chemler SR; Trauner D; Danishefsky SJ The B-Alkyl Suzuki-Miyaura Cross-Coupling Reaction: Development, Mechanistic Study, and Applications in Natural Product Synthesis. *Angew. Chem. Int. Ed* 2001, 40, 4544–4568. (b) Lennox AJJ; Lloyd-Jones GC Selection of Boron Reagents for Suzuki-Miyaura Coupling. *Chem. Soc. Rev* 2014, 43, 412–443. [PubMed: 24091429]
- (7). (a) For stereospecific transformations, see: Scott HK; Aggarwal VK Highly Enantioselective Synthesis of Tertiary Boronic Esters and their Stereospecific Conversion to other Functional Groups and Quaternary Stereocenters. *Chem. Eur. J* 2011, 17, 13124–13132. [PubMed: 22052475] (b) Leonori D; Aggarwal VK Stereospecific Couplings of Secondary and Tertiary Boronic Esters. *Angew. Chem. Int. Ed. Engl* 2015, 54, 1082–1096. [PubMed: 25414056] (c) Sandford C; Aggarwal VK Stereospecific Functionalizations of Secondary and Tertiary Boronic Esters. *Chem. Commun* 2017, 53, 5481–5494.
- (8). Green JC; Joannou MV; Murray SA; Zanghi JM; Meek SJ Enantio- and Diastereoselective Synthesis of Hydroxy Bis(boronates) via Cu-Catalyzed Tandem Borylation/1,2-Addition. *ACS Catal.* 2017, 7, 4441–4445. [PubMed: 29520326]
- (9). Jang H; Zhugralin AR; Lee Y; Hoveyda AH Highly Selective Methods for Synthesis of Internal (α -) Vinylboronates through Efficient NHC-Cu-Catalyzed Hydroboration of Terminal Alkynes. Utility in Chemical Synthesis and Mechanistic Basis for Selectivity. *J. Am. Chem. Soc* 2011, 133, 7859–7871. [PubMed: 21526827]
- (10). Semba K; Fujihara T; Terao J; Tsuji Y Copper-Catalyzed Borylative Transformations of Non-Polar Carbon–Carbon Unsaturated Compounds Employing Borylcopper as an Active Catalyst Species. *Tetrahedron* 2015, 71, 2183–2197.
- (11). (a). Lee Y; Jang H; Hoveyda AH Vicinal Diboronates in High Enantiomeric Purity through Tandem Site-Selective NHC-Cu-Catalyzed Boron-Copper Additions to Terminal Alkynes. *J. Am. Chem. Soc* 2009, 131, 18234–18235. [PubMed: 19968273] (b) Jung H-Y; Yun J Copper-Catalyzed Double Borylation of Silylacetylenes: Highly Regio- and Stereoselective Synthesis of Syn-Vicinal Diboronates. *Org. Lett* 2012, 14, 2606–2609. [PubMed: 22548477]
- (12). (a) For recent example of non-enantioselective carboboration of vinyl-B(pin), see: Yoshida H; Kageyuki I; Takaki K Copper-Catalyzed Three-Component Carboboration of Alkynes and Alkenes. *Org. Lett* 2013, 15, 952–955. [PubMed: 23384399] (b) Kageyuki I; Osaka I; Takaki K; Yoshida H Copper-Catalyzed B(Dan)-Installing Carboboration of Alkenes. *Org. Lett* 2017, 19, 830–833. [PubMed: 28156117]
- (13). (a) For recent examples of catalytic carboboration of olefins, see: Ito H; Kosaka Y; Nonoyama K; Sasaki Y; Sawamura M Synthesis of Optically Active Boron-Silicon Bifunctional Cyclopropane Derivatives through Enantioselective Copper(I)-Catalyzed Reaction of Allylic Carbonates with a Diboron Derivative. *Angew. Chem. Int. Ed* 2008, 47, 7424–7427. (b) Zhong C; Kunii S; Kosaka Y; Sawamura M; Ito H Enantioselective Synthesis of Trans-Aryl-and-Heteroaryl-Substituted Cyclopropylboronates by Copper(I)-Catalyzed Reactions of Allylic Phosphates with a Diboron Derivative. *J. Am. Chem. Soc* 2010, 132, 11440–11442. [PubMed: 20684557] (c) Kubota K; Yamamoto E; Ito H Copper(I)-Catalyzed Borylative *Exo*-Cyclization of Alkenyl Halides Containing Unactivated Double Bond. *J. Am. Chem. Soc* 2013, 135, 2635–2640. [PubMed: 23351005] (d) Su W; Gong T-J; Lu X; Xu M-Y; Yu C-G; Xu Z-Y; Yu H-Z; Xiao B; Fu Y Ligand-Controlled Regiodivergent Copper-Catalyzed Alkylboration of Alkenes. *Angew. Chem. Int. Ed* 2015, 54, 12957–12961. (e) Cheng Y; Mück-Lichtenfeld C; Studer A Transition Metal-Free 1,2-Carboboration of Unactivated Alkenes. *J. Am. Chem. Soc* 2018, 140, 6221–6225. [PubMed: 29741375] (f) Wang H-M; Zhou H; Xu Q-S; Liu T-S; Zhuang C-L; Shen M-H; Xu H-D Copper-Catalyzed Borylative Cyclization of Substituted N-(2-Vinylaryl)Benzaldimines. *Org. Lett* 2018, 20, 1777–1780. [PubMed: 29537288] (g) Chen B; Cao P; Liao Y; Wang M; Liao J

Enantioselective Copper-Catalyzed Methylboration of Alkenes. *Org. Lett* 2018, 20, 1346–1349. [PubMed: 29461840] (h)Logan KM; Sardini SR; White SD; Brown MK Nickel-Catalyzed Stereoselective Arylboration of Unactivated Alkenes. *J. Am. Chem. Soc* 2018, 140, 159–162. [PubMed: 29271650]

- (14). Lam and co-workers also observed poor diastereoselectivity (60:40) for a [5,5]-bicycle using methyl at the quaternary stereocenter, see ref. 4b.
- (15). 6a is somewhat unstable to silica gel, resulting in a difference between NMR and isolated yield.
- (16). (a)A syn relationship would likely lead to a boron-Wittig syn-elimination from the Cu-bound product. For example, see:Matteson DS; Moody RJ Deprotonation of 1,1-Diboronic Esters and Reactions of the Carbanions with Alkyl Halides and Carbonyl Compounds. *Organometallics*. 1982, 1, 20–28.(b)Matteson DS; Moody RJ; Jesthi PK The Reaction of Aldehydes and Ketones with a Boron-Substituted Carbanion, Bis(ethylenedioxyboryl)methide. A Simple Aldehyde Homologation. *J. Am. Chem. Soc* 1975, 97, 5608–5609.(c)Matteson DS; Jesthi PKJ Lithium bis(ethylenedioxyboryl)methide and its Reactions with Carbonyl Compounds and with the Chlorotriphenyl Derivatives of Germanium, Tin, and Lead. *Organomet. Chem* 1976, 110, 25–37. (d)Matteson DS; Moody RJ Carbanions from Deprotonation of gem-Diboronic Esters. *J. Am. Chem. Soc* 1977, 99, 3196–3197.(e)Coombs JR; Zhang L; Morken JP Synthesis of Vinyl Boronates from Aldehydes by a Practical Boron-Wittig Reaction. *Org. Lett* 2015, 17, 1708–1711. [PubMed: 25799147]
- (17). (a)For other stereoinvertive Cu-catalyzed 1,2-additions, see:Yang Y; Perry IB; Buchwald SL Copper-Catalyzed Enantioselective Addition of Styrene-Derived Nucleophiles to Imines Enabled by Ligand-Controlled Chemoselective Hydrocupration. *J. Am. Chem. Soc* 2016, 138, 9787–9790. [PubMed: 27454393] (b)Itoh T; Kanzaki Y; Shimizu Y; Kanai M Copper(I)-Catalyzed Enantio- and Diastereodivergent Borylative Coupling of Styrenes and Imines. *Angew. Chem. Int. Ed* 2018, 57, 8265–8269.
- (18). L8 was tested for products 3c, 3d, 3f, 5a, 6a, and 6b and it affords the same diastereomers as L1 and L7 in all cases (except 6a-b, <5% conversion observed).
- (19). Attempts to grow X-ray quality crystals of cyclopentane products **6a–c** proved unsuccessful.
- (20). (a). Martin JC; Arhart RJ Sulfuranes. III. A Reagent for the Dehydration of Alcohols. *J. Am. Chem. Soc* 1971, 93, 4327–4329.(b)Martin JC; Arhart RJ Sulfuranes. VI. Reactions Involving the Alkoxy Ligands of Dialkoxydiarylsulfuranes. Formation of Olefins and Ethers. *J. Am. Chem. Soc* 1972, 94, 5003–5010.(c)Martin JC; Arhart RJ Single-Step Syntheses of Epoxides and Other Cyclic Ethers by Reaction of a Diaryldialkoxysulfurane with Diols. *J. Am. Chem. Soc* 1974, 96, 4604–4611.(d)Engel PS; Keys DE; Kitamura A “Spring-Loaded” Biradicals. The Radical and Electron-Transfer Photochemistry of Bridgehead Cyclopropyl-Substituted 2,3-Diazabicyclo[2.2.2]oct-2-enes (DBO's). *J. Am. Chem. Soc* 1985, 107, 4964–4975.
- (21). For boron-Wittig elimination of a tertiary acyclic alcohol and a borane using KH, see: Kawashima T; Yamashita N; Okazaki R *J. Am. Chem. Soc* 1995, 117, 6142–6143.

Scheme 1.

Enantio- and Diastereoselective Cu-Catalyzed Multicomponent Carboborylative Cyclizations.

Scheme 2. Reaction Optimization^a

^aReactions performed under N_2 atmosphere. Yield and diastereomeric ratios (dr) determined by analysis of 400, 500, or 600 MHz 1H NMR spectra of crude reactions with hexamethyldisiloxane as internal standard. Enantiomeric ratios (er) determined by HPLC or SFC analysis; see the SI for details. nd = not determined. ^bCuCl (10 mol%), L7 (11 mol%), KOtBu (10 mol%), MeOH (2.0 equiv).

Scheme 3. Comparison to Alkenyl Boronic Esters in Ketone Desymmetrization^a^aSame conditions and analytical methods as in Scheme 2; see the SI for details.

Scheme 4.
Proposed Dual Catalytic Cycles.

Scheme 5. Stereochemical Analysis and Ligand Controlled Diastereoselectivity^a^aSee SI for details. ^bData given in Scheme 5B refer to NMR yields.

Scheme 6.
Utility of Bis-Boronate Products.

^aSee SI for details.

Table 1.

Reaction Scope^{a-b}

entry	starting material	product	result
1			with L7 : 70% yield, 13:1 dr >99:1 er
2			with L7 : 54% yield, >20:1 dr >99:1 er
3			with L7 : 49% yield, 5:1 dr 98:2 er, >99:1 er
4			with L7 : 60% NMR yield 32% yield, >20:1 dr 96:4 er
5			with L7 : 65% NMR yield, 3:2 dr 25% yield, >20:1 dr 94:6 er
6			with L7 : 46% yield, 17:1 dr 88:12 er
7			with L1 : 69% yield, 2:1 dr 74:26 er, 90:10 er
8			with L1 : 57% yield, >20:1 dr 89:11 er
9			with L1 : 64% yield, >20:1 dr 94:6 er [recryst: 97.5:2.5 er]
10			with L7 : 56% NMR yield 37% yield, >20:1 dr >99:1 er
11			with L7 : 42% yield, >20:1 dr >99:1 er
12			with L7 : 48% yield, 17:1 dr >99:1 er
13			with L1 or L7 : <2% conv

^aSame conditions and analytical methods as in Scheme 2; see the SI for details.^bYields of purified hydroxyl bis-boronate products.