CORRESPONDENCE

Estimation of the Number of People with Opioid Addiction in Germany

by Prof. Dr. phil. Ludwig Kraus, Dr. phil. Nicki-Nils Seitz, Dr. rer. medic. Bernd Schulte, Dr. med. Peter Cremer-Schaeffer, Dr. phil. Barbara Braun, PD Dr. phil. Uwe Verthein, and Dr. hum. biol. Tim Pfeiffer-Gerschel in issue 9/2019

Number of Addicts Underestimated

Thank you for calculating the number of persons with an opioid addiction—but less for the “It has not gotten worse.” For the number of persons with addictions was not determined, but rather only the number of those seeking out “addiction care facilities” (1). This results in an underestimation of the number.

Namely, the number is missing those whose addiction is maintained by medical prescription. An editorial in JAMA states: “In 2017, an estimated 11 to 12 million people in the United States (4.2% of the total population) misused opioids (including heroin). What most physicians do not recognize is that 92% of people who misuse opioids do so by taking prescription opioids, and that 75% of individuals who use heroin report that they started misusing opioids through the misuse of prescription opioids.” (2).

In the United States, this topic has been officially declared a “medical emergency.” A number of measures have been implemented, with dozens of articles appearing each year even in journals with international readership, such as JAMA and NEJM.

In Germany, only Schubert et al. (3) and Marschall et al. (4) have occupied themselves with the topic. The conclusion based on health insurance data: 4%–5% of insured persons receive such substances with “at least more than a one-time prescription per quarter”. Around 80% of these people receive it for “non-cancer pain”—predominantly as a permanent prescription. From 2010 to 2017, the prescribed dose for the defined daily dose (DDD) has increased by 21%.

Excluding the few cancer patients among those receiving opioid prescriptions, the experience of practicing physicians in Germany is that they also are obviously treating addicts: first, over time, addicts require increasingly higher doses; and, second, they can practically never be taken off the drug—not even stepwise over months.

DOI: 10.3238/arztebl.2019.0459a

References

1. Kraus L, Seitz NN, Schulte B, et al.: Estimation of the number of people with opioid addiction in Germany. *Dtsch Arztebl Int* 2019; 116: 137–43.
2. Ashburn MA, Fleisher LA. Editorial: Increasing evidence for the limited role of opioids to treat chronic noncancer pain. *JAMA* 2018; 320: 2427–8.
3. Schubert I, Ihle P, Sabatowski R: Increase in opiate prescription in Germany between 2000 and 2010—a study based on insurance data. *Dtsch Arztebl Int* 2013; 110: 45–51.
4. Marschall U, L'hoest H: Opioidtherapie in der Versorgungsrealität. Ein Beitrag zur Diskussion um ein weitverbreitetes Arzneimittel. *BARMER GEK Gesundheitswesen* aktuell 2011: 242–69.

emer. Prof. Dr. med. Heinz-Harald Abholz
Institut für Allgemeinmedizin (ifam),
Universitätsklinikum Düsseldorf und
Praxis Elsdorf, Germany
abholz@med.uni-duesseldorf.de

In Reply:

To address the question of whether the current estimate of persons with an opioid dependence is an underestimate, it should be noted that this is an estimate accounting for the use of “illegal opioid-containing substances” (1). This does not rule out the existence of a group of individuals who have become addicted to opioids, for instance, by means of pain therapy. Our conclusion that the size of the first group did not increase, and that the situation has not “gotten worse”, must therefore be seen in the context of this target group.

The prevalence of the group of individuals who have become addicted to prescription opioid analgesics is estimated to be around 1% of the general population in Germany (2). The prevalence of opioid analgesic disorder among non-cancer pain patients based on German medical health insurance data was estimated at 0.56% (3). Whether the size of this group has increased in the context of increased opioid analgesic prescriptions is unknown.

Changes in the scale of the first group are conceivable in two circumstances: due to an increase in the number of persons with opioid addiction induced by the consumption of illegal opioid-containing substances, or to an increase in persons with iatrogenic opioid addiction who move to the illegal market. Our estimate compared with previous data and trend analyses of drug-related deaths due to opioids (4) do not indicate such a development.

In the US, the majority of people with iatrogenic opioid addiction have switched to the black market (5). This development has not been observed in Germany until now. Nevertheless, in view of the increase in treatment with opioid-containing analgesics, the indication and duration must be critically examined.

DOI: 10.3238/arztebl.2019.0459b

References

1. Kraus L, Seitz NN, Schulte B, et al.: Estimation of the number of people with opioid addiction in Germany. *Dtsch Arztebl Int* 2019; 116: 137–43.
2. Aydin D: Konsumstörungen im Zusammenhang mit opioidhaltigen und nicht opioidhaltigen Schmerzmitteln – Prävalenz in der Bevölkerung und prädiktive Effekte. Bachelorarbeit. Fernuniversität in Hagen 2018.
3. Marschall U, L'hoest H, Radbruch L, Häuser W: Long-term opioid therapy for chronic non-cancer pain in Germany. *Eur J Pain* 2016; 20: 767–76.
4. Kraus L, Pfeiffer-Gerschel T, Seitz N-N, Kurz A: Analyse drogeninduzierter Todesfälle. Berlin 2018.
5. Pergolizzi JVJ, LeQuang JAT, R.Jr., Raffa RB, Group NR: Going beyond prescription pain relievers to understand the opioid epidemic: the role of illicit fentanyl, new psychoactive substances, and street heroin. *Postgraduate Med* 2018; 130: 1–8.

On behalf of the authors

Prof. Dr. phil. Ludwig Kraus
IFT Institut für Therapieforchung, München, Germany
kraus@ift.de

Conflict of interest statement

The authors of each contribution declare that no conflict of interest exists.