CORRESPONDENCE

Coronary Angiography with Pressure Wire and Fractional Flow Reserve

State of the Art in the Diagnosis of Coronary Stenosis

by Dr. med. Luise Gaede, Prof. Dr. med. Helge Möllmann, Prof. Dr. med. Tanja Rudolph, PD Dr. med. Johannes Rieber, PD Dr. med. Florian Boenner, and Dr. med. Monique Tröbs in issue 12/2019

Computed Tomography Angiography

We would like to add the following to the very informative and structured article (1): non-invasive, good quality imaging of coronary arteries with low radiation exposure is available to almost every patient, using computer tomography angiography (CTA)—for instance, with third-generation dual-source scanners. Based on our experience of examining more than 6000 patients, absolute contraindications no longer exist. CTA has long been recognized in the English-speaking world as a method of choice for detection of coronary heart disease (CHD). In Germany, CTA inexplicably has a subordinate role, despite unambiguous, evidence-based current data, and is not reimbursed by statutory health insurance.

In the meantime, computer programs using FFA-CT (fractional flow reserve from computed tomography) are available that use the standard available CTA datasets to provide reliable statements on the hemodynamic relevance of stenosis, which are comparable to those from invasive pressure measurements. Patent litigation is still in the way of wider dissemination outside of trials. However, there are currently technical limitations, for example, in the presentation of main stem stenosis (similar to conventional FFR) or heavy calcification (2).

Nevertheless, it can be stated that an examination of coronary arteries is possible for a large number of

patients that is fast, non-invasive, less burdensome, less expensive and provides reliable information about the hemodynamics of a stenosis. Taking into consideration the fact that, in contrast to cardiac catheter, CTA allows a distinction to be made between vulnerable and stable plaques, and that FFA-CT allows the entire coronary tree, rather than just individual segments, to be checked for significant stenosis, at least the perspective of diagnostic CTA / FFA-CT playing a significant role in a proposed revascularization therapy must be considered.

DOI: 10.3238/arztebl.2019.0460a

References

1. Gaede L, Möllmann H, Rudolph T, Rieber J, Boenner F, Tröbs M: Coronary angiography with pressure wire and fractional flow reserve—state of the art in the diagnosis of coronary stenosis. *Dtsch Arztebl Int* 2019; 116: 205–11.
2. Fairbairn TA, Nieman K, Akasaka T, et al.: Real-world clinical utility and impact on clinical decision-making of coronary computed tomography angiography-derived fractional flow reserve: lessons from the ADVANCE Registry. *Eur Heart J* 2018; 39: 3701–11.

Prof. Dr. med. Johannes Görich
Prof. Dr. med. Sebastian Buß
 Radiology practice, Heidelberg, Germany
 jgoerich@t-online.de

Conflict of interest statement

Prof. Görich has received speaking honoraria from Siemens Healthineers. Prof. Buß declares that no conflict of interest exists.

In Reply:

We thank our colleagues Görich and Buß for pointing out the issue of computed tomography angiography (CTA) of coronary arteries and the possibility to estimate FFR (fractional flow reserve) by simulations based on CT datasets (1, 2). Indeed, the largest study published to date showed that, in comparative trials, FFA-CT scores with respect to a threshold of 0.80 were 81% consistent with invasively measured FFR for assessment of coronary lesions (1).

Incidentally, simulation of FFR is also successful on the basis of invasive coronary angiogram with similar software solutions (“FFR angio”) (3). However, these methods are not widespread and at least in part still in the development stage. In addition, patients enrolled in the comparative studies are selected, and prospective randomized controlled trials that demonstrate

prognostic significance, as for invasive FFR, do not exist for either FFR-CT or FFR-angio.

These are certainly procedures whose further development, validation, and performance in clinical use should be followed with interest. However, it should be noted that FFR-CT in particular would occupy a slightly different place in the spectrum of cardiac examination procedures than coronary angiography using pressure wire measurement. CT angiography is already an examination procedure that is usually used early in the diagnostic sequence, especially in patients with a low-to-moderate pre-test probability of coronary heart disease. Invasive coronary artery diagnostics with the optional addition of pressure wire measurement are performed in non-invasive pathological examinations and for patients with high probability of pre-testing as well as for those with a clinical picture of an acute

coronary syndrome. In this respect, the—potential—areas of application overlap only partially.

DOI: 10.3238/arztebl.2019.0460b

References

1. Nørgaard BL, Leipsic J, Gaur S, et al.: Diagnostic performance of non-invasive fractional flow reserve derived from coronary computed tomography angiography in suspected coronary artery disease: the NXT trial (Analysis of Coronary Blood Flow Using CT Angiography: Next Steps). *J Am Coll Cardiol* 2014; 63: 1145–55.
2. Coenen A, Lubbers MM, Kurata A, et al.: Coronary CT angiography derived fractional flow reserve: methodology and evaluation of a point of care algorithm. *J Cardiovasc Comput Tomogr* 2016; 10: 105–13.

3. Fearon WF, Achenbach S, Engstrom T, et al.: Accuracy of fractional flow reserve derived from coronary angiography. *Circulation* 2019; 139: 477–84.
4. Gaede L, Möllmann H, Rudolph T, Rieber J, Boenner F, Tröbs M: Coronary angiography with pressure wire and fractional flow reserve—state of the art in the diagnosis of coronary stenosis. *Dtsch Arztebl Int* 2019; 116: 205–11.

On Behalf of the Authors

Dr. med. Monique Tröbs

Universitätsklinikum Erlangen, Germany

Department of Medicine 2—Cardiology and Angiology

monique.troebs@uk-erlangen.de

Conflict of interest statement

Dr. Tröbs has received consultant honoraria from Abbott Vascular.

CLINICAL SNAPSHOT

Giant Bulla Imitating Tension Pneumothorax

A 61-year-old man with a long history of exercise dyspnea came to the emergency department with acute respiratory distress at rest. His oxygen saturation was below 90 %, and there were no breathing sounds on the left side. Posteroanterior chest radiography in expiration with the patient standing erect showed almost complete absence of lung markings in the left thorax with residual lung and what looked like a pneumothorax line in the basal portion, together with displacement of the mediastinum to the right (*Figure 1*).

We assumed the diagnosis of left tension pneumothorax and inserted a thoracic drain. Repeat radiography showed no change. Computed tomography (coronal thorax–lung window) then revealed a giant bulla (*Figure 2*). Thoracic drain placement had not perforated the bulla.

The patient's respiratory distress was gradually relieved by means of oxygen administration. Later, elective resection of the bulla took place. The patient was then symptom-free at rest and could perform normal physical activities.

Patients with an extensive history of pulmonary emphysema and conspicuous findings on radiography should, provided their clinical status permits, be examined by thoracic CT for confident differentiation between pneumothorax and bullous lesions.

Dr. med. Matthias Schumann, Ahmad Hamed, Dr. med. Thomas Kohl, Klinik für Thorax- und Gefäßchirurgie, Zentralklinik Bad Berka, matthias.schumann@zentraklinik.de

Conflict of interest statement: The authors declare that no conflict of interest exists.

Translated from the original German by David Roseveare.

Cite this as: Schumann M, Hamed A, Kohl T: Giant bulla imitating tension pneumothorax. *Dtsch Arztebl Int* 2019; 116: 461. DOI: 10.3238/arztebl.2019.0461