Published in final edited form as:

Cardiol Young. 2019 July ; 29(7): 945–953. doi:10.1017/S1047951119001161.

Persistence of Right Ventricular Dysfunction and Altered Morphometry in Asymptomatic Preterm Infants through One Year of Age: Cardiac Phenotype of Prematurity

Collin T. Erickson¹, Meghna D. Patel^{2,3}, Swati Choudhry^{2,4}, Karl Stessy Bisselou¹, Tim Sekarski², Mary Craft¹, Ling Li¹, Afif El Khuffash⁵, Aaron Hamvas⁶, Shelby Kutty, MD^{1,7}, Gautam K. Singh², Philip T. Levy^{2,8}

¹Department of Pediatric Cardiology and Cardiovascular Surgery, University of Nebraska College of Medicine and Children's Hospital and Medical Center, Omaha, Nebraska ²Department of Pediatrics, Washington University School of Medicine, St. Louis, Missouri ³Department of Pediatrics, Stanford University School of Medicine, Stanford, CA. ⁴Section of Pediatric Cardiology, Texas Children's Hospital, Baylor College of Medicine, Houston, Texas ⁵Department of Neonatology, The Rotunda Hospital & School of Medicine, Royal College of Surgeons in Ireland, Dublin, Ireland. ⁶Department of Pediatrics, Northwestern University Feinberg School of Medicine, Chicago, IL. ⁷Division of Pediatric Cardiology, Johns Hopkins Medical Institutions, Baltimore, MD. ⁸Division of Newborn Medicine, Boston Children's Hospital and Department of Pediatrics, Harvard Medical School, Boston, Massachusetts, USA

Abstract

Introduction: Prematurity impacts myocardial development and may determine long-term outcomes. The objective of this study was to test the hypothesis that preterm neonates develop right ventricle dysfunction and adaptive remodeling by 32 weeks postmenstrual age that persists through 1 year corrected age.

Materials and Methods: A subset of 80 preterm infants (born <29 weeks) was selected retrospectively from a prospectively enrolled cohort and measures of right ventricle systolic function and morphology by two-dimensional echocardiography were assessed at 32 weeks postmenstrual age and at 1 year of corrected age. Comparisons were made to 50 term infants at 1 month and 1 year of age. Sub-analyses were performed in preterm-born infants with bronchopulmonary dysplasia and/or pulmonary hypertension.

Result: In both term and preterm infants, right ventricle function and morphology increased over the 1st year ($p < 0.01$). The magnitudes of right ventricle function measures were lower in preterm-born infants at each time period ($p < 0.01$ for all) and right ventricle morphology indices were wider

Correspondence Address: Philip T. Levy MD, Division of Newborn Medicine, Department of Pediatrics, Boston Children's Hospital, 300 Longwood Avenue | Hunnewell 436, Boston, Massachusetts 02115, Phone: (617) 919 6682, philip.levy@childrens.harvard.edu.

Conflicts of Interest: None.

Ethical Standards: The Institutional Review Board for human studies at Washington University and the University of Nebraska Medical Center approved the study. Written informed consents were obtained from the parents/guardians of the study subjects.

in all preterm infants by 1 year corrected age, irrespective of lung disease. Measures of a) right ventricle function were further decreased, and b) morphology increased through 1 year in preterm infants with bronchopulmonary dysplasia and/or pulmonary hypertension ($p<0.01$).

Conclusion: Preterm infants exhibit abnormal right ventricle performance with remodeling at 32 weeks postmenstrual age that persists through 1 year corrected age, suggesting a less developed intrinsic myocardial function response following preterm birth. The development of bronchopulmonary dysplasia and pulmonary hypertension leave a further negative impact on right ventricle mechanics over the first year of age.

Keywords

Right ventricular function; Prematurity; Strain Imaging; Echocardiography; Neonates

INTRODUCTION

Right ventricular mechanics in preterm neonates begin to undergo maturational changes during the early postnatal period that have prognostic implications for clinical status and long-term outcomes[1-3]. The determinants of right ventricle performances involve interactions between its contractility, afterload, preload, and morphology[4]. Preterm infants have maladaptive changes in right ventricle performance when compared to healthy term infants[2, 5, 6], where important morphological differences that explain the relative dysfunction are exacerbated as prematurity exposes the right ventricle to the stress of postnatal cardiovascular performance before development is concluded[7]. Recent work has validated emerging quantitative methods to characterize neonatal right ventricle performance,[1, 2] including: myocardial tissue deformation by two- dimensional speckle-tracking echocardiography[1, 2], fractional area of change[8], tricuspid annular plane systolic excursion[5], pulmonary artery acceleration time (estimate of right ventricle afterload)[9], right ventricle areas[8, 10] and linear dimensions[9]. Our group has also recently demonstrated that common cardiorespiratory complications of prematurity, including bronchopulmonary dysplasia and pulmonary hypertension, negatively impact right ventricle performance over the first year of age[1, 8, 9]. However, longitudinal comparisons of right ventricle performance in robust healthy birth cohorts and birth cohorts affected by prematurity from the neonatal period through 1 year of age are lacking, but are necessary prerequisites for clinical adoption in evaluating pathologic changes and progression.

We hypothesize that developmental and maturational disruption to the right ventricle from preterm birth results in dysfunction and remodeling as a distinct pathophysiology from bronchopulmonary dysplasia and pulmonary hypertension that is evident at 32 weeks postmenstrual age and persists through 1 year corrected age. Accordingly, the objective of this study was to identify and longitudinally track right ventricle performance using emerging measures of right ventricle mechanics in a cohort of infants born preterm (asymptomatic and those with bronchopulmonary dysplasia and/or pulmonary hypertension) and compare them with healthy infants born at term at 1 month and 1 year of age.

MATERIALS AND METHODS

Study Design and Population

To define the cardiac phenotype we retrospectively analyzed prospectively acquired data from two cohorts: (1) a cohort of preterm-born infants and (2) a cohort of full term-born infants. Each cohort was recruited at birth and longitudinally followed to 1 year of age. The cohort of 80 preterm infants were enrolled through the Prematurity and Respiratory Outcomes Program at Washington University School of Medicine/Saint Louis Children's Hospital neonatal intensive care unit between August 2011 and November 2013. The detailed design of the Prematurity and Respiratory Outcomes Program Study ([ClinicalTrials.gov: NCT01435187](https://clinicaltrials.gov/ct2/show/study/NCT01435187)) has been described[11, 12]. This was a retrospective secondary analysis of prospectively acquired data. Preterm subjects were eligible for inclusion in this study if they were born between 23 0/7 weeks and 28 6/7 weeks gestational age and had an echocardiogram at 32 weeks postmenstrual age and 1 year corrected age (Subjects were excluded if they had any suspected congenital anomalies of the airways, congenital heart disease (except age-appropriate atrial septal communications), intrauterine growth restriction (< 3rd percentile), or small for gestational age (birth weight < 10th centile for gestation). We have provided a list of terms (Supplementary Information, Appendix 1) that describe the definitions for length of gestation and age in the perinatal period based on the policy statement from the American Academy of Pediatrics.[13]

A cohort of 50 healthy, age, weight, and sex matched infants born at term (>37 weeks gestation at birth) were included as the control group from the University of Nebraska Medical Center from February 2013 to September 2016. Echocardiograms were obtained at 1 month and 1 year of age. The data was also prospectively acquired and retrospectively matched to the preterm cohort. Specific exclusion criteria for the control cohort consisted of multiple gestation pregnancy, maternal self-reported use of illegal drugs, maternal smoking, maternal diabetes, antepartum hemorrhage, pre-eclampsia, and/or proven chorioamnionitis. Infants with intrauterine growth restriction, perinatal acidosis (umbilical cord pH < 7.10), suspected congenital or chromosomal anomalies, neuro-muscular disorders, abnormal cardiac rhythms, heart failure, and infants readmitted to the hospital within the first year of age were also excluded. The Institutional Review Board for human studies at Washington University and the University of Nebraska Medical Center approved the study. Written informed consents were obtained from the parents/guardians of the study subjects.

Clinical demographics

For both cohorts, perinatal clinical information and demographic characteristic were obtained. In addition for the preterm infants, clinical cardio-respiratory characteristics during the two echocardiographic assessments were collected and included: systolic and diastolic blood pressure, heart rate, oxygen requirements, and oxygen saturation. The following clinical outcomes were also obtained in the preterm cohort: intraventricular hemorrhage classified according to Papile Classification[14], necrotizing enterocolitis with radiological evidence of pneumatosis, retinopathy of prematurity (stage 2 or higher), pulmonary hypertension[15], and bronchopulmonary dysplasia (using a modified definition

of the 2001 National Institutes of Health bronchopulmonary dysplasia workshop, Supplementary Information, Appendix 2)[12].

Echocardiography

Echocardiograms were performed with commercially available ultrasound imaging systems (Vivid 7 and E9; General Electric Medical Systems, Milwaukee, Wisconsin) at each center. Images were obtained at 32 weeks postmenstrual age and 1 year corrected age in infants born preterm and at 1 month and 1 year of age in infants born at term. One designated experienced pediatric cardiac sonographer at each center obtained all the echocardiographic images using a phased array transducer (7.5–12 megahertz)[16]. The echocardiographic images were acquired using a standardized image acquisition protocol based on the guidelines of the American Society of Echocardiography[17, 18]. The image data were acquired digitally and stored in raw Digital Imaging and Communications in Medicine cine-loop format for offline analysis for the following functional and morphometric parameters.

Deformation Analysis

Two-Dimensional Speckle Tracking Imaging—Myocardial mechanics were analyzed by the quantification of right ventricle free wall longitudinal strain (%) and systolic longitudinal strain rate (1/sec) using previously published image acquisition and data analysis protocols from our laboratories[16, 19]. Right ventricle segmental longitudinal strain was obtained from the segments at the apical, mid-ventricular, and basal levels of the right ventricle free wall from a focused right ventricle apical 4-chamber view[1, 20]. Peak strain for each index was measured as end-systolic strain at the closure of the pulmonic valve[21],(Supplementary Information, Figure S1). Two observers (PL and CE), who were blinded to the maternal and infant clinical and cardio-respiratory conditions, analyzed deformation using vendor customized commercially available software (EchoPAC; General Electric Medical Systems, Waukesha, WI, USA, version 112).

Other measurements of right ventricle performance

We assessed and compared additional indices of 1) right ventricle function: tricuspid annular plane systolic excursion,[5] fractional area of change [8]; 2) right ventricle morphology: right ventricle areas from the apical 4-chamber view in systole and diastole[8, 10], right ventricle linear dimensions at the basal and mid ventricular levels and in the longitudinal direction from the apical 4-chamber view at end-diastole[9, 17, 18]; and 3) right ventricle afterload[9]: pulmonary artery acceleration time and ratio of pulmonary acceleration time to right ventricle ejection time between the term and preterm infants at each time point[9]. All measures were generated according to guidelines of the American Society of Echocardiography[17, 18] and recent validated protocols in neonates[2].

Confounders

We evaluated the effects of common cardiorespiratory complications of prematurity, including bronchopulmonary dysplasia and pulmonary hypertension, on right ventricle structure and function.[1, 8, 9] We utilized a modified definition of the 2001 National Institutes of Health bronchopulmonary dysplasia workshop[12] (Supplementary

Information, Appendix 2) and a broad echocardiogram-based definition of pulmonary hypertension. [1, 15], in the assessment of their contributions on right ventricle performance indices at 32 weeks postmenstrual age, and at 1 year corrected age. Preterm infants were classified with pulmonary hypertension at 32 weeks postmenstrual age if they had one or more of the following conventional echocardiographic measures of pulmonary hemodynamics: 1) an estimated right ventricle systolic pressure > 40 mmHg, using the Doppler velocity of tricuspid regurgitation, 2) a ratio of right ventricle systolic pressure to systemic blood pressure $> 1/2$; 3) bidirectional or right to left shunt at the atrial level or patent ductus arteriosus; 4) septal wall flattening as defined as decreased septal curvature into the right ventricle at end systole, or 5) dilated/hypertrophy of the right ventricle and right atrial morphology. The independent effect of the risk factors for bronchopulmonary dysplasia and pulmonary hypertension, including gestational age, sex, total oxygen days, length of hospital stay, perinatal factors, neonatal morbidities (necrotizing enterocolitis, intraventricular hemorrhage, and retinopathy of prematurity), and presence of a patent ductus arteriosus at 32 weeks postmenstrual age[22] were also accounted for in the analysis.

Statistical Analyses

Continuous data were summarized as mean $\pm$ standard deviation or as medians (interquartile range) as appropriate. Categorical data were summarized as count and percent. Continuous variables of right ventricle longitudinal strain, longitudinal systolic strain rate, and segmental longitudinal strain at the base, mid-ventricular, and apical levels were tested for normality using the Shapiro-Wilk test and histogram of the data. Analysis of variance tests were used to compare the changes in measures of right ventricle function and structure amongst preterm infants with and without bronchopulmonary dysplasia and/or pulmonary hypertension at 32 weeks postmenstrual age with term born control infants. Student t-test and Bonferroni adjustment where appropriate were separately utilized to compare the patterns between uncomplicated preterm infants and those with bronchopulmonary dysplasia and/or pulmonary hypertension. All outcome variables with non-normal distributions were analyzed in simple comparisons using Wilcoxon rank sum tests or Kruskal-Wallis one-way analysis of variance for tests with more than two independent groups. Chi-square tests (or Fisher Exact test as appropriate) were used to assess the association between categorical variable. Univariate analysis was used to determine the best predictors to enter in the model and then backward step-wise regression was performed to assess the independent effect of gestational age, sex, total oxygen days, length of stay, and common neonatal morbidities, while adjusting for weight at examination. We performed stepwise regression to analyze the influence of pulmonary hypertension and bronchopulmonary dysplasia on right ventricle performance patterns at 32 weeks postmenstrual age and 1 year corrected age [15, 23]. All results were considered statistically significant for values of $p < 0.05$. The statistical analysis was performed using SAS v9.4 (SAS Institute, Cary, NC).

RESULTS

One hundred and thirty infants (80 preterm and 50 term) were included in this study (Figure 1). We compared the maternal and infant demographic characteristics and clinical data between preterm infants with and without bronchopulmonary dysplasia in Table 1, and with

and without pulmonary hypertension in Supplementary Information, Table S1. Clinical data not described in the Table 1 or Table S1 is detailed in the Supplementary Information.

Maturational patterns of RV performance in term and preterm infants

The magnitudes of right ventricle free wall longitudinal strain, fractional area of change, tricuspid annular plane systolic excursion, pulmonary artery acceleration time, and the ratio of pulmonary artery acceleration time to right ventricle ejection time increased from 32 weeks postmenstrual age to 1 year corrected age in the preterm infants and from 1 month of age to 1 year of age in the term infants ($p < 0.01$ for all measures, Figure 2A, Table 2). The magnitude of RV free wall systolic longitudinal strain rate in infants born at term increased from 1 month to 1 year of age ($p < 0.01$) but remained unchanged in infants born preterm over the same time period ($p = 0.5$), (Figure 2B). A significant base-to-apex (the magnitude was highest-to-lowest) segmental longitudinal strain gradient ($p < 0.001$) was observed in the right ventricle free wall in both the preterm and term infants, (Figure 3). Right ventricle morphology measures of systolic and diastolic areas, basal, mid-cavity, and apex-to-base linear dimensions also increased in preterm infants from 32 weeks postmenstrual age to 1 year corrected age ($p < 0.01$ for all measures), (Table 2).

Comparison of RV performance between term and preterm infants

All preterm infants had significantly lower values of right ventricle free wall longitudinal strain, longitudinal systolic strain rate, tricuspid annular plane systolic excursion, pulmonary artery acceleration time, pulmonary artery acceleration time / right ventricular ejection time ($p < 0.01$ for all measures) than those of term infants at 1 month and 1 year of age (Table 2, Figure 2). Segmental longitudinal strain at the basal, mid-ventricular, and apical levels of the right ventricle free wall were also decreased at 1 month and 1 year corrected age in preterm infants compared to term infants ($p < 0.01$ for all measures). As expected, the systolic and diastolic areas, the basal, mid-cavity, and the longitudinal diameter from the apex to the base were all smaller in the preterm infants at 1 month of age ($p < 0.01$ for all measures). At 1 year of age, all morphometric measures, except the longitudinal diameter from the apex to the base were larger in the preterm infants ($p < 0.01$ for all measures). All values for comparisons can be found in Table 2.

Bronchopulmonary dysplasia

Since measures of right ventricle function performed better in infants born at term, pairwise comparison was conducted using the Bonferroni method to detect any difference in right ventricle function in a) preterm infants with and without bronchopulmonary dysplasia and b) preterm infants with and without pulmonary hypertension at 32 weeks postmenstrual age. Adjustments for gestational age at birth, sex and heart rate revealed significantly lower values in the magnitudes of right ventricle free wall longitudinal strain, longitudinal systolic strain rate, fractional area of change, tricuspid annular plane systolic excursion, pulmonary artery acceleration time, pulmonary artery acceleration time / right ventricle ejection time, and segmental longitudinal strain at the basal, mid-ventricular, and apical levels of the right ventricle free wall at 32 weeks postmenstrual age in preterm infants with bronchopulmonary dysplasia ($p < 0.0001$ for all). The same patterns persisted at 1 year corrected age ($p < 0.0001$ for all). The systolic and diastolic areas were larger in the preterm infants with

bronchopulmonary dysplasia at both time points. The basal, mid-cavity, and longitudinal diameter from the apex to the base were similar between the preterm groups at 32 weeks postmenstrual, but the basal and mid-cavity diameter were significantly larger in the preterm infants with bronchopulmonary dysplasia by 1 year corrected age ($p < 0.01$ for both measures).

Pulmonary hypertension

In preterm infants with and without pulmonary hypertension there was also a decrease in right ventricle free wall longitudinal strain, free wall longitudinal systolic strain rate, fractional area of change, tricuspid annular plane systolic excursion, pulmonary artery acceleration time, pulmonary artery acceleration time / right ventricle ejection time, and segmental longitudinal strain at the basal, mid-ventricular, and apical levels of the right ventricle free wall at 1 year corrected age in those with pulmonary hypertension ($p < 0.0001$ for all). The base-to-apex segmental longitudinal strain gradients were preserved in all preterm infants at both time points, irrespective of the presence of pulmonary hypertension. These trends persisted even after adjusting for the presence of bronchopulmonary dysplasia (for all measure, $\beta > 2.4$, $p < 0.01$) and presence of patent ductus arteriosus (for all measures $\beta > 2.6$, $p < 0.01$). The dimensions and areas were similar between preterm infants with and without pulmonary hypertension at 32 weeks postmenstrual age. Right ventricle basal and mid-ventricular diameter, and both right ventricle areas were increased in preterm infants with pulmonary hypertension at 1 year corrected age ($p < 0.01$ for both). There was no difference in the longitudinal right ventricle dimension from the apex to the base at any time point between the groups. (Supplementary Information, Table S1).

Term vs. uncomplicated Preterm

We compared the measures between term infants and preterm infants without bronchopulmonary dysplasia and/or pulmonary hypertension at each time point. Adjustments for weight at exam, sex and heart rate revealed significant lower values of the magnitudes of right ventricle free wall longitudinal strain, free wall longitudinal systolic strain rate at both time points, and decreases in segmental longitudinal strain at the basal and mid-ventricular levels of the right ventricle free wall at 1 month, and all segmental longitudinal strain levels by 1 year of age. At 1 month, fractional area of change was similar between the term and preterm without bronchopulmonary dysplasia and/or pulmonary hypertension, but tricuspid annular plane systolic excursion was decreased ($p < 0.01$). At 1 year of age, tricuspid annular plane systolic excursion was similar, and fractional area of change was decreased in the preterm infants without bronchopulmonary dysplasia and/or pulmonary hypertension. All morphometric measures were decreased, as expected, in preterm infants at 1 month of age, but by 1 year of age, the systolic and diastolic areas, and the basal and mid-cavity diameters were significantly larger in the preterm infants without bronchopulmonary dysplasia and/or pulmonary hypertension ($p < 0.01$ for both measures). The longitudinal diameter from the apex to the base was similar between the groups.

DISCUSSION

In this study we observed right ventricle dysfunction in asymptomatic infants born preterm at 32 weeks postmenstrual age (~ 1 month of age) that persisted to 1 year corrected age when compared to healthy infants born at term over the same time period. In addition, right ventricle morphologic differences existed between preterm and term infants at 1 year of age, irrespective of the degree of lung disease during the neonatal period. Subclinical right ventricle dysfunction assessed using deformation, fractional area of change, and tricuspid annular plane systolic excursion, as well as alteration in right ventricle morphology appears to be a distinct pathology of prematurity that is present even after resolution of bronchopulmonary dysplasia and pulmonary hypertension.

Right ventricle mechanics undergo postnatal physiologic adaptations over the first year of age that have long-term influence on myocardial performance[5]. Preterm birth is associated with alterations in myocardial performance that appear in the neonatal period[5], persist through 1 year corrected age[1], early childhood[24], and are present at an early adult age with a disproportionate increase in ventricular mass and decrease in right ventricle function[3]. In neonates, strain analysis has been shown to be more sensitive in detecting early myocardial changes than conventional echocardiography in health and disease states[2]. Kwon et al. found a lower magnitude of right ventricle strain in children born preterm with a history of pulmonary hypertension during the neonatal period compared to non-pulmonary hypertension children born preterm[25]. Lewandowski showed reduced right ventricle strain in adults born preterm compared to adults born at term, which was gestational age dependent, but unrelated to the premature lung disease[3]. In the present study we observed decreased right ventricle free wall longitudinal and longitudinal systolic strain rate in preterm infants compared to term infants at 32 weeks postmenstrual age and 1 year corrected age. Taken all together, these studies from the early neonatal period through adulthood suggest that the factors that contribute to the right ventricle maladaptation following premature birth are complex and may differ substantially between infants with different antenatal and postnatal experiences.

The exact mechanism that explains why preterm infants have subclinical right ventricle dysfunction and alterations in right ventricle morphology that persists to 1 year of age is not yet known. We propose two theories that may explain this observation: 1) changes in postnatal loading conditions[26], and 2) loss of the third trimester myocardial structural programming[7]. The exposure of an immature preterm right ventricle to a sustained increase in hemodynamic load of postnatal circulation (increased right ventricle afterload), at a time in the development when the right ventricle primarily supports a low-resistance circulation, may induce myoarchitectural adaptation that can lead to changes in geometry, structure, and function, “a process known as ventricular remodeling”[27]. In this study, measures of right ventricle function were decreased and indices of right ventricle morphology (right ventricle areas and right ventricle linear dimensions/length at the base and mid-cavity regions) were increased in preterm infants by 1 year corrected age. Functional measures were further decreased and morphological measures increased at 32 weeks postmenstrual in preterm infants with bronchopulmonary dysplasia, suggesting early dysfunction and remodeling[5]. However, these observations do not explain the increased

incidence of subclinical right ventricle dysfunction and remodeling in a subset of premature infants without apparent clinical signs of lung disease of prematurity[1, 15, 28]. Since premature birth results in the loss of the normal maturational process that occurs in utero during the third trimester, the evidence of disproportionate increase in ventricular size and decrease in right ventricle function in the postnatal period[5], over and above what would be expected, indicates that loss of a primary myocardial structural programming event may also lead to postnatal ventricular remodeling[7]. Bensely et al. demonstrated that preterm birth may lead to an abrupt reduction in cardiomyocyte cell division, adversely impact upon the final number of cardiomyocytes, reduce right ventricle functional reserve, and impair the reparative capacity of the myocardium[7].

Strain is influenced by preload and afterload[1, 2, 29]. Strain rate correlates well with load independent measures of contractility and is a more accurate reflection of intrinsic myocardial contractile function[30-32]. The force-frequency relationship reflects alteration of the contractile force of cardiac myocytes through the neurohumoral effects, which are influenced by changes in the frequency of stimulation[31]. Both preclinical[31, 32] and human studies[30, 31] have demonstrated that strain rate has a force-frequency relationship behavior that is consistent with a measure of contractility that is mostly load independent within physiological ranges. We speculate that the decreased strain in preterm infants reflects the myocardium's response to postnatal-loading conditions, according to the Frank Starling phenomenon, while the decreases in strain rate possibly reflects a less mature myocardial programming mechanism stemming from preterm birth that alters contractility, according to the force-frequency relationship. While all measures of right ventricle function increased from 32 weeks postmenstrual to 1 year corrected age, strain rate was unchanged in preterm infants, and decreased in preterm infants with bronchopulmonary dysplasia and/or pulmonary hypertension, indicating a less developed intrinsic myocardial function response in preterm infants.

In the right ventricle free wall a base-to-apex segmental longitudinal strain gradient (highest-to-lowest magnitude of segmental longitudinal strain values) has been observed in extreme preterm population[1], late preterm infants[5], healthy term neonates[33], infants[6], children[34], and adults[35]. This physiologic pattern reflects the base-to-apex alignment of the dominant deep longitudinal layers of the right ventricle that allows for greater longitudinal shortening during contraction[34]. All preterm infants had a decreased magnitude of segmental longitudinal strain at the basal and mid-ventricular, and apical levels of the right ventricle free wall at 32 weeks postmenstrual age and 1 year corrected age. Focusing on the right ventricle free wall provides an assessment of the predominant wall contributing to right ventricle contractile function[35]. We have previously demonstrated that the presence of bronchopulmonary dysplasia and/or pulmonary hypertension during the neonatal period leaves a negative impact of global and regional strain values[1]. However, even those preterm infants without significant morbidity have decreased regional strain values, further suggesting that the postnatal period appears to be a critical period of cardiac development and may offer a window for interventions to prevent the long-term cardiovascular consequences of preterm birth[5]. We observed a difference of segmental longitudinal strain at the basal and mid-ventricular levels of the right ventricle free wall between preterm infants without pulmonary hypertension and/or bronchopulmonary

dysplasia compared to healthy term infants. Interestingly, the apical segmental longitudinal strain values were similar between these two groups. We have previously demonstrated that only the basal and mid-ventricular segments of the right ventricle free wall increase throughout maturation, while the apical segmental longitudinal strain was unchanged from birth to 1 year of age in healthy term infants[36]. Since the right ventricle is composed on superficial oblique fibers that are arranged in parallel with the AV groove and turn obliquely toward the cardiac apex and continue into the superficial myofibers of the left ventricle[37], the expected physiologic increase of segmental longitudinal strain at the right ventricle side of the apical free wall is likely masked by the contribution of the left ventricle to the contraction patterns of the right ventricle free wall[16]. Therefore, we would expect not to see a difference between the segmental longitudinal strain at the apical level, unless a significant disease process like pulmonary hypertension or bronchopulmonary dysplasia alters these “physiological” gradients.

Study Limitations

We acquired images of the right ventricle free wall from a right ventricle focused apical four-chamber view. Recently, several authors have outlined an approach for assessment of right ventricle function that utilizes three-chamber right ventricle-focused views to capture the inlet, apical, and outlet in one image view[38, 39]. Currently there is no consensus on the best approach, but more studies have utilized the right ventricle focused apical four chamber method[34]. Since the right ventricle lacks a middle layer of circumferential fibers, and the major contribution to ejection fraction and stroke volume during systole is provided by the dominant deep longitudinal fiber shortening, right ventricle circumferential and radial strain patterns were not assessed[40]. Future work is needed to investigate right ventricle circumferential, radial, and circumferential-longitudinal shear strain (rotational mechanics) in neonates with the use of three-dimensional deformational approaches[40]. Finally, we chose to exclude intrauterine growth restricted neonates from the analysis, since it has been documented that fetal growth restriction alone may induce cardiac remodeling that persists beyond the neonatal period[41].

CONCLUSION

This study tracks the maturational patterns of global and regional right ventricle performance by two- dimensional speckle-tracking echocardiography in extremely preterm infants and term infants over the first year of age. Infants born preterm exhibit decreased right ventricle performance at 1 month of age that persists to 1 year of age when compared to infants born at term. Right ventricle remodeling is evident by 1 year corrected age in preterm infants, irrespective of the degree of neonatal lung disease, suggesting a less developed intrinsic myocardial function response following preterm birth.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Financial Support: This study was supported by a grant from Premature and Respiratory Outcomes Program (NIH 1U01 HL1014650, U01 HL101794),

REFERENCES

1. Levy PT, El-Khuffash A, Patel MD, et al. Maturational patterns of systolic ventricular deformation mechanics by two-dimensional speckle tracking echocardiography in preterm infants over the first year of age. *J Am Soc Echocardiogr* 2017;30:685–698. [PubMed: 28433214]
2. Breatnach CR, Levy PT, James AT, Franklin O, El-Khuffash A. Novel echocardiography methods in the functional assessment of the newborn heart. *Neonatology* 2016;110:248–260. [PubMed: 27287615]
3. Lewandowski AJ, Bradlow WM, Augustine D, Davis EF, Francis J, Singhal A, et al. Right ventricular systolic dysfunction in young adults born preterm. *Circulation* 2013;128:713–720. [PubMed: 23940387]
4. El-Khuffash A, Mcnamara PJ. Hemodynamic Assessment and Monitoring of Premature Infants. *Clin Perinatol* 2017; 44:377–393. [PubMed: 28477667]
5. Aye CYL, Lewandowski AJ, Lamata P, al. Disproportionate cardiac hypertrophy during early postnatal development in infants born preterm. *Pediatr Res* 2017;82:36–46. [PubMed: 28399117]
6. Schubert U, Muller M, Abdul-Khaliq H, Norman M. Preterm birth is associated with altered myocardial function in infancy. *J Am Soc Echocardiogr* 2016;29:670–678. [PubMed: 27156903]
7. Bensley JG, Moore L, De Matteo R, Harding R, Black MJ. Impact of preterm birth on the developing myocardium of the neonate. *Pediatr Res*. 2017.
8. Levy PT, Dioneda B, Holland MR, et al. Right Ventricular Function in Preterm and Term Neonates: Reference Values for Right Ventricle Areas and Fractional Area of Change. *J Am Soc Echocardiogr*. 2015;28:559–569. [PubMed: 25753503]
9. Levy PT, Patel MD, Choudhry S, Hamvas A, Singh GK. Evidence of Echocardiographic Markers of Pulmonary Vascular Disease in Asymptomatic Preterm Infants at One year of Age *Journal of Pediatr* 2018
10. James AT, Corcoran JD, Franklin O, EL-Khuffash AF. Clinical utility of right ventricular fractional area change in preterm infants. *Early Hum Dev* 2016;92:19–23. [PubMed: 26619069]
11. Pryhuber GS, Maitre NL, Ballard RA, Cifelli D, Davis SD, Ellenberg JH, et al. Prematurity and respiratory outcomes program (PROP): study protocol of a prospective multicenter study of respiratory outcomes of preterm infants in the United States. *BMC Pediatr* 2016;10:15:37.
12. Poindexter BB, Feng R, Schmidt B, et al. Comparisons and Limitations of Current Definitions of Bronchopulmonary Dysplasia for the Prematurity and Respiratory Outcomes Program. *Ann Am Thorac Soc* 2016; 12:1822–1830.
13. Engle WA. Age Terminology During the Perinatal Period. *Pediatrics* 2004;114:1362–4. [PubMed: 15520122]
14. Papile LA, Burstein J, Burstein R, Koffler H. Incidence and evolution of subependymal and intraventricular hemorrhage: a study of infants with birth weights less than 1,500 gm. *J Pediatr* 1978; 92:529–354. [PubMed: 305471]
15. Mourani PM, Sontag MK, Younoszai A, et al. Early Pulmonary Vascular Disease in Preterm Infants at Risk for Bronchopulmonary Dysplasia. *Am J Respir Crit Care Med* 2015; 191:87–95. [PubMed: 25389562]
16. Levy PT, Holland MR, Sekarski TJ, Hamvas A, Singh GK. Feasibility and reproducibility of systolic right ventricular strain measurement by speckle-tracking echocardiography in premature infants. *J Am Soc Echocardiogr* 2013;26:1201–13. [PubMed: 23880052]
17. Lang RM, Badano LP, Mor-Avi V, et al. Recommendations for chamber quantification. *J Am Soc Echocardiogr* 2015;28:1–39. [PubMed: 25559473]
18. Mitchell C, Rahko PS, Blauwet LA, Canaday B, Finstuen JA, Foster MC, et al. Guidelines for Performing a Comprehensive Transthoracic Echocardiographic Examination in Adults:

Recommendations from the American Society of Echocardiography. *J Am Soc Echocardiogr* 2019;32; 2019: 1–64. [PubMed: 30282592]

19. Sanchez AA, Levy PT, Sekarski TJ, Hamvas A, Holland MR, Singh GK. Effects of frame rate on two-dimensional speckle tracking-derived measurements of myocardial deformation in premature infants. *Echocardiography* 2015;32:839–847. [PubMed: 25109389]
20. Cade WT, Tinius RA, Reeds DN, Patterson BW, Cahil AG. Maternal Glucose and Fatty Acid Kinetics and Infant Birth Weight in Obese Women With Type 2 Diabetes. *Pediatr Res*. 2017;82:768–775. [PubMed: 28604759]
21. Voigt JU, Pedrizzetti G, Lysyansky P, et al. Definitions for a common standard for 2D speckle tracking echocardiography: consensus document of the EACVI/ASE/Industry Task Force to Standardize Deformation Imaging. *J Am Soc Echocardiogr* 2015;28: 183–193. [PubMed: 25623220]
22. Schena F, Francescato G, Cappelleri A, et al. Association between Hemodynamically Significant Patent Ductus Arteriosus and Bronchopulmonary Dysplasia. *J Pediatr* 2015;166:1488–1492. [PubMed: 25882876]
23. Mourani PM, Abman SH. Pulmonary Hypertension and Vascular Abnormalities in Bronchopulmonary Dysplasia. *Clin Perinatol* 2015; 42:839–55. [PubMed: 26593082]
24. Mohlert LA, Hallberg J, Broberg O, et al. The Preterm Heart in Childhood: Left Ventricular Structure, Geometry, and Function Assessed by Echocardiography in 6-Year-Old Survivors of Periviable Births. *J Am Heart Assoc* 2018;7:e007742. [PubMed: 29353231]
25. Kwon HW, Kim H-S, An HS, et al. Long-Term Outcomes of Pulmonary Hypertension in Preterm Infants with Bronchopulmonary Dysplasia. *Neonatology* 2016;11:181–189.
26. Kang SJ, Kim M, Hwang SJ, Kim HJ. Progression of Right Ventricular Systolic Dysfunction Detected by Myocardial Deformation Imaging in Asymptomatic Preterm Children. *J Cardiovasc Ultrasound* 2015;25:98.
27. Haddad F, Hunt SA, Rosenthal DN, Murphy DJ. Right Ventricular Function in Cardiovascular Disease, Part II: Pathophysiology, Clinical Importance, and Management of Right Ventricular Failure. *Circulation* 2008;11:1436–1448.
28. Weismann CG, Asnes JD, Bazzi-Asaad A, Tolomeo C, Ehrenkranz RA, Bizzarro MJ. Pulmonary hypertension in preterm infants: results of a prospective screening program. *J Perinatol* 2017;37:572–577. [PubMed: 28206997]
29. Geyer H, Caracciolo G, Abe H, et al. Assessment of myocardial mechanics using speckle tracking echocardiography: fundamentals and clinical applications. *J Am Soc Echocardiogr* 2010;23:351–369. [PubMed: 20362924]
30. Breatnach CR, El-Khuffash A, James A, McCallion N, Franklin O. Serial measures of cardiac performance using tissue Doppler imaging velocity in preterm infants <29weeks gestations. *Early Hum Dev* 2017;108:33–39. [PubMed: 28388489]
31. Alvarez SV, Fortin-Pellerin E, Alhabdan M, Lomelin JS, Kantoch M, Atallah J et al. Strain Rate in Children and Young Piglets Mirrors Changes in Contractility and Demonstrates a Force-Frequency Relationship. *J Am Soc Echocardiogr*. 2017;30:797–806. [PubMed: 28602206]
32. Ferferieva V, Van den Bergh A, Claus P, et al. The relative value of strain and strain rate for defining intrinsic myocardial function. *Am J Physiol Heart Circ Physiol* 2012;302:H188–H195. [PubMed: 22081696]
33. Schubert U, Muller M, Norman M, Abdul-Khaliq H. Transition from fetal to neonatal life: Changes in cardiac function assessed by speckle-tracking echocardiography. *Early Hum Dev*. 2013; 89:803–8. [PubMed: 23948155]
34. Levy PT, Sanchez A, Machefsky A, Fowler S, Holland MR, Singh GK. Normal ranges of right ventricular systolic and diastolic strain measures in children: a systematic review and meta-analysis. *J Am Soc Echocardiogr* 2014;27:549–60. [PubMed: 24582163]
35. Fine NM, Chen L, Bastiansen PM, et al. Reference values for right ventricular strain in patients without cardiopulmonary disease: a prospective evaluation and meta-analysis. *Echocardiography* 2015;32:787–96. [PubMed: 25323591]

36. Erickson CT, Levy PT, Craft M, Li L, Danford D, Kutty S. Maturational patterns in right ventricular strain mechanics from the fetus to the young infant. *Early Hum Dev* 2019;129:23–32. [PubMed: 30616038]
37. Sadeghpour A, Alizadehasl A. The Right Ventricle: A Comprehensive Review From Anatomy, Physiology, and Mechanics to Hemodynamic, Functional, and Imaging Evaluation. *Arch Cardiovasc Imaging* 2015;3: e35717.
38. Jain A, Mohamed A, El-Khuffash A, et al. A comprehensive echocardiographic protocol for assessing neonatal right ventricular dimensions and function in the transitional period: normative data and z scores. *J Am Soc Echocardiogr* 2014;27: 1293–1304. [PubMed: 25260435]
39. Forsha D, Risum N, Kropf PA, et al. Right ventricular mechanics using a novel comprehensive three-view echocardiographic strain analysis in a normal population. *J Am Soc Echocardiogr* 2014;27:413–422. [PubMed: 24508364]
40. Atsumi A, Seo Y, Ishizu T, et al. Right Ventricular Deformation Analyses Using a Three-Dimensional Speckle-Tracking Echocardiographic System Specialized for the Right Ventricle. *J Am Soc Echocardiogr* 2016;29:402–411 [PubMed: 26879190]
41. Sarvari SI, Rodriguez-Lopez M, Nunez-Garcia M, et al. Persistence of Cardiac Remodeling in Preadolescents With Fetal Growth Restriction. *Circ Cardiovasc Imaging*. 2017;10

Figure 1:
Study subject flow diagram.

Figure 2:

Comparison of right ventricle free wall longitudinal systolic strain (A) and strain rate (B) in term and preterm infants at 1 month of age and 1 year of age. * $P < 0.01$ vs. Term; § $P < 0.01$ vs. Preterm without BPD; + $P < 0.001$ vs. first echocardiogram. **In the preterm-born cohort, echocardiograms were performed at 32 weeks postmenstrual age with a median of 38 days (interquartile range: 33 - 45). In the term-born cohort, echocardiograms were performed at a median of 32 days (interquartile range: 31 - 35). ***In the preterm-born cohort, echocardiograms were performed at 1 year corrected of 494 days (interquartile range: 452 - 521). These numbers reflect actual days of age, but corrected age in days for all preterm infants was a median of 381 days (interquartile range: 355- 424). In the term-born cohort, echocardiograms were performed at a median of 371 days (interquartile range: 365 - 380).

Figure 3:

Comparison of right ventricle segmental longitudinal systolic strain in term and preterm infants at 1 month and 1 year of age. * $P < 0.01$ vs. Term; § $P < 0.01$ vs. Preterm without BPD; + $P < 0.001$ vs. first echocardiogram. **In the preterm-born cohort, echocardiograms were performed at 32 weeks postmenstrual age with a median of 38 days (interquartile range: 33 - 45). In the term-born cohort, echocardiograms were performed at a median of 32 days (interquartile range: 31 - 35). *** In the preterm-born cohort, echocardiograms were performed at 1 year corrected of 494 days (interquartile range: 452 - 521). These numbers reflect actual days of age, but corrected age in days for all preterm infants was a median of 381 days (interquartile range: 355- 424). In the term-born cohort, echocardiograms were performed at a median of 371 days (interquartile range: 365 - 380).

Table 1:

Infant characteristics and clinical outcomes

Characteristics	Preterm cohort (n=80)	No BPD (n=32)	BPD (n=48)	P value
Birthweight (g)	890 [765,1010]	960 [825,1125]	863[717,980]	0.002
Birthweight strata (g)				
500 - 749 (n=16)	660 [590,690]	690 [655,705]	640 [578,685]	0.23
750 - 999 (n=41)	890 [825,955]	853 [800,945]	895 [835,940]	0.37
1000 - 1250 (n=23)	1125 [905,1265]	1140 [1050,1260]	1100 [1040,1220]	0.61
Gestational age	27 [26,28]	27 [26,28]	26 [25,27]	< 0.01
Male sex	39 (49%)	18 (56%)	21 (41%)	0.22
Multiple gestation	14 (18%)	3 (9%)	11 (23%)	0.17
Infant Race				0.66
White	38 (48%)	12 (38%)	26 (54%)	
Black	42 (52%)	20 (62%)	22 (46%)	
Asian	0	0	0	
Other	0	0	0	
Maternal Smoking	16 (20%)	5 (16%)	11 (23%)	0.48
Antenatal steroids	77 (96%)	32 (100%)	45 (94%)	0.16
Surfactant Replacement Therapy	80 (100%)	32 (100%)	48 (100%)	0.84
Cesarean section	55 (69%)	21 (66%)	34 (71%)	0.86
Maternal complications				
Gestational diabetes mellitus	4 (5%)	1 (3%)	3 (6%)	0.56
Gestational hypertension	22 (28%)	12 (38%)	10 (21%)	0.28
Prolonged rupture of membranes	38 (17%)	15 (15%)	23 (19%)	0.12
Chorioamnionitis	8 (10%)	4 (13%)	4 (8%)	0.51
Preeclampsia	22 (28%)	10 (31%)	12 (25%)	0.51
Pulmonary hypertension	12 (15%)	4 (12.5%)	8 (17%)	0.45
Presence of ductus arteriosus *	9 (11%)	3 (9%)	6 (13%)	0.36
Necrotizing enterocolitis	7 (9%)	2 (6%)	5 (10%)	0.56
Retinopathy of prematurity **	30 (38%)	6 (19%)	24 (50%)	0.02
Intraventricular hemorrhage ***	9 (11%)	2 (6%)	7 (15%)	0.54
Total oxygen days (Neonatal ICU)	87 [46,109]	38 [18,58]	100 [88,115]	< 0.001
Length of stay (Neonatal ICU)	93 [79,114]	80 [69,100]	106 [91,120]	< 0.001

Data are presented as median [interquartile range], or number (Percentage). Data are presented as column percentages

* Presence of ductus arteriosus at 32 weeks postmenstrual age

** Retinopathy of prematurity threshold (stage 2 of higher)

*** Intraventricular hemorrhage (Grade 3 or 4)

Table 2:

Echocardiographic data of term and preterm infants at 1 month of age and 1 year of age

Variables		First Echocardiogram (1 month of age)				Second Echocardiogram (1 year of age)		
	Term (n=50)	All Preterm (n=80)	Preterm w/out bronchopulmonary dysplasia (n=32)	Preterm with bronchopulmonary dysplasia (n=48)	Term (n=50)	All Preterm (n=80)	Preterm w/out bronchopulmonary dysplasia (n=32)	Preterm with bronchopulmonary dysplasia (n=48)
Age at echocardiogram (days)	32 (31, 35)	38 (33, 45)	33 (28, 41)	39 (36, 49)	371 (365, 380)	494 (452, 521) ^{**}	494 (450, 525) ^{**}	500 (468, 535) ^{**}
Weight at echocardiogram (kg)	4.6 (4.3, 4.9)	1.4 (1.2, 1.6) [*]	1.4 (1.3, 1.7) [*]	1.4 (1.1, 1.6) [*]	9.5 (8.7, 10.5) ⁺	10.1 (9.4, 11.1) ⁺	9.7 (9.1, 10.7) ⁺	10.1 (9.4, 11.2) ⁺
Body surface area (m ²)	0.27 (0.26, 0.28)	0.26 (0.24, 0.29)	0.26 (0.25, 0.30)	0.26 (0.24, 0.28)	0.46 (0.42, 0.48) ⁺	0.45 (0.44, 0.47) ⁺	0.45 (0.43, 0.47) ⁺	0.45 (0.44, 0.48) ⁺
Heart rate (beats/minutes)	155 (129, 166)	157 (147, 169)	162 (145, 170)	156 (147, 168)	127 (120-136) ⁺	129 (118, 141) ⁺	130 (121, 139) ⁺	127 (115, 143) ⁺
Right ventricle function								
Fractional area of change (%)	36 (31, 39)	32 (28, 36) [*]	36 (32-43)	30 (26, 33) [§]	44 (40, 48) ⁺	34 (32, 36) ⁺ *	39 (36, 43) ⁺ *	33 (31, 34) ⁺ §
Tricuspid annular plane systolic excursion (mm)	10 (9, 10)	8 (7, 9) [*]	9 (8, 10) [*]	8 (7, 9) [§]	18 (17, 19) ⁺	16 (15, 18) ⁺ *	18 (16, 19) ⁺	16 (15, 17) ⁺ §
Free wall longitudinal strain (%)	-26 (-25, -29)	-19 (-17, -21) [*]	-22 (-20, -23) [*]	-18 (-17, -19) [§]	-34 (-33, -37) ⁺	-27 (-26, -29) ⁺ *	-30 (-29, -31) ⁺ *	-27 (-26, -28) ⁺ §
Segmental longitudinal strain (%)								
Basal	-30 (-34, -28)	-24 (-22, -25) [*]	-26 (-24, -27) [*]	-23 (-22, -24) [§]	-38 (-35, -44) ⁺	-30 (-28, -31) ⁺ *	-30 (-28, -30) ⁺ *	-29 (-28, -30) ⁺ §
Mid-ventricular	-27 (-25, -30)	-22 (-21, -23) [*]	-23 (-22, -24) [*]	-21 (-20, -22) [§]	-36 (-34, -39) ⁺	-28 (-27, -29) ⁺ *	-29 (-28, -30) ⁺ *	-27 (-26, -28) ⁺ §
Apical	-20 (-16, -24)	-19 (-17, -21) [*]	-21 (-20, -22)	-18 (-17, -19) [§]	-28 (-21, -36) ⁺	-25 (-24, -27) ⁺ *	-26 (-27, -28) ⁺ *	-24 (-23, -25) ⁺ §
Free wall longitudinal strain rate (1/sec)	-2.7 (-2.4, -3.0)	-1.9 (-1.7, -2.3) [*]	-2.2 (-1.9, -2.4) [*]	-1.8 (-1.6, -2.1) [§]	-2.9 (-2.7, -3.4) ⁺	-1.8 (-1.6, -2.3) [*]	-2.3 (-1.8, -2.7) ⁺ *	-1.6 (-1.5, -1.9) ⁺ §
Right ventricle Morphology								

Variables	First Echocardiogram (1 month of age)				Second Echocardiogram (1 year of age)			
	Term (n=50)	All Preterm (n=80)	Preterm w/out bronchopulmonary dysplasia (n=32)	Preterm with bronchopulmonary dysplasia (n=48)	Term (n=50)	All Preterm (n=80)	Preterm w/out bronchopulmonary dysplasia (n=32)	Preterm with bronchopulmonary dysplasia (n=48)
Basal length (cm)	1.7 (1.5, 1.8)	1.3 (1.2, 1.4)*	1.2 (1.1, 1.3)*	1.3 (1.2, 1.5)*	2.1 (2.0, 2.3) [†]	2.4 (2.2, 2.6) ^{†*}	2.4 (2.1, 2.5) ^{†*}	2.5 (2.3, 2.7) ^{†*§}
Mid-cavity length (cm)	1.2 (1.0, 1.2)	1.1 (1.0, 1.2)*	1.0 (0.9, 1.2)	1.2 (1.0, 1.3)	1.5 (1.4, 1.6) [†]	2.0 (1.9, 2.2) ^{†*}	1.9 (1.6, 2.0) ^{†*}	2.1 (1.9, 2.3) ^{†*§}
Major length (cm)	3.5 (3.3, 3.6)	2.0 (1.9, 2.2)*	2.0 (1.8, 2.2)*	2 (1.9, 2.2)*	4.0 (3.7, 4.2) [†]	3.9 (3.7, 4.2) [†]	3.9 (3.6, 4.2) [†]	3.9 (3.8, 4.2) [†]
Systolic area (cm2)	2.5 (2.3, 2.8)	1.7 (1.6, 1.9)*	1.6 (1.5, 1.8)*	1.8 (1.6, 2.0)*§	3.6 (3.3, 4.0) [†]	4.5 (4.1, 4.7) ^{†*}	4.1 (3.9, 4.2) ^{†*}	4.6 (4.5, 4.8) ^{†*§}
Diastolic area (cm2)	3.8 (3.4, 4.3)	2.6 (2.3, 2.8)*	2.6 (2.4, 2.8)*	2.6 (2.3, 3.0)*§	6.4 (6.1, 6.8) [†]	6.7 (6.1, 7.1) ^{†*}	6.7 (6.4, 7.0) ^{†*}	6.9 (6.8, 7.2) ^{†*§}
RV Afterload								
Pulmonary artery acceleration time (mseconds)	74 (67, 81)	52 (39 - 60)*	56 (45, 66)*	43 (35, 53)*§	96 (92, 100) [†]	73 (65, 78) ^{†*}	79 (69, 82) ^{†*}	65 (60, 69) ^{†*§}
Pulmonary artery acceleration time / Right ventricle ejection time	0.34 (0.31, 0.38)	0.26 (0.21, 0.31)*	0.31 (0.23, 0.36)*	0.23 (0.20, 0.27)*§	0.36 (0.33, 0.38) [†]	0.31 (0.26, 0.34) ^{†*}	0.31 (0.30, 0.36) ^{†*}	0.27 (0.25, 0.29) ^{†*§}

All data presented as median (interquartile range)

* P < 0.01 vs. Term;

§ P < 0.01 vs. Preterm without bronchopulmonary dysplasia;

[†] P < 0.001 vs. first echocardiogram,

Note: right ventricle strain rate significantly decreased from the 1st echocardiogram to the 2nd echocardiogram for Preterm infants with bronchopulmonary dysplasia (P < 0.001)

** These numbers reflect actual days of age, but corrected age (days) for all preterm infants was 381 (355, 424), and for preterm without bronchopulmonary dysplasia was 381 (357, 428) and preterm infants with bronchopulmonary dysplasia was 398 (377, 428).