Published in final edited form as:

Orthop Nurs. 2019 ; 38(2): 150–156. doi:10.1097/NOR.0000000000000525.

Multimodal Pain Management for Major Joint Replacement Surgery

Victoria M. Goode, PhD, CRNA¹, Brett Morgan, DNP, CRNA¹, Virginia C. Muckler, DNP, CRNA, CHSE¹, Michael P. Cary Jr., PhD, RN¹, Christine E. Zdeb, BSN, RN, ONC¹, Michael Zychowicz, DNP, ANP, ONP, FAAN, FAANP¹

¹Duke University School of Nursing, Durham, NC

Abstract

Effective pain management for orthopedic major joint replacement is key to achieving earlier recovery, better functioning, and high rates of patient satisfaction. In an effort to decrease opioid dependency, practitioners are turning to multimodal pain management, which involves the use of multiple analgesic agents and techniques. In order to utilize this technique, a patient's history of and preoperative consumption of medications to treat pain impacts the success of this regimen. Multimodal pain management involves the use of NSAIDs, acetaminophen, NMDA antagonists, gabapentin, serotonin inhibitors, regional techniques and opioids as needed. It is necessary for the nurse to understand the mechanism of pain and how the multimodal adjuncts target the pain response in order to benefit the patient's perioperative course as well as their postoperative and discharge management.

Introduction

Several decades ago, major joint replacement of the knee, hip, and shoulder were considered medical breakthroughs. Now, hundreds of thousands of these orthopedic surgeries are performed annually. In 2014, there were more than 400,000 total knee replacements (Delanois et al., 2017), 230,000 total hip replacements (Gwam et al., 2017), and 45,000 total shoulder replacements (Trofa, Rajaei, & Smith, 2014) performed to treat osteoarthritis, relieve pain, restore function, and improve quality of life—increases of 100 to 300 percent since 1990. As the population continues to age, obesity rates rise, and technological innovations advance, the demand for major joint replacements are projected to skyrocket nationally (Padegimas et al., 2015; Kurtz, Ong, Lau, & Bozic, 2014). Despite the noted growth in these surgical procedures, the lack of attention paid to nursing interventions to relieve post-operative pain and reduce complications is surprising. Increasing evidence suggests poorly managed perioperative pain may cause post-operative complications, implant failure, and hospitalization. Therefore, effective pain management for orthopedic joint replacement is key to achieving earlier recovery, better functioning, and high rates of

Corresponding Author: Victoria M. Goode, PhD, CRNA, DUMC 3322 307 Trent Drive, Durham, NC 27710.

Conflicts of Interest

The Author(s) declare(s) that there is no conflict of interest.

patient satisfaction. This article provides an overview of multimodal pain management in patients who undergo major joint replacement surgery.

The Opioid Crisis

The Centers for Disease Control (CDC) estimates that more than 126 million adults, or approximately 56% of the adult population, suffer from some form of pain ranging in severity and occurrence (daily or near daily) ((Nahin, 2015; Makary, Overton, & Wang, 2017). Treating their pain has resulted in a record number of prescriptions (Nahin, 2015; Makary, Overton, & Wang, 2017; Roth, Boateng, Berken, Carlyle, & Vadivelu, 2018), including highly addictive opioids as an adjunct for pain relief in adults suffering from chronic pain. As a result, Bohnert, Guy, & Losby (2018), reported a 300% increase in opioid-related deaths from 1999 to 2016. This statistic includes illicit opioid use but also includes the use of opioids for the relief of chronic pain in non-surgical patients who have such diseases as rheumatoid arthritis and osteoarthritis. The danger posed by opioids has placed healthcare providers in the precarious position of balancing the drug's use for pain relief and its potential long-term impact on patients' lives.

In 2016, the CDC responded to the growing opioid crisis in the number of patients suffering from acute and chronic pain and the overuse of opioids by publishing national guidelines to aid in prescription practices (Dowell, Haegerich, & Chou 2016). According to the CDC guidelines, four key questions are important to consider when prescribing opioids as a treatment for pain: the effectiveness of opioid use, the perceived risks and benefits, an acceptable criterion for dosing patterns, and strategies to control the risk for long-term use and addiction.

In an effort to decrease opioid dependency, practitioners are turning to multimodal pain management, which involves the use of multiple analgesic agents and techniques that act on different pain mechanisms to provide pain relief and improve quality of life with less opioid consumption (Brooks, Freter, Bowles, & Amirault 2017). In addition to decreasing opioid use, multimodal pain management techniques used postoperatively reduce prolonged sedation for the orthopedic surgical patient. The techniques also alleviate many adverse side effects of opioids including nausea, vomiting, itching, respiratory depression and (Wick, Grant & Wu, 2017).

Mechanisms of Pain

Pain is a noxious stimulus that begins in injured nerve fiber endings and is transmitted via the central nervous system where it is perceived by the brain via primary nociceptor fibers (Type A, B or C) through the dorsal horn of the spinal cord where a response is mediated. Type A delta fibers and C fibers are the mediators of the pain response and sensation. This activity projects the response upward to the spinal cord and finally the thalamus. The activation of the pain response involves the efferent pathway where neurons take the painful sensation back from the level of the spinal cord to the site of injury at the peripheral nociceptors (Naglehout & Plaus 2014, Barash et al., 2013, Marks, Shah, Patkar, Masand, Park, & Pae, 2009, Parvizi & Bloomfield, 2013). The tissue injury mediates the release of

substances such as bradykinin, histamine, lactic acid, prostaglandins, serotonin, glutamate and most importantly, substance P as well as an excitatory ion channel (vanilloid receptor) secondary to tissue injury manifesting in an inflammatory response and the pain sensation (Naglehout & Plaus, 2014, Barash et al., 2013).

The nurses' understanding of the role of opioids, NSAIDS, acetaminophen, serotonin inhibitors, gabapentin, NMDA antagonists and local anesthetics in the multimodal pain management of the orthopedic surgical patient is crucial to positive patient outcomes (Naglehout & Plaus 2014, Barash et al., 2013, Marks, Shah, Patkar, Masand, Park, & Pae, 2009, Parvizi & Bloomfield, 2013). A patient's physiologic responses to pain include activation of the body's sympathetic nervous system resulting in increased heartrate, increased oxygen consumption, hypertension, increased cardiac output, changes in the rate and depth of respiration, slowing of the gastrointestinal and genitourinary tract leading to nausea, vomiting, and difficulty in urination and finally the increased secretion of cortisol. The use of non-steroidal anti-inflammatory drugs (NSAIDS) and aspirin interfere with the inflammatory process and the activation of prostaglandins thus limiting the magnitude of the inflammatory pain response. Serotonin works with inflammatory mediators as well and useful in the treatment of chronic pain. The use of local anesthetics hinder the ion channel's response to pain from the afferent pain fibers diminishing the perception of pain for the patient. Acute pain is most often addressed with opioids, which target five opioid receptors to modify pain signals and diminish pain perception (Naglehout & Plaus, 2014, Barash et al., 2013 Marks, Shah, Patkar, Masand, Park, & Pae, 2009, Parvizi & Bloomfield, 2013).

Preoperative Assessment

The use of opioids for surgery patients often begins well before surgery. Many joint replacement candidates delay surgery for as long as possible, often on the advice of their physicians. Instead, they opt for a regimen of anti-inflammatory and pain medication for weeks to years. A preoperative patient assessment is an opportunity for healthcare providers to identify baseline pain levels, current pain medication intake, successful past pain medication strategies, and past medical and surgical histories, all of which support our ability to educate the patient and are used by the anesthesia provider to formulate a personalized plan of care for perioperative pain management (American Society of Anesthesiologists (ASA), 2012). Ideally, healthcare providers will seek further detailed responses related to each question including: 1) pain levels at rest and with activity; 2) type of pain whether acute or chronic; 3) type, route, frequency, and quantity of pain medication consumed, 4) over-the-counter medications, medications prescribed to the patient, and any medications the patient takes that are not prescribed to the patient, 5) adverse effects associated with the pain regimen, 6) extent of pain relief various modalities achieve, and 7) alternative pain strategies such as acupuncture, chiropractic, yoga, aromatherapy, and hypnosis. The medical, surgical, and pharmacological intake that occurs during the preoperative assessment assists the provider to develop the clinical profile of the patient and guides the formulation of a perioperative plan of care including a multimodal pain management plan.

Armed with information regarding previous pain medication consumption, healthcare providers must consider whether their patients are opioid naïve or opioid tolerant (Food and Drug Administration, 2018) as they develop a multimodal pain regimen for proactive pain management rather than the traditional reactive approach. Studies have shown that opioid-tolerant patients or patients that receive opioids on a daily basis, often develop an exaggerated pain hypersensitivity known as opioid-induced hyperalgesia (Cooney & Broglio, 2017; Soffin et al., 2017). Pain management for the opioid-tolerant patient requires combined anesthetic techniques such as anesthetic infiltration or regional anesthesia. While neither will eliminate the need for opioids in the orthopedic surgical patient, they may result in decreased opioid consumption rates (Schwenk, Pozek, & Viscusi, 2018). More recently, consensus guidelines support the use of ketamine infused at subanesthetic doses as a perioperative pain adjunct for opioid-tolerant patients due to its potent analgesic properties (Schwenk, Pozek, & Viscusi, 2018).

Opioid-naïve patients, (patients not receiving opioids on a daily basis), report lower post-surgical opioid consumption rates (Schwenk, Pozek, & Viscusi, 2018). However, despite their relatively low opioid consumption rates immediately following surgery, there is evidence that the potential for opioid abuse remains. In a retrospective study of more than 1 million opioid-naïve patients from 2008-2016, Brat et al. (2018) noted that each opioid refill increased the potential for opioid misuse by more than 40%. Thus, healthcare providers must promote non-opioid pain control options and consider the need for opioid prescription refills with extreme caution.

Modalities and techniques for perioperative pain management-

Table 1 outlines the techniques utilized for multi-modal pain management therapies for the orthopedic surgery patient which often include NSAIDS such as aspirin, naproxen, ketorolac, celecoxib, meloxicam, diclofenac, and ibuprofen (Donahue, Bradbury, Zychowicz, & Muckler, 2018; Brooks, Fretter, Bowles, & Amirault, 2017; Song, 2017; Canata, Casale, & Chiey, 2016; Chou, et al., 2016; ASA 2012). Analgesia resulting from NSAID administration is due to inhibited synthesis and release of prostaglandins and inhibited cyclooxygenase 2 (COX-2) that results in a suppressed inflammatory response (Elmallah, et al., 2018). A 10-year database review from 2006 to 2016 of total hip and knee arthroplasties (N=1,540,462) showed that NSAIDS, as part of a multi-modal regimen, were effective and resulted in improved patient outcomes including decreased opioid consumption (Memtsoudis et al., 2018). Although non-opioid analgesics such as acetaminophen and ketorolac are commonly incorporated into multimodal pain management plans, healthcare providers must consider the gastrointestinal side effects such as bleeding, and an increased risk for renal failure, stroke, and heart attack that are associated with NSAIDS (U.S. Food and Drug Administration, 2015).

NMDA antagonists: Ketamine as part of a multi-modal pain management plan

Ketamine can be used as part of multi-modal pain management. Intravenous (IV) ketamine infusions and boluses have been successfully used in a multi-modal approach to manage pain for orthopedic surgery throughout the perioperative period (Chin & Lewis, 2018; Kadic,

Haren, Wilder-Smith, Bruhn, Driessen, de Waal Malefijt, 2016; Alviar, Hale, & Dungca, 2016; Martinez et al., 2014). Ketamine, a N-methyl-D-aspartate (NMDA) receptor antagonist and dissociative agent, is a potent analgesic yet lacks respiratory depressant effects (Kadic et al., 2016). Low-dose ketamine infusions have been shown to lower pain scores and decrease opioid consumption rates (Zhang, Shi, & Jia, 2018; Kadic et al., 2016; Martinez, et al., 2014). Thus, the use of these infusions is recommended for surgical pain in opioid-tolerant orthopedic patients, patients with sickle cell pain, and patients with sleep apnea (Schwenk, et al., 2018). The common dosage of an intraoperative IV ketamine infusion and bolus ranges from 0.1 to 0.5 mg/kg/hour and 0.3 to 0.5 mg/kg, respectively (Schwenk, et al., 2018; Vadivelu et al., 2016).

Gabapentin as part of a multi-modal pain management approach for orthopedic surgery

Oral gabapentin is often included as a component of multimodal pain regimens. Gabapentin is an anticonvulsant originally developed to treat epileptic seizures resistant to traditional therapies. While the mechanism is not fully understood, gabapentin is believed to alter neurotransmission by interrupting voltage-gated calcium channels in the dorsal horn of the spinal cord. This leads to decreased effects due to hyper-excitability in those regions caused by tissue damage (Hah et al., 2018).

Studies have shown that gabapentin is effective in decreasing the need for postoperative opioids in orthopedic surgical patients. Specific to procedures on lower extremities, research has shown that patients who receive gabapentin preoperatively as well as during the postoperative period report lower pain scores, require lower doses of opioids for breakthrough pain, and experience better range of motion and activity when compared to those patients who do not receive the medication (Clarke et al., 2014; Mardani-Kivi, Mobarakeh, Keyhani, Motlagh, & Ekhtiari, 2013). However, gabapentin has side effects that may limit its use with some patients. The literature suggests that the use of gabapentin to treat or prevent pain may be associated with dizziness, drowsiness, and confusion. This is more likely to occur in older patients. It appears that dose plays a large role in the development of these side effects, and using the lowest effective dose, specific to individual patients, decreases that risk (Fleet et al., 2018).

Continuous Peripheral Nerve Block for Perioperative Pain Control

A continuous peripheral nerve block (CPNB) is a regional anesthesia technique that involves the administration of local anesthetic to a targeted nerve through an indwelling percutaneous sheath that remains in place for an extended period of time (Ilfeld, 2017). As compared to a single-shot technique, indwelling catheters have demonstrated decreased post-operative pain scores, and lowered opioid consumption and opioid-related side effects for patients undergoing both upper- and lower-extremity orthopedic procedures. Researchers have also found that the use of indwelling catheters is associated with earlier ambulation and joint mobility (Arsoy et al., 2017; Ilfeld, 2017; Ullah, Samad, & Khan, 2014). Recently, the use of liposomal local anesthetics has become common. These longer-acting formulations allow for sustained release of local anesthetic at the nerve sheath for several days (Ulrich, Lavandero, Woods, & Early, 2014).

When caring for a patient receiving a CPNB, nurses should be aware of potential postoperative complications associated with their use. The most common complications include the development of a hematoma, falls, and infusion-induced local anesthetic toxicity (Elmallah et al., 2018; Elmallah et al., 2016). While the incidence of hematomas has decreased in recent years with the use of ultrasound-guided techniques for placing catheters, nurses should closely monitor patients with bleeding disorders or those on anticoagulant therapy (Gwam, 2018; Ilfeld, 2017). While it does not appear that patients receiving single-injection femoral nerve blocks are at a greater risk for falls, research has suggested that patients receiving a femoral CPNB for lower extremity surgery are four to five times more likely to fall during the post-operative period (Gwam, 2018; Ilfeld, 2017; Elmallah et al., 2018; Elmallah et al., 2016). Although infusion-induced local anesthetic toxicity is a rare adverse effect of CPNB, the ASA recommends that all nurses caring for patients receiving CPNB be able to recognize signs of local anesthetic toxicity (Ilfeld, 2017). These include changes in cognition, peri-oral paresthesia, audio-visual disturbances, cardiac rhythm disturbances, and potential respiratory and cardiovascular collapse (El-Boghdadly, Pawa, & Chin, 2018).

Post-operative multimodal pain management on the surgical unit

A multi-modal pain management regimen should continue as a patient transitions from the post-anesthesia care unit to the postoperative unit. This includes the considerations associated with PNBs, continued use of medications to treat neuropathic pain such as Keppra or gabapentin, acetaminophen, NSAIDs, and the sparingly use of opioids. Special consideration should be given to opioid-tolerant patients, whose postoperative course for pain management can be challenging.

Patients who receive PNBs demonstrate earlier mobilization, decreased length of stay, and greater participation in physical therapy (Gaffney, Pelt, Gililand & Peters, 2017). However, the need to establish fall-prevention strategies is essential as patients who receive a femoral nerve block are at increased risk for fall due to decreased quadriceps strength (Elmallah, et al., 2016). The nurse caring for patients with a PNB in place for more than 48 hours must be aware of the increased risk for both catheter site inflammation and infection (Nicolotti, Iotti, Fanelli, & Compagnone, 2016). Nurses should monitor PNBs carefully, observing for signs and symptoms of local anesthetic toxicity, and take measures to protect the insensate extremity from injury due to PNB (Bromberg, Dennis, & Gritsenko, 2017).

The continued post-operative use of oral medications such as acetaminophen, NSAIDs and gabapentin also play a role in postoperative pain management. Acetaminophen is considered safe and effective at a dosage of less than 3 grams per day (Malec & Shega, 2015), but should be used cautiously in patients with poor hepatic function. Gabapentin can be titrated in doses of 1,200 to 3,600 mg/day given in three doses (Lexicomp, 2018). Attention to the side effects associated with these medications should be considered especially in the elderly population (Devin et al., 2014; Gaffney et al., 2017; Malec & Shega, 2015). The scheduled dosing regimen of these pharmacologic adjuncts should reduce the need for opioids as well as limit the manifestation of physiologic derangements associated with opioid use.

When considering a plan that includes long-acting opioid medications, healthcare providers should use opioids sparingly in the early post-operative period. Some patients may require medication for the treatment of breakthrough acute pain. Nurses should be aware of the untoward side effects of this class of drugs, which include nausea, vomiting, constipation, ileus, respiratory depression, pruritis, delirium, and urinary retention (Parvizi & Bloomfield, 2013).

Alternative non-pharmacological pain control methods include the application of ice or the use of a cooling system to decrease postoperative pain (Ni et al., 2014). Patients can continue with around-the-clock application of ice after discharge to assist in pain management (Trasolini, McKnight, & Dorr, 2018).

Discharge Considerations

Pain management can affect length of stay (LOS) and discharge status. A multi-modal pain management strategy, initiated while the patient is in the hospital, should continue after discharge with the goal of decreasing opioid consumption. A study of shoulder surgery patients who received hospital-initiated PNBs demonstrated reduced opioid consumption for up to seven days post-operative (Beaussier et al., 2015). Patients who continue on multi-modal therapy after discharge reported significantly decreased pain at postoperative appointments and fewer pain-related hospital visits (Desai et al., 2017).

Nurses should develop realistic pain control expectations with patients. Opiates are useful, when judiciously used, to reduce a patient's pain and remove barriers to rehabilitation. Rather than a one-size-fits-all approach, opioid prescriptions at discharge should be tailored to patients' preoperative pain assessment and providers and nurses should take into consideration methods used to control preoperative pain, including opioid consumption during hospitalization (Sabatino, Kunkel, Ramkumar, Keeney, & Jevsevar, 2018). An interdisciplinary pain management team guided by the patient's surgeon can determine safe prescriptions at discharge (Genord, Frost, & Eid, 2017).

A post-discharge management plan may benefit from the inclusion of routine follow-up calls from nurses; these calls can reduce readmissions as well as aid in the identification of pain control needs and surgical complications while triaging the need for a return office visit to address those issues (Trasolini et al., 2018). Follow-up appointments should include the assessment of the patient's analgesic needs, the potential for surgical complications, and attempts to identify other etiologies of pain if present (Macintyre, Huxtable, Flint, & Dobbin, 2014).

Conclusion

The use of a multimodal pain approach in the orthopedic major joint replacement patient is an effective technique to guide care throughout the operative continuum. It is important for healthcare providers to have a clear understanding of the patient's previous history of pain management, if applicable, in order to utilize the technique. Nurses involved in care for this population need to have adequate education on the multimodal pain adjuncts including medications to aid in the plan of care for the patient.

Acknowledgments

The authors thank Jane Shealy, who provided editorial assistance for this manuscript.

References

- Alviar MJ, Hale T, Dungca M (2016). Pharmacologic interventions for treating phantom limb pain. *Cochrane Database of Systematic Reviews*, 10, CD006380. doi:10.1002/14651858.CD006380.pub3 [PubMed: 27737513]
- American Society of Anesthesiologists Task Force on Acute Pain Management. (2012). Practice guidelines for acute pain management in the perioperative setting: An updated report by the American Society of Anesthesiologists task force on acute pain management. *Anesthesiology*, 116(2), 248–273. doi:10.1097/ALN.0b013e31823c1030 [PubMed: 22227789]
- Arsoy D, Gardner MJ, Amanatullah DF, Huddleston JI III, Goodman SB, Maloney WJ, Bishop JA (2017). Continuous femoral nerve catheters decrease opioid-related side effects and increase home disposition rates among geriatric hip fracture patients. *Journal of Orthopaedic Trauma*, 31(6), e186–e189. doi:10.1097/bot.0000000000000854 [PubMed: 28538458]
- Barash PG (2013). *Clinical anesthesia* (7th ed.). Philadelphia, PA: Wolters Kluwer Health/Lippincott Williams & Wilkins.
- Bohnert A, Guy G, Losby J (2018). Opioid prescribing in the United States before and after the Centers for Disease Control and Prevention's 2016 Opioid Guidelines. *Annals of Internal Medicine*, 169(6), 367–375. doi:10.7326/M18-1243 [PubMed: 30167651]
- Brat GA, Agniel D, Beam A, Yorkgitis B, Bicket M, Homer M, Kohane I (2018). Postsurgical prescriptions for opioid naive patients and association with overdose and misuse: Retrospective cohort study. *British Medical Journal*, 360(j5790), 1–9. doi:10.1136/bmj.j5790
- Bromberg AL, Dennis JA, Gritsenko K (2017). Exparel/peripheral catheter use in the ambulatory setting and use of peripheral catheters postoperatively in the home setting. *Current Pain & Headache Reports*, 21(3), 13. doi:10.1007/s11916-017-0605-0 [PubMed: 28271332]
- Brooks E, Freter S, Bowles S, Amirault D (2017). Multimodal pain management in older elective arthroplasty patients. *Geriatric Orthopedic Surgery and Rehabilitation*, 8(3), 151–154. doi:10.1177/2151458517720297
- Canata GL, Casale V, Chiey A (2017). Pain management in total knee arthroplasty: Efficacy of a multimodal opiate-free protocol. *Joints*, 4(4), 222–227. doi:10.11138/jts/2016.4.4.222 [PubMed: 28217658]
- Chin KJ, Lewis S (2018). Opioid-free analgesia for posterior spinal fusion surgery using erector spinae plane (ESP) blocks in a multimodal anesthetic regimen. *Spine: An International Journal for the Study of the Spine*. Publish Ahead of Print. Published August 31, 2018. doi:10.1097/BRS.0000000000002855
- Chou R, Gordon DB, de Leon-Casasola OA, Resenberg JM, Bickler S, Brennan T, Wu CL (2016). Management of postoperative pain: A clinical practice guideline from the American Pain Society, the American Society of Regional Anesthesia and Pain Medicine, and the American Society of Anesthesiologists' Committee on Regional Anesthesia, Executive Committee, and Administrative Council. *Journal of Pain*, 17(2), 131–157. doi:10.1016/j.jpain.2015.12.008 [PubMed: 26827847]
- Clarke HA, Katz J, McCartney CJ, Stratford P, Kennedy D, Page MG, Kay J (2014). Perioperative gabapentin reduces 24 h opioid consumption and improves in-hospital rehabilitation but not post-discharge outcomes after total knee arthroplasty with peripheral nerve block. *British Journal of Anaesthesia*, 113(5), 855–864. doi:10.1093/bja/aeu20 [PubMed: 24980426]
- Cooney MF, Broglio K (2017). Acute pain management in opioid-tolerant individuals. *The Journal for Nurse Practitioners*, 13(6), 394–399. doi:10.1016/j.nurpra.2017.04.016
- Devin CJ, Lee DS, Armaghani SJ, Bible J, Shau DN, Martin PR, Ehrenfeld JM (2014). Approach to pain management in chronic opioid users undergoing orthopaedic surgery. *Journal of the American Academy of Orthopedic Surgery*, 22(10), 614–622. doi:10.5435/JAAOS-22-10-614
- Donahue RE, Bradbury GR, Zychowicz ME, Muckler VC (2018). Multimodal perioperative analgesia regimen to improve patient outcomes after total knee arthroplasty: A multidisciplinary quality

improvement project. *Journal of Perianesthesia Nursing*, 33(2), 138–152. doi:S1089-9472(16)30321-5 [PubMed: 29580593]

Dowell D, Haegerich TM (2016). Using the CDC guideline and tools for opioid prescribing in patients with chronic pain. *American Family Physician*, 93(12), 970–972. doi:d12732 [PubMed: 27304765]

El-Boghdadly K, Pawa A, Chin KJ (2018). Local anesthetic systemic toxicity: Current perspectives. *Local Regional Anesthesia*, 11, 35–44. doi:10.2147/lra.s154512 [PubMed: 30122981]

Elmallah RK, Cherian JJ, Pierce TP, Jauregui JJ, Harwin SF, Mont MA (2016). New and common perioperative pain management techniques in total knee arthroplasty. *The Journal of Knee Surgery*, 29(2), 169–178. doi:10.1055/s-0035-1549027 [PubMed: 25892004]

Elmallah R, Chughtai M, Khlopas A, Newman J, Stearns KL, Roche M, Mont MA (2018). Pain control in total knee arthroplasty. *The Journal of Knee Surgery*, 31(6), 504–513. [PubMed: 28719941]

Fleet JL, Dixon SN, Kuwornu PJ, Dev VK, Montero-Odasso M, Burneo J, Garg AX (2018). Gabapentin dose and the 30-day risk of altered mental status in older adults: A retrospective population-based study. *PloS One*, 13(3), e0193134. doi:10.1371/journal.pone.0193134 [PubMed: 29538407]

Food and Drug Administration (2015, 7). FDA drug safety communication: FDA strengthens warning that non-aspirin nonsteroidal anti-inflammatory drugs (NSAIDs) can cause heart attacks or strokes. Retrieved from U.S. Food and Drug Administration,

Food and Drug Administration. (2018). FDA blueprint for prescriber education for extended-release and long-acting opioid analgesics. Retrieved from <https://www.fda.gov/downloads/Drugs/DrugSafety/InformationbyDrugClass/UCM515636.pdf>

Gaffney CJ, Pelt CE, Gililland JM, Peters CL (2017). Perioperative pain management in hip and knee arthroplasty. *Orthopedic Clinics of North America*, 48(4), 407–419. doi:S0030-5898(17)30091-3 [PubMed: 28870302]

Gwam CU, Mistry JB, Delanois RE, Chughtai M, Khlopas A, Ramkumar PN, Mont MA (2017). Does patient perception of pain influence how patients rate their experience after total knee arthroplasty? *The Journal of Knee Surgery*, 30(7), 622–626. doi:10.1055/s-0037-1603794 [PubMed: 28666293]

Gwam CU, Mistry JB, Richards IV, Patel D, Patel NG, Thomas M, Delanois RE (2018). Does addition of adductor canal blockage to multimodal periarticular analgesia improve discharge status, pain levels, opioid use, and length of stay after total knee arthroplasty? *The Journal of Knee Surgery*, 31(2), 184–188. [PubMed: 28464196]

Hah J, Mackey SC, Schmidt P, McCue R, Humphreys K, Trafton J, Carroll I (2018). Effect of perioperative gabapentin on postoperative pain resolution and opioid cessation in a mixed surgical cohort: A randomized clinical trial. *JAMA Surgery*, 153(4), 303–311. doi:10.1001/jamasurg.2017.4915 [PubMed: 29238824]

Ilfeld BM (2017). Continuous peripheral nerve blocks: An update of the published evidence and comparison with novel, alternative analgesic modalities. *Anesthesia & Analgesia*, 124(1), 308–335. doi:10.1213/ane.0000000000001581 [PubMed: 27749354]

Kadic L, Haren FV, Wilder-Smith O, Bruhn J, Driessen JJ, de Waal Malefijt M (2016). The effect of pregabalin and s-ketamine in total knee arthroplasty patients: A randomized trial. *Journal of Anaesthesiology Clinical Pharmacology*, 32(4), 476–482.

Kurtz SM, Ong KL, Lau E, Bozic KJ (2014). Impact of the economic downturn on total joint replacement demand in the United States: Updated projections to 2021. *The Journal of Bone & Joint Surgery*, 96(8), 624–630. doi:10.2106/jbjs.m.00285 [PubMed: 24740658]

Makary MA, Overton HN, Wang P (2017). Overprescribing is major contributor to opioid crisis. *BMJ (Clinical Research Ed.)*, 359, j4792. doi:10.1136/bmj.j4792

Malec M, Shega JW (2015). Pain management in the elderly. *Medical Clinics of North America*, 99(2), 337–350. doi:10.1016/j.mcna.2014.11.007 [PubMed: 25700587]

Mardani-Kivi M, Mobarakeh MK, Keyhani S, Motlagh KH, Ekhtiari KS (2013). Is gabapentin effective on pain management after arthroscopic anterior cruciate ligament reconstruction? A triple blinded randomized controlled trial. *Archives of Bone Joint Surgery*, 1(1), 18–22. [PubMed: 25207278]

- Marks DM, Shah MJ, Patkar AA, Masand PS, Park GY, Pae CU (2009). Serotonin-norepinephrine reuptake inhibitors for pain control: Premise and promise. *Current Neuropharmacology*, 7(4), 331–336. doi:10.2174/157015909790031201 [PubMed: 20514212]
- Martinez V, Cymerman A, Ammar SB, Fiaud JF, Rapon C, Poindessous F, Fletcher D (2014). The analgesic efficiency of combined pregabalin and ketamine for total hip arthroplasty: A randomised, double blind, controlled study. *Anaesthesia*, 69(1), 46–52. doi:10.1111/anae.12495 [PubMed: 24320856]
- Memtsoudis SG, Poeran J, Zubizarreta N, Morwald EE, Mariano ER, Mazumdar M (2018). Association of multimodal pain management strategies with perioperative outcomes and resource utilization: A population-based study. *Anesthesiology*, 128(5), 891–902. doi:10.1097/ALN.0000000000002132 [PubMed: 29498951]
- Nagelhout JJ, Plaus KL (2014). *Nurse anesthesia* (5 ed.). St. Louis, MO: Elsevier/Saunders.
- Nahin RL (2015). Estimates of pain prevalence and severity in adults: United States, 2012. *The Journal of Pain: Official Journal of the American Pain Society*, 16(8), 769–780. doi:10.1016/j.jpain.2015.05.002 [PubMed: 26028573]
- Ni S, Jiang W, Guo L, Jin Y, Jiang T, Zhao Y, Zhao J (2018). Cryotherapy on postoperative rehabilitation of joint arthroplasty. *Knee Surgery, Sports Traumatology, Arthroscopy*, 2015(23), 3354–3361. doi:10.1007/s00167-014-3135-x
- Nicolotti D, Lotti E, Fanelli G, Compagnone C (2016). Perineural catheter infection: A systematic review of the literature. *Journal of Clinical Anesthesia*, 35, 123–128. doi:10.1016/j.jclinane.2016.07.025 [PubMed: 27871508]
- Padegimas EM, Maltenfort M, Lazarus MD, Ramsey ML, Williams GR, Namdari S (2015). Future patient demand for shoulder arthroplasty by younger patients: National projections. *Clinical Orthopaedics and Related Research*, 473(6), 1860–1867. doi:10.1007/s11999-015-4231-z [PubMed: 25758376]
- Parvizi J, Bloomfield MR (2013). Multimodal pain management in orthopedics: Implications for joint arthroplasty surgery. *Orthopedics*, 36(2 Suppl.), 7–14. doi:10.3928/01477447-20130122-51 [PubMed: 23379570]
- Roth B, Boateng A, Berken A, Carlyle D, Vadivelu N (2018). Post-operative weaning of opioids after ambulatory surgery: The importance of physician stewardship. *Current Pain & Headache Reports*, 22(6), 4. doi:10.1007/s11916-018-0694-4 [PubMed: 29350303]
- Schwenk ES, Pozek JJ, Viscusi ER (2018). Managing prolonged pain after surgery: Examining the role of opioids. *The Journal of Arthroplasty*. Retrieved from
- Soffin EM, Waldman SA, Stack RJ, Liguori GA (2017). An evidence-based approach to the prescription opioid epidemic in orthopedic surgery. *Anesthesia & Analgesia*, 125(5), 1704–1713. doi:10.1213/ANE.0000000000002433 [PubMed: 29049115]
- Song SJ (2017). Pain management and anesthesia in total knee arthroplasty. *Knee Surgery & Related Research*, 29(2), 77–79. doi:10.5792/ksrr.17.077 [PubMed: 28545170]
- Trasolini NA, McKnight BM, Dorr LD (2018). The opioid crisis and the orthopedic surgeon. *The Journal of Arthroplasty*, 33, 1–4. doi:10.1016/j.arth.2018.07.002 [PubMed: 28844631]
- Trofa D, Rajae SS, Smith EL (2014). Nationwide trends in total shoulder arthroplasty and hemiarthroplasty for osteoarthritis. *The American Journal of Orthopedics*, 43(4), 166–172. [PubMed: 24730001]
- Ullah H, Samad K, Khan FA (2014). Continuous interscalene brachial plexus block versus parenteral analgesia for postoperative pain relief after major shoulder surgery. *Cochrane Database of Systematic Reviews*, (2), Cd007080. doi:10.1002/14651858.CD007080.pub2 [PubMed: 24492959]
- Ulrich BT, Lavandero R, Woods D, Early S (2014). Critical care nurse work environments 2013: A status report. *Critical Care Nurse*, 34(4), 64–79. doi:10.4037/ccn2014731
- Vadivelu N, Schermer E, Kodumudi V, Belani K, Urman RD, Kaye AD (2016). Role of ketamine for analgesia in adults and children. *Journal of Anaesthesiology Clinical Pharmacology*, 32(3), 298–306.

- Wick E, Grant M, Wu C (2017). Postoperative multimodal analgesia pain management with nonopioid analgesics and techniques. *JAMA Surgery*, 152(7), 691–697. doi:10.1001/jamasurg.2017.0898 [PubMed: 28564673]
- Zhang J, Shi K, Jia H (2018). Ketamine and bupivacaine attenuate post-operative pain following total knee arthroplasty: A randomized clinical trial. *Experimental and Therapeutic Medicine*, 15(6), 5537–5543. doi:10.3892/etm.2018.6104 [PubMed: 29904433]

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 1.**Common Opioid Components of Multimodal Analgesic Approaches to Surgical Pain Control**

Modality	Medications	Mechanism of Action
NSAIDs	Aspirin, Ibuprofen, Ketorolac, Celecoxib, Naproxen	Inhibited synthesis and release of prostaglandins Inhibited COX-2
NMDA antagonists	Ketamine	Antagonize NMDA receptor; dissociative agent; potent analgesic properties
Gabapentinoids	Neurontin, Lyrica	Alters neurotransmission in the dorsal horn of the spinal cord
Continuous peripheral nerve block	Bupivacaine, Ropivacaine, Exparel, Lidocaine	Direct neural blockade at the site of peripheral nerve sheath

Note. NMDA = N -methyl-D-aspartate; NSAIDs = nonsteroidal anti-inflammatory drugs