Published in final edited form as:

Abdom Radiol (NY). 2018 July ; 43(7): 1590–1611. doi:10.1007/s00261-017-1383-1.

Magnetic resonance elastography: beyond liver fibrosis—a case-based pictorial review

Sudhakar K. Venkatesh¹, Michael L. Wells¹, Frank H. Miller², Kartik S. Jhaveri³, Alvin C. Silva⁴, Bachir Taouli⁵, Richard L. Ehman¹

¹Department of Radiology, Mayo Clinic College of Medicine, Mayo Clinic, 200, First Street SW, Rochester, MN 55905, USA

²Department of Radiology, Northwestern University, Chicago, IL, USA

³Joint Department of Medical Imaging, University Health Network, University of Toronto, Toronto, ON, Canada

⁴Department of Radiology, Mayo Clinic, Scottsdale, AZ, USA

⁵NYU Medical College, Icahn School of Medicine at Mount Sinai, New York, NY, USA

Abstract

Magnetic resonance elastography (MRE) has been introduced for clinical evaluation of liver fibrosis for nearly a decade. MRE has proven to be a robust and accurate technique for diagnosis and staging of liver fibrosis. As clinical experience with MRE grows, the possible role in evaluation of other diffuse and focal disorders of liver is emerging. Stiffness maps provide an opportunity to evaluate mechanical properties within a large volume of liver tissue. This enables appreciation of spatial heterogeneity of stiffness. Stiffness maps may reveal characteristic and differentiating features of chronic liver diseases and focal liver lesions and therefore provide useful information for clinical management. The objective of this pictorial review is to recapture the essentials of MRE technique and illustrate with examples, the utility of stiffness maps in other chronic liver disorders and focal liver lesions.

Keywords

Elastograms; Diffuse liver diseases; Focal liver lesions; Congestive hepatopathy; Infiltrative diseases; Primary sclerosing cholangitis; Amyloidosis; Sarcoidosis; Acute hepatitis

Magnetic resonance elastography (MRE) has been used for clinical evaluation of liver fibrosis for nearly a decade since its first clinical application in 2007 [1]. MRE has proven to be an accurate, reproducible, and reliable non-invasive technique for detection and staging of

Correspondence to: Sudhakar K. Venkatesh; Venkatesh.Sudhakar@mayo.edu.

Conflict of interest The authors confirm that this article content has no conflict of interest.

Ethical approval All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards.

Informed consent A waiver of informed consent was obtained from our institutional review board for the retrospective review of images.

liver fibrosis caused by many different chronic liver diseases [2–11]. The worldwide experience with this new technique is growing and the technique has been applied for both initial evaluation and treatment response assessment for a spectrum of chronic liver diseases [12–14]. There is also growing interest in the use of MRE for characterizing focal liver lesions and assessing their response to therapy [15–18].

MRE, unlike other elastography techniques provides a stiffness map (elastogram) that displays stiffnesses of tissues in large areas across the abdomen. This provides an opportunity for the assessment of stiffness distribution not only within the liver but also other abdominal organs. The volume of liver parenchyma assessed with a single slice of MRE is about 250 cm³ which is much higher with shear wave elastography (20 cm³), point shear wave elastography (0.5–1.0 cm³), transient elastography (4 cm³), and nearly 500 times that of a liver biopsy (10–50 mm³ or < 0.05 cm³) [19–21]. Assessing a large volume of liver provides an opportunity to appreciate the stiffness distribution within the liver parenchyma. The pattern of stiffness across the liver may provide clues for diagnosis of a specific diffuse disease or point towards a focal pathology. Knowledge of liver pathology that can cause diffuse or focal increased stiffness is important for interpretation of the clinical MRE. In this pictorial review, we demonstrate with examples, the different causes of increased liver stiffness, variability in the stiffness distribution, characteristic patterns of stiffness in chronic liver disease, and also causes for focal increased or decreased stiffness.

Magnetic resonance elastography (MRE) and generation of stiffness maps

The basic principle of MRE is to image the propagation of shear waves within the tissue. Shear waves are created by mechanically applied low frequency continuous vibrations. The propagating shear waves are detected and mathematically processed into quantitative images depicting tissue stiffness, known as stiffness maps or elastograms. MRE has three important steps (Fig. 1). The first step is to generate shear waves in the liver. This is achieved by conducting low frequency (60 Hz) vibrations produced by an active pneumatic mechanical wave driver placed outside the magnet room. The active driver is connected by a long plastic tube to a passive driver that is placed in contact with lower chest wall/anterior abdominal wall overlying the right lobe of the liver. The passive driver has a soft vibrating membrane in contact with anterior abdominal wall that produces shear waves, which propagate through the entire abdomen including the liver. The propagating shear waves cause tissue displacement typically in the order of microns. The second step is to image the pattern of tissue displacements with a motion-encoded phase-contrast MRE sequence. The pulse sequence uses trigger pulses to synchronize the motion-encoding gradients with the applied vibrations. By imaging the propagating waves at regular intervals in each wave cycle, snapshots of the propagating waves are obtained. The MRE sequence produces two kinds of images—the magnitude image showing the cross-sectional anatomy and the phase image containing motion information. The third step is to process the wave images to generate a quantitative map of tissue stiffness, known as stiffness map or elastogram. The processing takes place automatically after the scan is completed. For detailed description of MRE technique please see previous reviews on the technique [1, 22, 23].

The most commonly used clinical MRE sequence is two-dimensional gradient echo sequences (2D-GRE MRE). Other sequences are modified spin-echo (SE) and spin-echo echo-planar imaging (EPI) MRE. 2D-GRE MRE is not limited by obesity but loss of signal caused by the short T2* transverse relaxation time of iron overloaded liver may cause low signal-to-noise ratio (SNR) of the liver parenchyma. The low SNR results in poor visualization of the propagating shear waves. In such cases SE and SE-EPI MRE may be useful for obtaining better tissue signal [24].

Stiffness map (elastogram)

In 2D MRE, an elastogram is a two-dimensional image representing the stiffnesses of the tissues in a cross-sectional slice of the anatomy, typically with thickness of 5–10 mm. This quantitative map displays the mechanical property known as the *magnitude of complex shear modulus* which is measured in units of kilopascals (kPa). Detailed explanation of different mechanical properties of tissues used in evaluation of liver fibrosis can be found elsewhere [25, 26]. The stiffness maps can be either gray scale or color-coded (rainbow) maps with a scale (Fig. 1) provided for interpretation. Red and purple colors represent the higher and lower ends of the stiffness scale, respectively. Different vendors display slightly different hues of color in the color stiffness maps to maintain distinctions; however, all of them provide a color scale map of 0–8 kPa that is standardized across all vendors. Slight variation in the smoothness of the maps may be seen as the matrix size and/or resolution has changed and improved inversion algorithms have become available over the years.

Liver stiffness is measured by drawing regions of interest (ROI) either on gray scale map or color-coded map depending on the platform used (Fig. 2). A confidence map (Fig. 1) is a complex statistical signal-to-noise measure that is derived from a regional least squares fit of the phase difference wave data and it excludes noisy or discontinuous data. A checkerboard overlay is placed on the elastogram to depict regions of lowest confidence or less reliable measurement. ROIs should be drawn largely within the region of highest confidence. The ROI is drawn following the liver contour, avoiding the liver edge to exclude partial volume effects. Larger (geographical) ROIs are preferred (less sampling error). Left lobe measurements may be affected by cardiac pulsations and therefore are preferably not included in the ROIs whenever possible. The ROIs should exclude large vessels (> 3 mm), hot spots (a focal region of artefactual high stiffness) typically found immediately below the passive driver, regions of wave interference, and liver fissures and gallbladder fossa (Fig. 3). Mean stiffness from each ROI is obtained. For typical clinical use, when reporting estimates of liver fibrosis, the average stiffness ($\pm$ SD) from ROIs from four slices through the largest cross section of liver is calculated. Due to spatial heterogeneity of liver stiffness, a qualitative description of the distribution of fibrosis is often added to the report.

Liver stiffness

The normal (healthy, non-fibrotic) liver parenchyma stiffness should be less than 2.5 kPa [27]. Normal liver stiffness is dependent on: (a) tissue components including hepatocytes, bile ducts, interstitial space, intercellular matrix, and capsule; (b) tissue organization which is a regular pattern of cords and lobules in liver; (c) vascular component comprised

sinusoids, blood flow, and blood pressure. Interstitial pressure is typically low (0–2 mmHg) in normal liver parenchyma [28]. Liver is also wrapped by a thin capsule (Glisson's) that also contributes to the stiffness measured. Any external factor that compresses or puts strain on the liver can increase liver stiffness.

Pathological conditions that result in alterations in the tissue components, organizational structure of parenchyma, vascular components, and interstitial pressure would lead to altered liver stiffness.

Diffuse increased liver stiffness

Liver fibrosis

Chronic liver disease resulting in fibrosis and cirrhosis is the most common cause of significantly elevated liver stiffness and is of high clinical significance. Chronic liver diseases are characterized by hepatocyte loss, an increase in extracellular collagen content (liver fibrosis/cirrhosis), and disruption of lobules and nodule formation. The architectural disruption is accompanied by reduced portal venous flow and a compensatory increased arterial flow. Intrahepatic vascular shunts are also commonly present in advanced fibrosis and cirrhosis. The vascular changes lead to portal hypertension. Extracellular collagen deposition in liver fibrosis is the most significant contributing factor for diffuse liver stiffness. The collagen deposition typically occurs in small increments in early stages of fibrosis and becomes exponential in advanced stages of fibrosis [29–31]. Liver stiffness shows a similar trend of mild increase in stiffness in early stages and exponential increase in stiffness in advanced stages [3, 7, 8, 32].

Spatial heterogeneity of stiffness distribution in liver fibrosis

Liver parenchyma stiffness in chronic liver disease is not uniform with some regions showing more stiffness than other regions. This phenomenon is well known and likely reflects heterogeneous collagen deposition throughout the liver [33]. In fact, spatial heterogeneity is one of the major limitations of liver biopsy and ultrasound-based elastography techniques which sample a small volume of liver parenchyma, and therefore may not accurately reflect the overall burden of fibrosis. Heterogeneity of fibrosis deposition becomes prominent in advanced fibrosis and cirrhosis [34–36]. Stiffness maps correspondingly show more heterogeneity as the fibrosis stage advances. Stiffness maps can also be used to perform imaging guided biopsy in order to target the most diseased regions of the liver and to reduce sampling error.

Stiffness distribution may vary between patients even though biopsy determined fibrosis stage is the same (Fig. 4). The distribution of stiffness may also vary between etiologies (Fig. 5). There is very little data on fibrosis distribution versus etiology. Different cutoffs are used for staging liver fibrosis from different etiologies with ultrasound-based elastography techniques [37]. However, with MRE the data are still emerging and it is possible that different cutoffs may be established in the future, specifically for NAFLD [38].

As experience with clinical use of MRE grows, there is definite potential to find features that are characteristic of a chronic liver disease. In our experience we have observed that in

primary sclerosing cholangitis (Figs. 6, 7), increased stiffness may be found in the periphery of the liver particularly in early stages. As the disease advances, a segmental or lobar distribution of increased stiffness may progress in the advanced stage to involve the entire liver. When the entire liver is involved, the elastogram appearance of PSC may be indistinguishable from other diffuse liver diseases. Multifocal biliary strictures and peripheral increased liver stiffness may be diagnostic of primary sclerosing cholangitis.

MRE may demonstrate increased liver stiffness due to fibrosis even when there are no morphological signs of fibrosis or cirrhosis (Fig. 8). It is useful to perform MRE in patients who are clinically suspected to have liver fibrosis but conventional imaging does not show any morphological characteristics (Fig. 9).

Inflammation

Acute inflammatory processes such as acute infective hepatitis, acute autoimmune hepatitis and acute flare of chronic liver disease can cause diffusely increased liver stiffness. This increased stiffness is probably due to recruitment of inflammatory cells and associated edema and hyperemia in the inflamed liver which causes parenchymal expansion and stretching of the liver capsule. The increased liver stiffness may be severe sometimes simulating cirrhosis (Fig. 10). Often in clinical follow-ups, there may be an acute increase in liver stiffness due to flare. This is commonly seen in chronic hepatitis C and autoimmune hepatitis; correlation with laboratory findings is useful for confirmation. Staging of liver fibrosis should be avoided during acute inflammation. Follow-up MRE for liver fibrosis staging can be performed when acute inflammation has subsided (typically 3–6 months) or when laboratory tests particularly serum alanine aminotransferase (ALT), a marker of liver inflammation returns to normal. Other causes of acute inflammation include acute alcoholic hepatitis (Fig. 11) and follow-up MRE can be useful to show reduced stiffness to near normal levels in patients who show abstinence.

Chronic inflammation or subacute inflammation that frequently accompanies fibrosis in chronic liver diseases such as viral hepatitis, steatohepatitis, and autoimmune hepatitis can cause a mild increase in stiffness (typically raising liver stiffness to no more than 3 kPa) [2]. The contribution of chronic inflammation to overall liver stiffness is difficult to quantify and separate from fibrosis with current MRE technique and inversion. Advances in MRE technique may help in the future to separate the contributions of inflammation and fibrosis to overall liver stiffness [39]. Chronic inflammation and liver fibrosis is probably best viewed as a continuum in the progression of disease as inflammation results in liver fibrosis if untreated.

Biliary obstruction

Biliary obstruction especially occurring in the central hilar region can lead to diffuse increased liver stiffness (Fig. 12). This is likely due to increased interstitial pressure secondary to biliary obstruction and cholestasis. More peripheral biliary obstruction may result in lobar, segmental, or sub-segmental pattern of increased stiffness and can be observed in primary sclerosing cholangitis.

Passive congestion

Passive venous congestion of the liver due to obstruction in the hepatic venous outflow, congestive heart failure, congenital heart diseases with pulmonary stenosis, and tricuspid regurgitation can lead to congestive hepatopathy. This causes sinusoidal congestion that can lead to increased liver stiffness. Clinical history of congenital heart disease, imaging findings of dilated inferior vena cava and hepatic veins (Fig. 13) should provide the clue to the diagnosis. Congestive hepatopathy also leads to peripheral increased liver stiffness particularly in the right lobe (Fig. 25) which bears the most brunt of passive venous congestion [40]. Chronic congestive hepatopathy can lead to liver fibrosis and cirrhosis [41] and hepatic fibrosis can be difficult to differentiate from congestive hepatopathy alone with MRE (Fig. 13). Advances in MRE in the future may be able to quantify stiffness resulting from congestion alone and distinguish from other causes such as fibrosis and inflammation.

Diffuse infiltrative disorders

Diffuse infiltrative processes can also lead to increased liver stiffness. The infiltrative pathologies by themselves cause increased stiffness or can also lead to liver fibrosis which in turn will increase liver stiffness. Sarcoidosis is a systemic inflammatory disorder characterized by chronic inflammation with non-caseating granulomas. Patients with involvement of liver with sarcoidosis may be asymptomatic and detected incidentally (Fig. 14). Sarcoidosis can lead to scarring of the liver and ultimately to fibrosis simulating a cirrhotic liver (Fig. 15). History of sarcoidosis and no other chronic liver disease may help in arriving at correct diagnosis. Lymphoma can also cause diffuse increase in liver stiffness (Fig. 16) without any signs of fibrosis or cirrhosis. Presence of lymphadenopathy elsewhere is an important clue to differentiate from liver fibrosis alone.

Amyloidosis is a systemic disorder resulting in the deposition of gamma chains in multiple organs including the liver. Amyloid deposition in the liver causes extremely elevated liver stiffness (Fig. 17), probably the stiffest diffuse livers one would come across in clinical practice.

Liver metastases may diffusely infiltrate the entire liver. MRE may show diffusely elevated liver stiffness (Fig. 18) similar to cirrhosis and should be differentiated. MRI features of primary or other metastatic lesions and clinical history would be useful in distinction.

In summary, elevated liver stiffness may result from several different pathologies. Correlation with clinical features, liver function tests, and imaging findings are important when interpreting elevated liver stiffness.

Focal or segmental increased stiffness

Stiffness maps may reveal focal regions of increased stiffness within a normal liver parenchyma or within diffusely stiff liver secondary to fibrosis. These are often localized to few segments or may correspond to focal liver lesions. Examples include confluent fibrosis (Fig. 19), focal biliary stricture (Fig. 20), healing infective/inflammatory process by fibrosis (Fig. 21), and focal liver lesions (Figs. 22, 23). Malignant focal liver lesions may be recognized as region of increased stiffness in a normal or mildly stiff liver parenchyma, but

may not be distinguishable from surrounding liver if the parenchyma is fibrotic and therefore also abnormally stiff (Fig. 22).

Focal or segmental decreased stiffness in a diffusely stiff liver

Regions of decreased stiffness within abnormal liver parenchyma may be seen and they usually correspond to a region of normal or regenerating liver parenchyma. Regenerative nodules have similar or slightly higher stiffness than normal liver parenchyma and may appear as focal regions of decreased stiffness in a cirrhotic liver (Fig. 24), often surrounded by a rim of increased stiffness representing fibrosis (Fig. 25).

A focal lesion that has same stiffness as surrounding liver parenchyma may not be detectable on stiffness maps. Some benign lesions such as hemangiomas and hepatic adenomas may have similar stiffness as surrounding liver parenchyma and not detectable (stealth lesions) (Fig. 26) on stiffness maps. Currently, the stiffness maps are of low resolution (96×96 to 128×256) and therefore small focal lesions are not easily resolved. Also with 2D MRE, the entire liver is typically not assessed and therefore stiffness maps should not be used alone for detection of focal liver lesions. However, if new 3D MRE techniques are being used, nearly entire liver can be covered [38, 42, 43].

Liver tumors that undergo necrosis following locoregional treatment may demonstrate decreased stiffness when they undergo internal necrosis (Fig. 27). This finding may be useful for demonstrating response to treatment [18].

Summary/conclusion

MRE is well-established for identifying hepatic fibrosis and cirrhosis in chronic liver disease. Inflammation, congestion, biliary obstruction, and diffuse infiltrative diseases can also lead to increased liver stiffness. Peripheral distribution of elevated stiffness is often present in early PSC and passive hepatic congestion which may be useful for early detection. Regions of abnormal stiffness may have corresponding findings on conventional MRI sequences and the information may be complementary for detecting pathology.

Acknowledgments

Funding was provided by National Institutes of Health (Grant No. EB001981).

Funding No funding was received for this study.

References

1. Venkatesh SK, Yin M, Ehman RL (2013) Magnetic resonance elastography of liver: technique, analysis, and clinical applications. *J Magn Reson Imaging* 37:544–555 [PubMed: 23423795]
2. Singh S, Venkatesh SK, Wang Z, et al. (2015) Diagnostic performance of magnetic resonance elastography in staging liver fibrosis: a systematic review and meta-analysis of individual participant data. *Clin Gastroenterol Hepatol* 13(440–451):e6
3. Singh S, Venkatesh SK, Loomba R, et al. (2016) Magnetic resonance elastography for staging liver fibrosis in non-alcoholic fatty liver disease: a diagnostic accuracy systematic review and individual participant data pooled analysis. *Eur Radiol* 26:1431–1440 [PubMed: 26314479]

4. Singh S, Venkatesh SK, Keaveny A, et al. (2016) Diagnostic accuracy of magnetic resonance elastography in liver transplant recipients: a pooled analysis. *Ann Hepatol* 15:363–376
5. Jajamovich GH, Dyvome H, Donnerhack C, et al. (2014) Quantitative liver MRI combining phase contrast imaging, elastography, and DWI: assessment of reproducibility and postprandial effect at 3.0 T. *PLoS ONE* 9:e97355 [PubMed: 24840288]
6. Yasar TK, Wagner M, Bane O, et al. (2016) Interplatform reproducibility of liver and spleen stiffness measured with MR elastography. *J Magn Reson Imaging* 43:1064–1072 [PubMed: 26469708]
7. Venkatesh SK, Wang G, Lim SG, et al. (2014) Magnetic resonance elastography for the detection and staging of liver fibrosis in chronic hepatitis B. *Eur Radiol* 24:70–78 [PubMed: 23928932]
8. Ichikawa S, Motosugi U, Ichikawa T, et al. (2012) Magnetic resonance elastography for staging liver fibrosis in chronic hepatitis C. *Magn Reson Med Sci* 11:291–297 [PubMed: 23269016]
9. Cui J, Heba E, Hernandez C, et al. (2016) Magnetic resonance elastography is superior to acoustic radiation force impulse for the Diagnosis of fibrosis in patients with biopsy-proven nonalcoholic fatty liver disease: a prospective study. *Hepatology* 63:453–461 [PubMed: 26560734]
10. Bensamoun SF, Leclerc GE, Charleux F, et al. (2012) Magnetic resonance elastography (MRE): a non invasive technique to identify the cut-off values for alcoholic liver fibrosis. *J Hepatol* 56:S527
11. Wang J, Malik N, Yin M, et al. (2017) Magnetic resonance elastography is accurate in detecting advanced fibrosis in autoimmune hepatitis. *World J Gastroenterol* 23:859–868 [PubMed: 28223730]
12. Venkatesh SK, Yin M, Ehman RL (2013) Magnetic resonance elastography of liver: clinical applications. *J Comput Assist Tomogr* 37:887–896 [PubMed: 24270110]
13. Cui J, Ang B, Haufe W, et al. (2015) Comparative diagnostic accuracy of magnetic resonance elastography vs. eight clinical prediction rules for non-invasive diagnosis of advanced fibrosis in biopsy-proven non-alcoholic fatty liver disease: a prospective study. *Aliment Pharmacol Ther* 41:1271–1280 [PubMed: 25873207]
14. Loomba R, Sirlin CB, Ang B, et al. (2015) Ezetimibe for the treatment of nonalcoholic steatohepatitis: assessment by novel magnetic resonance imaging and magnetic resonance elastography in a randomized trial (MOZART trial). *Hepatology* 61:1239–1250 [PubMed: 25482832]
15. Venkatesh SK, Yin M, Glockner JF, et al. (2008) MR elastography of liver tumors: preliminary results. *AJR* 190:1534–1540 [PubMed: 18492904]
16. Garteiser P, Doblas S, Daire JL, et al. (2012) MR elastography of liver tumours: value of viscoelastic properties for tumour characterisation. *Eur Radiol* 22:2169–2177 [PubMed: 22572989]
17. Hennedige TP, Hallinan JT, Leung FP, et al. (2016) Comparison of magnetic resonance elastography and diffusion-weighted imaging for differentiating benign and malignant liver lesions. *Eur Radiol* 26:398–406 [PubMed: 26032879]
18. Gordie S, Ayache JB, Kennedy P, et al. (2017) Value of tumor stiffness measured with MR elastography for assessment of response of hepatocellular carcinoma to locoregional therapy. *Abdom Radiol* 42:1685–1694
19. Barr RG, Ferraioli G, Palmeri ML, et al. (2015) Elastography Assessment of liver fibrosis: society of radiologists in ultrasound consensus conference statement. *Radiology* 276:845–861 [PubMed: 26079489]
20. Yin M, Venkatesh SK (2017) Ultrasound or MR elastography of liver: which one shall I use? *Abdom Radiol*. 10.1007/s00261-017-1340-z
21. Bravo AA, Sheth SG, Chopra S (2001) Liver biopsy. *N Engl J Med* 344:495–500 [PubMed: 11172192]
22. Venkatesh SK, Ehman RL (2014) Magnetic resonance elastography of abdomen. *Abdominal Imaging* 40:745–759
23. Glaser KJ, Manduca A, Ehman RL (2012) Review of MR elastography applications and recent developments. *J Magn Reson Imaging* 36:757–774 [PubMed: 22987755]

24. Mariappan YK, Dzyubak B, Glaser KJ, et al. (2017) Application of modified spin-echo-based sequences for hepatic MR elastography: evaluation, comparison with the conventional gradient-echo sequence, and preliminary clinical experience. *Radiology* 282:390–398 [PubMed: 27509543]
25. Mariappan YK, Glaser KJ, Ehman RL (2010) Magnetic resonance elastography: a review. *Clin Anat* 23:497–511 [PubMed: 20544947]
26. Tang A, Cloutier G, Szeverenyi NM, et al. (2015) Ultrasound elastography and MR elastography for assessing liver fibrosis: part 1, principles and techniques. *AJR* 205:22–32 [PubMed: 25905647]
27. Venkatesh SK, Ehman RL (2014) Magnetic resonance elastography of liver. *Magn Reson Imaging Clin N Am* 22:433–446 [PubMed: 25086938]
28. Kurbel S, Sven K, Flam J, et al. (2007) Interstitial hydrostatic pressure: a manual for students. *Adv Physiol Educ* 31:116–117 [PubMed: 17327592]
29. Standish RA, Cholongitas E, Dhillon A, et al. (2006) An appraisal of the histopathological assessment of liver fibrosis. *Gut* 55:569–578 [PubMed: 16531536]
30. Lazzarini AL, Levine RA, Ploutz-Snyder RJ, et al. (2005) Advances in digital quantification technique enhance discrimination between mild and advanced liver fibrosis in chronic hepatitis C. *Liver Int* 25:1142–1149 [PubMed: 16343064]
31. Biagini G, Ballardini G (1989) Liver fibrosis and extracellular matrix. *J Hepatol* 8:115–124 [PubMed: 2646366]
32. Singh S, Venkatesh S, Wang Z, et al. (2014) Diagnostic performance of magnetic resonance elastography for the staging of liver fibrosis: a systematic review and collaborative individual participant data meta-analysis. *Am J Gastroenterol* 109:S144
33. Soloway RD, Baggenstoss AH, Schoenfield LJ, et al. (1971) Observer error and sampling variability tested in evaluation of hepatitis and cirrhosis by liver biopsy. *Am J Dig Dis* 16:1082–1086 [PubMed: 5135770]
34. Maharaj B, Maharaj RJ, Leary WP, et al. (1986) Sampling variability and its influence on the diagnostic yield of percutaneous needle biopsy of the liver. *Lancet* 1:523–525 [PubMed: 2869260]
35. Abdi W, Millan J, Mezey E (1979) Sampling variability on percutaneous liver biopsy. *Arch Intern Med* 139:667–669 [PubMed: 443970]
36. Regev A, Berho M, Jeffers LJ, et al. (2002) Sampling error and intraobserver variation in liver biopsy in patients with chronic HCV infection. *Am J Gastroenterol* 97:2614–2618 [PubMed: 12385448]
37. Sigrist RMS, Liao J, Kaffas AE, et al. (2017) Ultrasound elastography: review of techniques and clinical applications. *Theranostics* 7:1303–1329 [PubMed: 28435467]
38. Loomba R, Cui J, Wolfson T, et al. (2016) Novel 3D magnetic resonance elastography for the noninvasive diagnosis of advanced fibrosis in NAFLD: a prospective study. *Am J Gastroenterol* 111:986–994 [PubMed: 27002798]
39. Yin M, Glaser KJ, Manduca A, et al. (2017) Distinguishing between hepatic inflammation and fibrosis with MR elastography. *Radiology* 284:694–705 [PubMed: 28128707]
40. Babu AS, Wells ML, Teytelboym OM, et al. (2016) Elastography in chronic liver disease: modalities, techniques, limitations, and future directions. *Radiographics* 36:1987–2006 [PubMed: 27689833]
41. Wells ML, Fenstad ER, Poterucha JT, et al. (2016) Imaging findings of congestive hepatopathy. *Radiographics* 36:1024–1037 [PubMed: 27284758]
42. Morisaka H, Motosugi U, Glaser KJ, et al. (2017) Comparison of diagnostic accuracies of two- and three-dimensional MR elastography of the liver. *J Magn Reson Imaging* 45:1163–1170 [PubMed: 27662640]
43. Wang J, Glaser KJ, Zhang T, et al. (2017) Assessment of advanced hepatic MR elastography methods for susceptibility artifact suppression in clinical patients. *J Magn Reson Imaging*. 10.1002/jmri.25818

Fig. 1.

Steps in MRE of liver and generation of stiffness map. Top image showing the typical location of the passive driver on a T2W image in a patient with chronic hepatitis C cirrhosis. Magnitude and Phase maps (second row) are displayed on the scanner within seconds of acquiring an MRE slice. The color wave map, stiffness maps in gray scale, color map, and confidence maps (bottom row) are processed on the scanner and displayed within few minutes after completion of MRE sequence. The arrow shows the color scale from 0 to 8 kPa provided on the color map. Arrowheads point to the hashed out regions that are outside confidence limits. In this particular example, nearly the entire cross section of the liver is within confidence map.

Fig. 2.

Examples illustrating correct drawing of region of interest (ROI) for measuring liver stiffness. Top row showing magnitude (**A**), stiffness map (**B**), and confidence map (**C**) in a 57-year-old female with BMI of 33.6 kg/m² and non-alcoholic steatohepatitis. The ROI includes entire liver and mean stiffness is 7.2 kPa. Middle row showing magnitude (**D**), stiffness map (**E**), and confidence map (**F**) in a 56-year-old female with BMI of 42.8 kg/m² with alcoholic cirrhosis. Arrow points to the artifact due to cholecystectomy clips which is seen as an area of low stiffness and the confidence map hashes out that region. The geographical ROI covers most of the right lobe of the liver and mean stiffness was 6.8 kPa. Bottom row showing wave image (**G**), stiffness map (**H**), and confidence map (**I**) in a 58-year-old female with BMI of 38.4 kg/m² and fatty liver. Note that the confidence map hashes out most of the liver except for small peripheral region in right lobe which shows wave propagation (arrow, **G**). The ROI is drawn within the confidence map and is much smaller than two examples above. Mean stiffness was 1.8 kPa. Smaller regions of confidence may be seen in normal livers as healthy non-fibrotic liver parenchyma is soft and absorbs shear wave

energy causing rapid attenuation of waves deep in the liver. This should not be interpreted as failed MRE study.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Fig. 3.

Examples illustrating areas to avoid when drawing regions of interest. Top row showing an area of hot spot (dotted circle) in stiffness map (**A**) that corresponds to area of large wave amplitude on wave image (**B**). No focal lesions demonstrated on corresponding T2W image (**C**). Hot spots are usually seen under the passive driver. Second row shows area of low stiffness (**D**, arrow) that corresponds to a region separating two wave fronts (**E**) and fissure for ligamentum teres seen on magnitude image (**F**). Third row showing region of low stiffness in the region of gallbladder fossa (arrow, **G**), corresponding to an area of low amplitude waves (arrow, **H**) and hypointensity due to gallstones in corresponding anatomical image (**I**). Fourth row showing a focal region of increased stiffness (white circle, **J**) due to wave interference (white circle, **K**) with no corresponding anatomical abnormality on magnitude image (**L**). Fifth row demonstrating a focal area of decreased stiffness (white arrow, **M**) secondary to artifact, and wave interference (white arrow, **N**) with no corresponding focal abnormality on magnitude image (**O**). Changing the position of passive driver may help in removing the wave interference artifacts. Arrow heads (**E**, **H**) point to vascular pulsation artifacts.

Fig. 4. Heterogeneity of stiffness distribution and variability of mean stiffness in chronic liver disease of same etiology and same biopsy determined stage of fibrosis. Stiffness maps in four patients (A–D) with chronic hepatitis C and cirrhosis. Note heterogeneity of parenchyma stiffness and mean stiffness ranges from 5.8 to 8.2 kPa.

Fig. 5. Heterogeneity of stiffness distribution in different chronic liver diseases at same biopsy determined stage of fibrosis. Stiffness maps from biopsy-proven cirrhosis secondary to chronic hepatitis C (A), chronic hepatitis B (B), non-alcoholic steatohepatitis (C), and primary biliary cirrhosis (D).

Fig. 6. Primary sclerosing cholangitis in a patient with ulcerative colitis. Axial T2W image (**A**) shows dilated peripheral ducts in right anterior liver lobe with no corresponding perfusion change on post-contrast image (**B**). Stiffness maps (**C**, **D**) of two consecutive MRE sections shows peripheral increased stiffness, particularly in the right anterior lobe (arrows) with normal stiffness in central liver (arrowheads).

Fig. 7. Primary sclerosing cholangitis showing correlate of peripherally increased liver stiffness (**A**) to hyperintensity on T2W (**B**) and DWI (**C**), hyper perfusion in arterial phase (**D**) and delayed enhancement in the delayed phase (**E**) and decreased uptake of Eovist on hepatobiliary phase (**F**).

Fig. 8.

Non-alcoholic steatohepatitis with biopsy-proven stage 2 fibrosis. Axial in-phase (**A**) and opposed-phase (**B**) images show drop off signal consistent with fatty change in the liver. Note no morphological signs of advanced fibrosis or cirrhosis on T2W (**C**) image. Stiffness map (**D**) shows elevated liver stiffness with a mean of 3.8 kPa consistent with stage 2 fibrosis.

Fig. 9.

Chronic hepatitis C patient on follow-up. Biopsy performed 7 years earlier showed stage 1 fibrosis. Conventional T2W (**A**) and opposed-phase image (**B**) do not show any signs of chronic liver disease. Spleen size is also normal. Stiffness map (**C**) shows liver stiffness of 6.2 kPa consistent with advanced fibrosis/cirrhosis.

Fig. 10.

Acute inflammation. A 66-year-old male with no prior history of liver disease presented with jaundice and fatigue with elevated serum ALT of 1773 (Normal, < 55 IU/L), AST 1244 (Normal, < 48 IU/L). MRI and MRE were performed. Axial T2W image (**A**) and intravenous gadolinium enhanced portal venous phase (**B**) do not reveal any significant abnormality in liver. MRE (**C**) shows significantly elevated mean liver stiffness of 8.6 kPa. Biopsy showed severe inflammatory infiltrate, marked interface activity, severe lobular inflammatory with several necroinflammatory foci with focal bridging necrosis. Periportal fibrosis (stage 2) was also present. Findings were consistent with autoimmune hepatitis and patient responded to oral steroids and azathioprine therapy and currently on follow-up.

Fig. 11.

A 56-year-old female with history of alcoholism. At initial presentation showing elevated liver stiffness measuring 14 kPa (**A**). After abstinence and follow-up at 7 months shows liver stiffness returning to nearly normal (2.6 kPa) (**B**).

Fig. 12.

A 75-year-old female with biopsy-proven hilar cholangiocarcinoma. MRCP (A) showing the hilar tumor (arrow). Axial T2W image (B) and post-contrast-enhanced T1-weighted image (C) showing subtle differences in signal intensity of right and left lobes and intrahepatic biliary dilatation right more than left. MRE (D) shows diffuse increase in liver stiffness with right lobe stiffer than left lobe which may be related to cholestasis.

Fig. 13.

Congestive hepatopathy. Top row—A 36-year-old male with history of pulmonary atresia and post-Fontan surgery. Axial T2W (**A**) image showing dilated hepatic veins (arrow) and MRE (**B**) showing stiff liver predominantly in the peripheral distribution and right lobe more involved. Bottom row—A 32-year-old patient with history of long-standing constrictive pericarditis following congenital heart surgery. Axial T2W (**C**) image showing hepatomegaly with dilated hepatic veins (arrows). MRE (**D**) shows increased stiffness somewhat more in periphery. Liver biopsy confirmed cirrhosis in this patient.

Fig. 14.

59-year-old diabetic male with elevated alkaline phosphatase and presumed fatty liver: There is mild hepatomegaly but otherwise normal MRI of the liver. Note enlarged spleen (**A**) with heterogeneous enhancement (**B**, arrow). There were enlarged portal lymph nodes (**C**, arrow) and thoracic lymphadenopathy and bone lesion (**D**, arrows). MRE (**E**) helped show liver is abnormal which would otherwise appear normal on routine MRI. PET (**F**) showed diffusely increased FDG uptake in liver and spleen. Patient underwent biopsy that was consistent with sarcoidosis.

Fig. 15.

45-year-old male with sarcoidosis diagnosed 3 years earlier. T2W (**A**) and T1W (**B**) images showing nodular liver suggestive of cirrhosis. Note granulomas in the spleen (arrows). MRE (**C**) shows increased liver stiffness with a mean of 4 kPa.

Fig. 16.

Lymphoma. A 19-year-old female with common variable immunodeficiency with history of splenectomy and presenting with enlarged liver. Coronal post-contrast-enhanced T1W image (A) showing hepatomegaly and enlarged mesenteric lymph nodes, axial DWI (B) showing multiple enlarged retroperitoneal lymph nodes as well (B, arrows). MRE (C) showed diffuse increased stiffness of liver with a mean of 5.9 kPa. Histology confirmed lymphoma and no fibrosis.

Fig. 17.

47-year-old female with proven amyloidosis. Coronal T2W image (A) showing hepatosplenomegaly. Stiffness map (B) showing elevated stiffness in both liver and spleen. Stiffness map with 0–20 scale (C) showing the heterogeneous distribution of amyloid deposition in the liver.

Fig. 18.

Neuroendocrine metastases diffusely studding both lobes of the liver on axial T2W (**A**) and post-contrast T1W (**B**) images. MRE (**C**) shows diffusely increased stiffness of the liver with focal areas of increased stiffness corresponding to regions of coalescing metastases.

Fig. 19. Focal confluent fibrosis. Chronic biliary cirrhosis showing a region of hyperintensity on T2W (**A**) and diffusion-weighted images (**B**) that correspond to focally increased stiffness region on the elastograms (**C**, **D**). The focally increased stiffness is better demonstrated on 0–20 kPa scale (arrow, **D**).

Fig. 20.
Focal biliary stricture causing focal region of elevated stiffness in primary sclerosing cholangitis. T2W (A) and MRCP (B) showing focal dilatation of segment 3 ducts (circle). Stiffness map (C) shows focal increased stiffness over same region.

Fig. 21.

Healing liver abscess in a liver transplant. Focal area of increased liver stiffness corresponds to hyperintensity (arrow) on T2W image (**A**) and enhancing region (arrow) on post-gadolinium-enhanced portal venous image (**B**). Note central core of higher stiffness within the abscess in the stiffness map (**C**).

Fig. 22.

Hepatocellular carcinomas in two patients. Top row (**A–C**) showing a large HCC with arterial phase (**A**) hyperenhancement and portal venous washout (**B**) with corresponding heterogeneous but increased stiffness region (arrow) on the stiffness map (**C**). Note only mildly elevated stiffness of the surrounding liver parenchyma. A small HCC in another patient (bottom row, **D–F**) shows arterial phase hyperenhancement (arrow, **A**) with no washout in portal venous phase (arrow, **B**) and is not distinguishable from surrounding cirrhotic liver parenchyma (circle) on the stiffness map (**F**).

Fig. 23.

Malignant focal lesions. Top row T2W (**A**), postcontrast T1W (**B**), and stiffness map (**C**) showing an intrahepatic cholangiocarcinoma with satellite lesions and intrahepatic metastases. Large primary mass (arrow) corresponds to stiff region on stiffness map. Note the satellite lesion and metastases (arrowheads) corresponding to regions of increased stiffness in right lobe. Note also increased left lobe stiffness due to other metastases (metastases are not shown). Bottom row T2W (**D**), hepatobiliary phase image (**E**), and stiffness map (**F**) showing two large metastases (arrows) from neuroendocrine carcinoma of the pancreas (not shown). The focal regions of increased stiffness correspond to metastases within normal liver parenchyma.

Fig. 24.

Primary sclerosing cholangitis on follow-up. Two adjacent large hypointense regions (*) in right lobe on T2W image (A) that show no restricted diffusion on DWI (B) and only mild significant post-contrast hyperenhancement in arterial phase (C) and other phases (not shown). These regions have normal or slightly elevated stiffness on stiffness map (D, arrows) suggestive of regenerative nodules. These lesions showed mild increase in size during 2-year follow-up and no other suspicious features.

Fig. 25.

Chronic autoimmune hepatitis with multiple regenerative nodules. Axial T2W (A), DWI (B), post-contrast-enhanced delayed phase T1W (C) images and stiffness map (D) show multiple regenerative nodules studded throughout the liver. Regenerative nodules show lower stiffness as compared to background fibrotic parenchyma. Arrows point to a single regenerative nodule in posterior right lobe that is surrounded by fibrotic band.

Fig. 26.

Hepatic adenomas as stealth lesions on stiffness maps. Axial T2W (A), DWI (B), and hepatobiliary phase image following Eovist (C) showing multiple lesions in liver in a patient with hepatic adenomatosis. The stiffness map (D) shows normal stiffness with no discrete lesions. Note difference from diffuse metastatic disease which is associated with increases in liver stiffness.

Fig. 27.

A 47-year-old male with a biopsy-proven large HCC involving the left lobe of the liver. The mass is mildly hyperintense on T2W (**A**) at initial diagnostic MRI and shows partial internal necrosis on the T2W (**B**) on MRI performed 3 weeks after chemoembolization. Post-contrast-enhanced arterial phase image (**C**, **E**) and their corresponding MRE slices (**D**, **F**). Residual enhancing nodule (arrow, **D**) appears stiff on the elastogram with remainder necrotic region appearing softer (dotted outline). The completely necrotic inferior aspect appears soft (*, dotted outline) similar to gallbladder (GB).