Published in final edited form as:

Psychopharmacology (Berl). 2019 January ; 236(1): 265–272. doi:10.1007/s00213-018-5031-4.

Increased amygdalar metabotropic glutamate receptor 7 mRNA in a genetic mouse model of impaired fear extinction

Richard M. O'Connor^{1,2,3}, Cian P. McCafferty^{1,4}, Javier A. Bravo⁵, Nicolas Singewald⁶, Andrew Holmes⁷, John F. Cryan¹

¹Department of Anatomy and Neuroscience and APC Microbiome Institute, University College Cork, Cork, Ireland ²Department of Neuroscience, Icahn School of Medicine at Mount Sinai, S10-20 Hess CSM, 1470 Madison Avenue, New York, NY 10029, USA ³Present address: Department of Neuroscience, Icahn School of Medicine, Mount Sinai Hospital, New York, NY, USA ⁴Present address: Department of Neurology, Yale University School of Medicine, New Haven, CT, USA ⁵Grupo de NeuroGastroBioquímica, Laboratorio e Química Biológica & Bioquímica de Sistemas, Instituto de Química, Facultad de Ciencias, Pontificia Universidad Católica de Valparaíso, Valparaíso, Chile ⁶Department of Pharmacology and Toxicology, Institute of Pharmacy and Center for Molecular Biosciences Innsbruck (CMBI), University of Innsbruck, Innsbruck, Austria ⁷Laboratory of Behavioral and Genomic Neuroscience, National Institute on Alcohol Abuse and Alcoholism, Rockville, MD, USA

Abstract

Rationale—Post-traumatic stress disorder (PTSD) is a devastating anxiety-related disorder which develops subsequent to a severe psychologically traumatic event. Only ~ 9% of people who experience such a trauma develop PTSD. It is clear that a number of factors, including genetics, influence whether an individual will develop PTSD subsequent to a trauma. The 129S1/SvImJ (S1) inbred mouse strain displays poor fear extinction and may be useful to model this specific aspect of PTSD. The metabotropic glutamate receptor 7 (mGlu₇ receptor) has previously been shown to be involved in cognitive processes and anxiety-like behaviour placing it in a key position to regulate fear extinction processes. We sought to compare mGlu₇ receptor mRNA levels in the S1 strain with those in the robustly extinguishing C57BL/6J (B6) inbred strain using in situ hybridisation (ISH) in three brain regions associated with fear extinction: the amygdala, hippocampus and prefrontal cortex (PFC).

Results—Compared to the B6 strain, S1 mice had increased mGlu₇ receptor mRNA levels in the lateral amygdala (LA) and basolateral amygdala (BLA) subdivisions. An increase was also seen in the hippocampal CA1 and CA3 subregions of S1 mice. No difference in mGlu₇ receptor levels

Richard M. O'Connor: richard.oconnor@mssm.edu.

This article belongs to a Special Issue on Psychopharmacology of Extinction.

Compliance with ethical standards

All experiments were conducted in accordance with the European Directive 86/609/EEC and the local Animal Care and Use Committees.

Conflict of interest The authors declare that they have no conflict of interest.

were seen in the central nucleus (CeA) of the amygdala, dentate gyrus (DG) of the hippocampus or prefrontal cortex.

Conclusions—These data show altered mGlu₇ receptor expression in key brain regions associated with fear extinction in two different inbred mouse strains which differ markedly in their fear extinction behaviour. Altered mGlu₇ receptor levels may contribute to the deficit fear extinction processes seen in fear extinction in the S1 strain.

Keywords

mGlu₇ receptor expression; Post-traumatic stress disorder; Trauma

Introduction

Post-traumatic stress disorder (PTSD) is a chronic anxiety disorder that develops subsequent to a severe trauma (DSM-V 2013). With only ~ 9% of individuals who experience a trauma developing full blown PTSD (Breslau et al. 1998; Kessler et al. 1995), it is clear that different susceptibility factors predispose individuals to PTSD with genetics likely to be influential (Kendler 2001; Maren and Holmes 2016; Stein et al. 2002). Our knowledge of the molecular underpinnings of PTSD remain incomplete with only a limited number of genetic models available (Depino and Gross 2007; Stiedl et al. 2003; Tovote et al. 2004). The inbred 129S1/SvImJ (S1) mouse strain has been shown to display impaired fear extinction without abnormal fear acquisition under most experimental conditions (Fitzgerald et al. 2014; Hefner et al. 2008; Whittle et al. 2013) and could be a potential model of the genetics underlying PTSD (Camp et al. 2009; Mi et al. 2018).

Recently, research on the metabotropic glutamate receptors (mGlu_s) has come to the fore in anxiety and cognition research (Peterlik et al. 2016) with early work on mGlu₇ receptor showing great promise (Gee et al. 2014; Goddyn et al. 2008; Moloney et al. 2015; O'Connor and Cryan 2010; O'Connor and Cryan 2013; O'Connor et al. 2010; O'Connor et al. 2013a; O'Connor et al. 2013b; O'Mahony et al. 2010; Palucha-Poniewiera and Pilc 2013; Toth et al. 2012). The advent of mGlu₇ receptor knockout mice paved the way for early research on mGlu₇ receptor's role in both anxiety and cognition. Interestingly, mGlu₇ receptor knockouts display impaired cognitive abilities in many paradigms including spatial memory acquisition (Callaerts-Vegh et al. 2006; Goddyn et al. 2015; Holscher et al. 2004), impaired extinction learning (Callaerts-Vegh et al. 2006; Goddyn et al. 2015; Goddyn et al. 2008) as well as deficits to acquiring conditioned responses (Goddyn et al. 2015; Goddyn et al. 2008). Furthermore, mGlu₇ receptor knockouts display an attenuated response to anxiogenic stimuli as measured in many different behavioural tests including the elevated-plus maze, staircase test, open-field test, marble-burying test and stress-induced hyperthermia (SIH) (Callaerts-Vegh et al. 2006; Cryan et al. 2003; Stachowicz et al. 2008).

The development of the first selective mGlu₇ receptor agonist, AMN082 (Mitsukawa et al. 2005), has allowed modulation of mGlu₇ receptor function in adult animals. However, the results obtained have been somewhat paradoxical when examined in light of some of the previous knockout studies. AMN082 administration impaired fear acquisition (Siegl et al. 2008) and the short-term extinction of fear memories (Toth et al. 2012) but interestingly

facilitated the extinction of fear memories when tested over a longer time frame (Dobi et al. 2013; Fendt et al. 2008; Toth et al. 2012) possibly due to AMN082 induced impaired amygdalar plasticity (Fendt et al. 2013; Fendt et al. 2008). Moreover, AMN082 rescued the fear extinction deficits observed in the S1 mouse (Whittle et al. 2013). In relation to anxiety-like behaviour, AMN082 administration produced data similar to knockout animals with administration resulting in an anxiolytic phenotype in tests such as stress-induced hyperthermia (SIH) and the four-plate test (Stachowicz et al. 2008). Furthermore, siRNA-induced reduction of mGlu₇ receptor led to potent anxiolysis in mice (O'Connor et al. 2013b). Interestingly, a negative allosteric modulator of the mGlu₇ receptor, ADX71743, induced anxiolysis as measured by the reduction of visceral hypersensitivity in a stress sensitive rat strain (Moloney et al. 2015). In contrast, neither genetic nor pharmacological reduction of mGlu₇ receptor activity impacts anxiety or fear learning in a mouse model of a neurodevelopmental disorder (*MECP2* Duplication syndrome) (Fisher et al. 2017).

Given the apparent similar effects resulting from mGlu₇ receptor agonism and inhibition it is worth noting reports that a metabolite of AMN082, N-benzhydrylethane-1,2-diamine (Met-1), at physiologically relevant levels, inhibits the serotonin transporter, noradrenaline transporter and dopamine transporter (Sukoff Rizzo et al. 2011) possibly inducing AMN082's observed effects in vivo. However, the AMPA receptor antagonist 2,3-dihydroxy-6-nitro-7-sulfamoyl-benzo[f]quinoxaline-2,3-dione (Bradley et al. 2012) and an mGlu₇ receptor antagonist MMPIP both attenuated antidepressant-like effects of AMN082 (O'Connor and Cryan 2013) administration pointing to a glutamatergic mechanism.

Although much of the data has been somewhat paradoxical with knockout and pharmacological upregulation producing similar results, it is clear that mGlu₇ receptor is involved in regulating emotional behaviour and cognitive processes, including fear extinction. The S1 mouse strain represents a model of impaired regulation of fear memories seen in PTSD (Holmes and Singewald 2013). To assess whether differences in mGlu₇ receptor levels contribute to these fear extinction deficits, the present study used in situ hybridisation (ISH) to compare mGlu₇ receptor mRNA levels between this strain and the B6 strain, which shows robust fear extinction. Expression levels were analysed in three different brain regions critical for fear extinction, namely the prefrontal cortex (PFC), hippo-campus and amygdala.

Methods

Animals

Subjects were male S1 and B6 mice obtained at ~ 8 weeks of age from the Jackson Laboratory (Bar Harbour, ME). Animals were naïve from the same cohort tested in (Camp et al. 2009) where S1 mice showed robust deficits to fear extinction. Mice were housed by strain (2 per cage) in a temperature (22 ± 3 °C) and humidity ($45 \pm 15\%$) controlled vivarium under a 12-h light/dark cycle (lights on 0600 h). Cage dimensions were 37 cm × 16 cm × 13 cm. All experiments were conducted in accordance with the European Directive 86/609/EEC and the local Animal Care and Use Committees.

Sample collection

Mice were sacrificed by decapitation. The brain was immediately extracted, snap-frozen in cold isopentane and stored at -80°C before being processed for in situ hybridisation.

In situ hybridisation

The procedure was carried out using an oligodeoxynucleotide (cDNA) probe complementary to mGlu₇ receptor mRNA (1336–1292 pb access number NM_177328.3), labelled with a digoxigenin (DIG) oligonucleotide tailing kit (Roche, Molecular Biochemicals, Mannheim, Germany). Frozen 10- μm -thick sections were initially post-fixed in 4% paraformaldehyde and hybridised over night at 37°C in formamide 50%, saline and sodium citrate buffer (SSC) 4 $\times$, sheared salmon DNA 6.25 mg/mL, tRNA 125 $\mu\text{g/mL}$ and cDNA probe at fixed concentration of 100 pmol/mL. On the following day, sections were washed in saline-sodium citrate buffer, and the DIG molecules attached to the hybridised probes were detected with an anti-DIG antibody, conjugated with an alkaline phosphatase (Roche, Molecular Biochemicals, Mannheim, Germany). Finally, a substrate for the alkaline phosphatase (NBT/BCIP) (Sigma, St. Louis, MO, USA) was added. Specificity of the hybridization was evaluated by the use of 100-fold excess of the unlabelled oligodeoxynucleotide. For quantitative analysis, densitometric measurements of each hippocampal areas were analysed using FujiFilm's Science Lab Multi Gauge v2.2 software. All pictures were analysed in grey scale by measuring the density of pixels in a given area.

Statistical analysis

Data was analysed using Student's *t* test using GraphPad prism 6 software. A *p* value of 0.05 was chosen as the threshold for significance.

Results

Amygdala

mGlu₇ receptor mRNA expression was found to be significantly higher in the amygdala (LA + BLA) in the S1 strain when compared to B6 (Student's *t* test $p < 0.001$, $t = 3.557$, $df = 10$) (Fig. 1a, e, f). mGlu₇ receptor mRNA expression was found to be significantly higher in the LA in the S1 strain when compared to B6 (Student's *t* test $p < 0.001$, $t = 4.304$, $df = 10$) (Fig. 1b, e, f). mGlu₇ receptor mRNA expression was found to be significantly higher in the basolateral amygdala in the S1 strain when compared to B6 (Student's *t* test $p < 0.05$, $t = 2.318$, $df = 10$) (Fig. 1c, e, f). No significant difference in mGlu₇ receptor mRNA was found in the shell of the CeA/medial intercalated cell clusters between the strains (Student's *t* test $p > 0.05$, $t = 1.604$, $df = 10$) (Fig. 1d, e, f).

Hippocampus

mGlu₇ receptor mRNA expression was found to be significantly higher in the hippocampus (CA1 + CA3) in the S1 strain when compared to B6 (Student's *t* test $p < 0.001$, $t = 2.770$, $df = 10$) (Fig. 2a, e, f). mGlu₇ receptor mRNA expression was found to be significantly higher in the CA1 region of the hippocampus in the S1 strain when compared to B6 (Student's *t* test $p < 0.05$, $t = 2.660$, $df = 10$) (Fig. 2b, e, f). mGlu₇ receptor mRNA expression was found to

be significantly higher in the CA3 region of the hippocampus in the S1 strain when compared to B6 (Student's t test $p < 0.05$, $t = 3.839$, $df = 10$) (Fig. 2c, e, f). No significant difference in mGlu₇ receptor mRNA levels was found in the dentate gyrus of the hippocampus between the strains (Student's t test $p > 0.05$, $t = 0.4827$, $df = 10$) (Fig. 2d, e, f).

Infralimbic cortex, anterior cingulate gyrus and prelimbic region

No significant difference in mGlu₇ receptor mRNA was found in the infralimbic cortex (Student's t test $p > 0.05$, $t = 0.5937$, $df = 10$), anterior cingulate gyrus (Student's t test $p > 0.05$, $t = 0.5757$, $df = 10$) or the prelimbic region between strains (Student's t test $p > 0.05$, $t = 0.5496$, $df = 10$) (Fig. 3).

Discussion

The current study compared mGlu₇ receptor expression levels between the S1 inbred mouse strain, impaired in its ability to extinguish fear, and the B6 inbred strain which has normal extinction patterns (Hefner et al. 2008; Whittle et al. 2013) in key brain regions associated with fear extinction namely the amygdala, hippocampus, prefrontal cortex. In the amygdala, we found increased mGlu₇ receptor mRNA expression in the S1 strain and when further divided into different sub-regions, increased expression was found in the LA and the BLA, but not in the CeA. Similarly, increased expression levels of mGlu₇ receptor were also seen in the hippocampus of S1 mice compared to the B6 strain, with an increase seen in the CA1 and CA3 regions but not the DG. Interestingly, no difference in mGlu₇ receptor levels was seen in the medial prefrontal cortex, prelimbic region or anterior cingulated gyrus.

Previous work on mGlu₇ receptor has demonstrated its critical role in both maintaining correct cognitive functioning (Callaerts-Vegh et al. 2006; Dobi et al. 2013; Fendt et al. 2008; Goddyn et al. 2015; Goddyn et al. 2008; Holscher et al. 2004, 2005; Klakotskaia et al. 2013; Moloney et al. 2015; Toth et al. 2012), regulating anxiety-like behaviour (Callaerts-Vegh et al. 2006; Cryan et al. 2003; Moloney et al. 2015; O'Connor et al. 2013b; Stachowicz et al. 2008) and appropriately responding to psychologically stressful events (Gee et al. 2014; Moloney et al. 2015; Peterlik et al. 2016; Peterlik et al. 2017). Situated at the interface of cognition and emotion, mGlu₇ receptor is in a key position to regulate processes that may contribute to the pathophysiology underlying PTSD (O'Connor et al. 2010).

It is interesting to find that the S1 strain has increased mGlu₇ receptor mRNA in the amygdala and hippocampus since both global mGlu₇ receptor ablation and knockdown result in an attenuation to the extinction of aversive memories (Callaerts-Vegh et al. 2006; Fendt et al. 2008) and systemic pharmacological mGlu₇ receptor activation facilitates the extinction of aversive memories (Fendt et al. 2008; Toth et al. 2012). It might be expected that a strain showing poor extinction such as the S1 would have lower mGlu₇ receptor levels in neuronal circuits that regulate fear extinction. There are a number of different possibilities that may help explain elevated amygdalar and hippocampal mGlu₇ receptor levels including the mGlu₇ receptor's role as a presynaptic inhibitor of both GABA and glutamate release and possible increased compensatory mGlu₇ receptor expression in response to additional deficits in the signalling cascades that mediate the extinction of fear memories.

Evidence to date has ascribed a role to many different members of the mGlu receptor family in the acquisition, extinction and recall of fear memories. However, each mGlu receptor seems to occupy a specific niche in these processes. For example, intra-amygdalar pharmacological activation of mGlu_{2/3} receptors did not affect the rate of fear extinction acquisition but did enhance the retention of fear memories (Kim et al. 2015). Furthermore, systemically blocking mGlu₅ receptor activity facilitated the spontaneous recovery of fear memories but not extinction per se, whereas the other group 1 mGlu receptor member, mGlu₁ receptor, does not seem to play a role in fear extinction (Mao et al. 2013). The other group III mGlu receptors appear to be more involved in the acquisition of fear memories, given mutant mice lacking mGlu₄ receptor had enhanced acquisition of cued freezing responses (Davis et al. 2013) whereas mGlu₈ receptor knockout mice exhibit deficits in fear acquisition but no changes to fear extinction (Fendt et al. 2010). Thus, most members of the mGlu receptor family play key roles in fear memory formation but specifically regulating the extinction of such memories appears unique to mGlu₇ receptor based on the currently available evidence.

It is worth noting that the mGlu₇ receptor agonist AMN082 was able to rescue the fear extinction deficits seen in the S1 mouse (Whittle et al. 2013). The amygdala contains a large population of GABAergic neurons which are activated by glutamate (Rainnie et al. 1991) and it has been suggested that glutamate-induced increases in GABA release may contribute to the extinction of fear memories (Rainnie et al. 1991). Increased amygdalar mGlu₇ receptor levels would result in potentiation of the negative feedback that inhibits further release of glutamate (Summa et al. 2013). This would lead to a reduction in the glutamatergic innervation of GABAergic neurons, resulting in an overall decrease of inhibition levels, which may contribute to the poor extinction observed in the S1 strain.

Preceding extinction, the hippocampus contributes to the contextual retrieval of fear memory through its inputs to the BLA (Corcoran and Maren 2001; Herry et al. 2008). These inputs are coupled to a subset of neurons that are activated when a conditioned stimulus and an unconditioned stimulus are presented together. Following extinction, these neurons are no longer active. If the increased mGlu₇ receptor levels seen in the S1 strain are on hippocampal GABAergic neurons, the subsequent reduced GABA release could lead to an exaggerated activation of these fear neurons, preventing fear extinction. Recent data indicates that correct mGlu₇ receptor function is required for hippocampal plasticity (Klar et al. 2015). Currently, we can only speculate on the differing contributions of glutamatergic and GABAergic mGlu₇ receptor levels to fear extinction processes. Experiments localising mGlu₇ receptor on either GABAergic or glutamatergic neurons in different regions of the brain critical to fear extinction will reveal more about how mGlu₇ receptor influences the extinction of fear memories and may help to explain why a strain exhibiting profoundly impaired fear extinction displays increased mGlu₇ receptor in brain regions associated with fear extinction.

It has been well established that the mPFC plays a prominent role in both the acquisition and extinction of fear memories (Corcoran and Quirk 2007; Sotres-Bayon and Quirk 2010). Furthermore, there are reciprocal projections between the mPFC and the amygdala (Conde et al. 1995; Dzhaiani et al. 1987; Herry et al. 2008; McDonald et al. 1996) with cortical

neurons providing inhibitory control over BLA neurons during the extinction of fear memories (Bukalo et al. 2015; Marek et al. 2013). Interestingly, in contrast to the hippocampus and the amygdala, no difference in mGlu₇ receptor levels was seen between the B6 and S1 strains in various subregions of the mPFC. Thus suggesting the impaired fear extinction seen in the S1 strain may not result from any differences in cortical derived inhibition of amygdalar activity driven via a mGlu₇ receptor-dependent mechanism.

In sum, the current study shows mGlu₇ receptor mRNA levels to be different between two different inbred mouse strains that differ markedly in their ability to extinguish conditioned fear with the S1 mouse having increased mGlu₇ receptor mRNA levels in the amygdala and hippocampus. This strongly suggests a role for mGlu₇ receptor in the neuronal processes associated with fear extinction. Furthermore, based on pharmacological data showing activation of mGlu₇ receptor facilitates the extinction of aversive memories (Fendt et al. 2008; Toth et al. 2012), it is more likely, at least in C57 mice which display robust fear extinction, that changes to receptor function and not any changes to expression contributes to fear memory extinction. Moreover, it further confirms the utility of the S1 mouse as a tool to understand the molecular basis of extinction (Gunduz-Cinar et al. 2018) and ways to boost extinction memories (Bukalo et al. 2014; Singewald et al. 2015). This could be of great relevance to our understanding of the neuronal circuitry involved in PTSD and increases our knowledge of what may be the underlying molecular disturbances taking place in this devastating disorder.

Funding information

JFC is supported by Science Foundation Ireland (SFI) (Grant Numbers 07/CE/BI368 and 12/RC/2273); the Irish Health Research Board. RMOC was supported by a postgraduate scholarship awarded by the Irish Research Council for Science Engineering and Technology (IRCSET). AH is supported by the NIAAA IRP. NS is supported by the Austrian Science Fund FWF SFB F4410.

References

- Bradley SR, Uslaner JM, Flick RB, Lee A, Groover KM, Hutson PH (2012) The mGluR7 allosteric agonist AMN082 produces antidepressant-like effects by modulating glutamatergic signaling. *Pharmacol Biochem Behav* 101:35–40 [PubMed: 22138407]
- Breslau N, Kessler RC, Chilcoat HD, Schultz LR, Davis GC, Andreski P (1998) Trauma and posttraumatic stress disorder in the community: the 1996 Detroit area survey of trauma. *Arch Gen Psychiatry* 55:626–632 [PubMed: 9672053]
- Bukalo O, Pinard CR, Holmes A (2014) Mechanisms to medicines: elucidating neural and molecular substrates of fear extinction to identify novel treatments for anxiety disorders. *Br J Pharmacol* 171:4690–4718 [PubMed: 24835117]
- Bukalo O, Pinard CR, Silverstein S, Brehm C, Hartley ND, Whittle N, Colacicco G, Busch E, Patel S, Singewald N, Holmes A (2015) Prefrontal inputs to the amygdala instruct fear extinction memory formation. *Sci Adv* 1(6). 10.1126/sciadv.1500251
- Callaerts-Vegh Z, Beckers T, Ball SM, Baeyens F, Callaerts PF, Cryan JF, Molnar E, D'Hooge R (2006) Concomitant deficits in working memory and fear extinction are functionally dissociated from reduced anxiety in metabotropic glutamate receptor 7-deficient mice. *J Neurosci* 26:6573–6582 [PubMed: 16775145]
- Camp M, Norcross M, Whittle N, Feyder M, D'Hanis W, Yilmazer-Hanke D, Singewald N, Holmes A (2009) Impaired Pavlovian fear extinction is a common phenotype across genetic lineages of the 129 inbred mouse strain. *Genes Brain Behav* 8:744–752 [PubMed: 19674120]

- Conde F, Maire-Lepoivre E, Audinat E, Crepel F (1995) Afferent connections of the medial frontal cortex of the rat. II Cortical and sub-cortical afferents. *J Comp Neurol* 352:567–593 [PubMed: 7722001]
- Corcoran KA, Maren S (2001) Hippocampal inactivation disrupts contextual retrieval of fear memory after extinction. *J Neurosci* 21:1720–1726 [PubMed: 11222661]
- Corcoran KA, Quirk GJ (2007) Activity in prelimbic cortex is necessary for the expression of learned, but not innate, fears. *J Neurosci* 27:840–844 [PubMed: 17251424]
- Cryan JF, Kelly PH, Neijt HC, Sansig G, Flor PJ, van Der Putten H (2003) Antidepressant and anxiolytic-like effects in mice lacking the group III metabotropic glutamate receptor mGluR7. *Eur J Neurosci* 17:2409–2417 [PubMed: 12814372]
- Davis MJ, Iancu OD, Acher FC, Stewart BM, Eiwaz MA, Duvoisin RM, Raber J (2013) Role of mGluR4 in acquisition of fear learning and memory. *Neuropharmacology* 66:365–372 [PubMed: 22884897]
- Depino AM, Gross C (2007) Simultaneous assessment of autonomic function and anxiety-related behavior in BALB/c and C57BL/6 mice. *Behav Brain Res* 177:254–260 [PubMed: 17204340]
- Dobi A, Sartori SB, Busti D, Van der Putten H, Singewald N, Shigemoto R, Ferraguti F (2013) Neural substrates for the distinct effects of presynaptic group III metabotropic glutamate receptors on extinction of contextual fear conditioning in mice. *Neuropharmacology* 66:274–289 [PubMed: 22643400]
- DSM-V (2013) Diagnostic and statistical manual of mental disorders: DSM-V. American Psychiatric Association, Washington D.C
- Dzhaiani NN, Egorova VV, Iezuitova NN, Nikitina AA, Timofeeva NM (1987) Structural-functional characteristics of the small intestine in adult and growing rats after various surgical procedures. *Dokl Akad Nauk SSSR* 295:1501–1504 [PubMed: 3678031]
- Fendt M, Schmid S, Thakker DR, Jacobson LH, Yamamoto R, Mitsukawa K, Maier R, Natt F, Husken D, Kelly PH, McAllister KH, Hoyer D, van der Putten H, Cryan JF, Flor PJ (2008) mGluR7 facilitates extinction of aversive memories and controls amygdala plasticity. *Mol Psychiatry* 13:970–979 [PubMed: 17712315]
- Fendt M, Burki H, Imobersteg S, van der Putten H, McAllister K, Leslie JC, Shaw D, Holscher C (2010) The effect of mGlu8 deficiency in animal models of psychiatric diseases. *Genes Brain Behav* 9:33–44 [PubMed: 19740090]
- Fendt M, Imobersteg S, Peterlik D, Chaperon F, Mattes C, Wittmann C, Olpe HR, Mosbacher J, Vranesic I, van der Putten H, McAllister KH, Flor PJ, Gee CE (2013) Differential roles of mGlu(7) and mGlu(8) in amygdala-dependent behavior and physiology. *Neuropharmacology* 72:215–223 [PubMed: 23664812]
- Fisher NM, Gogliotti RG, Vermudez SAD, Stansley BJ, Conn PJ, Niswender CM (2017) Genetic reduction or negative modulation of mGlu7 does not impact anxiety and fear learning phenotypes in a mouse model of MECP2 duplication syndrome. *ACS Chem Neurosci*. 10.1021/acscchemneuro.7b00414
- Fitzgerald PJ, Whittle N, Flynn SM, Graybeal C, Pinard CR, Gunduz-Cinar O, Kravitz AV, Singewald N, Holmes A (2014) Prefrontal single-unit firing associated with deficient extinction in mice. *Neurobiol Learn Mem* 113:69–81 [PubMed: 24231425]
- Gee CE, Peterlik D, Neuhauser C, Bouhelal R, Kaupmann K, Laue G, Uschold-Schmidt N, Feuerbach D, Zimmermann K, Ofner S, Cryan JF, van der Putten H, Fendt M, Vranesic I, Glatthar R, Flor PJ (2014) Blocking metabotropic glutamate receptor subtype 7 (mGlu7) via the Venus flytrap domain (VFTD) inhibits amygdala plasticity, stress, and anxiety-related behavior. *J Biol Chem* 289:10975–10987 [PubMed: 24596089]
- Goddyn H, Callaerts-Vegh Z, Stroobants S, Dirix T, Vansteenwegen D, Hermans D, van der Putten H, D'Hooge R (2008) Deficits in acquisition and extinction of conditioned responses in mGluR7 knockout mice. *Neurobiol Learn Mem* 90:103–111 [PubMed: 18289889]
- Goddyn H, Callaerts-Vegh Z, D'Hooge R (2015) Functional Dissociation of Group III Metabotropic Glutamate Receptors Revealed by Direct Comparison between the Behavioral Profiles of Knockout Mouse Lines. *Int J Neuropsychopharmacol* 18:pyv053 [PubMed: 25999589]

- Gunduz-Cinar O, Brockway E, Lederle L, Wilcox T, Halladay LR, Ding Y, Oh H, Busch EF, Kaugars K, Flynn S, Limoges A, Bukalo O, MacPherson KP, Masneuf S, Pinard C, Sibille E, Chesler EJ, Holmes A (2018) Identification of a novel gene regulating amygdala-mediated fear extinction. *Mol Psychiatry* 10.1038/s41380-017-0003-3
- Hefner K, Whittle N, Juhasz J, Norcross M, Karlsson RM, Saksida LM, Bussey TJ, Singewald N, Holmes A (2008) Impaired fear extinction learning and cortico-amygdala circuit abnormalities in a common genetic mouse strain. *J Neurosci* 28:8074–8085 [PubMed: 18685032]
- Herry C, Cioocchi S, Senn V, Demmou L, Muller C, Luthi A (2008) Switching on and off fear by distinct neuronal circuits. *Nature* 454:600–606 [PubMed: 18615015]
- Holmes A, Singewald N (2013) Individual differences in recovery from traumatic fear. *Trends Neurosci* 36:23–31 [PubMed: 23260015]
- Holscher C, Schmid S, Pilz PK, Sansig G, van der Putten H, Plappert CF (2004) Lack of the metabotropic glutamate receptor subtype 7 selectively impairs short-term working memory but not long-term memory. *Behav Brain Res* 154:473–481 [PubMed: 15313036]
- Holscher C, Schmid S, Pilz PK, Sansig G, van der Putten H, Plappert CF (2005) Lack of the metabotropic glutamate receptor subtype 7 selectively modulates Theta rhythm and working memory. *Learn Mem* 12:450–455 [PubMed: 16204199]
- Kendler KS (2001) Twin studies of psychiatric illness: an update. *Arch Gen Psychiatry* 58:1005–1014 [PubMed: 11695946]
- Kessler RC, Sonnega A, Bromet E, Hughes M, Nelson CB (1995) Posttraumatic stress disorder in the National Comorbidity Survey. *Arch Gen Psychiatry* 52:1048–1060 [PubMed: 7492257]
- Kim J, An B, Kim J, Park S, Park S, Hong I, Lee S, Park K, Choi S (2015) mGluR2/3 in the Lateral Amygdala is Required for Fear Extinction: Cortical input synapses onto the lateral amygdala as a target site of the mGluR2/3 action. *Neuropsychopharmacology* 40:2916–2928 [PubMed: 26081171]
- Klakotskaia D, Ramsey AK, Fowler SW, Serfozo P, Simonyi A, Schachtman TR (2013) Effects of group II and III metabotropic glutamate receptor ligands on conditioned taste aversion learning. *Behav Brain Res* 253:9–16 [PubMed: 23827202]
- Klar R, Walker AG, Ghose D, Grueter BA, Engers DW, Hopkins CR, Lindsley CW, Xiang Z, Conn PJ, Niswender CM (2015) Activation of metabotropic glutamate receptor 7 is required for induction of long-term potentiation at SC-CA1 synapses in the Hippocampus. *J Neurosci* 35:7600–7615 [PubMed: 25972184]
- Mao SC, Chang CH, Wu CC, Orejarena MJ, Manzoni OJ, Gean PW (2013) Inhibition of spontaneous recovery of fear by mGluR5 after prolonged extinction training. *PLoS One* 8:e59580 [PubMed: 23555716]
- Marek R, Strobel C, Bredy TW, Sah P (2013) The amygdala and medial prefrontal cortex: partners in the fear circuit. *J Physiol* 591:2381–2391 [PubMed: 23420655]
- Maren S, Holmes A (2016) Stress and fear extinction. *Neuropsychopharmacology* 41:58–79 [PubMed: 26105142]
- McDonald AJ, Mascagni F, Guo L (1996) Projections of the medial and lateral prefrontal cortices to the amygdala: a Phaseolus vulgaris leucoagglutinin study in the rat. *Neuroscience* 71:55–75 [PubMed: 8834392]
- Mi B, Liu G, Zhou W, Lv H, Liu Y, Liu J (2018) Identification of genes and pathways in the synovia of women with osteoarthritis by bioinformatics analysis. *Mol Med Rep* 17:4467–4473 [PubMed: 29344651]
- Mitsukawa K, Yamamoto R, Ofner S, Nozulak J, Pescott O, Lukic S, Stoeckl N, Mombereau C, Kuhn R, McAllister KH, van der Putten H, Cryan JF, Flor PJ (2005) A selective metabotropic glutamate receptor 7 agonist: activation of receptor signaling via an allosteric site modulates stress parameters in vivo. *Proc Natl Acad Sci U S A* 102:18712–18717 [PubMed: 16339898]
- Moloney RD, Golubeva AV, O'Connor RM, Kalinichev M, Dinan TG, Cryan JF (2015) Negative allosteric modulation of the mGlu7 receptor reduces visceral hypersensitivity in a stress-sensitive rat strain. *Neurobiol Stress* 2:28–33 [PubMed: 26844237]

- O'Connor RM, Cryan JF (2010) Role of Metabotropic Glutamate Receptors in CNS Disorders In: Siehler S, Milligan G (eds) *G Protein-Coupled Receptors: Structure, Signaling, and Physiology*. Cambridge University Press, Cambridge, pp 321–379
- O'Connor RM, Cryan JF (2013) The effects of mGlu(7) receptor modulation in behavioural models sensitive to antidepressant action in two mouse strains. *Behav Pharmacol* 24:105–113 [PubMed: 23455446]
- O'Connor RM, Finger BC, Flor PJ, Cryan JF (2010) Metabotropic glutamate receptor 7: at the interface of cognition and emotion. *Eur J Pharmacol* 639:123–131 [PubMed: 20371242]
- O'Connor RM, Pusccheddu MM, Dinan TG, Cryan JF (2013a) Impact of early-life stress, on group III mGlu receptor levels in the rat hippocampus: effects of ketamine, electroconvulsive shock therapy and fluoxetine treatment. *Neuropharmacology* 66:236–241 [PubMed: 22609536]
- O'Connor RM, Thakker DR, Schmutz M, van der Putten H, Hoyer D, Flor PJ, Cryan JF (2013b) Adult siRNA-induced knockdown of mGlu7 receptors reduces anxiety in the mouse. *Neuropharmacology* 72:66–73 [PubMed: 23603202]
- O'Mahony CM, Bravo JA, Dinan TG, Cryan JF (2010) Comparison of hippocampal metabotropic glutamate receptor 7 (mGlu7) mRNA levels in two animal models of depression. *Neurosci Lett* 482:137–141 [PubMed: 20638442]
- Palucha-Poniewiera A, Pilc A (2013) A selective mGlu7 receptor antagonist MMPIP reversed antidepressant-like effects of AMN082 in rats. *Behav Brain Res* 238:109–112 [PubMed: 23085340]
- Peterlik D, Flor PJ, Uschold-Schmidt N (2016) The emerging role of metabotropic glutamate receptors in the pathophysiology of chronic stress-related disorders. *Curr Neuropharmacol* 14:514–539 [PubMed: 27296643]
- Peterlik D, Stangl C, Bludau A, Grabski D, Strasser R, Schmidt D, Flor PJ, Uschold-Schmidt N (2017) Relief from detrimental consequences of chronic psychosocial stress in mice deficient for the metabotropic glutamate receptor subtype 7. *Neuropharmacology* 115:139–148 [PubMed: 27188844]
- Rainnie DG, Asprodini EK, Shinnick-Gallagher P (1991) Inhibitory transmission in the basolateral amygdala. *J Neurophysiol* 66:999–1009 [PubMed: 1684384]
- Siegl S, Flor PJ, Fendt M (2008) Amygdaloid metabotropic glutamate receptor subtype 7 is involved in the acquisition of conditioned fear. *Neuroreport* 19:1147–1150 [PubMed: 18596617]
- Singewald N, Schmuckermair C, Whittle N, Holmes A, Ressler KJ (2015) Pharmacology of cognitive enhancers for exposure-based therapy of fear, anxiety and trauma-related disorders. *Pharmacol Ther* 149:150–190 [PubMed: 25550231]
- Sotres-Bayon F, Quirk GJ (2010) Prefrontal control of fear: more than just extinction. *Curr Opin Neurobiol* 20:231–235 [PubMed: 20303254]
- Stachowicz K, Branski P, Klak K, van der Putten H, Cryan JF, Flor PJ, Andrzej P (2008) Selective activation of metabotropic G-protein-coupled glutamate 7 receptor elicits anxiolytic-like effects in mice by modulating GABAergic neurotransmission. *Behav Pharmacol* 19:597–603 [PubMed: 18690114]
- Stein MB, Jang KL, Taylor S, Vernon PA, Livesley WJ (2002) Genetic and environmental influences on trauma exposure and posttraumatic stress disorder symptoms: a twin study. *Am J Psychiatry* 159:1675–1681 [PubMed: 12359672]
- Stiedl O, Meyer M, Kishimoto T, Rosenfeld MG, Spiess J (2003) Stress-mediated heart rate dynamics after deletion of the gene encoding corticotropin-releasing factor receptor 2. *Eur J Neurosci* 17:2231–2235 [PubMed: 12786991]
- Sukoff Rizzo SJ, Leonard SK, Gilbert A, Dollings P, Smith DL, Zhang MY, Di L, Platt BJ, Neal S, Dwyer JM, Bender CN, Zhang J, Lock T, Kowal D, Kramer A, Randall A, Huselton C, Vishwanathan K, Tse SY, Butera J, Ring RH, Rosenzweig-Lipson S, Hughes ZA, Dunlop J (2011) The metabotropic glutamate receptor 7 allosteric modulator AMN082: a monoaminergic agent in disguise? *J Pharmacol Exp Ther* 338:345–352 [PubMed: 21508084]
- Summa M, Di Prisco S, Grilli M, Usai C, Marchi M, Pittaluga A (2013) Presynaptic mGlu7 receptors control GABA release in mouse hippocampus. *Neuropharmacology* 66:215–224 [PubMed: 22564442]

- Toth I, Dietz M, Peterlik D, Huber SE, Fendt M, Neumann ID, Flor PJ, Slattery DA (2012) Pharmacological interference with metabotropic glutamate receptor subtype 7 but not subtype 5 differentially affects within- and between-session extinction of Pavlovian conditioned fear. *Neuropharmacology* 62:1619–1626 [PubMed: 22079159]
- Tovote P, Meyer M, Beck-Sickinger AG, von Horsten S, Ove Ogren S, Spiess J, Stiedl O (2004) Central NPY receptor-mediated alteration of heart rate dynamics in mice during expression of fear conditioned to an auditory cue. *Regul Pept* 120:205–214 [PubMed: 15177939]
- Whittle N, Schmuckermair C, Gunduz Cinar O, Hauschild M, Ferraguti F, Holmes A, Singewald N (2013) Deep brain stimulation, histone deacetylase inhibitors and glutamatergic drugs rescue resistance to fear extinction in a genetic mouse model. *Neuropharmacology* 64:414–423 [PubMed: 22722028]

Fig. 1.

In situ hybridisation of mGlu₇ receptor in the amygdala of B6 and S1 mice. **a–c** Densitometric analysis showing the S1 mice had increased levels of mGlu₇ receptor in the amygdala (LA + BLA), lateral amygdala (LA) and basolateral amygdala (BLA), respectively. **d** No significant difference in mGlu₇ receptor expression of the CeA of the amygdala was found between the two strains. **e** Representative image of mGlu₇ receptor mRNA staining in the amygdala of B6 mice. **f** Representative image of mGlu₇ receptor mRNA staining in the amygdala of S1 mice.

*Significant using Student's *t* test when compared to B6. *n* = 6

Fig. 2.

In situ hybridisation of mGlu₇ receptor in the hippocampus of B6 and S1 mice. **a-c** Densitometric analysis showing the S1 mice had increased levels of mGlu₇ receptor in the hippocampus (CA1 + CA3), CA1 and CA3 regions of the hippocampus. **d** No significant difference in mGlu₇ receptor expression of the dentate gyrus of the hippocampus was found between the two strains. **e** Representative image of mGlu₇ receptor mRNA staining in the hippocampus of B6 mice. **f** Representative image of mGlu₇ receptor mRNA staining in the hippocampus of S1 mice.

*Significant using Student's *t* test when compared to B6. *n* = 6

Fig. 3.

In situ hybridisation of mGlu7 receptor in the prefrontal cortex of B6 and S1 mice. **a–c** No significant difference in mGlu7 receptor expression in the prefrontal cortex was found between the two strains. **d** Representative image of mGlu7 receptor mRNA staining in the prefrontal cortex of B6 mice. **e** Representative image of mGlu7 receptor mRNA staining in the prefrontal cortex of S1 mice.

CgL = cingulate gyrus, PrL = prelimbic region, IL = infralimbic. $n = 6$