Published in final edited form as:

Curr Treat Options Cardiovasc Med. ; 21(8): 38. doi:10.1007/s11936-019-0741-4.

Optimal non-invasive strategies to reduce recurrent atherosclerotic cardiovascular disease risk

Maeve Jones-O'Connor, MB BCh BAO^{1,2}, Pradeep Natarajan, MD MMSc^{1,2,3,4}

¹Department of Medicine, Massachusetts General Hospital, Boston, MA

²Department of Medicine, Harvard Medical School, Boston, MA

³Cardiovascular Research Center and Center for Genomic Medicine, Massachusetts General Hospital, Boston, MA

⁴Program in Medical and Population Genetics, Broad Institute of Harvard & MIT, Cambridge, MA

Abstract

Purpose of review—Cardiovascular disease (CVD) remains the leading cause of death worldwide, with coronary artery disease (CAD) responsible for the vast majority of these deaths. Incidence is increasing in developing countries, and prevalence is increasing globally as populations age. Once coronary artery disease is manifest, recurrent event risk remains high.

Recent findings—Multiple therapeutic avenues have had significant recent developments, including diet, low density lipoprotein management, triglycerides, hypoglycemic agents, anti platelet agents and oral anti-coagulants.

Summary—Combined approaches involving specific, tailored lifestyle and pharmacological interventions will provide the most effective strategy for reducing the risk of recurrent CVD events. Here, we review risk prediction, non-invasive non-pharmacologic and pharmacologic approaches to mitigate residual coronary artery disease risk.

INTRODUCTION

Despite many advances in cardiovascular medicine, cardiovascular disease (CVD) remains the leading cause of mortality worldwide. In 2016, nearly 18 million people died of cardiovascular diseases globally. Of these, 9.4 million were caused by coronary artery disease (CAD). In addition, the burden of mortality from CAD is spreading beyond traditional high-income areas. In 2000, it was the 8th leading cause of death in low-income countries. By 2016, it progressed to the 3rd leading cause [1]

In addition to this burden of mortality, CVD also generates a high economic cost. In the USA, the direct cost of CVD was \$213.8 billion in 2014–2015. This is over double the cost

Please address correspondence to: Pradeep Natarajan, MD MMSc, 185 Cambridge St, CPZN 3.184, Boston, MA 02114, Tel: 617-724-3526, Fax: 617-726-2203, pnatarajan@mgh.harvard.edu, Twitter: @pnatarajanmd.

Disclosures

P.N. reports grant support from Amgen and Boston Scientific, and consulting income from Apple. M.J.-O. reports no relevant disclosures.

from 1996–1997, of \$103.5 billion. Although risk factor modification, widespread use of primary revascularisation, and secondary prevention strategies have helped reduce cardiovascular mortality and morbidity in recent times, these costs are anticipated to continue to precipitously increase. A further doubling of the total economic cost of CVD is predicted between 2015 to 2035, from \$318 billion to \$749 billion [2].

RECURRENT RISK PREDICTION

The recently revised ACC/AHA ASCVD risk calculator is widely used and is intended for use in predicting risk of a first cardiovascular event [3]. However, this tool is commonly used in a secondary prevention setting as well, with a relative lack of specific risk calculation strategies in this cohort. Recently, a clinical tool for calculating recurrent CVD risk has been developed in TRS-2oP (Table 1) [4]. It has been validated in the trial setting, and recently has been shown to demonstrate good correlation in multiple cohorts. It is particularly useful in identifying patients at high risk [5–7].

DIET

Previous data suggests that a “higher quality” diet, rich in fruits, vegetables and whole grains, and lower in alcohol and trans-fats, conferred benefit in prevention of major chronic diseases, including CAD [8,9]. As such, this strategy is supported by the 2013 ACC/AHA guideline on lifestyle management to reduce CVD risk [10].

Results of the DART trial also suggest increased intake of fatty fish conferred benefit in terms of reducing all-cause mortality in men post myocardial infarction [11]. However, results of the follow up trial, DART-2, did not replicate these findings, and indeed showed a trend towards increased mortality in a subgroup taking fish oil supplements [12].

Other studies have demonstrated that a “Mediterranean” diet, lower in dietary fat overall and relatively higher in oleic and alpha-linolenic acids confers benefit in prevention of cardiovascular disease [13, 14]. The PREDIMED study is awaited to objectively assess this diet in secondary prevention. [15]

Other dietary interventions for health have come to public attention recently, notably intermittent fasting and its variations. A recent randomised, crossover trial has demonstrated benefit of the fasting-mimicking diet (FMD) in reducing numerous cardiovascular risk markers, including BMI, systemic blood pressure, total and LDL-cholesterol, triglycerides and C-reactive protein [16].

Epidemiologic studies link the consumption of whole food plant-based diets with improvements in CVD risk factors [17–20]. Analysis of trials comparing plant based strategies for weight loss suggest plant-based dietary strategies to reduce recurrent CVD risk [21]. Plant-based choices may also be linked to an overall improvement in health-related behaviors, separately conferring benefit.

Indeed, older studies suggested 'intensive lifestyle changes', involving a combination of 'whole-food', vegetarian diets, exercise, stress management and medications, had been suggested to halt or even reverse prevalent CAD [22–27].

PHYSICAL ACTIVITY

Physical activity confers a plethora of health benefits. Comprehensive cardiac rehabilitation programs help deliver these benefits to patients with symptomatic CAD, as well as providing education and resources. Individual trials demonstrated reduced hospital admissions and potentially reduced all-cause mortality [28–31]. Recently, Cochrane reviewers found that exercise-based CR reduces cardiovascular mortality in comparison to a no-exercise control (risk ratio (RR) 0.74, (95% CI 0.64 to 0.86)) among individuals referred after myocardial infarction. The risk for hospital admission was decreased in CR groups (RR 0.82, 95% CI 0.70 to 0.96)[32].

Exercise also directly modifies CV risk factors. One study demonstrated reductions in small LDL-cholesterol and overall concentration of LDL-cholesterol in a high-volume, high-intensity treatment group [33]. High-volume, high-intensity exercise is also associated with increased HDL efflux as measured by global radiolabelled efflux [34].

Recent data also suggests that physical activity independently confers a survival advantage for those with CAD. High-volume physical activity (150 min of moderate-intensity or 60 min of vigorous activity per week) was associated with a 36% lower all-cause mortality risk versus inactivity (HR: 0.64; 95% CI: 0.50 to 0.83). A change from inactive to high physical activity conferred a 32% (HR: 0.68; 95% CI: 0.47 to 0.97) lower risk of cardiovascular mortality [35].

The 2018 US Physical Activity Guidelines recommends that adults should do >150–300 min/week of moderate-intensity, or 75–150 min/week of vigorous-intensity aerobic physical activity, or an equivalent combination [36].

SMOKING

The 1964 Surgeon General's report on the health effects of smoking clearly articulated the CVD risks of smoking [37]. The 1983 Surgeon General's report further outlined evidence that multiple pathogenic pathways are activated by smoking, including endothelial dysfunction, inflammation, lipid alterations, increase in prothrombotic state and myocardial oxygen supply-demand mismatch [38]. Recent smoking statistics from the CDC give overall prevalence rates of 17.5% and 13.5% in men and women in the general population [39]

A recently published review of three decades of data from the Framingham cohort demonstrated mitigated but persistently elevated cardiovascular risk in smokers across all decades of follow-up, despite improvements in risk factor modulation across that time [40].

Smoking cessation confers a significant protective effect in those with CAD and recent acute coronary syndrome. In those with known CAD, observational studies suggest a pooled reduction in overall mortality of up to 35% [41, 42]. In follow up analyses of OASIS-5,

patients post-ACS who successfully quit were 43% less likely to experience myocardial infarction (MI) at 6 months (95% CI, 0.36 to 0.89; $P=0.0145$) [43].

LOW-DENSITY LIPOPROTEIN (LDL) CHOLESTEROL

Circulating cholesterol levels were linked to atherosclerotic coronary artery disease since the original Framingham Heart Study in 1957. Over time, low-density lipoprotein cholesterol (LDL-C) levels emerged as being directly related to elevated coronary artery disease risk [44–49]. More recently, LDL particle number (LDL-P) has been observed to be strongly associated with primary and secondary CAD risk compared to conventional LDL-C measurements [50,51]. Particle number may also more accurately predict risk in those with discordant LDL-P and LDL-C, such as in type 2 diabetes and metabolic syndrome [52, 53]. Also, recent data suggest the subfractions of small dense LDL and oxidised LDL particles are proposed as being more atherogenic than others [54, 55].

HMG co-a reductase inhibitors (statins)

Statin medications have long been used to reduce circulating LDL-C levels via inhibition of HMG Co-A reductase, the rate-limiting enzyme for cholesterol synthesis. They were known to reduce cardiovascular morbidity and mortality in stable coronary artery disease, and their secondary prevention benefit in reducing recurrent CAD events and mortality was demonstrated in multiple randomized controlled clinical trials, including MIRACL, PROVE-TIMI 22 and IDEAL trials (Table 2). PROVE-TIMI 22 and IDEAL also suggested superiority of higher intensity therapy in reducing residual risk [56–58]. As such, LDL-C lowering with statins is a cornerstone of CVD risk reduction.

Ezetimibe

Ezetimibe acts via inhibition of gastrointestinal absorption of dietary cholesterol. The IMPROVE-IT compared the combination of ezetimibe with simvastatin against simvastatin monotherapy among individuals with ASCVD. At a median of 6 years, there were decreased rates of the combined primary end point of cardiovascular death, nonfatal myocardial infarction (MI), unstable angina requiring readmission, revascularisation or non-fatal stroke (Kaplan-Meier event rate of 32.7% in treatment group vs. 34.7%; HR 0.93 (95% CI 0.89–0.99, $p=0.016$). Notably, similar effects were seen with rates of fatal MI (14.8% v 13.1%, $p=0.002$) and nonfatal MI (14.4% vs. 12.8%) [59]. Recently, subgroup analyses among diabetics demonstrated greater absolute risk reduction with ezetimibe and potentially greater relative risk reduction among diabetics younger than 75 years [60].

PCSK9 monoclonal antibodies

ODYSSEY Outcomes studied alirocumab, a humanised monoclonal antibody inhibitor of PCSK9, in patients with recent (1–12 months) acute coronary syndrome on maximal statin therapy but with persistent LDL-C > 70 mg/dl. Primary outcome was defined as a composite of major adverse cardiac events (MACE); median follow-up was 2.8 years. Overall rate of MACE was 11.1% in the placebo group and 9.5% in the treatment group (ARR 1.6%). Rates of CAD events were 14.3% vs. 12.7% (ARR 1.6%; HR 0.88 [95% CI 0.81–0.95]) and major CAD events were 8.4% vs. 9.5% (ARR 1.1%; HR 0.88, 95% CI 0.8–0.96). Notably, there

was a greater treatment effect regarding MACE, CAD death, CV death and all cause death in patients with baseline LDL-C >100 mg/dL [61,62]. Furthermore, in an analysis pooling total recurrent incident events (instead of only considering first events), demonstrated even greater clinical benefit (i.e., ~2-fold reduction in total CVD events) with alirocumab versus placebo [63].

FOURIER evaluated evolocumab, another monoclonal antibody targeting PCSK9, in patients with prior myocardial infarction, stroke or symptomatic peripheral arterial disease on statin therapy but with LDL-C >70 mg/dl. Primary outcome was a composite of cardiovascular death, MI, stroke, hospitalisation for unstable angina or coronary revascularisation. This was seen in 9.8% of treatment group, and 11.3% of the placebo group (ARR 2%, $p < 0.0001$). Absolute benefit was greater in patients with significant residual CAD, recent MI and multiple prior MIs, and with higher baseline high sensitivity C-reactive protein (hs-CRP) [64,65].

In the 2018 ACC/AHA Cholesterol Guidelines, PCSK9 monoclonal antibodies are recommended to achieve LDL-C >50% lowering to <100 mg/dl when “very high risk” ASCVD is present [66].

PCSK9 siRNA

Another avenue for manipulation of the PCSK9 pathway is via small interfering RNA molecules, notably inclisiran. ORION-1 was a Phase 2 trial that evaluated inclisiran dosed bi-annually in patients with LDL-C levels above target despite statin therapy, of high CVD risk. The average reductions in LDL-C were significantly greater after a single dose of inclisiran (27.9–41.9% reduction) than with placebo (2.1% increase) [67]. More recent studies demonstrated the safety and durability of effect in patients with type 2 diabetes [68].

PCSK9 gene editing

In recent years, it has been demonstrated that clustered regularly interspaced short palindromic repeats (CRISPR)/CRISPR-associated (Cas) system mediated genetic editing effectively and efficiently induced loss-of-function mutations in endogenous PCSK9. In one study, there was response within days to administration of CRISPR-Pcsk9 adenoviruses in hepatocytes, with resultant 35%–40% reduction in total cholesterol compared to control group. This reduction in LDL-C was comparable to rates seen in PCSK9 knockout mice. [69,70]

PCSK9 vaccination

Another novel method of PCSK9 modulation is via PCSK9 antibody formation. A recent study in a male, ApoE-deficient mouse model demonstrated durable formation of anti-PCSK9 antibodies, resulting in decreased PCSK9-LDLR interactions and resultant lower plasma total cholesterol, VLDL-cholesterol and triglycerides. However, there was less effect on LDL-C levels. [71]

Bempedoic acid

Bempedoic acid is a first-in-class small molecule adenosine triphosphate (ATP)-citrate lyase inhibitor that acts upstream of 3-hydroxy-3-methylglutaryl-coenzyme A (HMG-CoA) reductase to inhibit cholesterol biosynthesis and increase **LDL receptor** expression. Phase II trials had demonstrated its tolerability and efficacy in lowering LDL-C and hs-CRP levels [72–75]. CLEAR Tranquillity assessed bempedoic acid with ezetimibe in CVD patients intolerant of high intensity statins. Results demonstrated that bempedoic acid with ezetimibe decreased mean LDL-cholesterol concentration by –28.5% (95% CI: –34.4%, –22.5%; $p<0.001$) compared to ezetimibe alone at 12 weeks [76].

TRIGLYCERIDES

Epidemiological studies have suggested that triglycerides (TG) are associated with residual CVD events independent of achieved LDL-C concentrations. Elevated TG levels are also elevated levels of remnant circulating cholesterol particles, which are also present in triglyceride-rich lipoproteins. Many epidemiological cohort studies support TG as an independent risk factor for ASCVD, and recent meta-analysis suggest dose dependent increases in CVD risk and mortality with increasing TG levels. In addition, genetic analyses have suggested a causal link between TG-rich lipoprotein concentrations and CVD risk [77–80]. Recent genetic analyses suggest that the correlation of TG-rich lipoprotein with CAD risk may be explained by apolipoprotein B, the primary determinant of atherogenic lipoproteins including LDL [81,82].

Eicosapentaenoic acid (EPA)

Given the plausibility that omega-3 fatty acids may mediate some of the CVD risk reduction with consumption of fatty fish, a number of studies have evaluated the relationship between pharmacologic omega-3 fatty acids and CVD risk. Omega-3 fatty acid supplements are primarily comprised of eicosapentaenoic acid (EPA) and docosahexaenoic acid (DHA), but have largely not shown an association of CVD risk reduction [83].

The JELIS trial examined high-dose EPA with statin against statin monotherapy and demonstrated an ARR of 0.7% and RRR of 19% in major coronary events, with effect was significant only in those with baseline CAD [84]. This was further supported by the REDUCE-IT trial, which assessed reduction in MACE with supplementation of 4g/day EPA. It showed an ARR of 6.2% and RRR of 25% in primary outcome of MACE for the secondary prevention cohort [85]. Whether the apparent treatment effect for EPA is explained only by reduction in TG-rich lipoproteins or by other potential cardioprotective pleiotropic effects requires further study.

The pending STRENGTH trial will examine whether a proprietary blend of omega-3 carboxylic acids in high-risk primary and secondary prevention will lower CVD event risk [86].

LIPOPROTEIN(a)

Lipoprotein(a) (Lp(a)) consists of an apolipoprotein B-containing LDL-like particle linked to a plasminogen-like glycoprotein apo(a). Observational studies showed that Lp(a) levels are associated with higher risk of incident cardiovascular disease in primary prevention settings [87,88].

AIM-HIGH, JUPITER and LIPID trials all demonstrated that baseline Lp(a) levels were significantly associated with risk of future CAD events. A meta-analysis showed that those with Lp(a) levels in the highest quintile increased risk of MACE by 40% (OR 1.4; 95% CI 1.15–1.71, $p=0.001$). This analysis suggested the risk only remained significant with baseline elevated LDL-C ($>130\text{mg/dL}$) [89–93]. However, genome-wide association studies have shown *LPA* genetic variants are associated with recurrent CVD risk, independent of levels of LDL-C [94].

A recent genetic analysis assessed by how much Lp(a) would theoretically need to be reduced in order to produce a meaningful level of risk reduction. It suggested a reduction of 102 mg/dL would achieve similar levels of risk reduction as reducing LDL-C by 39 mg/dL. It also suggested reductions of 50mg/dL and greater were associated with $>10\%$ estimated risk reductions in the short term [95].

One issue with including Lp(a) in risk reducing strategies is the current lack of direct therapies. Lp(a) levels are variably affected by other lipid-lowering interventions. Indeed, high dose, high potency statins have been associated with increased Lp(a) levels [96,97]. Other interventions have been shown to reduce Lp(a) levels, such as niacin, IL-6 antagonists and PCSK9 inhibitors [98–100].

Therapies that are under investigation as directed therapies to reduce Lp(a) include antisense nucleotides to apo(a). IONIS-APO(a)Rx phase I and II trials have investigated antisense oligonucleotides, have suggested reductions in circulating concentration of Lp(a) of up to 90% with no apparent acute side effects. Phase III trials evaluating the effect of this reduction on cardiovascular outcomes are awaited [101,102].

DIABETES MELLITUS

Diabetes mellitus is one of the most well-established CVD risk factors. However, the DCCT trial demonstrated that intensive blood glucose control did not reduce risk of macrovascular complications. The ABCD and UKPDS studies some years later corroborated this, and did not detect significant changes related to cardiovascular mortality with improvements in glycemic control [103,104].

Since data suggested that use of rosiglitazone may increase CV risk, cardiovascular outcomes trials (CVOTs) have been required for all new glucose-lowering agents. Recent CVOTs that demonstrated safety but not protective benefit include SAVOR-TIMI 53 (saxagliptin), EXAMINE (alogliptin), EXSCEL (exanetide), ELIXA (lixisenatide) and TOSCA.IT (pioglitazone plus sulfonylureas) [105–110].

There are now five CVOTs to-date that, however, demonstrated benefit in reducing CV risk in diabetic patients. These include SUSTAIN-6 and LEADER, which evaluated semaglutide and liraglutide of the GLP-1 agonist class, and EMPA-REG OUTCOME, CANVAS and DECLARE-TIMI 58, which evaluated empagliflozin, canagliflozin and dapagliflozin of the SGLT2 inhibitor class [111–116].

GLP-1 receptor agonists (GLP1-RA)

SUSTAIN-6 CVOT examined semaglutide. Overall, 60% of participants had cardiovascular disease at baseline. After a median of 2.1 years, primary outcome of MACE occurred in 6.6% of the semaglutide group and 8.9% in the placebo group (HR, 0.74; 95% CI, 0.58 to 0.95). It did not demonstrate benefit in reducing risk of cardiovascular death or nonfatal MI [111].

The LEADER CVOT examined liraglutide. It demonstrated benefit in reducing rates of MACE (13.0% vs. 14.9%; HR 0.87, 95% CI 0.78–0.97). In addition, there were significantly lower rates of cardiovascular and all cause death [112].

SGLT2 Inhibitors (SGLT2i)

EMPA-REG OUTCOMES was the first CVOT to demonstrate cardiovascular benefit with glucose-lowering agents. It evaluated empagliflozin. Over 75% of patients had established CAD, with ~50% having prior MI. Rates of MACE were significantly lower in the treatment group (10.5% versus 12.1%; HR 0.86, 95% confidence interval 0.74–0.99). There was also significant difference in the rates of hospitalization for heart failure and all-cause mortality. The effect of the empagliflozin was seen early on Kaplan-Meier event curves [113].

The CANVAS CVOTs assessed canagliflozin. They found a significant reduction in MACE in treatment group compared to placebo (26.9 vs. 31.5 participants with an event per 1000 patient years; HR 0.86; 95% CI, 0.75 to 0.97; $P=0.02$ for superiority). Rates of hospitalisation for heart failure were again significantly reduced. However, they noted an increased risk of fracture (15.4 vs. 11.9 events per 1000 patient years; HR, 1.26; 95% CI, 1.04 to 1.52) and lower limb amputation (6.3 vs. 3.4 events per 1000 patient years, HR 1.97; 95% CI, 1.41 to 2.75) in the treatment group [114, 115].

DECLARE-TIMI 58 assessed cardiovascular outcomes for dapagliflozin. It did not find a significant difference in MACE rate in the general cohort (8.8% vs. 9.4% in placebo; HR, 0.93; 95% CI, 0.84–1.03; $P=0.17$), or in those with prior CAD. However, it did corroborate lower risk of hospitalisation for heart failure. It did not find significant increases in rates of lower limb amputation or fracture [116].

A 2018 ACC Expert Consensus document recommends initiation of either of a GLP-1 receptor agonist or SGLT-2 inhibitor in patients with type 2 diabetes mellitus, in addition to metformin and other risk factor modification strategies. Liraglutide is the preferred agent of the GLP-1 RA class, and empagliflozin is the preferred agent of the SGLT2i class. [117]

HIGH-DENSITY LIPOPROTEIN (HDL) CHOLESTEROL

Epidemiologic data suggests HDL-C concentrations are inversely related to cardiovascular risk. However, recent analyses demonstrated that increases in HDL-C caused by genetic polymorphisms do not confer a risk advantage in preventing future myocardial infarction, implying lack of a causal relationship [118].

Previously, AIM-HIGH assessed addition of niacin to a high intensity statin regime. It demonstrated no additional risk reduction benefit despite significant increases in HDL-C [119]. Later, HPS2-THRIVE assessed the risk reduction effects of niacin in a heterogeneous cohort of high-risk patients with CVD. Again, there was no risk reduction benefit seen despite increased serum HDL-C [120]. Additional evidence against HDL-C in preventing cardiovascular events was suggested from dalOUTCOMES trial, which assessed the cholesterol ester transfer protein (CETP) inhibitor, dalcetrapib. Despite a 31% increase in HDL-cholesterol levels in their treatment group with no difference seen in rates of the primary endpoint of MACE [121].

ACCELERATE assessed another CETP inhibitor, evacetrapib, in patients with recent ACS, known ASCVD or diabetes mellitus. Despite a marked increase in HDL-C (133.2% vs. 1.6%), there was no difference in rates of primary endpoint of MACE (12.9% vs. 12.8%; HR 1.01; 95% CI, 0.91 to 1.11; P=0.91) [122].

The REVEAL trial assessed anacetrapib in patients with known ASCVD on high-intensity statins. Mean HDL-C levels increased in the treatment group by 43 mg/dL (104% vs. placebo), and non-HDL-C notably also declined by 17mg/dL (-18% vs. placebo). Median follow up was 4.1 years. There was ARR of 1% in first occurrence of MACE (10.8% vs. 11.8%; RR, 0.91; 95% CI 0.85–0.97). Overall, it is thought the trial effect is driven by the reduction in non-HDL-C rather than increase in HDL-C [123].

Whether other properties of HDL, such as reverse cholesterol transport, may influence ASCVD risk requires further study. Observational data in patients without cardiovascular disease demonstrated that those with the highest quartile of HDL cholesterol efflux capacity (a measure of reverse cholesterol transport) had a 67% reduction in composite ASCVD rates compared to the lowest quartile at 9.7 years of follow up [124].

PLATELETS AND THROMBOSIS

The final common pathway for a CAD event is coagulation and thrombosis. The anti-platelet agent aspirin is known to reduce mortality in acute coronary syndromes [125]. Dual anti-platelet therapy, combining aspirin with P2Y₁₂ inhibitors is critical in maintaining stent patency after PCI. Recently, the DAPT and PEGASUS TIMI-54 trials have demonstrated longer durations of DAPT beyond 1 year of treatment reduce risk of stent thrombosis and MACE [126]. In PEGASUS TIMI-54, there was a slight but significant increase in the rates of major bleeding only in the higher dose ticagrelor group [127]. More recent follow up analyses in of this trial have demonstrated this benefit persists at 5 years in the trial setting, with slight attenuation of bleeding risk as time progressed [128]. Scores, such as the DAPT score, may identify candidates for a more prolonged duration of P2Y₁₂ inhibition [129].

Oral vitamin K antagonists (VKORs) help reduce residual cardiovascular risk in patients with myocardial infarction, at a cost of increased bleeding rates. However, direct acting oral anticoagulants (DOACs) have been developed with reduced bleeding risk compared to VKORs. COMPASS investigated the benefit and safety of rivaroxaban 2.5 mg twice daily in secondary prevention. The rivaroxaban plus aspirin group demonstrated the lowest MACE event rate (4.1%) compared to rivaroxaban alone (4.9%) and aspirin alone (5.4%). Rivaroxaban plus aspirin compared to aspirin alone had a HR of 0.76 (95% CI 0.66–0.86; $p < 0.001$). Major bleeding occurred more in the combination group than with aspirin alone, but there were no significant difference in rates of fatal bleeding [130]. Recently, the FDA approved the use rivaroxaban 2.5mg twice daily in adjunct to aspirin for secondary prevention; this strategy has not yet been incorporated into professional guidelines [131].

INFLAMMATION

Manipulating inflammation to reduce ASCVD risk has been proposed based on epidemiologic and basic biomedical investigations [132]. Elevated concentrations of hs-CRP, an inflammatory biomarker, is associated with elevated CVD risk independent of established cardiovascular risk factors [133].

Recently, the CANTOS trial tested the clinical efficacy of this strategy in humans using canakinumab, a monoclonal antibody targeting interleukin-1 beta (IL-1B). IL-1B is a central cytokine in the interleukin-6 (IL-6) pathway, which has been suggested by genetic analyses to be causative in atherosclerosis, and high levels are associated with increased risk of cardiovascular events [134–136].

CANTOS evaluated patients with a history of prior MI and hs-CRP level of $> 2 \text{ mg/dL}$. All treatment groups experienced significant reductions in hs-CRP levels. After a median of 3.7 years, there was a significant difference in the rates of primary endpoint of MACE in the 150mg (HR 0.85, 95% CI 0.74–0.98; $p = 0.021$) and 300mg groups (HR 0.86, 95% CI 0.75–0.99; $p = 0.031$). There was an increased rate of deaths related to infection in the pooled treatment groups [137]. However, FDA recently indicated that current evidence for canakinumab is insufficient to support a label for CVD risk reduction [138]. The CIRT trial did not observe an improvement in CVD risk among a similar population with low-dose methotrexate [139]. These data indicate that specific inflammatory pathways, such as IL-1B/IL-6, may be more favourable for risk modulation than others.

Conclusion:

Despite increasingly durable and intensive LDL-C-lowering therapies, significant residual risk remains after people experience cardiovascular events. Strategies to improve the delivery and adherence of lifestyle modification have an opportunity to address this risk. Additionally, orthogonal strategies to lower recurrent CVD risk among key subgroups can help facilitate tailored therapies to optimize medical strategies to reduce recurrent CVD risk.

Funding

P.N. is supported by a grant from the National Heart, Lung, and Blood Institute (K08 HL140203) and Hassenfeld Scholar Award from the Massachusetts General Hospital.

References:

1. Global Health Estimates 2016: Disease burden by Cause, Age, Sex, by Country and by Region, 2000–2016. Geneva, World Health Organization; 2018.
2. American Heart Association (AHA). Heart Disease and Stroke Statistics-2019 Update: A Report From the American Heart Association. American Heart Association Council on Epidemiology and Prevention Statistics Committee and Stroke Statistics Subcommittee. *Circulation*. 2019 131:10.1161/CIR.0000000000000659. [Epub ahead of print]
3. Yadlowsky S, Hayward RA, Sussman JB, McClelland RL, Min Y-I, Basu S. Clinical Implications of Revised Pooled Cohort Equations for Estimating Atherosclerotic Cardiovascular Disease Risk. *Annals of Internal Medicine*. 2018;169(1):20–29. doi:10.7326/M17-3011 [PubMed: 29868850]
4. Bohula Erin A., Bonaca Marc P., Braunwald Eugene, et al. Atherothrombotic Risk Stratification and the Efficacy and Safety of Vorapaxar in Patients With Stable Ischemic Heart Disease and Previous Myocardial Infarction. *Circulation*. 2016;134(4):304–313. doi:10.1161/CIRCULATIONAHA.115.019861 [PubMed: 27440003] *Established a new predictive clinical calculator for estimation of residual risk
5. Bohula EA, Morrow DA, Giugliano RP, et al. Atherothrombotic Risk Stratification and Ezetimibe for Secondary Prevention. *Journal of the American College of Cardiology*. 2017;69(8):911–921. doi:10.1016/j.jacc.2016.11.070 [PubMed: 28231942]
6. Williams BA, Chagin KM, Bash LD, et al. External validation of the TIMI risk score for secondary cardiovascular events among patients with recent myocardial infarction. *Atherosclerosis*. 2018;272:80–86. doi:10.1016/j.atherosclerosis.2018.03.026 [PubMed: 29579671]
7. Yejin Mok, Ballew Shoshana H., Bash Lori D., et al. International Validation of the Thrombolysis in Myocardial Infarction (TIMI) Risk Score for Secondary Prevention in Post-MI Patients: A Collaborative Analysis of the Chronic Kidney Disease Prognosis Consortium and the Risk Validation Scientific Committee. *Journal of the American Heart Association*. 2018;7(14):e008426. doi:10.1161/JAHA.117.008426 [PubMed: 29982232]
8. Dehghan M, Mente A, Teo KK, Gao P, Sleight P et al. Relationship Between Healthy Diet and Risk of Cardiovascular Disease Among Patients on Drug Therapies for Secondary Prevention: A Prospective Cohort Study of 31 546 High-Risk Individuals From 40 Countries. *Circulation*. 2012;126:2705–2712 [PubMed: 23212996]
9. Yu E, Malik VS, Hu FB. Cardiovascular Disease Prevention by Diet Modification: JACC Health Promotion Series. *Journal of the American College of Cardiology*. 2018;72(8):914–926. doi: 10.1016/j.jacc.2018.02.085 [PubMed: 30115231]
10. American College of Cardiology/American Heart Association. 2013 AHA/ACC Guideline on Lifestyle Management to Reduce Cardiovascular Risk - A Report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. *Circulation*. 2014 6 24;129(25 Suppl 2):S76–99. doi: 10.1161/01.cir.0000437740.48606.d1. Epub 2013 Nov 12. [PubMed: 24222015]
11. Burr ML, Fehily AM, Gilbert JF, Rogers S, Holliday RM. Effects of changes in fat, fish, and fibre intakes on death and myocardial reinfarction: diet and reinfarction trial (DART). *Lancet*. 1989;2(8666):757–61. [PubMed: 2571009]
12. Burr ML. Secondary prevention of CHD in UK men: the Diet and Reinfarction Trial and its sequel. *Proc Nutr Soc*. 2007 2;66(1):9–15. [PubMed: 17343767]
13. deLorgeril M, Renaud S, Salen P, Monjaud I, Mamelle N. Mediterranean alpha-linolenic acid-rich diet in secondary prevention of coronary heart disease. *Lancet*. 1994;343:8911:1454–1459 [PubMed: 7911176]
14. Estruch R, Ros E, Salas-Salvadó J, et al. Primary Prevention of Cardiovascular Disease with a Mediterranean Diet Supplemented with Extra-Virgin Olive Oil or Nuts. *N Engl J Med*. 2018 6 21;378(25):e34. [PubMed: 29897866]
15. Itsiopoulos C, Kucianski T, Mayr HL, et al. The AUStralian MEDiterranean Diet Heart Trial (AUSMED Heart Trial): A randomized clinical trial in secondary prevention of coronary heart disease in a multiethnic Australian population: Study protocol. *American Heart Journal*. 2018. doi: 10.1016/j.ahj.2018.05.010.

16. Wei M, Brandhorst S, Shelehchi M, et al. Fasting-mimicking diet and markers/risk factors for aging, diabetes, cancer, and cardiovascular disease. *Science Translational Medicine*. 2017;9(377).
17. Crowe FL, Appleby PN, Travis RC, Key TJ. Risk of hospitalization or death from ischemic heart disease among British vegetarians and nonvegetarians: results from the EPIC-Oxford cohort study. *Am J Clin Nutr*. 2013;97(3):597–603. doi:10.3945/ajcn.112.044073 [PubMed: 23364007]
18. Tonstad S, Butler T, Yan R, Fraser GE. Type of Vegetarian Diet, Body Weight, and Prevalence of Type 2 Diabetes. *Diabetes Care*. 2009;32(5):791–796. doi:10.2337/dc08-1886 [PubMed: 19351712]
19. Rizzo NS, Sabaté J, Jaceldo-Siegl K, Fraser GE. Vegetarian dietary patterns are associated with a lower risk of metabolic syndrome: the adventist health study 2. *Diabetes Care*. 2011;34(5):1225–1227. doi:10.2337/dc10-1221 [PubMed: 21411506]
20. Yokoyama Y, Nishimura K, Barnard ND, et al. Vegetarian diets and blood pressure: a meta-analysis. *JAMA Intern Med*. 2014;174(4):577–587. doi:10.1001/jamainternmed.2013.14547 [PubMed: 24566947]
21. Barnard ND, Levin SM, Yokoyama Y. A systematic review and meta-analysis of changes in body weight in clinical trials of vegetarian diets. *J Acad Nutr Diet*. 2015;115(6):954–969. doi:10.1016/j.jand.2014.11.016 [PubMed: 25620754]
22. Ornish D, Scherwitz LW, Doody RS, et al. Effects of stress management training and dietary changes in treating ischemic heart disease. *JAMA*. 1983;249(1):54–59. [PubMed: 6336794]
23. Esselstyn CB, Ellis SG, Medendorp SV, Crowe TD. A strategy to arrest and reverse coronary artery disease: a 5-year longitudinal study of a single physician's practice. *J Fam Pract*. 1995;41(6):560–568. [PubMed: 7500065]
24. Ornish D, Scherwitz LW, Billings JH, et al. Intensive lifestyle changes for reversal of coronary heart disease. *JAMA*. 1998;280(23):2001–2007. [PubMed: 9863851]
25. Ornish D, Brown SE, Scherwitz LW, et al. Can lifestyle changes reverse coronary heart disease? The Lifestyle Heart Trial. *Lancet*. 1990;336(8708):129–133. [PubMed: 1973470]
26. Gould KL, Ornish D, Scherwitz L, et al. Changes in myocardial perfusion abnormalities by positron emission tomography after long-term, intense risk factor modification. *JAMA*. 1995;274(11):894–901. [PubMed: 7674504]
27. Esselstyn CB. Updating a 12-year experience with arrest and reversal therapy for coronary heart disease (an overdue requiem for palliative cardiology). *Am J Cardiol*. 1999;84(3):339–341, A8. [PubMed: 10496449]
28. Goel K, Lennon RJ, Tilbury RT, et al. Impact of cardiac rehabilitation on mortality and cardiovascular events after percutaneous coronary intervention in the community. *Circulation*. 2011 5 31;123(21):2344–52. [PubMed: 21576654]
29. Dunlay SM, Pack QR, Thomas RJ, Killian JM, Roger VL. Participation in cardiac rehabilitation, readmissions, and death after acute myocardial infarction. *Am J Med*. 2014;127(6):538–46. [PubMed: 24556195]
30. Kotseva K et al. 2012. Use and effects of cardiac rehabilitation in patients with coronary heart disease: results from the EUROASPIRE III survey. *Eur J Prev Cardiol*. 2012;20(5), 817–826. [PubMed: 22718794]
31. Gomadam PS, Douglas CJ, Sacrinty MT. Degree and Direction of Change of Body Weight in Cardiac Rehabilitation and Impact on Exercise Capacity and Cardiac Risk Factors. *Am J Cardiol*. 2016 2 15;117(4):580–584. [PubMed: 26768674]
32. Anderson L, Thompson DR, Oldridge N, Zwisler A, Rees K, Martin N, Taylor RS. Exercise-based cardiac rehabilitation for coronary heart disease. *Cochrane Database of Systematic Reviews* 2016, 10.1002/14651858.CD001800.pub3
33. Kraus WE, Houmard JA, Duscha BD. Effects of the amount and intensity of exercise on plasma lipoproteins. *N Engl J Med*. 2002; 347(19) 1483–1492. [PubMed: 12421890]
34. Sarzynski MA, Ruiz-Ramie JJ, Barber JL, Slentz CA, Apolzan JW, McGarrah RW, et al. Effects of increasing exercise intensity and dose on multiple measures of HDL (High-Density Lipoprotein) function. *Arterioscler Thromb Vasc Biol*. 2018;38:943–52 [PubMed: 29437573]

35. Moholdt T, Lavie CJ, Nauman J. Sustained physical activity, not weight loss, associated with improved survival in coronary heart disease. *J Am Coll Cardiol*. 2018; 71:1094–101 [PubMed: 29519349]
36. Piercy KL, Troiano RP, Ballard RM, et al. The Physical Activity Guidelines for Americans. *JAMA*. 2018;320(19):2020–2028. doi:10.1001/jama.2018.14854 [PubMed: 30418471]
37. U.S. Department of Health, Education and Welfare. Smoking and Health: Report of the Advisory Committee to the Surgeon General of the Public Health Service. Washington D.C.: U.S. Department of Health, Education and Welfare; 1964
38. U.S. Department of Health and Human Services. The Reports of the Surgeon General: The Health Consequences of Smoking: Cardiovascular Disease. Washington D.C.: Department of Health and Human Services; 1983
39. Centers for Disease Control and Prevention. Current Cigarette Smoking Among Adults—United States, 2016. *Morbidity and Mortality Weekly Report* 2018;67(2):53–9 [PubMed: 29346338]
40. Burke GM, Genuardi M, Shappell H, D'Agostino RB, Magnani JW. Temporal Associations Between Smoking and Cardiovascular Disease, 1971 to 2006 (from the Framingham Heart Study). *American Journal of Cardiology*. 2017;(10):1787. [PubMed: 28865894]
41. Critchley JA, Capewell S. Mortality risk reduction associated with smoking cessation in patients with coronary heart disease: a systematic review. *JAMA*. 2003;290(1):86–97. doi:10.1001/jama.290.1.86 [PubMed: 12837716]
42. Iestra JA, Kromhout D, van der Schouw YT, Grobbee DE, Boshuizen HC, van Staveren WA. Effect Size Estimates of Lifestyle and Dietary Changes on All-Cause Mortality in Coronary Artery Disease Patients: A Systematic Review. *Circulation*. 2005;112:924–934 [PubMed: 16087812]
43. Chow CK, Jolly S, Rao-Melacini P, Fox KA, Anand SS, Yusuf S. Association of diet, exercise, and smoking modification with risk of early cardiovascular events after acute coronary syndromes. *Circulation*. 2010 2 16;121(6):750–8 [PubMed: 20124123]
44. Hajar R Risk Factors for Coronary Artery Disease: Historical Perspectives. *Heart Views* 2017;18(3):109–114. [PubMed: 29184622]
45. Gofman JW, Young W, Tandy R. Ischemic heart disease, atherosclerosis, and longevity. *Circulation*. 1966;34:679–97 [PubMed: 5921763]
46. Kannel WB, Castelli WP, Gordon T. Cholesterol in the prediction of atherosclerotic disease. New perspectives based on the Framingham study. *Ann Intern Med*. 1979;90:85–91 [PubMed: 217290]
47. Klag MJ, Ford DE, Mead LA, He J, Whelton PK, Liang KY, et al. Serum cholesterol in young men and subsequent cardiovascular disease. *N Engl J Med*. 1993;328:313–8 [PubMed: 8419817]
48. Ip S, Lichtenstein AH, Chung M, Lau J, Balk EM. Systematic review: association of low-density lipoprotein subfractions with cardiovascular outcomes. *Ann Intern Med*. 2009;150:474–84 [PubMed: 19349632]
49. Shiffman D, Louie JZ, Caulfield MP, Nilsson PM, Devlin JJ, Melander O. LDL subfractions are associated with incident cardiovascular disease in the Malmö Prevention Project Study. *Atherosclerosis*. 2017 8;263:287–292 [PubMed: 28728064]
50. Otvos JD, Collins D, Freedman DS, et al. Low-density lipoprotein and high-density lipoprotein particle subclasses predict coronary events and are favorably changed by gemfibrozil therapy in the Veterans Affairs High-Density Lipoprotein Intervention Trial. *Circulation*. 2006;113(12):1556–1563. doi:10.1161/CIRCULATIONAHA.105.565135 [PubMed: 16534013]
51. Cromwell WC, Otvos JD, Keyes MJ, et al. LDL particle number and risk of future cardiovascular disease in the Framingham Offspring Study—Implications for LDL management. *Journal of Clinical Lipidology*. 2007;1(6):583–592. doi:10.1016/j.jacl.2007.10.001 [PubMed: 19657464]
52. Otvos JD, Mora S, Shalaurova I, Greenland P, Mackey RH, Goff DC. Clinical implications of discordance between low-density lipoprotein cholesterol and particle number. *Journal of Clinical Lipidology*. 2011;5(2):105–113. doi:10.1016/j.jacl.2011.02.001 [PubMed: 21392724]
53. Cromwell WC, Otvos JD. Heterogeneity of Low-Density Lipoprotein Particle Number in Patients With Type 2 Diabetes Mellitus and Low-Density Lipoprotein Cholesterol <100 mg/dl. *American Journal of Cardiology*. 2006;98(12):1599–1602. doi:10.1016/j.amjcard.2006.07.036 [PubMed: 17145217]

54. Carmena R, Duriez P, Fruchart J-C. Atherogenic lipoprotein particles in atherosclerosis. *Circulation*. 2004;109(23 Suppl 1):III2–7. doi:10.1161/01.CIR.0000131511.50734.44 [PubMed: 15198959]
55. Ai M, Otokozawa S, Asztalos BF, et al. Small dense LDL cholesterol and coronary heart disease: results from the Framingham Offspring Study. *Clin Chem*. 2010;56(6):967–976. doi:10.1373/clinchem.2009.137489 [PubMed: 20431054]
56. Schwartz GG, Olsson AG, Ezekowitz MD, Ganz P, Oliver MF et al. Effects of atorvastatin on early recurrent ischemic events in acute coronary syndromes: the MIRACL study: a randomized controlled trial. *JAMA*. 2001 4 4;285(13):1711–8. [PubMed: 11277825]
57. Cannon CP, Braunwald E, McCabe CH, Rader DJ, Rouleau JL et al. Intensive versus moderate lipid lowering with statins after acute coronary syndromes. *N Engl J Med*. 2004 4 8;350(15):1495–504. [PubMed: 15007110]
58. Pedersen TR, Faergeman O, Kastelein JJ, Olsson AG, Tikkanen MJ et al. High-dose atorvastatin vs usual-dose simvastatin for secondary prevention after myocardial infarction: the IDEAL study: a randomized controlled trial. *JAMA*. 2005 11 16;294(19):2437–45. [PubMed: 16287954]
59. Cannon CP, Blazing MA, Giugliano RP, McCagg A, White JA, et al. Ezetimibe Added to Statin Therapy after Acute Coronary Syndromes. *N Eng J Med*; 2015 372(25);2387–2397**Established risk reduction benefit of ezetimibe in combination with statin therapy
60. Giugliano RP, Cannon CP, Blazing MA, Nicolau JC, Corbalan R et al. Benefit of Adding Ezetimibe to Statin Therapy on Cardiovascular Outcomes and Safety in Patients With Versus Without Diabetes Mellitus: Results from IMPROVE-IT (Improved Reduction of Outcomes: Vytorin Efficacy International Trial). *Circulation*; 2018; 137:1571–1582 [PubMed: 29263150]
61. Schwartz GG, Szarek M, Bhatt DL, Bittner V, Diaz R, Edelberg J, Goodman GG, Hanotin C, Harrington R, Jukema JW, Lecorps G, Moryusef A, Pordy R, Roe MT, White HD, Zeiher A, Steg G. Evaluation of Cardiovascular Outcomes After an Acute Coronary Syndrome During Treatment With Alirocumab - ODYSSEY OUTCOMES. *American College of Cardiology - 67th Scientific Sessions*; 2018; Orlando
62. Schwartz GG, Steg G, Szarek M, Bhatt DL et al. Alirocumab and Cardiovascular Outcomes after Acute Coronary Syndrome. *New Eng J Med*, 2018 DOI: 10.1056/NEJMoa1801174
63. Szarek M, White HD, Schwartz GG, et al. Alirocumab Reduces Total Nonfatal Cardiovascular and Fatal Events: The ODYSSEY OUTCOMES Trial. *Journal of the American College of Cardiology*. 2019;73(4):387–396. doi:10.1016/j.jacc.2018.10.039 [PubMed: 30428396] *Later data from ODYSSEY OUTCOMES cohort demonstrating even greater impact in reducing cardiovascular events, in particular related to recurrent events
64. Sabatine MS, Giugliano RP, Keech AC, et al., on behalf of the FOURIER Steering Committee and Investigators. Evolocumab and Clinical Outcomes in Patients with Cardiovascular Disease. *N Engl J Med* 2017;376:1713–22. [PubMed: 28304224]
65. Bohula EA, Giugliano RP, Leiter LA, et al. Inflammatory and Cholesterol Risk in the FOURIER Trial (Further Cardiovascular Outcomes Research With PCSK9 Inhibition in Patients With Elevated Risk). *Circulation* 2018;3 12:[Epub ahead of print].
66. Grundy SM, Stone NJ, Bailey AL, et al. 2018 AHA/ACC/AACVPR/AAPA/ABC/ACPM/ADA/AGS/APhA/ASPC/NLA/PCNA Guideline on the Management of Blood Cholesterol: Executive Summary: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *Journal of the American College of Cardiology*. 11 2018. doi:10.1016/j.jacc.2018.11.002
67. Ray KK, Landmesser U, Leiter LA, Kallend D, Dufour R, Karakas M, et al. Inclisiran in patients at high cardiovascular risk with elevated LDL cholesterol. *N Engl J Med*. 2017;376(15):1430–40. [PubMed: 28306389]
68. Leiter L, Teoh H, Kallend D, Wright RS, Landmesser U, Wijngaard PL, Kastelein JJ, Ray KK. Inclisiran Lowers LDL-C and PCSK9 Irrespective of Diabetes Status without Worsening Glycemia. *Diabetes*. 2018; 67(Supplement 1). 10.2337/db18-337-OR. American Diabetes Association 2018.
69. Ding Q, Strong A, Patel KM, et al. Permanent alteration of PCSK9 with in vivo CRISPR-Cas9 genome editing. *Circ Res*. 2014;115(5):488–92. [PubMed: 24916110]

70. Rashid S, Curtis DE, Garuti R, et al. Decreased plasma cholesterol and hypersensitivity to statins in mice lacking Pcsk9. *Proc Natl Acad Sci USA*. 2005;102(15):5374–9. [PubMed: 15805190]
71. Kawakami R, Nozato Y, Nakagami H, et al. Development of vaccine for dyslipidemia targeted to a proprotein convertase subtilisin/kexin type 9 (PCSK9) epitope in mice. *PLoS One*. 2018;13(2):e0191895 10.1371/journal.pone.0191895 [PubMed: 29438441]
72. Ballantyne CM, Davidson MH, Macdougall DE, Bays HE, Dicarlo LA, et al. Efficacy and safety of a novel dual modulator of adenosine triphosphate-citrate lyase and adenosine monophosphate-activated protein kinase in patients with hypercholesterolemia: results of a multicenter, randomized, double-blind, placebo-controlled, parallel-group trial. *J Am Coll Cardiol*, 62 (2013), pp. 1154–1162 [PubMed: 23770179]
73. Gutierrez MJ, Rosenberg NL, Macdougall DE, Hanselman JC, Margulies JR, et al. Efficacy and safety of ETC-1002, a novel investigational low-density lipoprotein-cholesterol-lowering therapy for the treatment of patients with hypercholesterolemia and type 2 diabetes mellitus. *Arterioscler. Thromb. Vasc. Biol*, 34 (2014), pp. 676–683 [PubMed: 24385236]
74. Thompson PD, Rubino J, Janik MJ, MacDougall DE, McBride SJ, et al. Use of ETC-1002 to treat hypercholesterolemia in patients with statin intolerance. *J Clin Lipidol*, 9 (2015), pp. 295–304 [PubMed: 26073387]
75. Thompson PD, MacDougall DE, Newton RS, Margulies JR, Hanselman JC, et al. Treatment with ETC-1002 alone and in combination with ezetimibe lowers LDL cholesterol in hypercholesterolemic patients with or without statin intolerance. *J Clin Lipidol*, 10 (2016), pp. 556–567 [PubMed: 27206943]
76. Ballantyne CM, Banach M, Mancini GBJ, Lepor NE, Hanselman JC, Zhao X, Leiter LA. Efficacy and safety of bempedoic acid added to ezetimibe in statin-intolerant patients with hypercholesterolemia: A randomized, placebo-controlled study. *Atherosclerosis*. 2018 10;277:195–203. [PubMed: 29910030]
77. Liu J, Zeng FF, Liu ZM, Zhang CX, Ling WH et al. Effects of blood triglycerides on cardiovascular and all-cause mortality: a systematic review and meta-analysis of 61 prospective studies. *Lipids Health Dis*. 2013 10 29;12:159. [PubMed: 24164719]
78. Varbo A, Benn M, Tybjaerg-Hansen A, Jørgensen AB, Frikke-Schmidt R, Nordestgaard BG. Remnant cholesterol as a causal risk factor for ischemic heart disease. *J Am Coll Cardiol*. 2013;61:427–36 [PubMed: 23265341]
79. Triglyceride Coronary Disease Genetics Consortium and Emerging Risk Factors Collaboration. Sarwar N, Sandhu MS, Ricketts SL, Butterworth AS, Di Angelantonio E, et al. Triglyceride-mediated pathways and coronary disease: Collaborative analysis of 101 studies. *Lancet*. 2010;375:1634–9. [PubMed: 20452521]
80. Jørgensen AB, Frikke-Schmidt R, West AS, Grande P, Nordestgaard BG, Tybjaerg-Hansen A. Genetically elevated non-fasting triglycerides and calculated remnant cholesterol as causal risk factors for myocardial infarction. *Eur Heart J*. 2013;34:1826–33. [PubMed: 23248205]
81. Ference BA, Kastelein JJP, Ray KK, et al. Association of Triglyceride-Lowering LPL Variants and LDL-C-Lowering LDLR Variants With Risk of Coronary Heart Disease. *JAMA*. 2019;321(4):364–373. doi:10.1001/jama.2018.20045 [PubMed: 30694319]
82. Ference BA, Kastelein JJP, Ginsberg HN, et al. Association of Genetic Variants Related to CETP Inhibitors and Statins With Lipoprotein Levels and Cardiovascular Risk. *JAMA*. 2017;318(10): 947–956. doi:10.1001/jama.2017.11467 [PubMed: 28846118]
83. Abdelhamid AS, Brown TJ, Brainard JS, Biswas P, Thorpe GC, et al. Omega-3 fatty acids for the primary and secondary prevention of cardiovascular disease. *Cochrane Database of Systematic Reviews* 2018, 10.1002/14651858.CD003177.pub4.
84. Yokoyama M, Origasa H, Matsuzaki M, Matsuzawa Y, Saito Y et al. Effects of eicosapentaenoic acid on major coronary events in hypercholesterolaemic patients (JELIS): a randomised open-label, blinded endpoint analysis. *Lancet*. 2007 3 31;369(9567):1090–8. [PubMed: 17398308]
85. Bhatt DL, Steg PG, Miller M, et al. Cardiovascular Risk Reduction with Icosapent Ethyl for Hypertriglyceridemia. *New England Journal of Medicine*. 2019;380(1):11–22. doi:10.1056/NEJMoa1812792 [PubMed: 30415628] **Large-scale randomised clinical trial demonstrating clinical benefit in terms of reduction of cardiovascular events with high-dose EPA supplementation

86. Nicholls SJ, Lincoff AM, Bash D, Ballantyne CM, Barter PJ et al. Assessment of omega-3 carboxylic acids in statin-treated patients with high levels of triglycerides and low levels of high-density lipoprotein cholesterol: Rationale and design of the STRENGTH trial. *Clin Cardiol*. 2018 10;41(10):1281–1288. [PubMed: 30125052]
87. Kamstrup PR, Tybjaerg-Hansen A, Steffensen R, Nordestgaard BG. Genetically elevated lipoprotein(a) and increased risk of myocardial infarction. *JAMA*. 2009 6 10;301(22):2331–9 [PubMed: 19509380]
88. Emerging Risk Factors Collaboration, Erqou S, Kaptoge S, Perry PL, Di Angelantonio E, et al. Lipoprotein(a) concentration and the risk of coronary heart disease, stroke, and nonvascular mortality. *JAMA*. 2009 7 22;302(4):412–23 [PubMed: 19622820]
89. Puri R, Ballantyne CM, Hoogeveen RC, Shao M, Barter P, et al. Lipoprotein(a) and coronary atheroma progression rates during long-term high-intensity statin therapy: Insights from SATURN. *Atherosclerosis*. 2017 8;263:137–144 [PubMed: 28641153]
90. Albers JJ, Slee A, O'Brien KD, Robinson JG, Kashyap ML, et al. Relationship of apolipoproteins A-1 and B, and lipoprotein(a) to cardiovascular outcomes: the AIM-HIGH trial (Atherothrombosis Intervention in Metabolic Syndrome with Low HDL/High Triglyceride and Impact on Global Health Outcomes). *J Am Coll Cardiol*. 2013 10 22; 62(17):1575–9. [PubMed: 23973688]
91. Nestel PJ, Barnes EH, Tonkin AM, Simes J, Fournier M et al. Plasma lipoprotein(a) concentration predicts future coronary and cardiovascular events in patients with stable coronary heart disease. *Arterioscler Thromb Vasc Biol*. 2013 12;33(12):2902–8 [PubMed: 24092750]
92. Khera AV, Everett BM, Caulfield MP, Hantash FM, Wohlgemuth J, Ridker PM, Mora S. Lipoprotein(a) concentrations, rosuvastatin therapy, and residual vascular risk: an analysis from the JUPITER Trial (Justification for the Use of Statins in Prevention: an Intervention Trial Evaluating Rosuvastatin). *Circulation*. 2014 2 11; 129(6):635–42. [PubMed: 24243886]
93. O'Donoghue ML, Morrow DA, Tsimikas S, Sloan S, Ren AF et al. Lipoprotein(a) for risk assessment in patients with established coronary artery disease. *J Am Coll Cardiol*. 2014 2 18;63(6):520–7 [PubMed: 24161323]
94. Wei WQ, Li X, Feng Q, Kubo M, Kullo IJ et al. LPA Variants Are Associated With Residual Cardiovascular Risk in Patients Receiving Statins. *Circulation*. 2018 10 23;138(17):1839–1849. [PubMed: 29703846] *Described Lp(a) in terms of contribution to residual risk of patients already on directed medical therapy
95. Burgess S, Ference BA, Staley JR, Freitag DF, Mason AM, et al. Association of LPA Variants With Risk of Coronary Disease and the Implications for Lipoprotein(a)-Lowering Therapies: A Mendelian Randomization Analysis. *JAMA Cardiol*. 2018 7 1;3(7):619–627 [PubMed: 29926099] *Quantified risk associated with elevated Lp(a) in relation to LDL-C reductions
96. Murphy SA, Cannon CP, Wiviott SD, McCabe CH, Braunwald E. Reduction in recurrent cardiovascular events with intensive lipid-lowering statin therapy compared with moderate lipid-lowering statin therapy after acute coronary syndromes from the PROVE IT-TIMI 22 (Pravastatin or Atorvastatin Evaluation and Infection Therapy-Thrombolysis In Myocardial Infarction 22) trial. *J Am Coll Cardiol*. 2009 12 15;54(25):2358–62 [PubMed: 20082923]
97. Capoulade R, Chan KL, Yeang C, Mathieu P, Bossé Y et al. Oxidized Phospholipids, Lipoprotein(a), and Progression of Calcific Aortic Valve Stenosis. *J Am Coll Cardiol*. 2015 9 15;66(11):1236–1246 [PubMed: 26361154]
98. Chennamsetty I, Kostner KM, Claudel T, et al. Nicotinic acid inhibits hepatic APOA gene expression: studies in humans and in transgenic mice. *J Lipid Res*. 2012;53(11):2405–12. [PubMed: 22930813]
99. Müller N, Schulte DM, Türk K, et al. IL-6 blockade by monoclonal antibodies inhibits apolipoprotein (a) expression and lipoprotein (a) synthesis in humans. *J Lipid Res*. 2015;56(5): 1034–42. [PubMed: 25713100]
100. Raal FJ, Giugliano RP, Sabatine MS, Koren MJ, Langslet G et al. Reduction in lipoprotein(a) with PCSK9 monoclonal antibody evolocumab (AMG 145): a pooled analysis of more than 1,300 patients in 4 phase II trials. *J Am Coll Cardiol*. 2014 4 8;63(13):1278–1288 [PubMed: 24509273]
101. Tsimikas S, Viney NJ, Hughes SG, Singleton W, Graham MJ, et al. Antisense therapy targeting apolipoprotein(a): a randomised, double-blind, placebo-controlled phase 1 study. *Lancet* 2015;386:1472–1483. [PubMed: 26210642]

102. Viney NJ, Capelleveen JCV, Geary RS, Xia S, Tami JA, et al. Antisense oligonucleotides targeting apolipoprotein(a) in people with raised lipoprotein(a): two randomised, double-blind, placebo-controlled, dose-ranging trials. *Lancet* 2016;388:2239–2253. [PubMed: 27665230]
103. The Diabetes Control and Complications Trial Research Group. The effect of intensive treatment of diabetes on the development and progression of long-term complications in insulin-dependent diabetes mellitus. *N Engl J Med.* 1993;329:977–986 [PubMed: 8366922]
104. UK Prospective Diabetes Study (UKPDS) Group. Intensive blood-glucose control with sulphonylureas or insulin compared with conventional treatment and risk of complications in patients with type 2 diabetes (UKPDS 33) *Lancet.* 1998;352:837–853 [PubMed: 9742976]
105. Food and Drug Administration Guidance for industry: diabetes mellitus—evaluating cardiovascular risk in new antidiabetic therapies to treat type 2 diabetes. [Accessed Nov 8, 2018]. Available from: <https://www.fda.gov/downloads/drugs/guidancecomplianceregulatoryinformation/guidances/ucm071627.pdf>
106. Scirica BM, Bhatt DL, Braunwald E, et al. SAVOR-TIMI 53 Steering Committee and Investigators Saxagliptin and cardiovascular outcomes in patients with type 2 diabetes mellitus. *N Engl J Med.* 2013;369(14):1317–1326 [PubMed: 23992601]
107. White WB, Cannon CP, Heller SR, et al. EXAMINE Investigators Alogliptin after acute coronary syndrome in patients with type 2 diabetes. *N Engl J Med.* 2013;369(14):1327–1335. [PubMed: 23992602]
108. Holman RR, Bethel MA, Mentz RJ, et al. EXSCEL Study Group Effects of once-weekly exenatide on cardiovascular outcomes in type 2 diabetes. *N Engl J Med.* 2017;377(13):1228–1239 [PubMed: 28910237]
109. Pfeiffer MA, Claggett B, Diaz R, et al. ELIXA Investigators Lixisenatide in patients with type 2 diabetes and acute coronary syndrome. *N Engl J Med.* 2015;373(23):2247–2257 [PubMed: 26630143]
110. Vaccaro O, Masulli M, Nicolucci A, et al. Thiazolidinediones Or Sulfonylureas Cardiovascular Accidents Intervention Trial (TOSCA. IT) study group; Italian Diabetes Society Effects on the incidence of cardiovascular events of the addition of pioglitazone versus sulfonylureas in patients with type 2 diabetes inadequately controlled with metformin (TOSCA.IT): a randomised, multicentre trial. *Lancet Diabetes Endocrinol.* 2017;5(11):887–897 [PubMed: 28917544]
111. Marso SP, Bain SC, Consoli A, et al. SUSTAIN-6 Investigators Semaglutide and cardiovascular outcomes in patients with type 2 diabetes. *N Engl J Med.* 2016;375(19):1834–1844. [PubMed: 27633186]
112. Marso SP, Daniels GH, Brown-Frandsen K, et al. LEADER Steering Committee; LEADER Trial Investigators Liraglutide and cardiovascular outcomes in type 2 diabetes. *N Engl J Med.* 2016;375(4):311–322 [PubMed: 27295427] **First to demonstrate cardiovascular benefit in terms of both MACE, reduced CV death and all cause death from the GLP2-RA class of agents
113. Zinman B, Wanner C, Lachin JM, et al. EMPA-REG OUTCOME Investigators Empagliflozin, cardiovascular outcomes, and mortality in type 2 diabetes. *N Engl J Med.* 2015;373(22):2117–2128. [PubMed: 26378978] **First in CVOTs to demonstrate cardiovascular benefit from hypoglycaemic agents in terms of reduced MACE and reduced rates of admissions for heart failure
114. Neal B, Perkovic V, Mahaffey KW, et al. CANVAS Program Collaborative Group Canagliflozin and cardiovascular and renal events in type 2 diabetes. *N Engl J Med.* 2017;377(7):644–657. [PubMed: 28605608]
115. Neal B, Perkovic V, Mahaffey KW, de Zeeuw D, Fulcher G et al. Canagliflozin and Cardiovascular and Renal Events in Type 2 Diabetes. *N Engl J Med.* 2017 8 17;377(7):644–657 [PubMed: 28605608]
116. Wiviott SD, Raz I, Bonaca MP, Mosenzon O, Kato ET et al. Dapagliflozin and Cardiovascular Outcomes in Type 2 Diabetes. *N Engl J Med.* 2018 11 10.
117. Das SR, Everrett BM, et al. 2018 ACC Expert Consensus Decision Pathway on Novel Therapies for Cardiovascular Risk Reduction in Patients With Type 2 Diabetes and Atherosclerotic Cardiovascular Disease: A Report of the American College of Cardiology Task Force on Expert Consensus Decision Pathways. *JACC.* 11 2018;25566. doi:10.1016/j.jacc.2018.09.020

118. Voight BF, Peloso GM, Orho-Melander M, Frikke-Schmidt R, Barbalic M et al. Plasma HDL cholesterol and risk of myocardial infarction: a mendelian randomisation study. *Lancet*. 2012 8 11;380(9841):572–80. [PubMed: 22607825]
119. Investigators AIM-HIGH. Niacin in Patients with Low HDL Cholesterol Levels Receiving Intensive Statin Therapy. *N Engl J Med*. 2011 12 15;365(24):2255–67. doi: 10.1056/NEJMoal107579. [PubMed: 22085343]
120. HPS2-THRIVE Collaborative Group, Landray MJ, Haynes R, Hopewell JC, Parish S, Aung T et al. Effects of extended-release niacin with laropiprant in high-risk patients. *N Engl J Med*. 2014 7 17;371(3):203–12. doi: 10.1056/NEJMoal1300955 [PubMed: 25014686]
121. Schwartz GG, Olsson AG, Abt M, Ballantyne CM, Barter PJ et al. Effects of dalcetrapib in patients with a recent acute coronary syndrome. *N Engl J Med*. 2012 11 29;367(22):2089–99 [PubMed: 23126252]
122. Lincoff AM, Nicholls SJ, Riesmeyer JS, Barter PJ, Brewer HB et al. Evacetrapib and Cardiovascular Outcomes in High-Risk Vascular Disease. *N Engl J Med*. 2017 5 18;376(20):1933–1942 [PubMed: 28514624]
123. HPS3/TIMI55–REVEAL Collaborative Group, Bowman L, Hopewell JC, Chen F, Wallendszus K, Stevens W et al. Effects of Anacetrapib in Patients with Atherosclerotic Vascular Disease. *N Engl J Med*. 2017 9 28;377(13):1217–1227 [PubMed: 28847206] *Trial evaluating CETP inhibitors that demonstrated a modest reduction in MACE, but not thought to be driven directly by increasing HDL-C levels
124. Rohatgi A, Khera A, Berry JD, et al. HDL cholesterol efflux capacity and incident cardiovascular events. *N Engl J Med*. 2014;371(25):2383–93.. [PubMed: 25404125]
125. ISIS-2 Collaborative Group. Randomised Trial of Intravenous Streptokinase, Oral Aspirin, Both, or Neither among 17187 Cases of Suspected Acute Myocardial Infarction. *The Lancet*. 1988 332(8607):349–360
126. Mauri L, Kereiakes DJ, Yeh RW, Driscoll-Shempp P, Cutlip DE et al. Twelve or 30 months of dual antiplatelet therapy after drug-eluting stents. *N Engl J Med*. 2014 12 4;371(23):2155–66. doi: 10.1056/NEJMoal1409312. Epub 2014 Nov 16. [PubMed: 25399658]
127. Bonaca MP, Bhatt DL, Cohen M, Steg PG, et al. Long-term use of ticagrelor in patients with prior myocardial infarction. *N Engl J Med*. 2015 5 7;372(19):1791–800 [PubMed: 25773268]
128. Bonaca MP, Storey RF, Theroux P3, Steg PG, Bhatt DL. et al. Efficacy and Safety of Ticagrelor Over Time in Patients With Prior MI in PEGASUS-TIMI 54. *J Am Coll Cardiol*. 2017 9 12;70(11):1368–1375 [PubMed: 28882235] **Demonstrated significant benefit in reducing rate of MACE and residual risk with longer course of dual antiplatelet therapy with ticagrelor plus aspirin
129. Yeh RW, Secemsky EA, Kereiakes DJ, et al. Development and Validation of a Prediction Rule for Benefit and Harm of Dual Antiplatelet Therapy Beyond 1 Year After Percutaneous Coronary Intervention. *JAMA*. 2016;315(16):1735–1749. doi:10.1001/jama.2016.3775 [PubMed: 27022822] *Developed a clinical tool for predicting those who may benefit from longer duration of DAPT and those at higher bleeding risk
130. Eikelboom JW, Connolly SJ, Bosch J, Dagenais GR, Hart RG, et al. Rivaroxaban with or without Aspirin in Stable Cardiovascular Disease. *N Engl J Med*. 2017 10 5;377(14):1319–1330 [PubMed: 28844192] **Demonstrated significant benefit in reducing rate of MACE and residual risk with rivaroxaban and aspirin in combination, with no increase in rates of fatal bleeding
131. Janssen Pharmaceuticals. U.S. FDA Approves XARELTO® (rivaroxaban) to Reduce the Risk of Major Cardiovascular Events in Patients with Chronic Coronary Artery Disease (CAD) or Peripheral Artery Disease (PAD). Janssen Global Services. 2018 10 11; internet page. Available at: <https://bit.ly/2A74vIT> [date accessed: 2019 Feb 11]
132. Libby P History of Discovery: Inflammation in Atherosclerosis. *Arterioscler Thromb Vasc Biol*. 2012;32(9):2045–2051. doi:10.1161/ATVBAHA.108.179705 [PubMed: 22895665]
133. Ridker PM, Rifai N, Rose L, Buring JE, Cook NR. Comparison of C-reactive protein and low-density lipoprotein cholesterol levels in the prediction of first cardiovascular events. *N Engl J Med*. 2002;347(20):1557–1565. doi:10.1056/NEJMoal021993 [PubMed: 12432042]

134. The Interleukin-6 Receptor Mendelian Randomisation Analysis (IL6R MR) Consortium. The interleukin-6 receptor as a target for prevention of coronary heart disease: a mendelian randomisation analysis. *Lancet* 2012;379:1214–1224 [PubMed: 22421340]
135. Ridker PM, Cushman M, Stampfer MJ, Tracy RP, Hennekens CH. Inflammation, aspirin, and the risk of cardiovascular disease in apparently healthy men. *N Engl J Med* 1997;336:973–979 [PubMed: 9077376]
136. Ridker PM, Hennekens CH, Buring JE, Rifai N. C-reactive protein and other markers of inflammation in the prediction of cardiovascular disease in women. *N Engl J Med* 2000;342:836–843 [PubMed: 10733371]
137. Ridker PM, Everett BM, Thuren T, MacFadyen JG, Chang WH et al. Antiinflammatory Therapy with Canakinumab for Atherosclerotic Disease. *N Engl J Med*. 2017 9 21;377(12):1119–1131 [PubMed: 28845751] *Demonstrated benefit in MACE reduction with directed anti-inflammatory therapy
138. Miller J FDA snubs Novartis bid to repurpose inflammation drug for heart attacks. Reuters. 2018 10 18; internet page. Available at: <https://reut.rs/2QXmHnt>. [Date accessed: 2019 Feb 11]
139. Ridker PM, Everett BM, Pradhan A, MacFadyen JG, Solomon DH et al. Low-Dose Methotrexate for the Prevention of Atherosclerotic Events. *N Engl J Med*. 2018 11 10. doi: 10.1056/NEJMoa1809798. [Epub ahead of print]

Table 1.

Clinical variables included in TRS 2oP Score

Clinical variable	HR (95% CI)	Points allocated
CHF	2.03 (1.68–2.46)	1
Prior CVA	1.83 (1.39–2.4)	1
Hypertension	1.61 (1.34–1.93)	1
Diabetes mellitus	1.49 (1.27–1.75)	1
Current smoking	1.47 (1.23–1.75)	1
Prior CABG	1.44 (1.2–1.73)	1
Age >75	1.4 (1.11–1.75)	1
Renal dysfunction [*]	1.36 (1.12–1.65)	1
Peripheral vascular disease [†]	1.36 (1.13–1.64)	1

Bohula et al., 2016

^{*} Renal dysfunction defined as eGFR <60mL/min/1.73 m²;[†] Peripheral vascular disease defined by clinical history of claudication, ankle-brachial index <0.85, previous revascularisation or amputation

Table 2.

Event rates in contemporary secondary prevention trials

Medication class	Trial	Recruitment period	Primary outcome* rate (control)	Primary outcome rate (ivn)	Median follow up (years)	Approximate annual event rate†(placebo)
Statins						
	TNT	April 1998-Dec1999	10.90%	8.70%	4.9	2%
	PROVE-IT/TIMI-22	Nov 2000-Dec 2001	26.30%	22.40%	2	13%
	LIPID [‡]	Apr 1990-Sept 1992	14.10%	11.00%	6	2%
Ezetimibe						
	IMPROVE-IT [§]	2005–2010	3.47%	3.27%	7	3%
PCSK9 monoclonal antibodies						
	ODYSSEY	Nov 2012-Nov 2015 (main cohort)	11.10%	9.50%	2.8	4%
	FOURIER	Feb 2013-June 2015 [§]	11.30%	9.80%	2.16	5%
	SPIRE-1,2 [¶]	Oct 2013-Nov 2016	3.16%	3.59%	0.8	3.16%
Eicosapentaenoic acid						
	REDUCE-IT ^{**}	Nov 2011-Aug 2016	19.30%	25.50%	4.9	4%
	JELIS ^{**}	1996–1999	10.70%	8.70%	4.6	2%
Niacin						
	HPS2 THRIVE ^{††}	Apr 2007-July 2010	13.70%	13.20%	3.9	4%
	AIM HIGH	2006–2010	16.20%	16.40%	3	5%
SGLT2 inhibitors						
	EMPA-REG	Sept2010-Apr2013	12.10%	10.50%	3.1	4%
	CANVAS ^{‡‡}	Dec 2009-Mar 2011	3.15%	2.69%	2.4	3.15%
	DECLARE-TIMI58	2013 onwards	9.40%	8.80%	4.2	2%
GLP-1 agonist						
	SUSTAIN-6	Feb 2013-Dec 2013	8.90%	6.60%	2.1	4%
	LEADER	Sept 2010-April 2012	14.90%	13.00%	2.8	5%
CETP inhibitors						
	dalOUTCOMES	April 2008-July 2010	8.30%	8.00%	2.6	3%
	ACCELERATE	Oct 2012-Dec 2013	12.80%	12.90%	2.16	6%
	REVEAL	Aug 2011-Oct 2013	11.80%	10.80%	4.1	3%
P2Y12 inhibitors						
	DAPT	Aug 2009-July 2011	5.90%	4.30%	2.5	2%
	PLATO ^{§§}	Oct 2006-July 2008	11.70%	9.80%	1	12%
	TRITON-TIMI38 ^{§§}	Nov 2004-July 2007	12.10%	9.90%	1.25	10%
	PEGASUS-TIMI54 ^{¶¶}	Oct2010-May2013	9.04%	7.81%	2.75	3%

Medication class	Trial	Recruitment period	Primary outcome* rate (control)	Primary outcome rate (ivn)	Median follow up (years)	Approximated annual event rate†(placebo)
Direct oral anticoagulants						
	COMPASS***	Mar 2013-May 2016	5.40%	4.1%	1.9	3%
Anti inflammatory agents						
	CANTOS§,†,†,†	Apr 2011-Mar 2014	4.50%	3.86%	3.7	4.50%
	CIRT¶	Apr 2013-Mar 2018	4.31%	4.13%	2.3	4.30%

* Primary outcome as a combined rate of major adverse cardiovascular events, unless otherwise stated

† Approximated annual event rate: reported primary outcome rate / median follow up

‡ Primary outcome in LIPID was all-cause mortality

§ Kaplan-Meier event rate at median follow up listed for IMPROVE-IT

¶ Event rates originally reported as events per 100 patient years; approximated to annual event rate here

*** Established cardiovascular disease subgroups listed

†† Primary outcome in HPS2-THRIVE was first major cardiovascular event

††† Event rates originally reported as events per 1000 patient years; approximated to annual event rate here

§§ Event rates reported at follow up duration listed

¶¶ Intervention group taken as pooled ticagrelor group in PEGASUS-TIMI 58

*** Intervention group taken as rivaroxaban plus aspirin for COMPASS

††† Intervention rate reported for 150mg subgroup under intervention group rates