The predictive value of peritubular capillaries C3d deposition in IgA glomerulonephritis

Mihaela Gherghiceanu^a*, M. Penescu^b, E. Mandache^a

^a "Victor Babes" National Institute of Pathology, Bucharest, Romania

^b "Carol Davila" Clinical Hospital of Nephrology, Bucharest, Romania

Received: July 28, 2004; Accepted: January 28, 2005

Abstract

The IgA glomerulonephritis (IgAGN) is one of the most common primary glomerulonephritis and has a variable and difficult to predict evolution toward the end-stage nephrosclerosis. The deposition of C3d complement component in peritubular capillaries (PTCs) indicates a variant type of acute rejection while C3d deposition in primary glomerulonephritis (GN) is poorly documented. The aim of this study is to examine C3d expression in peritubular capillaries (PTCs) as a predictive marker and its correlation with the severity of renal injury in IgA glomerulonephritis. Polyclonal FITC conjugated rabbit anti-human C3c and C3d antibodies were used for direct immunofluorescent evaluation of the C3c and C3d deposits in 24 kidney biopsies with IgA glomerulonephritis. The study revealed that the C3d deposits in peritubular capillaries were associated with known predictors for rapid progression of IgAGN: glomerular sclerosis (63.6%), atrophic tubules (90.9%) and interstitial sclerosis (81.8%). The intensity of the C3c glomerular immunofluorescent deposits was related with active lesions. Thus, the predictive value of C3d deposition on PTCs in IgAGN is worth to be taken into consideration as an unfavorable outcome of the disease and request further long run investigations.

Keywords: IgA glomerulonephritis • complement system • peritubular capillaries • C3c • C3d

Introduction

The IgA glomerulonephritis is one of the most common primary types of glomerulonephritis. Because IgAGN has a long natural history and variable evolution to end-stage renal disease it is difficult to predict its outcome.

The activation of the complement system (CS) plays an important role in the initiation and evolution of the inflammatory lesions in IgA glomeru-

lonephritis. Complement activation in IgAGN by polymeric IgA through the alternative and lectin pathways enhance the proinflammatory effects of IgA deposition and therefore play an unfavorable role in the disease progression [1, 2]. The evidence that complement C3 activation occurs in IgAGN patients with deteriorating course but not in those with a stable course may provide a new approach to stratify patients for treatment or no treatment [3].

The key of CS activation is the cleavage of C3 and the formation of soluble C3a and C3b, which contain a thioester bond within the C3d region. C3 is the most

* Correspondence to: Dr. M. GHERGHICEANU,
Ultrastructural Pathology Laboratory,
"Babes" National Institute of Pathology,
99-101 Splaiul Independentei, 050096, Bucharest, Romania.
Tel./Fax: 0040 21 411 51 05.
E-mail: mgherghiceanu@yahoo.com

abundant protein of the CS (90-120 mg/dl) and supports the activation through classical (by C1q, C reactive protein), lectin (MBL) and alternative pathways [4, 5, 6].

C3 is permanently activated on a low-level as a result of spontaneous hydrolysis of the thioester group in native C3 ("C3 tick-over" of the alternative pathway) that leads to C3b deposition on the cells surface or on the circulating biological particles. Recently it was demonstrated that C3 is continually low-level activated via the classical pathway ("C1q tick-over") mediated by soluble immunoglobulins, mostly by IgG [7]. This low-level mechanism of continuous activation of the C3 seems to facilitate the antigen recognition and to allow the binding of C3b on nonself surfaces. The self-surfaces are protected by the presence of membrane associated complement inhibitors [8, 9].

The C3 cleavage is the central point of the CS activation. Once C3b is formed, its actions are determined by the intervention or non-intervention of the complement regulator proteins (Factor I, CR1, CR2, Factor H, MCP, and DAF). The absence of this regulator intervention on C3b makes possible the amplification, formation of C5 convertase and of the membrane attack complex (MAC). The presence of C3d implies the action of the regulator proteins of the complement. The regulator proteins promote the cleavage of C3b and lead to the formation of more breakdown products (iC3b, C3f, C3c, C3dg, C3g), leaving the small C3d fragment covalently attached by thioester bound to the antigenic surface. The decay-accelerating factor is expressed in the glomerular vascular pole in normal kidney (DAF prevents B binding and CS amplification) and membrane cofactor protein is expressed in normal glomerular capillaries (MCP direct factor I to cleave C3b) [9, 10].

The significance of C3c and C3d deposits is poorly documented in native kidney glomerulonephritis. One of the first studies [11] showed deposits of C3dg and C3d in capillary basement membranes and mesangial matrix in patients with minimal change disease. Studying the clinical-pathological significance of the glomerular deposition of C3c and C3d in IgAGN Nakagawa et al [12] showed that C3c is associated with the active phase of IgAGN. Muso et al [13] found that C3d mesangial deposition in lupus nephritis reveals a significant correlation with the disease activity. The deposition of complement factor C3d in peritubular capillaries indicates a variant type of acute rejection characterized by a worse clinical

outcome [14] but the importance of C3d deposits present in peritubular capillaries in native kidney glomerulonephritis is still unclear.

The aim of this study is to evaluate the correlation between the C3d expression in peritubular capillaries and the severity of renal injury in IgAGN. Our study analyzed the deposition of C3 breakdown products (C3c and C3d) and C3d possible implication in IgAGN prognosis.

Materials and methods

Study subjects

Twenty-four renal biopsies from patients with IgAGN were studied. The renal biopsies that proved to have enough material for all three required techniques were selected from the samples received for routine diagnosis during the last year. The cases with Henoch-Schönlein purpura and secondary forms of IgAGN were excluded.

Two biopsies with minimal change disease (MCD) and two renal biopsies performed for microscopic hematuria showing normal structure in light and electron microscopy were used as controls. Clinical data were expressed as means $\pm$ SD.

Kidney biopsies were performed with disposable COLT-ABI Sterylab 16-gauge needles. Renal biopsy fragments were divided into three parts: the two extremities were fixed in glutaraldehyde for light and electron microscopy and the middle part was frozen for immunofluorescence. Diagnosis was made by standard examination of renal biopsy specimens by light microscopy, immunofluorescence, and electron microscopy.

Light and electron microscopy

Light and electron microscopy were performed on Epon embedded tissue. Slides containing 1 μ thick sections were stained with toluidine blue for optic evaluation. The ultra-thin sections (60 nm) were examined in a Philips 301 transmission electron microscope after uranyl acetate and lead citrate staining.

Immunofluorescence

Frozen sections (3 μ) were air dried at room temperature for 30 min. Two sections were performed for each anti-

Table 1 Histological classification of IgA glomerulonephritis used in this study.

Histological class		Renal lesions
I	Minimal glomerular changes	none (0) or minimal (1) mesangial proliferation
II	Diffuse endocapillary proliferation	moderate (2) or intense (3) mesangial proliferation
III	Focal segmental GN	segmental and focal adhesion and mesangial proliferation
IV	Diffuse endo- and extracapillary proliferation	diffuse mesangial proliferation with: - crescents in less than 50% glomeruli - IVa, - crescents in more than 50% glomeruli - IVb

body. The slides were placed in PBS 10% pH 7,4 for 15 min. and then incubated with polyclonal FITC conjugated rabbit anti-human antibodies for 30 min., washed twice in PBS 10% for 15 min., and mounted with glycerin. Polyclonal FITC conjugated rabbit anti-human C3c and C3d (DAKO) antibodies were used. Anti-C3c antibody reacts with both C3c and C3c part of native C3, C3b, or iC3b. The manufacturer claims that there is no reaction with C3d and C3a. FITC conjugated anti-C3d antibody reacts with C3d and C3d epitopes on C3 complement molecules. Immunofluorescence for IgA, IgG, IgM, C1q, albumin and fibrin was performed for each renal biopsy.

Data evaluation

We have assessed the pattern (linear, fine granular, granular), the site of deposition (glomerular capillaries, mesangium, vascular pole, arterioles, arteries, peritubular capillaries, tubular epithelial cells, tubular basement membranes, Bowman capsule) and the distribution (segmental or global and focal or diffuse) of immunofluorescent deposits. The binding of conjugated antibodies was evaluated in a grading of intensity and distribution. The assessing system for immunofluorescent deposits was the following: 0 - no deposits, 1 - weak glomerular deposits, 2 - moderate glomerular deposits, 3 - intense glomerular deposits. 0,5 points were added for associated arteriolar deposits. One point was added for the presence of C3d deposits in PTCs. The assessing system for morphological evaluation (mesangial proliferation, interstitial inflammation, glomerular sclerosis, interstitial fibrosis, crescents) was referred to their extension or number of glomeruli involved: 0 - absent (0%), 1 - weak, segmental and focal (<25%), 2 - moderate, segmental and diffuse (25-50%), 3 - global and diffuse (>50%). The following histopathologic classification (Table 1) was used for renal

biopsies evaluation: class I - minimal glomerular changes; class II - diffuse mesangial proliferation; class III - focal and segmental glomerulonephritis; class IV - diffuse endocapillary proliferation with crescents (adapted after Droz D. and Lanz B. [15]).

Results

The biopsies were analyzed with all three techniques: immunofluorescence, light, and electron microscopy.

Non IgA glomerulonephritis

Two renal biopsies performed for microscopic hematuria showed normal characteristics in light and electron microscopy and were used as controls. The immunofluorescence was negative and the renal histology was normal. The biopsies with MCD were negative for C3c in the glomerular area and weakly positive in the vascular pole and preglomerular arterioles (Fig. 1). The C3d was intensely positive and fine granular in the glomerular capillaries, vascular pole and arterioles (Fig. 2) but there was no histological correspondence such as electron-dense deposits for C3 presence. There was no C3d deposition in PTCs in these cases.

IgA glomerulonephritis

The IgAGN were divided into the following two groups: group A without C3d deposits in PTCs and group B with C3d deposits in PTCs.

Fig. 1 Minimal change disease. Weak positive reaction for C3c in the vascular pole (a) and small granular deposits in the glomerulus (g).

Fig. 2 Intense positive, fine granular C3d deposits in the glomerular capillaries (g) and coarse deposits in vascular pole and arterioles (a).

Fig. 3 Strong fluorescent, arborescent deposits of IgA in two glomeruli and an arteriole (>) with subendothelial hyaline deposits (insert, toluidine blue).

Fig. 4 Severe endocapillary proliferation associated with crescentic extracapillary proliferation – class IVb IgAGN. 1μ plastic section stained with toluidine blue.

The group B with C3d in PTCs had a higher level of blood C3, proteinuria and hematuria than group A without C3d in PTCs. Clinical data are expressed in table 2 as a mean±SD value for all analyzed kidney biopsies with IgAGN.

The study revealed that the C3d deposits present in peritubular capillaries were associated with more severe renal lesions in IgAGN. Table 3 shows the classification and morphological evaluation of renal biopsies with IgAGN in group A and group B. In group B, with C3d deposits in PTCs, the percentage of cases with severe renal lesions was higher than in group A (table 4). In group B we found more sclerotic lesions and the percentage of cases with crescents was higher than in group A. A distinctive fea-

ture for IgAGN with C3d deposits in PTCs was the presence of the electron dense round-oval hyaline deposits located in the glomerular vascular pole or afferent arteriolar wall. These deposits corresponded in rare cases with fluorescent IgA deposits in the same location (Fig. 3).

Kidney biopsies showed mesangial immunofluorescent deposits of IgA and confirmed the diagnosis (Fig.3). The immunofluorescence demonstrated in IgAGN that the vascular poles and the arterioles were intensely or moderately labeled by C3c and C3d, having a granular and transmural pattern in almost all cases. C3d stained intensely the mesangial area and the glomerular capillary walls. C3c was negative in the glomeruli

Fig. 5 Strong positive labeling for C3c in the mesangial area (g), arterial wall (a) and weak reaction in the cytoplasm of tubular epithelial cells (double arrows).

Fig. 6 Immunofluorescence for C3d showing strong labeling in the glomerulus (g), PTCs (double arrows), and arterial wall (a).

Fig. 7 IgAGN with positive labeling for C3c limited to arteriolar walls (a), glomerular parietal epithelial cells and some tubular epithelial cells (double arrows).

Fig. 8 C3d deposits in two glomeruli (g), arteriolar walls (a) and PTCs (double arrows).

when minimal histological changes and no inflammatory cells were present. The cases with crescentic proliferation (Fig. 4) showed C3c positive in the cytoplasm of proximal tubular epithelial cells (Fig. 5) and C3d in PTCs and interstitial space (Fig. 6). C3c was moderate or intense positive in the mesangial area for cases with mild or intense mesangial proliferation, PMNs and trombocytes in the capillary lumina.

The expression of C3c was sometimes restricted to the arteriolar walls, parietal epithelial cells and tubular epithelial cells in those cases with mesangial proliferation only (Fig. 7), while C3d showed immunofluorescent deposits in glomeruli and PTCs (Fig. 8).

In the cases with sclerous glomeruli and interstitial fibrosis C3d was intensely positive all over the glomerular area. The Bowman capsule was segmentally stained in the cases with capsular-glomerular adherences or hyaline deposits in glomerular capillaries. C3d deposits were more intense and wider distributed than C3c deposits in all cases and stained the glomeruli even when C3c was negative and the glomerular lesions were reduced.

The distribution of renal lesional classes in group A and group B (Fig. 9) shows that the PTCs C3d deposition was essentially associated with class III and class IV. Figure 10 shows in a relevant manner that the morphological and immunofluorescent features were more severe in group B.

Table 2 Mean values of clinical data.

Mean $\pm$ SD	Group A, n=13	Group B, n=11
Age	33.91 $\pm$ 18.72	36.5 $\pm$ 14.12
Blood pressure	133.88 $\pm$ 26.90	137.5 $\pm$ 24.03
Creatinine (mg/dl)	1.30 $\pm$ 0.08	1.62 $\pm$ 0.54
Urea (mg/dl)	59 $\pm$ 25.44	84.97 $\pm$ 65.58
C3 (mg/dl)	87 $\pm$ 7.88	97.4 $\pm$ 14.41
Proteinuria (mg/24h)	2.32 $\pm$ 3.96	3.025 $\pm$ 2.40
Hematuria (Er/min)	4957.14 $\pm$ 5155.53	7875 $\pm$ 5807.68

Discussion

The current study investigated the correlation of C3c and C3d deposits with the severity of renal injury in IgAGN. Our data showed that the presence of C3d deposits in glomeruli is unspecific but C3d deposits in PTCs are correlated with a high degree of crescents, glomerular sclerosis, and interstitial fibrosis. The extents of glomerular and interstitial sclerosis, and the presence of crescents, are identified as markers of poor prognosis [16]. Therefore, the presence of C3d deposits in PTCs could be an additional marker for unfavorable evolution.

The glomerular staining for C3c was found to be more intense in the glomerular lesions associated with acute inflammatory aspects such as: mesangial cellular proliferation and intraglomerular inflammation. Therefore, the staining for C3c was correlated with the inflammatory active phase of the IgA glomerular disease. The C3c deposits indicate an ongoing activation of CS with new immune complex formation [17] while C3d deposits persisted in spite of the lack of this ongoing complement activation. Schultze [17] showed that C3c deposits were reduced by over 85% within 24 hours in three experimental models of complement-dependent glomerulonephritis. Our results also confirm the study of Nakagawa et al [12] showing that C3c is associated with the active phase of IgAGN.

After a number of successive cleavages, the C3c fragment enters the fluid phase leaving the smaller fragment C3d attached to antigens. C3d showed more intense and wider distributed deposits than C3c in almost all cases but the glomerular staining

for C3d seems to be nonspecific. It is possible that the difference of intensity between C3c and C3d intraglomerular staining underlines an evolutionary stage: the higher the difference, the older the disease on a cumulative process. This hypothesis is in accordance with the observation that C3d could bind laminin and fibronectin [18–20].

The presence of C3c without C3d suggests that there is only native C3, which does not have the C3d fragment exposed. The C3c alone was present with focal distribution in the cytoplasm of tubular epithelial cells in cases with active lesions only. The presence of C3c protein in tubular epithelial cells suggests that C3 is locally synthesized during the acute phase of the inflammatory process in glomerulonephritis [21, 22]. In pathological conditions, local synthesis of C3 in tubular and glomerular cells increases and leads to the development of lesions [23, 24]. Tang et al [24] found that donor kidney contributed with 4-5% of the total circulating C3 and with 9,6% in allograft rejection episodes. This finding indicates that the kidney is probably a major extra hepatic source of C3 with a significant potential to increase its local concentration during inflammatory injuries. This could explain why there is no correlation between the level of C3 in blood and the glomerular deposition of C3 in glomerulonephritis and why C3 had a mean blood level higher in group B than in group A.

The presence of C3d outside the C3c stained areas suggests that there is only C3d small fragments. C3d alone was limited to PTCs and tubular basement membranes. The C3c in tubular epithelial cells and C3d in PTCs and interstitial spaces were found

Fig. 9 Distribution of renal lesional classes (table 1) and number of patients in Group A and Group B.

Fig. 10 Morphological and immunofluorescent findings in group A and group B.

Table 3 Immunofluorescent and morphological evaluation of renal biopsies of IgAGN in Group A and Group B.

Crt. no.	IgAGN class	Mesangial IgA deposits	Glomerular IgG deposits	Glomerular IgM deposits	Glomerular C1q deposits	C3c deposits	C3d deposits	Mesangial proliferation	Glomerular inflammation	Crescents	Glomerular sclerosis	Interstitial inflammation	Interstitial sclerosis	Atrophic tubules	Vascular hyaline deposits
Group A - without C3d in PTCs															
1.	I	2	1	0	0	0.5	2.5	0	0	0	0	0	0	0	0
2.	I	3	1	1	0	2.5	3.5	0	1	0	0	2	0	0	0
3.	I	2	1	0	0	0.5	1.5	0	0	0	0	0	0	0	0
4.	I	3	2	2	0	2	2.5	0	0	0	0	0	0	0	0
5.	I	3	2	1	1	1	3.5	1	0	0	0	0	0	0	0
6.	I	3	2	1	0	0.5	3.5	1	1	0	0	0	0	0	0
7.	I	2	1	0	0	0.5	3.5	1	0	0	0	0	0	0	0
8.	II	3	0	0	0	0.5	2.5	2	1	0	0	0	0	0	0
9.	II	3	2	1	0	1.5	2.5	2	1	0	0	1	0	0	0
10.	III	3	1	2	0	2.5	2.5	1	0	0	1	1	1	1	0
11.	III	3	2	2	0.5	0.5	3.5	1	0	0	1	0	1	1	0
12.	III	2	2	1	1	3.5	3.5	2	3	0	3	2	1	1	0
13.	IVb	2	2	1	1	2.5	3.5	2	2	3	1	1	0	1	0
Group B- with C3d in PTCs															
1.	II	3	1	0	0	1.5	4.5	3	1	0	0	1	1	1	0
2.	III	2	1	2	0	1	4.5	2	1	0	0	0	0	0	0
3.	III	3	2	2	2	3.5	4.5	2	1	0	3	1	1	1	1
4.	III	3	1	2	0.5	3.5	4.5	2	0	0	2	2	0	2	1
5.	III	3	1	1	2	3.5	4.5	2	3	0	3	3	2	2	3
6.	III	3	1	2	1	3.5	4.5	1	0	0	3	3	2	2	2
7.	IVa	3	1	0	0	2.5	4.5	3	2	1	0	3	2	1	1
8.	IVa	3	1	2	1	4.5	4.5	3	1	2	2	1	1	1	2
9.	IVb	3	2	1	1.5	4.5	4.5	3	2	3	2	2	1	1	2
10.	IVb	3	2	1	1	3.5	4.5	3	2	3	0	3	2	1	1
11.	IVb	2.5	2.5	2.5	2	3.5	4.5	3	1	3	2	3	1	1	2

Table 4 Occurrence of renal lesions in group A and group B.

	Mesangial proliferation	Glomerular inflammation	Crescents	Glomerular sclerosis	Interstitial inflammation	Interstitial sclerosis	Atrophic tubules	Vascular hyaline deposits
Group A	69.23%	46.15%	7.69%	30.76%	38.46%	23.07%	30.76%	0
Group B	100%	81.81%	45.45%	63.63%	90.90%	81.81%	90.90%	81.81%

together in some cases. The C3c deposition in tubular epithelial cells was found in correlation with interstitial inflammation. This fact could mean that C3 synthesized by tubular epithelial cells is secreted in the interstitial space where it may stimulate an antigen specific response through interstitial dendritic cells and could attract the inflammatory cells [25].

The presence of C3c and C3d in the vascular pole and arterioles could indicate a C3 activation or degradation process in these sites. In our study, C3 breakdown products are present in the vascular pole and glomerular arterioles not only in glomerulonephritis but also in the normal kidney. C3 has a protective function by preventing local accumulation of immune complexes. This point of view is sustained by the fact that C3 deficiencies lead to an impaired clearance of immune complexes and their accumulation.

The complement system, through the adjuvant effect of C3d, may select antigens for recognition by the acquired immune system [4, 26, 27]. Research papers accumulated in the last years show that C3d, in addition to its established role as an adjuvant, could have an immunosuppressive activity [28]. A functionally active complement system is present within the anterior chamber of the normal eye [29] and this could be the same in kidney. The local activation of C3 in the vascular territory could play a crucial role in both the immune control and the clearance of immune complexes. An active complement could be normal in this specialized unit composed by glomerulus, juxtaglomerular apparatus, and arterioles having a role in the physiology of the immune system. The presence of C3d in normal histological cases suggests this and it is sustained by the fact that C3d can be found in the glomerular basement membranes in the absence of signs for ongoing local complement activation or immune complex deposition [19]. It is also possible that this C3d deposition should be favored by the existence of some hereditary collagen type IV defects. Such

defects, visualized in electron microscopy, are present in at least 50% of IgA glomerulonephritis [30].

In conclusion, the active lesions in IgAGN are correlated with C3c glomerular deposition. The C3c presence in tubular epithelial cells and C3d deposits in PTCs are correlated with sclerotic lesions. Thus, this last pattern of staining could be predictive for the progression of kidney lesions and the deterioration of the renal function. Therefore, although the C3d deposition in PTCs during the IgAGN evolution seems to have prognostic significance, it still remains to be confirmed on further long run studies.

Acknowledgements

The VIASAN Program, grants 037/2001 and 035/2001 from the Ministry of Education and Research, Romania supported this work.

References

1. **Roos A., Bouwman L.H., van Gijlswijk-Janssen D.J., Faber-Krol M.C., Stahl G.L., Daha M.R.,** Human IgA activates the complement system via the mannan-binding lectin pathway, *J. Immunol.*, **167**: 2861-2868, 2001
2. **Lhotta K., Wurzner R., Konig P.,** Glomerular deposition of mannose-binding lectin in human glomerulonephritis, *Nephrol. Dial. Transplant.*, **14**: 881-886, 1999
3. **Floege J., Feehally J.,** IgA Nephropathy: Recent Developments, *J. Am. Soc. Nephrol.*, **11**:2395-2403, 2000
4. **Asghar S., Pasch M.C.,** Complement as a promiscuous signal transduction device, *Lab. Invest.* **78**: 1203-1225, 1998
5. **Wolbink G.J., Brouwer M.C., Buysmann S., ten Berge I.J.M., Hack C.E.,** CPR-mediated Activation of Complement in Vivo, *J. Immunol.*, **157**: 473-479, 1996
6. **Sahu A., Lambris J.D.,** Structure and biology of the complement protein C3, a connecting link between innate and acquired immunity, *Immunol. Rev.*, **180**: 35-48, 2001

7. **Manderson A.P., Pickering M.C., Botto M., Walport M.J., Parish C.R.,** Continual low-level activation of the classical complement pathway, *J. Exp. Med.*, **194**: 747-756, 2001
8. **Sheerin N.S., Sacks S.,** Complement and complement inhibitors: their role in autoimmune and inflammatory diseases, *Curr. Opin. Nephrol. Hypertens.*, **7**: 305-310, 1998
9. **Cosio F.G., Sedmak D.D., Nahman N.S.,** Localization of decay accelerating factor in normal and diseased kidneys, *Kidney Int.*, **36**: 100-107, 1989
10. **Sheerin N.S., Springall T., Abe K., Sacks S.,** Protection and injury: the differing roles of complement in the development of glomerular injury, *Eur. J. Immunol.*, **31**: 1255-1260, 2001
11. **Lesavre P., Monteiro RC, Chevailler A, Berger J, Halbwachs-Mecarelli L, Noel LH.,** Recent immunologic data on human glomerulonephritis. *Adv. Nephrol. Necker Hosp.*, **17**: 57-75, 1988
12. **Nakagawa H., Suzuki S., Haneda M., Gejyo F., Kikkawa R.,** Significance of glomerular deposition of C3c and C3d in IgA nephropathy, *Am. J. Nephrol.*, **20**: 122-128, 2000
13. **Muso E., Sekita K., Doi T., Kuwahara T., Yoshida H., Tamura T., Kawai C., Hamashima Y.,** Immunopathological correlation between mesangial C3d-deposition and C3d-fixing circulating immune complexes in lupus nephritis, *Clin. Immunol. Immunopathol.*, **32**: 351-358, 1984
14. **Kuypers D.R., Lerut E., Evenepoel P., Maes B., Vanrenterghem Y., Van Damme B.,** C3d deposition in peritubular capillaries indicates a variant of acute renal allograft rejection characterized by a worse clinical outcome, *Transplantation*, **76**: 102-108, 2003
15. **Droz D.** La biopsie rénale dans les glomérulonéphrites à dépôts mésangiaux d'IgA: maladie de Berger et purpura rhumatoïde. In: Droz D., Lantz B. La biopsie rénale. INSERM/Paris, 1996, pp. 161-163
16. **Radford M.G. Jr., Donadio J.V. Jr., Bergstralh E.J., Grande J.P.,** Predicting renal outcome in IgA nephropathy, *J. Am. Soc. Nephrol.*, **8**: 199-207, 1997
17. **Schulze M., Pruchno C.J., Burns M., Baker P.J., Johnson R.J., Couser W.G.,** Glomerular C3c localization indicates ongoing immune deposit formation and complement activation in experimental glomerulonephritis, *Am. J. Pathol.*, **142**: 179-187, 1993
18. **Hautanen A., Keski-Oja J.,** Interaction of fibronectin with complement component C3, *Scand. J. Immunol.*, **17**: 225-230, 1983
19. **Leivo I., Engvall E.,** C3d fragment of complement interacts with laminin and binds to basement membranes of glomerulus and trophoblast, *J. Cell. Biol.*, **103**: 1091-1100, 1986
20. **Johnson E., Gauperaa T., Eskeland T.,** Fibronectin binds to complement-coated agarose beads and increases their association to mouse macrophages, *Scand. J. Immunol.*, **22**: 315-20, 1985
21. **Welch T.R., Beischel L.S., Witte D.P.,** Differential expression of complement C3 and C4 in the human kidney, *J. Clin. Invest.*, **92**: 1451-1458, 1993
22. **Welch T.R., Frenzke M., Witte D.,** Evidence of a role for local complement expression in a murine model of progressive glomerulonephritis, *Pediatr. Res.*, **48**: 200-205, 2000
23. **Sacks S., Zhou W., Sheerin N.S.,** Complement synthesis in the injured kidney: does it have a role in immune complex glomerulonephritis?, *J. Am. Soc. Nephrol.*, **7**: 2314-2319, 1996
24. **Tang S., Zhou W., Sheerin N.S., Vaughan R.W., Sacks S.H.,** Contribution of renal secreted complement C3 to the circulating pool in humans, *J. Immunol.*, **162**: 4336-4341, 1999
25. **Sacks S., Zhou W.,** The effect of locally synthesized complement on acute renal allograft rejection, *Mol. Med.*, **81**: 404-410, 2003
26. **Dempsey P.W., Dempsey P.W., Allison M.D., Akkaraju S., Goodnow C., Fearon D.T.,** C3d of Complement as a Molecular Adjuvant: Bridging Innate and Acquired Immunity, *Science*, **271**: 348-350, 1996
27. **Carroll M.C.,** The role of complement and complement receptors in induction and regulation of immunity. *Annu. Rev. Immunol.*, **16**: 545-568, 1998
28. **Bennett M., Leanderson T.,** Was it there all the time?, *Scand. J. Immunol.*, **57**: 499-505, 2003
29. **Sohn J.H., Bora P., Suk H.J., Molina H., Kaplan H.J., Bora N.,** Tolerance is dependent on complement C3 fragment iC3b binding to antigen-presenting cells, *Nature Medicine*, **9**: 206-212, 2003
30. **Mandache E., Gherghiceanu M.,** Ultrastructural defects of the glomerular basement membranes associated with primary glomerular nephropathies, *Ultrastructural Pathology*, **2**: 103 – 108, 2004