Published in final edited form as:

Ann Epidemiol. 2019 May ; 33: 24–29. doi:10.1016/j.annepidem.2019.02.009.

Racial and Sex Differences in Biological and Chronological Heart Age in The Coronary Artery Risk Development in Young Adults (CARDIA) Study

Rachel Zmora^a, Pamela J Schreiner^a, Duke Appiah^b, Donald M Lloyd-Jones^c, Jamal S Rana^d, Cora E Lewis^e

^aUniversity of Minnesota, Minneapolis, MN

^bTexas Tech University Health Sciences Center, Lubbock, TX

^cNorthwestern University, Chicago, IL

^dDepartment of Cardiology and Division of Research, Kaiser Permanente Northern California, Oakland, CA

^eUniversity of Alabama at Birmingham, Birmingham, AL

Abstract

Purpose—Calculation of a biological heart age offers an alternative to absolute risk for characterizing cardiovascular risk by describing risk relative to an individual with normal health. We examined risk factors contributing to differences between biological and chronological heart age in young adults.

Methods—The Coronary Artery Risk Development in Young Adults (CARDIA) study included 2,264 black and white men and women who attended exam years 10 through 25. We estimated biological heart age using the non-laboratory-based Framingham 10-year cardiovascular disease (CVD) risk calculator. Trends in risk factors were examined cross-sectionally and longitudinally.

Results—Biological heart ages for black participants were 5.6 years older than their chronological ages over 15 years (p-value <0.001). In longitudinal analyses, urinary albumin-creatinine ratio (UACR) and alcohol intake were statistically significantly related to higher biological compared to chronological heart age while physical activity and education were statistically significantly related to negative heart age differences (p-values <0.001). Trends were similar in cross-sectional analyses at all time points.

Conclusions—Most risk factors driving biological heart age, including race, education, physical activity, and UACR, contributed to heart age differences cross-sectionally and longitudinally suggesting that risk factors related to adverse biological aging are important at younger and older ages.

Keywords

cardiovascular diseases; prevention and control; aging

Introduction

Biological heart age represents the functional age of the cardiovascular system of a person based on his or her cardiovascular risk factor profile^{1, 2}. Therefore, biological age offers another method for describing cardiovascular risk in individuals or groups beyond chronological age or probability of cardiovascular disease (CVD) event over time². When compared to chronological age, biological age can quantify both the excess or favorable risk in an individual and be used to identify lifestyle factors that may contribute to these differences.

Biological heart age is not a new concept, but thus far has not been widely utilized. The non-laboratory-based Framingham 10-year CVD risk calculator was designed to provide an individual risk of experiencing a CVD event¹. An analysis of data from the Behavioral Risk Factor Surveillance System (BRFSS) assessed biological heart age based on this risk score^{1, 2}. The authors of the BRFSS article found that biological heart age was significantly higher than chronological heart age for many adults in the US. They observed the greatest differences between biological and chronological age in men, Blacks, and those with lower socioeconomic status (SES), as indicated by education and household income². This analysis was limited by the lack of longitudinal data and the need to estimate systolic blood pressure, which is not captured by the survey².

The Coronary Artery Risk Development in Young Adults (CARDIA) Study offers an opportunity to reexamine the association between chronological and biological heart age over time with an emphasis on important demographic subgroups, specifically age, race, sex, and education. These subgroups were the recruitment strata for the cohort; therefore, the study has adequate distributions of these attributes³. Additionally, the relatively young age of CARDIA participants in earlier examinations may identify risk factors or trends that are not apparent at older ages when competing comorbidities or medication use may obscure trends.

CARDIA is able to provide directly measured data for biological heart age calculations both cross-sectionally and longitudinally. Using data from CARDIA, we calculated biological heart age in study participants at each exam starting at age 30. We examined differences in biological and chronological heart age and determined which subgroups had the largest differences between biological and chronological age across exam years. We also assessed if established cardiovascular risk factors not included in the Framingham risk calculator were also associated with biological heart age.

Methods

Study Design

The CARDIA study is a prospective cohort that enrolled 5,115 participants between the ages of 18 and 30 from 1985 to 1986. Enrollment was balanced across race, sex, age and education. Participants were from Birmingham, AL; Chicago, IL; Oakland, CA and Minneapolis, MN. Informed consent was obtained from all participants. The study design has been previously described in detail³.

Biological heart age was calculated at four consecutive exams to assess trends in risk factors. At exam years 10, 15, 20 and 25, there was high participant retention with 79, 74, 72, and 72% of surviving participants returning for follow-up exams, respectively. Of these participants, 3,790, 3,566, 3,453, and 3,406 individuals had sufficient data to calculate biological heart age at the four exam years, respectively. A total of 2,489 individuals had complete data at all four exams. Individuals who were less than 30 years old at exam year 10 were excluded from analysis because their chronological age did not fall within the parameters of the calculator. Additionally, pregnant women were excluded due to the potential impact on risk factors. One participant withdrew from the study. The final analyses included 2,264 participants. Using data from these four exams captured changes in biological heart age during early middle age.

Biological Heart Age

Biological heart age was derived from the non-laboratory-based Framingham 10-year cardiovascular risk score, which was chosen for its relevance in and out of clinical settings. The Framingham risk score was developed for adults between the ages of 30 and 74 years old based on sex, systolic blood pressure (SBP), height, weight, smoking status, anti-hypertensive medication use, and diabetes status^{1, 2}. Using the Framingham risk score, biological heart age represents the age at which someone would have the same 10-year cardiovascular as someone of the same sex with normal health characteristics. Using this method, normal health was defined as a non-smoker with a SBP of 125 mmHg, BMI of 22.5 kg/m², who does not have diabetes or use antihypertensive medication¹. (The cardiovascular disease risk calculator may be accessed here: <https://www.framinghamheartstudy.org/fhs-risk-functions/cardiovascular-disease-10-year-risk/>).

Heart age difference was defined as biological heart age minus chronological heart age. A biological heart age less than the chronological age therefore resulted in a negative heart age difference that was indicative of someone with lower than expected risk for their age. Conversely, a biological heart age greater than the chronological heart age, which resulted in a positive heart age difference, indicated greater cardiovascular risk.

Thus, a hypothetical 40-year-old woman, with a SBP of 125 mmHg and BMI of 22.5 kg/m² who did not smoke, have diabetes, or require treatment for hypertension would have a biological heart age of 40. Her biological and chronological ages would be the same, and her heart age difference would be 0. If her SBP were 110 mmHg instead of 125 mmHg but the rest of her risk factors remained normal, then her biological heart age would be 35, and her heart age difference would be -5. If instead she was taking medication for hypertension, but all other characteristics remained normal, then her biological heart age would be 45 and her heart age difference would be 5. *Variable Ascertainment*

Components of biological heart age were assessed throughout follow up. Sex was self-reported. Participant age at each exam was calculated using date of birth, which was assessed by self-report. Smoking status and the use of anti-hypertensive medications were ascertained by self-report at each exam. Serum glucose concentrations were measured using the hexokinase method at Linco Research (St Charles, Missouri, USA) through exam year 20 after which they were conducted at the Collaborative Studies Clinical Laboratory,

University of Minnesota. Hemoglobin A1c (HbA1c) was determined at the University of Minnesota (Minneapolis, Minnesota, USA) using the Tosoh G7 high-performance liquid chromatography instrument at exam years 25 and 30. Diabetes status was based on fasting glucose (≥ 126 mg/dL), 2-hour oral glucose tolerance test (OGTT ≥ 200 mg/dL), HbA1c ($\geq 6.5\%$), or the use of medication for diabetes when available. Study staff measured height, weight, and SBP using standard protocols³.

Additional CVD risk factors of interest were also assessed throughout the CARDIA study. Race was assessed by self-report at baseline and confirmed at exam year 2. Alcohol consumption, education, and marital status were obtained by self-report at all exams. Education was recorded as highest grade completed. Marital status was categorized as married or cohabiting, single, or separated, divorced or widowed. Physical activity was characterized using a modified Baecke questionnaire³. Plasma total cholesterol, high-density lipoprotein cholesterol (HDL-c) and triglyceride levels were measured using enzymatic methods; HDL-c levels were measured after dextran-sulfate-magnesium precipitation of other lipoproteins. Low-density lipoprotein cholesterol (LDL-c) levels were estimated with the Friedewald equation for individuals with fasting triglyceride values less than 400 mg/dL³⁻⁶. Urinary albumin-creatinine ratio was measured at exam years 10 through 25 with similar methods using a nephelometry-based assay by Dade Behring using a BN-II instrument. Detailed information on CARDIA study protocols have been described previously³.

Statistical Analysis

Heart age difference was the outcome of interest. Continuous demographic and health characteristics were assessed using univariate analyses to examine normality and determine if transformation was required. Data were presented as means and standard deviations for continuous variables and counts and percentages for categorical variables.

Linear regression was used to assess the associations between Heart age difference and individual risk factors. For each of the four exam years analyzed, multiple linear regression models were used to assess risk factors cross-sectionally. Repeated measures linear regression models with a Toeplitz correlation structure were used to examine longitudinal trends in the difference between biological and chronological age.

Variables included in the heart age derivation, specifically sex, SBP, height, weight, smoking status, anti-hypertensive medication use, and diabetes status, were not included in the models to avoid over-adjustment. Both cross-sectional and longitudinal linear regression models were conducted in three stages. First, crude models were run, then the models were adjusted for race, study center, total cholesterol, and urinary albumin-creatinine ratio, and finally for alcohol intake, physical activity, education, and marital status. All analyses were conducted using SAS 9.4, and statistical significance assessed as two-tailed type 1 errors of 0.05.

Results

Descriptive characteristics of CARDIA participants included and excluded from these analyses are in table 1. Compared to the CARDIA participants who were not included in the analysis, the analytic sample had fewer black participants (p-value < 0.001, data not shown). The participants in the analytic sample also were less likely to be treated for hypertension and had lower mean SBP at exam year 0 compared to those not included in analysis (p-value < 0.001 and 0.0155 respectively, data not shown). Participants included in the analysis were more likely to be married and had more years of education on average compared to CARDIA participants not included in the analysis (p-values < 0.001, data not shown). Prevalence of diabetes, mean alcohol intake, and mean body mass index were lower in participants analyzed, but these differences were not statistically significant (p-values 0.7738, 0.0667, and 0.3028 respectively, data not shown).

Cross-sectional trends are presented in table 2. At exam year 10, on average, participants had lower biological ages than chronological ages, largely driven by low SBP and the low prevalence of diabetes and hypertension medication use. By exam year 15, biological age was greater than chronological age and this trend continued through exam years 20 and 25.

Mean systolic blood pressure, body mass index, and total cholesterol increased across the exams. The means for systolic blood pressure remained below 120 mmHg, body mass index remained below 30 kg/m² and total cholesterol remained below 200 mg/dL. Prevalence of diabetes and antihypertensive medication use were higher at exam year 25 compared to exam year 10. At exam year 10, prevalence of diabetes and antihypertensive medication use were 2.1 and 2.6 % respectively. At exam year 25, 11.9 and 27.1% of participants reported diabetes and antihypertensive medication use respectively. Fewer participants reported being current smokers at exam year 25 (14.4%) than exam year 10 (21.5%).

Cross-sectional associations between heart age difference and many risk factors were largely consistent among the exams (Table 3). Total cholesterol and UACR were positively associated with heart age difference. Total cholesterol was statistically significantly associated with heart age difference at exam years 15, 20, and 25 (all p-values < 0.0001). UACR was statistically significantly associated with heart age difference at exam years 15, 20, and 25 (all p-values < 0.0001).

Race was associated with significant excess heart age after adjusting for all other variables and at all exams. At exam year 10, biological heart age exceeded chronological age for black participants by 2.6 years on average compared to whites (p-value < 0.001). By exam year 25, this difference was 4.7 years (p-value < 0.001). Higher physical activity and education were associated with lower biological heart age compared to chronological age. Education was highly statistically significant at all four exams (all p-values < .0001) with each additional year of education associated with between 0.5 and 0.8 lower biological heart ages on average. Physical activity was not statistically significant at exam year 10 but was significant at all subsequent exams (p-values = 0.0043, 0.0004, and 0.0153 respectively). Overall, marital status was statistically significant at exam years 15 and 25, but not in the remaining exams (p-value = 0.0567, 0.0182, 0.0987, and 0.0098 respectively). Being single was

associated with a positive heart age difference at exam years 10 and 25 (p-values = 0.0265 and 0.0053 respectively) and being divorced, separated, or widowed was associated positive heart age difference at exam year 15 (p-value = 0.0084) compared to being married (Table 3).

Longitudinal trends in heart age difference are shown in table 4. After adjusting for other risk factors, black participants had on average biological heart ages 4.8 years older than their chronological ages over 15 years of examinations. This difference was highly statistically significant (p-value < .0001). Total cholesterol was associated with -0.01 less years of aging per unit increment (p-value = 0.0415). UACR and physical activity were associated with 0.32 and -0.45 additional years of aging for each standard deviation increment (p-values < 0.0001) across the exams analyzed. Alcohol intake was associated with a 0.02 higher heart age difference (p-value < 0.0001). Each additional year of education was associated with an average of 0.37 lower years of biological age compared to chronological age, which was statistically significant after adjusting for other risk factors (p-values < .0001). Being single or never married compared to married was associated with greater heart age difference (p-value = 0.0058), but being divorced was not a significant predictor of heart age difference (p-value = 0.3735) (Table 4).

Discussion

Black participants, men, and those with lower levels of education had positive heart age differences, which correspond to higher biological heart age relative to their actual age. The results of these analyses are consistent with previous findings in BRFSS², NHANES⁷, and other analyses of the prevalence of cardiovascular risk factors in the US⁸⁻¹⁰. Of particular importance, these analyses support the presence the persistent racial disparities in cardiovascular disease risk present in the United States.

Additionally, these analyses demonstrated that biomarkers of disease states and lifestyle factors not included in the Framingham cardiovascular risk calculator were associated with positive heart age differences. Although these risk factors did not directly contribute to the excess heart age, the results suggest that subpopulations within the analytical sample had different risk profiles.

At year 10, the mean systolic blood pressure, which is present in the Framingham risk score, and mean total cholesterol, which is not present in the non-laboratory Framingham CVD risk equation, were not in ranges considered abnormal^{11,12}. However, these risk factors remain associated with excess heart age. Together these findings suggest targeting individual biological risk factors might not be sufficient to reduce biological heart age and cardiovascular risk or that there is a continuum of risk within the range of values considered normal.

We observed an unexpected finding in longitudinal analyses that total cholesterol was inversely associated with heart age difference. In cross-sectional analyses, total cholesterol was positively associated with heart age difference at exam years 10 and 15. At exam year 20, this association was not statistically significant. By exam year 25, total cholesterol was

inversely related to heart age difference, which appears to drive the significant inverse association in the longitudinal analyses. Neither the Framingham risk score nor the models presented were adjusted for lipid-lowering medication use, which increased from less than 1% in all race and sex subgroups at exam year 10 to between 10.4% for white women and 20.0% for white men at exam year 25. Additionally, by exam year 25, these medications were more likely to be the highly effective HMG CoA reductase inhibitor class rather than older, less effective medications. Previous analyses of CARDIA data showed that lipid-lowering medication use and obesity, which was present in the Framingham risk score, had opposing relationships with mean total cholesterol over time resulting in stable or lower total cholesterol throughout exams¹³.

The ability to clearly communicate cardiovascular risk is important in clinical settings. Absolute risk, as presented in Framingham and other risk scores may be difficult for individuals without health backgrounds to understand^{14,15}. The use of biological heart age compared to risk score to characterize and inform individual risk of cardiovascular disease demonstrated greater improvement in risk scores in a trial on motivating changes in lifestyle factors¹⁶. Therefore, understanding the impact of cardiovascular risk factors on biological heart age may improve the ability to communicate risk¹⁷.

Limitations

The entire CARDIA cohort had more Black participants (51.6% compared to 40.1%) and fewer never smokers (55.9 compared to 60.7%) than the analytic cohort, which was comprised of participants present at year 10 who also attended the next three study visits. Therefore, the sample analyzed may not be generalizable to the entire cohort. However, the results presented are expected to be conservative.

The Framingham 10-year cardiovascular disease (CVD) risk calculator has a maximum biological heart age value of 86. At exam year 25, 73 participants reached the maximum calculated biological heart age. As a result, the true biological heart ages for these individuals may be underestimated. Additionally, current blood pressure guidelines suggest a target systolic blood pressure below the value considered normal in the calculator¹¹. Finally, in the calculator, continuous risk factors were modeled as linear variables in the model.

Strengths

CARDIA is a longitudinal cohort with high participant retention. Participants were enrolled as young adults and have been followed through middle age. Variable ascertainment within study visits is also high. Biomarkers and risk factors were assessed at all study visits using standard protocols. Direct measurement of SBP and diabetes are also more accurate than self-report, which improves our analyses relative to previous analysis using BRFSS data. Overall, the study methods provide very good quality data.

Conclusions

These analyses of a population-based cohort demonstrated that overall, biological heart age exceeds chronological heart age in a sample of young and middle-aged adults. During exam years 10, 15, 20, and 25, 34.9, 43.1, 53.9, and 45.6%, respectively, of the available cohort

had biological heart ages that exceeded their chronological age. Across all exam years examined here, 62.5% of participants had a positive heart age difference.

Modifiable risk factors, both in the Framingham risk calculator and additional risk factors examined in these analyses, offer an opportunity to potentially reduce biological age. These analyses show that in a sample of black and white men and women, average biological heart age exceeds chronological age when individuals are in their early 40s. These results emphasize the importance of prevention prior to this transition. Overall, biological heart age provides valuable information about cardiovascular health using easy-to-obtain variables that can be obtained at home or in clinical settings.

Acknowledgements

The Coronary Artery Risk Development in Young Adults Study (CARDIA) is supported by contracts HHSN268201300025C, HHSN268201300026C, HHSN268201300027C, HHSN268201300028C, HHSN268201300029C, and HHSN268200900041C from the National Heart, Lung, and Blood Institute (NHLBI), the Intramural Research Program of the National Institute on Aging (NIA), and an intra-agency agreement between NIA and NHLBI (AG0005).

List of Abbreviations and Acronyms

BMI	body mass index
BRFSS	Behavioral Risk Factor Surveillance System
CARDIA	Coronary Artery Risk Development in Young Adults
CVD	cardiovascular disease
HDL-c	high-density lipoprotein cholesterol
LDL-c	low-density lipoprotein cholesterol
SBP	systolic blood pressure
SES	socioeconomic status
UACR	urinary albumin-creatinine ratio

References

1. D'Agostino RB Sr, Vasan RS, Pencina MJ, Wolf PA, Cobain M, Massaro JM, Kannel WB. General cardiovascular risk profile for use in primary care: the Framingham Heart Study. *Circulation*. 2008 2 12;117(6):743–53. [PubMed: 18212285]
2. Yang Q, Zhong Y, Ritchey M, Cobain M, Gillespie C, Merritt R, Hong Y, George MG, Bowman BA. Vital Signs: Predicted Heart Age and Racial Disparities in Heart Age Among U.S. Adults at the State Level. *MMWR Morb Mortal Wkly Rep*. 2015 9 4;64(34):950–8. [PubMed: 26335037]
3. Friedman GD, Cutter GR, Donahue RP, Hughes GH, Hulley SB, Jacobs DR Jr, Liu K, Savage PJ. CARDIA: study design, recruitment, and some characteristics of the examined subjects. *J Clin Epidemiol*. 1988;41(11):1105–16. [PubMed: 3204420]
4. Warnick GR. 1986 Enzymatic methods for quantification of lipoprotein lipids. *Meth Enzymol* 129:101–123. [PubMed: 3724535]
5. Warnick GR, Benderson J, and Albers JJ. Dextran sulfate-Mg²⁺ precipitation procedure for quantitation of high-density lipoprotein cholesterol. *Clin Chem* 28:1379–1388

6. Friedewald WT, Levy RI, Frederickson DS. Estimation of the concentration of low-density lipoprotein cholesterol in plasma, without use of the preparative ultracentrifuge. *Clin Chem.* 1972; 18(6):499–502. [PubMed: 4337382]
7. Hirsch JR, Waits G, Li Y, Soliman EZ. Racial Differences in Heart Age and Impact on Mortality. *J Natl Med Assoc.* 2018 4;110(2):169–175. [PubMed: 29580451]
8. Ford ES, Greenlund KJ, Hong Y. Ideal cardiovascular health and mortality from all causes and diseases of the circulatory system among adults in the United States. *Circulation* 2012;125:987–95. [PubMed: 22291126]
9. Yang Q, Cogswell ME, Flanders WD, et al. Trends in cardiovascular health metrics and associations with all-cause and CVD mortality among US adults. *JAMA* 2012;307:1273–83. [PubMed: 22427615]
10. Appiah D, Capistrant BD. Cardiovascular Disease Risk Assessment in the United States and Low- and Middle-Income Countries Using Predicted Heart/Vascular Age. *Sci Rep.* 2017;7(1):16673 [PubMed: 29192146]
11. Whelton PK, Carey RM, Aronow WS, Casey DE Jr, Collins KJ, Dennison Himmelfarb C, DePalma SM, Gidding S, Jamerson KA, Jones DW, MacLaughlin EJ, Muntner P, Ovbigele B, Smith SC Jr, Spencer CC, Stafford RS, Taler SJ, Thomas RJ, Williams KA Sr, Williamson JD, Wright JT Jr. 2017 ACC/AHA/AAPA/ABC/ACPM/AGS/APhA/ASH/ASPC/NMA/PCNA Guideline for the Prevention, Detection, Evaluation, and Management of High Blood Pressure in Adults: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *Hypertension.* 2017 11 13 pii: HYP.0000000000000065.
12. Stone NJ, Robinson JG, Lichtenstein AH, Bairey Merz CN, Blum CB, Eckel RH, Goldberg AC, Gordon D, Levy D, Lloyd-Jones DM, McBride P, Schwartz JS, Shero ST, Smith SC Jr, Watson K, Wilson PW; American College of Cardiology/American Heart Association Task Force on Practice Guidelines. 2013 ACC/AHA guideline on the treatment of blood cholesterol to reduce atherosclerotic cardiovascular risk in adults: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. *J Am Coll Cardiol.* 2014 7 1;63(25 Pt B):2889–934. [PubMed: 24239923]
13. Schreiner PJ, Jacobs DR Jr, Wong ND, Kiefe CI. Twenty-Five Year Secular Trends in Lipids and Modifiable Risk Factors in a Population-Based Biracial Cohort: The Coronary Artery Risk Development in Young Adults (CARDIA) Study, 1985–2011. *J Am Heart Assoc.* 2016 7 5;5(7). pii: e003384.
14. Soureti A, Hurling R, Murray P, van Mechelen W, Cobain M. Evaluation of a cardiovascular disease risk assessment tool for the promotion of healthier lifestyles. *Eur J Cardiovasc Prev Rehabil* 2010;17:519–23. [PubMed: 20195154]
15. Groenewegen K, den Ruijter H, Pasterkamp G, Polak J, Bots M, Peters SA. Vascular age to determine cardiovascular disease risk: a systematic review of its concepts, definitions, and clinical applications. *Eur J Prev Cardiol* 2015 [Epub ahead of print].
16. Lopez-Gonzalez AA, et al. Effectiveness of the Heart Age tool for improving modifiable cardiovascular risk factors in a Southern European population: a randomized trial. *European journal of preventive cardiology.* 2015;22:389–396 [PubMed: 24491403]
17. Webster R, Heeley E. Perceptions of risk: understanding cardiovascular disease. *Risk Manag Healthc Policy.* 2010;3:49–60. [PubMed: 22312218]

Table 1:

Characteristics of CARDIA Cohort and Analytic Sample at Exam Year 0

Variable	Entire Cohort (N=5,114)		Analytic Sample (N=2,264)	
Age, mean (SD), years *	24.8	(3.7)	25.9	(3.1)
Male, N (%)	2327	(45.5)	1006	(44.4)
Black, N (%) *	2637	(51.6)	908	(40.1)
Center, N (%)				
Birmingham	1178	(23.0)	530	(23.4)
Chicago	1108	(21.7)	510	(22.5)
Minneapolis	1402	(27.4)	619	(27.3)
Oakland	1426	(27.9)	605	(26.7)
Education, mean (SD), years *	13.8	(2.3)	14.4	(2.2)
Marital status, N (%) *				
Married/cohabiting	1137	(22.3)	607	(26.8)
Single	3530	(69.1)	1456	(64.4)
Separated/divorced/widowed	440	(8.6)	199	(8.8)
Physical activity, mean (SD), exercise units	420.1	(300.7)	419.2	(297.3)
Alcohol, mean (SD), mL/day	12.1	(22.0)	11.9	(19.6)
Current smoker, N (%) *	1546	(30.5)	547	(24.3)
Diabetes, N (%)	43	(0.9)	20	(0.9)
Hypertension medication, N (%)	115	(2.3)	46	(2.0)
Systolic blood pressure, mean (SD), mmHg *	110.4	(10.9)	110.0	(10.7)
Body mass index, mean (SD), kg/m²	24.5	(5.0)	24.4	(4.7)
Total cholesterol, mean (SD), mg/dL	176.7	(33.5)	177.7	(32.5)
HDL-c, mean (SD), mg/dL	53.2	(13.2)	53.3	(12.8)
LDL-c, mean (SD), mg/dL	109.1	(31.2)	110.0	(30.4)

Note: Variables in bold are in the heart age calculator;

* denotes variables that were statistically significantly different in participants who were and were not included in the analytic sample (p-value < 0.05)

Table 2:

Cross-Sectional Biological and Lifestyle Risk Factors in CARDIA for Exam Year 10 through Year 25

Variable	Year 10	Year 15	Year 20	Year 25
Chronological age, mean, years	35.9	41.0	46.0	50.9
Biological heart age, mean, years	35.4	42.0	49.7	53.1
Heart age difference, mean, years	-0.6	1.0	3.7	2.2
Education, mean, years	15.0	15.2	15.2	15.3
Marital status, %				
Married/cohabiting	54.7	62.5	65.5	64.6
Single	29.5	20.0	16.6	15.1
Separated/divorced/widowed	15.8	17.4	18.0	20.4
Physical Activity, mean (SD), exercise units	337.4	351.8	346.6	346.2
Alcohol, mean, mL/day	10.7	10.7	11.5	11.9
Current smoker, %	21.5	18.7	16.8	14.4
Diabetes, %	2.1	3.1	8.7	11.9
Hypertension medication, %	2.6	7.1	16.9	27.1
Systolic blood pressure, mean, mmHg	109.2	112.7	116.3	119.5
Body mass index, mean, kg/m²	27.2	28.4	29.2	29.7
Total cholesterol, mean, mg/dL	178.8	185.1	186.3	192.9
Urine albumin/creatinine ratio, mean, mg/g	13.2	14.3	12.8	12.0

Note: Variables in bold are in the heart age calculator

Cross-Sectional Multivariable Adjusted Biological and Lifestyle Risk Factors Associated with Heart Age Difference in CARDIA for Exam Year 10 through Year 25

Table 3:

Variable	Year 10		Year 15		Year 20		Year 25	
	Heart Age Difference	p-value	Heart Age Difference	p-value	Heart Age Difference	p-value	Heart Age Difference	p-value
Black (compared to white)	2.56	<.0001	3.56	<.0001	4.99	<.0001	4.73	<.0001
Total cholesterol, mg/dL	0.03	<.0001	0.03	<.0001	0.01	0.0563	-0.04	<.0001
UACR, mg/g	0.24	0.0656	1.46	<.0001	1.45	<.0001	1.83	<.0001
Alcohol, mL/day	0.05	<.0001	0.06	<.0001	0.04	<.0001	0.04	<.0001
PA, mean (SD), exercise units	0.21	0.1217	-0.48	0.0043	-0.78	0.0004	-0.57	0.0153
Education, years	-0.54	<.0001	-0.62	<.0001	-0.83	<.0001	-0.77	<.0001
Single*	0.69	0.0265	0.71	0.0989	1.05	0.0777	1.86	0.0053
Separated/divorced/ widowed*	0.60	0.1248	1.22	0.0084	0.96	0.1062	-0.29	0.6237

Note: PA = physical activity,

* compared to married; estimates after adjusting for all other variables and study center; heart age difference is measured in years; per 1 standard deviation (SD) increment where SD = 135.0, 127.6, 57.2, and 40.1 mg/g for UACR and SD = 269.5, 277.5, 275.4, and 279.5 exercise units for physical activity at the four exams respectively

Table 4:

Adjusted Longitudinal Trends in Biological and Social Risk Factors Associated with Heart Age Difference in CARDIA from Exam Year 10 through Year 25

Variable	Univariate		Multivariable	
	Heart Age Difference	p-value	Heart Age Difference	p-value
Black (compared to white)	5.61	<.0001	4.79	<.0001
Total Cholesterol, mg/dL	−0.01	0.0116	−0.01	0.0415
UACR, mg/g	0.33	<.0001	0.32	<.0001
Alcohol, mL/day	0.02	<.0001	0.02	<.0001
Physical activity, exercise units	−0.55	<.0001	−0.45	<.0001
Education, years	−0.53	<.0001	−0.37	<.0001
Marital status *				
Single	1.08	<.0001	0.73	0.0058
Separated/divorced/widowed	0.58	0.0071	0.20	0.3735

Note:

* compared to married; estimates after adjusting for all other variables, also adjusted for center and exam year; heart age difference is measured in years; per 1 standard deviation (SD) increment where SD = 98.0 mg/g for UACR and SD = 275.5 exercise units for physical activity