Published in final edited form as:

J Org Chem. 2018 September 21; 83(18): 11359–11368. doi:10.1021/acs.joc.8b01590.

Synthesis of Halogenated Anilines by Treatment of *N,N*-Dialkylaniline *N*-Oxides with Thionyl Halides

Hayley Reed, Tyler R. Paul, William J. Chain*

Department of Chemistry & Biochemistry University of Delaware Newark, Delaware 19716, United States

Abstract

The special reactivity of *N,N*-dialkylaniline *N*-oxides allows practical and convenient access to electron-rich aryl halides. A complementary pair of reaction protocols allow for the selective *para*-bromination or *ortho*-chlorination of *N,N*-dialkylanilines in up to 69% isolated yield. The generation of a diverse array of halogenated anilines is made possible by a temporary oxidation level increase of *N,N*-dialkylanilines to the corresponding *N,N*-dialkylaniline *N*-oxides and the excision of the resultant weak *N*-*O* bond via treatment with thionyl bromide or thionyl chloride at low temperature.

Graphical Abstract

Halogenated aromatic rings are ubiquitous in various disciplines of the organic chemistry field. Aryl halides are critically important synthetic building blocks for use in cross-coupling chemistries,¹ lithium–halogen exchange reactions,² and classical Grignard reactions.³ Collectively, these methods represent some of the most highly utilized means of constructing carbon–carbon bonds in synthesis. Aryl halides are also common substructures in synthetic and naturally occurring biologically active molecules and constitute some end-user industrial materials as well.⁴ Classical methods of preparing aryl halides (e.g., electrophilic aromatic substitution) are prone to complications in electron-rich settings;^{5,6} aniline and phenol substrates are highly reactive toward electrophilic halogenation reagents, often compromising the regioselectivity or stoichiometry of the resultant reactions.⁷ Thus, new synthetic methods for the regioselective halogenation of electron-rich aromatics are of high utility in synthetic organic chemistry.

* Corresponding Author: wchain@udel.edu.

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acs.joc.8b01590.

¹H and ¹³C NMR spectra of all compounds (PDF)

The authors declare no competing financial interest.

We have a longstanding interest in the manipulations of anilines at elevated oxidation levels (Scheme 1).⁸ The functionalization of anilines by group transfer is known at the *N*-arylhydroxylamine oxidation level (Scheme 1, eqs 1 and 2),⁹ and a small number of examples of *ortho*-chlorination reactions have been reported on that platform (Scheme 1, eq3).¹⁰ We report here the practical and convenient syntheses of 2-chloro-*N,N*-dialkylanilines and 4-bromo-*N,N*-dialkylanilines as part of our ongoing interests in the reactivity of aniline *N*-oxides (Scheme 1, eq 4). In our previous work, we demonstrated that *N,N*-dimethylaniline *N*-oxides can be acylated with a variety of electrophiles, and the resultant *O*-acylated materials can undergo a group transfer event by which a variety of C–O, C–N, and C–C bonds are generated.^{8a} The materials so derived constitute a diverse array of useful materials that are otherwise unavailable by traditional synthetic methods (e.g., electrophilic aromatic substitution). The mechanistic pathways of these group-transfer events are ambiguous; possibilities include [3,3]-sigmatropic rearrangements in which the N–O bond is excised as a C–X bond is formed or stepwise processes in which the N–O bond is excised homolytically followed by a radical recombination in which the C–X bond is formed. We envisioned that monohalogenation of the aniline platform could be achieved by treatment of aniline *N*-oxides with thionyl halides or phosphorus oxyhalides; transfer of the halogen atom would result in expulsion of sulfur dioxide or metaphosphoryl halide.

We quickly determined that treatment of the aniline *N*-oxide **1a** with phosphorus oxychloride could generate the corresponding aryl chloride but only in trace yields. However, treatment of aniline *N*-oxide **1a** with thionyl halides in dichloromethane followed by base cleanly afforded the corresponding aryl halides at low temperature (Table 1, entry 1). For example, treatment of *N,N*-dimethylaniline *N*-oxide **1a** with 1 equiv of thionyl bromide in dichloromethane at –78 °C for 4 h followed by addition of 4 equiv of triethylamine and warming to room temperature afforded the corresponding 4-bromo-*N,N*-dimethylaniline **2a** in 31% yield after an aqueous workup. The regioisomeric aryl chloride **3a** could be obtained in 25% yield by treatment of **1a** with thionyl chloride under otherwise identical conditions (Table 1, entry 1). Isolated yields were diminished when the reaction temperature was warmed above or cooled below –78 °C (Table 1, entries 2 and 3). Unlike our previous work with aniline *N*-oxides, we found the halogenation reactions to be highly sensitive to solvent. Reactions conducted in tetrahydrofuran were routinely more efficient; treatment of **1a** with thionyl halides in THF at –78 °C followed by addition of 4 equiv of triethylamine and warming to room temperature afforded a 55% yield of the aryl bromide **2a** and a 35% yield of the aryl chloride **3a** (Table 1, entry 5). Interestingly, addition of 1 equiv of thionyl chloride in two portions over a period of 4 h under otherwise identical reaction conditions markedly improved the isolated yield of the aryl chloride **3a** (49%, Table 1, entry 6). No such advantage was apparent for reactions with thionyl bromide. Most interestingly, we found the halogenation reactions to be highly regioselective; treatment of *N,N*-dimethylaniline *N*-oxides with thionyl bromide results in the corresponding 4-bromo-*N,N*-dimethylanilines exclusively, while treatment with thionyl chloride results in predominately the corresponding 2-chloro-*N,N*-dimethylanilines (3.6–6.6:1 2-chloro/4-chloro).

Utilizing our optimized reaction conditions, we generated a variety of halogenated anilines, and we found the selectivity profiles to be general. Treatment of diversely substituted *N,N*-

dialkylaniline *N*-oxides (including *N,N*-dimethyl-1-naphthylamine *N*-oxide, and *N*-methyl-tetrahydroquinoline *N*-oxide) with thionyl bromide under the optimized reaction conditions routinely afforded 4-bromo-*N,N*-dialkylanilines **2a–t** (including 4-bromo-*N,N*-dimethyl-1-naphthylamine **2d** and 7-bromo-*N*-methyltetrahydroquinoline **2t**) in up to 69% yield (Table 2). In all cases, we observed no evidence for the formation of the corresponding 2-bromo-substituted equivalents; however, if we blocked that position by subjecting 4-substituted substrates to our reaction conditions, we could observe 2-bromo-substituted products in reduced yields (Table 2, **2k–o**).

Treatment of a similar set of *N,N*-dialkylaniline *N*-oxides (including *N,N*-dimethyl-1-naphthylamine *N*-oxide and *N*-methyl-tetrahydroquinoline *N*-oxide) with thionyl chloride under the optimized reaction conditions routinely afforded 2-chloro-*N,N*-dialkylanilines **3a–j** (including 2-chloro-*N,N*-dimethyl-1-naphthylamine **3d** and 9-chloro-*N*-methyl-tetrahydroquinoline **3o**) in up to 65% yield with up to 6.7:1 (2-Cl/4-Cl) selectivity (Table 3), though some 2-substituted anilines are problematic in chlorination reactions (e.g., **3k**).

Together, these reactions comprise a complementary set of synthetic tools to access regioselectively monohalogenated anilines that are otherwise difficult to obtain. The elevated reactivity of aniline *N*-oxides allowed us to overcome the difficulties described in the chlorination of the equivalent *N*-arylhydroxylamines. In contrast to the few reported examples of halogenation of *N*-arylhydroxylamines, and consistent with our prior work,⁸ these halogenation reactions are general and conducted at low temperature, which renders the reactions more amenable to sensitive substrates. However, the elevated reactivity of aniline *N*-oxides comes at the cost of mechanistic ambiguity. The manipulations described above employing *N*-arylhydroxylamines almost certainly proceed by [3,3]-sigma-tropic rearrangement pathways;^{9–11} however, reactions with aniline *N*-oxides may proceed by multiple mechanisms. While the 2-haloaniline products are consistent with a sigmatropic rearrangement, the 4-haloaniline products must arise by a radical pathway or a nucleophilic aromatic substitution-type mechanism. We conducted bromination reactions in the presence of 2,6-di-*tert*-butyl-4-methylphenol (BHT) under otherwise identical conditions in order to probe the involvement of radical species. We observed no significant change in the isolated yield of 4-bromoaniline product; for example, treatment of **1a** with 1 equiv of thionyl bromide and 20 mol % of BHT in tetrahydrofuran at –78 °C for 4 h, followed by treatment with triethylamine (4 equiv), afforded the 4-bromo-*N,N*-dimethylaniline **2a** in 55% yield (cf. Table 2, 55% yield). Analogous experiments with other radical traps or tool compounds such as (2,2,6,6-tetramethylpiperidin-1-yl)-oxyl (TEMPO) afforded no conclusive data. Interestingly, bromination reactions conducted in the presence of LiCl additives and chlorination reactions conducted in the presence of LiBr additives result in scrambled halogenation products. These results are consistent with simple halogen exchange of the reagent, halogen exchange with intermediates such as those shown in Scheme 1, eq 4, or the involvement of nucleophilic aromatic substitution-type mechanisms. Efforts to fully elucidate the mechanistic pathway of these reactions is still underway but in any case, the synthetic building blocks available by these methods are widely utilized for a myriad of applications in synthetic organic chemistry.

The special reactivity of *N,N*-dialkylaniline *N*-oxides allows for the efficient, controlled, and regioselective halogenation of aromatic systems without exogenous activation of halogen sources, Lewis acids, or other exotic reagents. We are working to further probe the mechanisms of these transformations, and exploit further chemistries enabled by the weak *N*-*O* bond within aniline *N*-oxides.

EXPERIMENTAL SECTION

General Information.

Commercial reagents and solvents were used as received with the following exceptions. Triethylamine, dichloromethane, ethyl ether, dimethyl sulfoxide, tetrahydrofuran, hexane, toluene, *N,N*-dimethylformamide, and benzene were purified by the method of Pangborn et al.¹² Thionyl chloride (SOCl₂) was purified by distillation over calcium hydride prior to use. Thionyl bromide was used without further purification. All reactions were performed in single-neck oven- or flame-dried round-bottom flasks fitted with rubber septa under a positive pressure of nitrogen, unless otherwise noted. Air- and moisture-sensitive liquids were transferred via syringe or stainless steel cannula. Organic solutions were concentrated by rotary evaporation at or below 35 °C at 10 Torr (diaphragm vacuum pump) unless otherwise noted. Proton (¹H) and carbon (¹³C) nuclear magnetic resonance (NMR) spectra were recorded on Bruker AV400 CryoPlatform QNP or Bruker AVIII600 SMART NMR spectrometers at 23 °C. Proton chemical shifts are expressed in parts per million (ppm, δ scale) downfield from tetramethylsilane and are referenced to residual protium in the NMR solvent (CHCl₃: δ 7.26, C₆H₅D₅: δ 7.16). Carbon chemical shifts are expressed in parts per million (ppm, δ scale) downfield from tetramethylsilane and are referenced to the carbon resonance of the NMR solvent (CDCl₃: δ 77.16, C₆D₆: δ 128.06). Data are represented as follows: chemical shift, multiplicity (s = singlet, d = doublet, t = triplet, q = quartet, m = multiplet, br = broad, app = apparent), integration, and coupling constant (*J*) in hertz (Hz). Accurate mass measurements were obtained using an Agilent 1100 quaternary LC system coupled to an Agilent 6210 LC/MSD-TOF fitted with an ESI or an APCI source, or Thermo Q-Exactive Orbitrap using electrospray ionization (ESI) or a Waters GCT Premier spectrometer using chemical ionization (CI). Compounds were isolated using flash column chromatography¹³ with silica gel (60-Å pore size, 40–63 μ m, standard grade, Silicycle) or basic alumina (60-Å pore size, 50–200 μ m, Brockmann I, Acros Organics). Analytical thin-layer chromatography (TLC) was performed using glass plates precoated with silica gel (0.25 mm, 60-Å pore size, 5–20 μ m, Silicycle) impregnated with a fluorescent indicator (254 nm). TLC plates were visualized by exposure to ultraviolet light (UV), then were stained by submersion in aqueous ceric ammonium molybdate solution (CAM), ethanolic phosphomolybdic acid solution (PMA), acidic ethanolic *p*-anisaldehyde solution (anisaldehyde), or aqueous potassium permanganate solution (KMnO₄), followed by brief heating on a hot plate (215 °C, 10–15 s).

General Procedure for the Formation of *N,N*-Dimethylanilines.

1-(*N,N*-Dimethylamino)naphthalene *N*-oxide (**4d**), 4-methoxy-*N,N*-dimethylaniline *N*-oxide (**4p**), and 4-methyl-*N,N*-dimethylaniline *N*-oxide (**4q**) were obtained using the procedure of

Lewis et al.⁸ All other anilines were obtained using the following procedure modified from Chandrasekharam et al.¹⁴

Glacial acetic acid (7.3 mL, 130 mmol, 5.4 equiv) was added dropwise to a mixture of 3-methoxyaniline (3.0 g, 24 mmol, 1 equiv), paraformaldehyde (3.9 g, 130 mmol, 5.4 equiv), and sodium cyanoborohydride (7.9 g, 130 mmol, 5.4 equiv) in tetrahydrofuran (100 mL) at 23 °C. The resultant mixture was heated to 50 °C and stirred at that temperature for 18 h. The reaction mixture was cooled to 23 °C and then was partitioned between saturated aqueous sodium bicarbonate solution (100 mL) and diethyl ether (40 mL). The layers were separated, and the aqueous layer was extracted with diethyl ether (2 × 40 mL). The combined organic layers were washed sequentially with water (2 × 60 mL) and saturated aqueous sodium chloride solution (3 × 40 mL). The combined organic layers were dried over anhydrous sodium sulfate, and the dried solution was concentrated. The resultant oily residue was purified by flash column chromatography (silica gel, starting with 5% ethyl acetate–hexanes, grading to 10% ethyl acetate–hexanes) to afford **4b** as a yellow oil (2.8 g, 18 mmol, 77% yield).

3-Methoxy-*N,N*-dimethylaniline (4b).—TLC 10% ethyl acetate–hexanes, R_f = 0.40 (UV, KMnO₄). ¹H NMR (400 MHz, CDCl₃) δ : 7.16 (t, J = 8.1 Hz, 1H), 6.37 (dd, J_1 = 8.3, J_2 = 2.3 Hz, 1H), 6.30 (m, 2H), 3.80 (s, 3H), 2.94 (s, 6H). ¹³C NMR (101 MHz, CDCl₃) δ : 160.7, 152.1, 129.8, 105.8, 101.4, 99.2, 55.2, 40.7. HRMS: ESI⁺ [M + H]⁺ calcd for C₉H₁₄ON 152.1075, found 152.1071.

3-Methyl-*N,N*-dimethylaniline (4c).—Obtained as a yellow oil (2.8 g, 75%). TLC 10% ethyl acetate–hexanes, R_f = 0.59 (UV, KMnO₄). ¹H NMR (400 MHz, CDCl₃) δ : 7.16 (m, 1H), 6.59 (m, 3H), 2.95 (s, 6H), 2.35 (s, 3H). ¹³C NMR (101 MHz, CDCl₃) δ : 150.9, 138.8, 129.0, 117.7, 113.6, 110.0, 40.8, 22.0. HRMS: ESI⁺ [M + H]⁺ calcd for C₉H₁₄N 136.1126, found 136.1122.

Methyl 3-*N,N*-Dimethylaminobenzoate (4e).—Obtained as a yellow oil (3.1 g, 87%). TLC 10% ethyl acetate–hexanes, R_f = 0.27 (UV, KMnO₄). ¹H NMR (400 MHz, CDCl₃) δ : 7.41–7.36 (m, 2H), 7.29 (m, 1H), 6.90 (m, 1H), 3.90 (s, 3H), 2.99 (s, 6H). ¹³C NMR (101 MHz, CDCl₃) δ : 167.9, 150.5, 130.9, 129.1, 117.6, 116.8, 113.3, 52.2, 40.7. HRMS: ESI⁺ [M + H]⁺ calcd for C₁₀H₁₄O₂N 180.1025, found 180.1018.

3-Bromo-*N,N*-dimethylaniline (4f).—Obtained as an orange-yellow oil (3.2 g, 92%). TLC 10% ethyl acetate–hexanes, R_f = 0.53 (UV, KMnO₄). ¹H NMR (400 MHz, CDCl₃) δ : 7.08 (t, J = 8.0 Hz, 1H), 6.83 (m, 2H), 6.63 (m, 1H), 2.94 (s, 6H). ¹³C NMR (101 MHz, CDCl₃) δ : 151.7, 130.3, 123.5, 119.1, 115.1, 111.0, 40.5. HRMS: ESI⁺ [M + H]⁺ calcd for C₈H₁₁NBr 200.0075, found 200.0071.

3-Chloro-*N,N*-dimethylaniline (4g).—Obtained as a yellow oil (2.50 g, 68%). TLC 10% ethyl acetate–hexanes, R_f = 0.56 (UV, KMnO₄). ¹H NMR (400 MHz, CDCl₃) δ : 7.14 (m, 1H), 6.69 (m, 2H), 6.59 (dd, J_1 = 8.5, J_2 = 2.4 Hz, 1H), 2.95 (s, 6H). ¹³C NMR (101 MHz, CDCl₃) δ : 151.6, 135.1, 130.1, 116.2, 112.3, 110.5, 40.5. HRMS: ESI⁺ [M + H]⁺ calcd for C₈H₁₁NCl 156.0580, found 156.0575.

2-Fluoro-N,N-dimethylaniline (4h).—Obtained as a red-orange oil (2.9 g, 77%). TLC 10% ethyl acetate–hexanes, R_f = 0.49 (UV, KMnO_4). ^1H NMR (400 MHz, CDCl_3) δ : 7.08–6.98 (m, 2H), 6.95–6.84 (m, 2H), 2.84 (s, 6H). ^{13}C NMR (101 MHz, CDCl_3) δ : 155.3 (d, J = 245.1 Hz), 140.9 (d, J = 8.6 Hz), 124.4 (d, J = 3.6 Hz), 121.3 (d, J = 7.8 Hz), 118.4 (d, J = 3.4 Hz), 116.2 (d, J = 20.9 Hz), 43.0 (d, J = 4.0 Hz). HRMS: ESI^+ $[\text{M} + \text{H}]^+$ calcd for $\text{C}_8\text{H}_{11}\text{NF}$ 140.0876, found 140.0869.

3-(Trifluoromethyl)-N,N-dimethylaniline (4i).—Obtained as a yellow oil (2.3 g, 65%). TLC 10% ethyl acetate–hexanes, R_f = 0.51 (UV, KMnO_4). ^1H NMR (400 MHz, CDCl_3) δ : 7.32 (t, J = 8.0 Hz, 1H), 6.94 (d, J = 7.6 Hz, 1H), 6.89 (m, 1H), 6.85 (dd, J_1 = 8.4 Hz, J_2 = 2.7 Hz, 1H), 3.00 (s, 6H). ^{13}C NMR (101 MHz, CDCl_3) δ : 150.5, 131.4 (q, J = 31.4 Hz), 129.5, 124.7 (q, J = 272.4 Hz), 115.2 (d, J = 1.5 Hz), 112.7 (q, J = 3.9 Hz), 108.5 (q, J = 4.0 Hz), 40.5. HRMS: ESI^+ $[\text{M} + \text{H}]^+$ calcd for $\text{C}_9\text{H}_{11}\text{NF}_3$ 190.0844, found 190.0834.

3-(N,N-Dimethylamino)benzonitrile (4j).—Obtained as a yellow oil (3.5 g, 95%). TLC 10% ethyl acetate–hexanes, R_f = 0.28 (UV, KMnO_4). ^1H NMR (400 MHz, CDCl_3) δ : 7.27 (m, 1H), 6.95 (m, 1H), 6.88 (m, 2H), 2.98 (s, 6H). ^{13}C NMR (101 MHz, CDCl_3) δ : 150.3, 129.8, 119.9, 119.5, 116.3, 114.8, 112.8, 40.3. HRMS: ESI^+ $[\text{M} + \text{H}]^+$ calcd for $\text{C}_9\text{H}_{11}\text{N}_2$ 147.0922, found 147.0917.

Methyl 4-N,N-Dimethylaminobenzoate (4k).—Obtained as a white solid (2.4 g, 63%). Mp: 93–95 °C. TLC 10% ethyl acetate–hexanes, R_f = 0.25 (UV, KMnO_4). ^1H NMR (400 MHz, CDCl_3) δ : 7.91 (d, J = 8.8 Hz, 2H), 6.65 (d, J = 8.7 Hz, 2H), 3.86 (d, J = 0.8 Hz, 3H), 3.04 (s, 6H). ^{13}C NMR (101 MHz, CDCl_3) δ : 167.8, 153.4, 131.4, 116.9, 110.8, 51.7, 40.2. HRMS: ESI^+ $[\text{M} + \text{H}]^+$ calcd for $\text{C}_{10}\text{H}_{14}\text{O}_2\text{N}$ 180.1025, found 180.1012.

4-Fluoro-N,N-dimethylaniline (4l).—Obtained as a yellow solid (2.9 g, 77%). Mp: 34–35 °C. TLC 10% ethyl acetate–hexanes, R_f = 0.34 (UV, KMnO_4). ^1H NMR (400 MHz, CDCl_3) δ : 6.95 (m, 2H), 6.68 (m, 2H), 2.90 (s, 6H). ^{13}C NMR (101 MHz, CDCl_3) δ : 155.70 (d, J = 235.1 Hz), 147.61 (d, J = 1.8 Hz), 115.50 (d, J = 22.0 Hz), 114.04 (d, J = 7.4 Hz), 41.53. HRMS: ESI^+ $[\text{M} + \text{H}]^+$ calcd for $\text{C}_8\text{H}_{11}\text{NF}$ 140.0876, found 140.0869.

4-Bromo-N,N-dimethylaniline (4m).—Obtained as a white solid (3.2 g, 90%). Mp: 29–31 °C. TLC 10% ethyl acetate–hexanes, R_f = 0.45 (UV, KMnO_4). ^1H NMR (400 MHz, CDCl_3) δ : 7.30 (d, J = 9.1 Hz, 2H), 6.59 (d, J = 9.1 Hz, 2H), 2.92 (s, 6H). ^{13}C NMR (101 MHz, CDCl_3) δ : 149.6, 131.8, 114.2, 108.6, 40.7. HRMS: ESI^+ $[\text{M} + \text{H}]^+$ calcd for $\text{C}_8\text{H}_{11}\text{NBr}$ 200.0075, found 200.0070.

4-Chloro-N,N-dimethylaniline (4n).—Obtained as a white solid (3.0 g, 83%). Mp: 30–32 °C. TLC 10% ethyl acetate–hexanes, R_f = 0.57 (UV, KMnO_4). ^1H NMR (400 MHz, CDCl_3) δ : 7.17 (d, J = 9.1 Hz, 2H), 6.64 (d, J = 9.1 Hz, 2H), 2.93 (s, 6H). ^{13}C NMR (101 MHz, CDCl_3) δ : 149.3, 128.9, 121.5, 113.7, 40.8. HRMS: ESI^+ $[\text{M} + \text{H}]^+$ calcd for $\text{C}_8\text{H}_{11}\text{NCl}$ 156.0580, found 156.0577.

4-(N,N-Dimethylamino)benzonitrile (4o).—Obtained as an orange-brown solid (2.4 g, 65%). Mp: 72–74 °C. TLC 10% ethyl acetate–hexanes, R_f = 0.29 (UV, KMnO_4). ^1H NMR

(400 MHz, CDCl_3) δ : 7.47 (d, $J = 9.0$ Hz, 2H), 6.64 (d, $J = 9.0$ Hz, 2H), 3.04 (s, 6H). ^{13}C NMR (101 MHz, CDCl_3) δ : 152.5, 133.5, 120.9, 111.5, 97.4, 40.1. HRMS: $\text{ESI}^+ [\text{M} + \text{H}]^+$ calcd for $\text{C}_9\text{H}_{11}\text{N}_2$ 147.0922, found 147.0917.

2-Methyl-N,N-dimethylaniline (4p).—Obtained as a yellow oil (2.5 g, 65%). TLC 10% ethyl acetate–hexanes, $R_f = 0.65$ (UV, KMnO_4). ^1H NMR (400 MHz, CDCl_3): δ 7.20–7.13 (m, 2H), 7.07–7.02 (m, 1H), 6.96 (td, $J_1 = 7.4$ Hz, $J_2 = 1.3$ Hz, 1H), 2.71 (s, 6H), 2.34 (s, 3H). ^{13}C NMR (101 MHz, CDCl_3): δ 152.8, 132.2, 131.2, 126.5, 122.6, 118.4, 44.3, 18.5. HRMS: $\text{ESI}^+ [\text{M} + \text{H}]^+$ calcd for $\text{C}_9\text{H}_{14}\text{N}$ 136.1126, found 136.1120.

Procedures for the Formation of *N*-Alkyl-*N*-methylanilines.

Asymmetric anilines were obtained using the following procedure modified from Chandrasekharam et al.¹²

N-Butyl-N-methylaniline (4q).—Glacial acetic acid (4.8 mL, 84 mmol, 3.0 equiv) was added dropwise to a mixture of *N*-methylaniline (3.0 g, 28 mmol, 1 equiv), butyraldehyde (7.6 mL, 84 mmol, 3.0 equiv), and sodium cyanoborohydride (5.3 g, 84 mmol, 3.0 equiv) in tetrahydrofuran (100 mL) at 23 °C. The resultant mixture was heated to 50 °C and stirred at that temperature for 18 h. The reaction mixture was cooled to 23 °C and then was partitioned between saturated aqueous sodium bicarbonate solution (100 mL) and diethyl ether (40 mL). The layers were separated, and the aqueous layer was extracted with diethyl ether (2 $\times$ 40 mL). The combined organic layers were washed sequentially with water (2 $\times$ 60 mL) and saturated aqueous sodium chloride solution (3 $\times$ 40 mL). The combined organic layers were dried over anhydrous sodium sulfate, and the dried solution was concentrated. The resultant oily residue was purified by flash column chromatography (silica gel, starting with 1% ethyl acetate–hexanes, grading to 5% ethyl acetate–hexanes) to afford 4q as a yellow oil (2.3 g, 14 mmol, 50% yield). TLC 10% ethyl acetate–hexanes, $R_f = 0.75$ (UV, KMnO_4). ^1H NMR (400 MHz, CDCl_3): δ 7.28–7.19 (m, 2H), 6.76–6.64 (m, 3H), 3.37–3.27 (m, 2H), 2.93 (s, 3H), 1.57 (m, 2H), 1.36 (m, 2H), 0.96 (t, $J = 7.3$ Hz, 3H). ^{13}C NMR (101 MHz, CDCl_3): δ 149.4, 129.2, 115.8, 112.1, 52.6, 38.4, 28.9, 20.5, 14.2. HRMS: $\text{ESI}^+ [\text{M} + \text{H}]^+$ calcd for $\text{C}_{11}\text{H}_{18}\text{N}$ 164.1439, found 164.1433.

N-Isopentyl-N-methylaniline (4r).—Glacial acetic acid (4.8 mL, 84 mmol, 3.0 equiv) was added dropwise to a mixture of *N*-methylaniline (3.0 g, 28 mmol, 1 equiv), isovaleraldehyde (9.2 mL, 84 mmol, 3.0 equiv), and sodium cyanoborohydride (5.3 g, 84 mmol, 3.0 equiv) in tetrahydrofuran (100 mL) at 23 °C. The resultant mixture was heated to 50 °C and stirred at that temperature for 18 h. The reaction mixture was cooled to 23 °C and then was partitioned between saturated aqueous sodium bicarbonate solution (100 mL) and diethyl ether (40 mL). The layers were separated, and the aqueous layer was extracted with diethyl ether (2 $\times$ 40 mL). The combined organic layers were washed sequentially with water (2 $\times$ 60 mL) and saturated aqueous sodium chloride solution (3 $\times$ 40 mL). The combined organic layers were dried over anhydrous sodium sulfate, and the dried solution was concentrated. The resultant oily residue was purified by flash column chromatography (silica gel, starting with 1% ethyl acetate–hexanes, grading to 5% ethyl acetate–hexanes) to afford 4r as a yellow oil (2.2 g, 12 mmol, 43% yield). TLC 10% ethyl acetate–hexanes, $R_f =$

0.80 (UV, KMnO_4). ^1H NMR (400 MHz, CDCl_3): δ 7.27–7.20 (m, 2H), 6.69 (m, 3H), 3.37–3.29 (m, 2H), 2.92 (s, 3H), 1.61 (m, 1H), 1.46 (m, 2H), 0.96 (d, J = 6.6 Hz, 6H). ^{13}C NMR (101 MHz, CDCl_3): δ 149.4, 129.3, 115.9, 112.2, 51.1, 38.3, 35.1, 26.3, 22.8. HRMS: ESI^+ $[\text{M} + \text{H}]^+$ calcd for $\text{C}_{12}\text{H}_{20}\text{N}$ 178.1596, found 178.1589.

N-Cyclohexyl-N-methylaniline (4s).—Glacial acetic acid (9.2 mL, 160 mmol, 5.0 equiv) was added dropwise to a mixture of aniline (3.0 g, 32 mmol, 1 equiv), cyclohexanone (16.6 mL, 178 mmol, 5.6 equiv), and sodium cyanoborohydride (10.1 g, 161 mmol, 5.0 equiv) in tetrahydrofuran (100 mL) at 23 °C. The resultant mixture was heated to 50 °C and stirred at that temperature for 18 h. The reaction mixture was cooled to 23 °C and then was partitioned between saturated aqueous sodium bicarbonate solution (100 mL) and diethyl ether (40 mL). The layers were separated, and the aqueous layer was extracted with diethyl ether (2 $\times$ 40 mL). The combined organic layers were washed sequentially with water (2 $\times$ 60 mL) and saturated aqueous sodium chloride solution (3 $\times$ 40 mL). The combined organic layers were dried over anhydrous sodium sulfate, and the dried solution was concentrated. The resultant oily residue was purified by flash column chromatography (silica gel, starting with 5% ethyl acetate–hexanes, grading to 10% ethyl acetate–hexanes) to afford 5s as a yellow oil (2.7 g, 15 mmol, 47% yield). TLC 10% ethyl acetate–hexanes, R_f = 0.59 (UV, KMnO_4). ^1H NMR (400 MHz, CDCl_3): δ 7.16 (dd, J_1 = 8.4 Hz, J_2 = 7.4 Hz, 2H), 6.66 (t, J = 7.3 Hz, 1H), 6.62–6.57 (m, 2H), 3.52 (s, 1H), 3.27 (m, 1H), 2.07 (m, 2H), 1.77 (m, 2H), 1.70–1.62 (m, 1H), 1.45–1.31 (m, 2H), 1.29–1.09 (m, 3H). ^{13}C NMR (101 MHz, CDCl_3): δ 147.5, 129.4, 116.9, 113.2, 51.8, 33.6, 26.1, 25.2. HRMS: ESI^+ $[\text{M} + \text{H}]^+$ calcd for $\text{C}_{12}\text{H}_{18}\text{N}$ 176.1439, found 176.1430.

Glacial acetic acid (1.6 mL, 28 mmol, 4.9 equiv) was added dropwise to a mixture of 5s (1.0 g, 5.7 mmol, 1 equiv), paraformaldehyde (0.86 g, 29 mmol, 5.1 equiv), and sodium cyanoborohydride (1.8 g, 29 mmol, 5.1 equiv) in tetrahydrofuran (30 mL) at 23 °C. The resultant mixture was heated to 50 °C and stirred at that temperature for 18 h. The reaction mixture was cooled to 23 °C and then was partitioned between saturated aqueous sodium bicarbonate solution (30 mL) and diethyl ether (20 mL). The layers were separated, and the aqueous layer was extracted with diethyl ether (2 $\times$ 20 mL). The combined organic layers were washed sequentially with water (2 $\times$ 30 mL) and saturated aqueous sodium chloride solution (3 $\times$ 20 mL). The combined organic layers were dried over anhydrous sodium sulfate, and the dried solution was concentrated. The resultant oily residue was purified by flash column chromatography (silica gel, starting with 5% ethyl acetate–hexanes, grading to 10% ethyl acetate–hexanes) to afford 4s as a yellow oil (0.69 g, 3.6 mmol, 64% yield). TLC 10% ethyl acetate–hexanes, R_f = 0.61 (UV, KMnO_4). ^1H NMR (400 MHz, CDCl_3): δ 7.25–7.20 (m, 2H), 6.78 (d, J = 8.6 Hz, 2H), 6.71–6.66 (m, 1H), 3.61–3.53 (m, 1H), 2.78 (s, 3H), 1.87–1.75 (m, 4H), 1.69 (d, J = 13.0 Hz, 1H), 1.53–1.29 (m, 5H), 1.19–1.08 (m, 1H). ^{13}C NMR (101 MHz, CDCl_3): δ 150.27, 129.21, 116.29, 113.21, 58.21, 31.28, 30.17, 26.34, 26.07. HRMS: ESI^+ $[\text{M} + \text{H}]^+$ calcd for $\text{C}_{13}\text{H}_{20}\text{N}$ 190.1596, found 190.1589.

N-Methyl-1,2,3,4-tetrahydroquinoline (4t).—Glacial acetic acid (6.6 mL, 120 mmol, 5.1 equiv) was added dropwise to a mixture of 1,2,3,4-tetrahydroquinoline (3.0 g, 22 mmol, 1 equiv), paraformaldehyde (3.48 g, 120 mmol, 5.1 equiv), and sodium cyanoborohydride

(7.3 g, 120 mmol, 5.1 equiv) in tetrahydrofuran (100 mL) at 23 °C. The resultant mixture was heated to 50 °C and stirred at that temperature for 18 h. The reaction mixture was cooled to 23 °C and then was partitioned between saturated aqueous sodium bicarbonate solution (100 mL) and diethyl ether (40 mL). The layers were separated, and the aqueous layer was extracted with diethyl ether (2 × 40 mL). The combined organic layers were washed sequentially with water (2 × 60 mL) and saturated aqueous sodium chloride solution (3 × 40 mL). The combined organic layers were dried over anhydrous sodium sulfate, and the dried solution was concentrated. The resultant oily residue was purified by flash column chromatography (silica gel, starting with 5% ethyl acetate–hexanes, grading to 10% ethyl acetate–hexanes) to afford **4t** as a yellow oil (2.5 g, 17 mmol, 76% yield). TLC 10% ethyl acetate–hexanes, R_f = 0.71 (UV, KMnO₄). ¹H NMR (400 MHz, CDCl₃): δ 7.08 (t, J = 7.7 Hz, 1H), 6.98–6.94 (m, 1H), 6.61 (m, 2H), 3.27–3.18 (m, 2H), 2.89 (s, 3H), 2.78 (t, J = 6.4 Hz, 2H), 1.99 (p, J = 6.3 Hz, 2H). ¹³C NMR (101 MHz, CDCl₃): δ 146.9, 128.9, 127.2, 123.0, 116.3, 111.1, 51.4, 39.3, 27.9, 22.6. HRMS: ESI⁺ [M + H]⁺ calcd for C₁₀H₁₄N 148.1126, found 148.1117.

General Procedure for the Oxidation of *N,N*-Dialkylanilines.

N,N-Dimethylaniline *N*-oxide (**1a**), 3-methoxy-*N,N*-dimethylaniline *N*-oxide (**1b**), 3-methyl-*N,N*-dimethylaniline *N*-oxide (**1c**), 1-(*N,N*-dimethylamino)naphthalene *N*-oxide (**1d**), methyl 3-(*N,N*-dimethylamino)benzoate (**1e**), 4-fluoro-*N,N*-dimethylaniline *N*-oxide (**1i**), 4-chloro-*N,N*-dimethylaniline *N*-oxide (**1n**), 4-methoxy-*N,N*-dimethylaniline *N*-oxide (**1u**), and 4-methyl-*N,N*-dimethylaniline *N*-oxide (**1v**) were previously synthesized and reported by Lewis et al.⁸ All other *N,N*-dimethylaniline *N*-oxides were synthesized using the same procedure. A solution of 3-bromo-*N,N*-dimethylaniline **4f** (1.0 g, 5.0 mmol) in dichloromethane (25 mL) was added dropwise to a solution of m-CPBA (77%, 0.99 g, 6.0 mmol, 1.2 equiv) in dichloromethane (25 mL each) at 23 °C. The resultant solution was allowed to stir at 23 °C for 60 min. The solution was concentrated in vacuo to give a crude product, which was further purified using flash chromatography (basic alumina, starting with dichloromethane grading to 2% methanol–dichloromethane) to yield 3-bromo-*N,N*-dimethylaniline *N*-oxide **1f** (0.68 g, 63%) as a white solid.

3-Bromo-*N,N*-dimethylaniline N-Oxide (1f).—Mp: 117–119 °C. TLC 20% ethyl acetate–hexanes, R_f = 0.00 (UV, KMnO₄). ¹H NMR (400 MHz, CDCl₃): δ 8.28 (m, 1H), 7.87 (m, 1H), 7.55 (m, 1H), 7.34 (t, J = 8.1 Hz, 1H), 3.57 (s, 6H). ¹³C NMR (101 MHz, CDCl₃): δ 155.8, 132.3, 130.5, 124.1, 123.0, 118.8, 63.7. HRMS: ESI⁺ [M + H]⁺ calcd for C₈H₁₁ONBr 216.0024, found 216.0017.

3-Chloro-*N,N*-dimethylaniline N-Oxide (1g).—Obtained as a tan solid (0.99 g, 90%). Mp: 115–117 °C. TLC 20% ethyl acetate–hexanes, R_f = 0.00 (UV, KMnO₄). ¹H NMR (400 MHz, CDCl₃): δ 8.13 (m, 1H), 7.80 (m, 1H), 7.45–7.35 (m, 2H), 3.57 (s, 6H). ¹³C NMR (101 MHz, CDCl₃): δ 155.8, 135.2, 130.2, 129.4, 121.3, 118.2, 63.7. HRMS: ESI⁺ [M + H]⁺ calcd for C₈H₁₁ONCl 172.0529, found 172.0521.

2-Fluoro-*N,N*-dimethylaniline N-Oxide (1h).—Obtained as a orange-brown solid (0.96 g, 84%). Mp: 75–77 °C. TLC 20% ethyl acetate–hexanes, R_f = 0.00 (UV, KMnO₄). ¹H

NMR (400 MHz, CDCl₃) δ : 8.71 (m, 1H), 7.42 (m, 1H), 7.32 (m, 1H), 7.15 (m, 1H), 3.65 (d, J = 1.7 Hz, 6H). ¹³C NMR (101 MHz, CDCl₃) δ : 153.5 (d, J = 246.3 Hz), 141.3 (d, J = 9.9 Hz), 131.4 (d, J = 7.8 Hz), 125.3 (d, J = 3.6 Hz), 125.0, 116.5 (d, J = 22.5 Hz), 62.5 (d, J = 5.7 Hz). HRMS: ESI⁺ [M + H]⁺ calcd for C₈H₁₁ONF 156.0825, found 156.0817.

3-(Trifluoromethyl)-N,N-dimethylaniline N-Oxide (1i).—Obtained as a white solid (0.86 g, 79%). Mp: 131–132 °C. TLC 20% ethyl acetate–hexanes, R_f = 0.00 (UV, KMnO₄). ¹H NMR (400 MHz, CDCl₃) δ : 8.40 (m, 1H), 8.18 (m, 1H), 7.70 (m, 1H), 7.62 (m, 2H), 3.62 (s, 6H). ¹³C NMR (101 MHz, CDCl₃) δ : 155.3, 131.9 (q, J = 33.3 Hz), 130.0, 126.1 (q, J = 3.6 Hz), 123.5, 123.4 (q, J = 272.7 Hz), 117.9 (q, J = 3.8 Hz), 63.8. HRMS: ESI⁺ [M + H]⁺ calcd for C₉H₁₁ONF₃: 206.0793, found 206.0786.

3-(N,N-Dimethylamino)benzonitrile N-Oxide (1j).—Obtained as a white solid (0.85 g, 77%). Mp: 140–142 °C. TLC 20% ethyl acetate–hexanes, R_f = 0.00 (UV, KMnO₄). ¹H NMR (400 MHz, CDCl₃) δ : 8.44 (m, 1H), 8.25 (m, 1H), 7.72 (m, 1H), 7.62 (t, J = 8.0 Hz, 1H), 3.60 (s, 6H). ¹³C NMR (101 MHz, CDCl₃) δ : 155.5, 132.9, 130.3, 124.8, 124.5, 117.6, 113.6, 63.8. HRMS: ESI⁺ [M + H]⁺ calcd for C₉H₁₁ON₂ 163.0871, found 163.0863.

N-Methyl 4-(N,N-Dimethylamino)benzoate N-Oxide (1k).—Obtained as a white solid (0.64 g, 58%). Mp: 145–146 °C. TLC 20% ethyl acetate–hexanes, R_f = 0.00 (UV, KMnO₄). ¹H NMR (400 MHz, CDCl₃) δ : 8.15 (d, J = 8.9 Hz, 2H), 8.09 (d, J = 8.9 Hz, 2H), 3.94 (s, 3H), 3.61 (s, 6H). ¹³C NMR (101 MHz, CDCl₃) δ : 165.9, 158.2, 131.0, 130.8, 120.5, 63.6, 52.6. HRMS: ESI⁺ [M + H]⁺ calcd for C₁₀H₁₄O₃N 196.0974, found 196.0966.

4-Bromo-N,N-dimethylaniline N-Oxide (1m).—Obtained as a white solid (0.78 g, 71%). Mp: 158–160 °C. TLC 20% ethyl acetate–hexanes, R_f = 0.00 (UV, KMnO₄). ¹H NMR (400 MHz, CDCl₃) δ : 7.90 (d, J = 9.0 Hz, 2H), 7.59 (d, J = 9.0 Hz, 2H), 3.57 (s, 6H). ¹³C NMR (101 MHz, CDCl₃) δ : 153.8, 132.3, 123.0, 122.1, 63.7. HRMS: ESI⁺ [M + H]⁺ calcd for C₈H₁₁ONBr 216.0024, found 216.0016.

4-(N,N-Dimethylamino)benzonitrile N-Oxide (1o).—Obtained as a white solid (0.76 g, 69%). Mp: 139–140 °C. TLC 20% ethyl acetate–hexanes, R_f = 0.00 (UV, KMnO₄). ¹H NMR (400 MHz, CDCl₃) δ : 8.18 (d, J = 8.9 Hz, 2H), 7.80 (d, J = 9.0 Hz, 2H), 3.60 (s, 6H). ¹³C NMR (101 MHz, CDCl₃) δ : 158.2, 133.4, 121.6, 117.6, 113.5, 63.6. HRMS: ESI⁺ [M + H]⁺ calcd for C₉H₁₁ON₂: 163.0871, found 163.0863.

2-Methyl-N,N-dimethylaniline N-Oxide (1p).—Obtained as a yellow solid (0.39 g, 68%). Mp: 69–71 °C. TLC 20% ethyl acetate–hexanes, R_f = 0.00 (UV, KMnO₄). ¹H NMR (400 MHz, CDCl₃) δ : 8.29–8.24 (m, 1H), 7.30–7.23 (m, 3H), 3.65 (s, 6H), 2.79 (s, 3H). ¹³C NMR (101 MHz, CDCl₃) δ : 152.6, 134.1, 130.0, 129.2, 127.1, 120.9, 61.8, 22.5. HRMS: ESI⁺ [M + H]⁺ calcd for C₉H₁₄ON 152.1075, found 152.1066.

N-Butyl-N-dimethylaniline N-Oxide (1q).—Obtained as a yellow solid (0.44 g, 79%). TLC 20% ethyl acetate–hexanes, R_f = 0.00 (UV, KMnO₄). ¹H NMR (400 MHz, CDCl₃) δ : 7.93–7.88 (m, 2H), 7.49–7.43 (m, 2H), 7.40–7.36 (m, 1H), 3.68–3.56 (m, 2H), 3.52 (s, 3H), 1.96–1.85 (m, 2H), 1.26 (dddd, J = 17.1, 10.0, 4.1, 2.0 Hz, 2H), 0.86 (t, J = 7.2 Hz, 3H). ¹³C

NMR (101 MHz, CDCl₃): δ 152.6, 129.1, 128.7, 121.0, 73.5, 62.4, 25.6, 20.0, 14.0. HRMS: ESI⁺ [M + H]⁺ calcd for C₁₁H₁₈ON 180.1388, found 180.1377.

N-Isopentyl-N-dimethylaniline N-Oxide (1r).—Obtained as a yellow solid (0.38 g, 79%). TLC 20% ethyl acetate–hexanes, R_f = 0.00 (UV, KMnO₄). ¹H NMR (400 MHz, CDCl₃): δ 7.92–7.87 (m, 2H), 7.49–7.43 (m, 2H), 7.41–7.36 (m, 1H), 3.63 (m, 2H), 3.52 (s, 3H), 1.94–1.77 (m, 2H), 1.54 (m, 2H), 1.17 (m, 1H), 0.85 (dd, J = 13.5, 6.6 Hz, 6H). ¹³C NMR (101 MHz, CDCl₃): δ 152.6, 129.1, 128.7, 121.0, 72.4, 62.4, 32.0, 26.3, 22.7, 22.6. HRMS: ESI⁺ [M + H]⁺ calcd for C₁₂H₂₀ON 194.1545, found 194.1535.

N-Cyclohexyl-N-dimethylaniline N-Oxide (1s).—Obtained as a yellow solid (0.36 g, 66%). TLC 20% ethyl acetate–hexanes, R_f = 0.00 (UV, KMnO₄). ¹H NMR (400 MHz, CDCl₃): δ 7.89–7.84 (m, 2H), 7.43 (td, J = 7.3, 6.6, 1.4 Hz, 2H), 7.38–7.33 (m, 1H), 3.46 (s, 3H), 3.39 (m, 1H), 2.24–2.19 (m, 1H), 1.91 (m, 1H), 1.76 (m, 2H), 1.63–1.54 (m, 2H), 1.33–1.20 (m, 2H), 1.18–1.09 (m, 2H). ¹³C NMR (101 MHz, CDCl₃): δ 153.2, 128.8, 128.5, 121.5, 79.7, 57.5, 26.7, 26.6, 25.5, 25.4, 25.1. HRMS: ESI⁺ [M + H]⁺ calcd for C₁₃H₂₀ON 206.1545, found 206.1534.

N-Methyl-1,2,3,4-tetrahydroquinoline N-Oxide (1t).—Obtained as a tan solid (0.32 g, 57%). Mp: 118–120 °C. TLC 20% ethyl acetate–hexanes, R_f = 0.00 (UV, KMnO₄). ¹H NMR (600 MHz, CDCl₃): δ 8.15 (dd, J_1 = 8.4 Hz, J_2 = 1.2 Hz, 1H), 7.35–7.31 (m, 1H), 7.24 (dt, J_1 = 7.5 Hz, J_2 = 1.2 Hz, 1H), 7.12 (dd, J_1 = 7.7 Hz, J_2 = 1.4 Hz, 1H), 3.87–3.80 (m, 2H), 3.53 (s, 3H), 3.03–2.96 (m, 1H), 2.90 (m, 1H), 2.48 (m, 1H), 2.17–2.09 (m, 1H). ¹³C NMR (101 MHz, CDCl₃): δ 149.96, 129.51, 128.64, 128.36, 128.14, 122.09, 69.50, 62.46, 26.23, 19.85. HRMS: ESI⁺ [M + H]⁺ calcd for C₁₀H₁₄ON 164.1075, found 164.1065.

General Procedure for the Reaction of *N,N*-Dialkylaniline *N*-oxides with SOBr₂.

Thionyl bromide (54 μ L, 0.70 mmol, 1.0 equiv) was added dropwise to solution of the *N,N*-dimethylaniline *N*-oxide (96 mg, 0.70 mmol, 1 equiv) in tetrahydrofuran (4 mL) at –78 °C. The resultant mixture was stirred at –78 °C for 4 h, whereupon triethylamine (365 μ L, 2.80 mmol, 4.0 equiv) was added. The cooling bath was removed, and the resultant mixture was allowed to warm to 23 °C and was stirred at that temperature for 45 min. The resultant mixture was diluted with saturated aqueous sodium bicarbonate solution (4 mL), the layers were separated, and the aqueous layer was extracted with dichloromethane (3 $\times$ 5 mL). The combined organic layers were dried over anhydrous sodium sulfate, and the dried solution was concentrated. The resultant oily residue was purified by flash column chromatography (silica gel, starting with hexanes, grading to 1% ethyl acetate–hexanes) to afford 2a as a white solid (77 mg, 0.38 mmol, 55% yield).

4-Bromo-N,N-dimethylaniline (2a).—Mp: 29–31 °C. TLC: 10% ethyl acetate–hexanes, R_f = 0.45 (UV, KMnO₄). ¹H NMR (400 MHz, CDCl₃): δ 7.30 (d, J = 9.1 Hz, 2H), 6.59 (d, J = 9.1 Hz, 2H), 2.92 (s, 6H). ¹³C NMR (101 MHz, CDCl₃): δ 149.6, 131.8, 114.2, 108.6, 40.7. HRMS: ESI⁺ [M + H]⁺ calcd for C₈H₁₁NBr 200.0075, found 200.0069.

4-Bromo-3-methoxy-N,N-dimethylaniline (2b).—Obtained as a tan solid (65 mg, 39%). Mp: 67–68 °C. TLC 10% ethyl acetate–hexanes, R_f = 0.55 (UV, KMnO₄). ¹H NMR (400 MHz, CDCl₃) δ : 7.31 (d, J = 8.7 Hz, 1H), 6.25 (d, J = 2.8 Hz, 1H), 6.21 (dd, J_1 = 8.7, J_2 = 2.8 Hz, 1H), 3.89 (s, 3H), 2.95 (s, 6H). ¹³C NMR (101 MHz, CDCl₃) δ : 156.4, 151.4, 133.1, 106.2, 98.0, 97.2, 56.1, 40.8. HRMS: ESI⁺ [M + H]⁺ calcd for C₉H₁₃ONBr 230.0181, found 230.0175.

4-Bromo-3-methyl-N,N-dimethylaniline (2c).—Obtained as a white solid (103 mg, 66%). Mp: 51–53 °C. TLC 10% ethyl acetate–hexanes, R_f = 0.41 (UV, KMnO₄). ¹H NMR (400 MHz, CDCl₃) δ : 7.32 (d, J = 8.8 Hz, 1H), 6.59 (d, J = 3.1 Hz, 1H), 6.44 (dd, J_1 = 8.8, J_2 = 3.1 Hz, 1H), 2.91 (s, 6H), 2.35 (s, 3H). ¹³C NMR (101 MHz, CDCl₃) δ : 150.0, 138.0, 132.5, 115.0, 112.0, 111.5, 40.8, 23.5. HRMS: ESI⁺ [M + H]⁺ calcd for C₉H₁₃NBr 214.0231, found 214.0225.

4-Bromo-1-(N,N-dimethylamino)naphthalene (2d).—Obtained as a yellow oil (85 mg, 56%). TLC 10% ethyl acetate–hexanes, R_f = 0.46 (UV, KMnO₄). ¹H NMR (400 MHz, CDCl₃) δ : 8.29–8.17 (m, 2H), 7.67 (d, J = 8.0 Hz, 1H), 7.56 (m, 2H), 6.93 (d, J = 8.0 Hz, 1H), 2.88 (s, 6H). ¹³C NMR (101 MHz, CDCl₃) δ : 151.1, 132.9, 130.2, 129.7, 127.7, 127.2, 126.0, 124.7, 116.5, 114.8, 45.3. HRMS: ESI⁺ [M + H]⁺ calcd for C₁₂H₁₃NBr 250.0231, found 250.0225.

Methyl 2-Bromo-5-(N,N-dimethylamino)benzoate (2e).—Obtained as a yellow oil (91 mg, 50%). TLC 10% ethyl acetate–hexanes, R_f = 0.26 (UV, KMnO₄). ¹H NMR (400 MHz, CDCl₃) δ : 7.42 (d, J = 8.9 Hz, 1H), 7.06 (d, J = 3.2 Hz, 1H), 6.65 (dd, J_1 = 8.9, J_2 = 3.2 Hz, 1H), 3.92 (s, 3H), 2.95 (s, 6H). ¹³C NMR (101 MHz, CDCl₃) δ : 167.6, 149.4, 134.5, 132.3, 116.5, 114.6, 106.9, 52.5, 40.5. HRMS: ESI⁺ [M + H]⁺ calcd for C₁₀H₁₃O₂NBr 258.0130, found 258.0135.

3,4-Dibromo-N,N-dimethylaniline (2f).—Obtained as a white solid (109 mg, 58%). Mp: 64–65 °C. TLC 10% ethyl acetate–hexanes, R_f = 0.47 (UV, KMnO₄). ¹H NMR (400 MHz, CDCl₃) δ : 7.37 (d, J = 9.0 Hz, 1H), 6.92 (d, J = 3.0 Hz, 1H), 6.50 (dd, J_1 = 9.0, J_2 = 3.0 Hz, 1H), 2.92 (s, 6H). ¹³C NMR (101 MHz, CDCl₃) δ : 150.4, 133.4, 125.2, 116.9, 112.9, 110.0, 40.5. HRMS: ESI⁺ [M + H]⁺ calcd for C₈H₁₀NBr₂ 277.9180, found 277.9185.

4-Bromo-3-chloro-N,N-dimethylaniline (2g).—Obtained as a white solid (101 mg, 61%). Mp: 48–49 °C. TLC 10% ethyl acetate–hexanes, R_f = 0.50 (UV, KMnO₄). ¹H NMR (400 MHz, CDCl₃) δ : 7.37 (d, J = 8.9 Hz, 1H), 6.75 (d, J = 3.0 Hz, 1H), 6.46 (dd, J_1 = 9.0, J_2 = 3.0 Hz, 1H), 2.93 (s, 6H). ¹³C NMR (101 MHz, CDCl₃) δ : 150.5, 134.7, 133.5, 113.7, 112.3, 107.7, 40.5. HRMS: ESI⁺ [M + H]⁺ calcd for C₈H₁₀NBrCl 233.9685, found 233.9690.

4-Bromo-2-fluoro-N,N-dimethylaniline (2h).—Obtained as a yellow oil (49 mg, 31%). TLC 10% ethyl acetate–hexanes, R_f = 0.55 (UV, KMnO₄). ¹H NMR (400 MHz, CDCl₃) δ : 7.15 (m, 2H), 6.75 (m, 1H), 2.82 (s, 6H). ¹³C NMR (101 MHz, CDCl₃) δ : 154.8 (d, J = 249.9 Hz), 140.2 (d, J = 8.5 Hz), 127.3 (d, J = 3.6 Hz), 119.5 (d, J = 40.4 Hz), 119.4 (d, J =

12.2 Hz), 112.0 (d, $J = 9.4$ Hz), 42.8 (d, $J = 4.2$ Hz). HRMS: ESI⁺ [M + H]⁺ calcd for C₈H₁₀NBrF: 217.9980, found 217.9984.

4-Bromo-3-(trifluoromethyl)-N,N-dimethylaniline (2i).—Obtained as a yellow oil (92 mg, 48%). TLC 10% ethyl acetate–hexanes, $R_f = 0.47$ (UV, KMnO₄). ¹H NMR (400 MHz, CDCl₃) δ : 7.70 (d, $J = 2.4$ Hz, 1H), 7.38 (dd, $J_1 = 8.6$, $J_2 = 2.3$ Hz, 1H), 6.96 (d, $J = 8.6$ Hz, 1H), 2.80 (s, 6H). ¹³C NMR (101 MHz, CDCl₃) δ : 149.2, 135.2, 130.1 (q, $J = 30.4$ Hz), 123.3 (q, $J = 273.5$ Hz), 116.1, 111.2 (q, $J = 5.8$ Hz), 104.5 (q, $J = 1.9$ Hz), 40.5. HRMS: ESI⁺ [M + H]⁺ calcd for C₉H₁₀NBrF₃ 267.9949, found 267.9952.

2-Bromo-5-(N,N-dimethylamino)benzonitrile (2j).—Obtained as a white solid (77 mg, 49%). Mp: 70–72 °C. TLC 5% ethyl acetate–hexanes, $R_f = 0.21$ (UV, KMnO₄). ¹H NMR (400 MHz, CDCl₃) δ : 7.41 (d, $J = 9.1$ Hz, 1H), 6.87 (d, $J = 3.2$ Hz, 1H), 6.73 (dd, $J_1 = 9.1$, $J_2 = 3.2$ Hz, 1H), 2.97 (s, 6H). ¹³C NMR (101 MHz, CDCl₃) δ : 149.2, 133.3, 118.2, 117.6, 116.9, 115.7, 109.9, 40.4. HRMS: ESI⁺ [M + H]⁺ calcd for C₉H₁₀N₂Br 225.0027, found 225.0020.

Methyl 3-Bromo-4-(N,N-dimethylamino)benzoate (2k).—Obtained as a yellow oil (60 mg, 33%). TLC 10% ethyl acetate–hexanes, $R_f = 0.57$ (UV, KMnO₄). ¹H NMR (400 MHz, CDCl₃) δ : 8.20 (d, $J = 2.0$ Hz, 1H), 7.89 (dd, $J_1 = 8.5$, $J_2 = 2.0$ Hz, 1H), 7.01 (d, $J = 8.5$ Hz, 1H), 3.87 (s, 3H), 2.88 (s, 6H). ¹³C NMR (101 MHz, CDCl₃) δ : 166.0, 155.8, 135.7, 129.7, 124.5, 119.3, 116.9, 52.2, 43.7. HRMS: ESI⁺ [M + H]⁺ calcd for C₁₀H₁₃O₂NBr 258.0130, found 258.0123.

2-Bromo-4-fluoro-N,N-dimethylaniline (2l).—Obtained as a yellow oil (39 mg, 26%). TLC 10% ethyl acetate–hexanes, $R_f = 0.56$ (UV, KMnO₄). ¹H NMR (400 MHz, CDCl₃) δ : 7.31 (dd, $J_1 = 8.1$, $J_2 = 2.9$ Hz, 1H), 7.06 (m, 1H), 6.99 (m, 1H), 2.75 (s, 6H). ¹³C NMR (101 MHz, CDCl₃) δ : 158.3 (d, $J = 245.5$ Hz), 148.4 (d, $J = 3.0$ Hz), 121.1 (d, $J = 8.5$ Hz), 120.9 (d, $J = 25.0$ Hz), 119.7 (d, $J = 9.5$ Hz), 114.8 (d, $J = 21.6$ Hz), 44.7. HRMS: ESI⁺ [M + H]⁺ calcd for C₈H₁₀NBrF 217.9981, found 217.9981.

2,4-Dibromo-N,N-dimethylaniline (2m).—Obtained as a yellow oil (51 mg, 26%). TLC 10% ethyl acetate–hexanes, $R_f = 0.62$ (UV, KMnO₄). ¹H NMR (400 MHz, CDCl₃) δ : 7.68 (d, $J = 2.4$ Hz, 1H), 7.36 (dd, $J_1 = 8.6$, $J_2 = 2.3$ Hz, 1H), 6.94 (d, $J = 8.6$ Hz, 1H), 2.77 (s, 6H). ¹³C NMR (101 MHz, CDCl₃) δ : 151.2, 136.1, 131.1, 121.7, 119.8, 115.4, 44.2. HRMS: ESI⁺ [M + H]⁺ calcd for C₈H₁₀NBr₂ 279.9180, found 279.9158.

2-Bromo-4-chloro-N,N-dimethylaniline (2n).—Obtained as a yellow oil (58 mg, 37%). TLC 10% ethyl acetate–hexanes, $R_f = 0.54$ (UV, KMnO₄). ¹H NMR (400 MHz, CDCl₃) δ : 7.55 (d, $J = 2.5$ Hz, 1H), 7.23 (dd, $J_1 = 8.6$, $J_2 = 2.4$ Hz, 1H), 7.00 (d, $J = 8.6$ Hz, 1H), 2.78 (s, 6H). ¹³C NMR (101 MHz, CDCl₃) δ : 150.8, 133.4, 128.2, 128.2, 121.2, 119.5, 44.3. HRMS: ESI⁺ [M + H]⁺ calcd for C₈H₁₀NBrCl 233.9685, found 233.9680.

3-Bromo-4-(N,N-dimethylamino)benzonitrile (2o).—Obtained as a yellow oil (20 mg, 10%). TLC 5% ethyl acetate–hexanes, $R_f = 0.23$ (UV, KMnO₄). ¹H NMR (400 MHz, CDCl₃) δ : 7.79 (d, $J = 2.0$ Hz, 1H), 7.51 (dd, $J_1 = 8.5$ Hz, $J_2 = 2.0$ Hz, 1H), 7.01 (d, $J = 8.4$

Hz, 1H), 2.91 (s, 6H). ^{13}C NMR (101 MHz, CDCl_3) δ : 155.7, 137.8, 132.1, 119.9, 118.3, 116.6, 105.4, 43.5. HRMS: $\text{ESI}^+ [\text{M} + \text{H}]^+$ calcd for $\text{C}_9\text{H}_{10}\text{N}_2\text{Br}$ 225.0027, found 225.0030.

4-Bromo-2-methyl-N,N-dimethylaniline (2p).—Obtained as a yellow oil (77 mg, 51%). TLC 10% ethyl acetate–hexanes, R_f = 0.63 (UV, KMnO_4). ^1H NMR (600 MHz, CDCl_3): δ 7.28 (d, J = 2.1 Hz, 1H), 7.24 (dd, J_1 = 8.5 Hz, J_2 = 2.3 Hz, 1H), 6.88 (d, J = 8.5 Hz, 1H), 2.67 (s, 6H), 2.29 (s, 3H). ^{13}C NMR (101 MHz, CDCl_3): δ 152.0, 134.5, 133.8, 129.3, 120.2, 115.2, 44.2, 18.3. HRMS: $\text{ESI}^+ [\text{M} + \text{H}]^+$ calcd for $\text{C}_9\text{H}_{13}\text{NBr}$ 214.0231, found 214.0222.

4-Bromo-N-butyl-N-dimethylaniline (2q).—Obtained as a yellow oil (100 mg, 59%). TLC 10% ethyl acetate–hexanes, R_f = 0.71 (UV, KMnO_4). ^1H NMR (400 MHz, CDCl_3): δ 7.27 (d, J = 9.2 Hz, 2H), 6.54 (d, J = 9.0 Hz, 2H), 3.30–3.24 (m, 2H), 2.89 (s, 3H), 1.58–1.48 (m, 2H), 1.34 (m, 2H), 0.94 (t, J = 7.3 Hz, 3H). ^{13}C NMR (101 MHz, CDCl_3): δ 148.3, 131.8, 113.7, 107.6, 52.6, 38.5, 28.8, 20.4, 14.1. HRMS: $\text{ESI}^+ [\text{M} + \text{H}]^+$ calcd for $\text{C}_{11}\text{H}_{17}\text{NBr}$ 242.0544, found 242.0534.

4-Bromo-N-isopentyl-N-dimethylaniline (2r).—Obtained as a yellow oil (96 mg, 53%). TLC 10% ethyl acetate–hexanes, R_f = 0.74 (UV, KMnO_4). ^1H NMR (400 MHz, CDCl_3): δ 7.27 (d, J = 9.3 Hz, 2H), 6.54 (d, J = 9.1 Hz, 2H), 3.32–3.25 (m, 2H), 2.88 (s, 3H), 1.65–1.51 (m, 2H), 1.46–1.38 (m, 2H), 0.94 (d, J = 6.6 Hz, 6H). ^{13}C NMR (101 MHz, CDCl_3): δ 148.3, 131.9, 113.8, 107.7, 51.1, 38.4, 35.0, 26.3, 22.8. HRMS: $\text{ESI}^+ [\text{M} + \text{H}]^+$ calcd for $\text{C}_{12}\text{H}_{19}\text{NBr}$ 256.0701, found 256.0690.

4-Bromo-N-cyclohexyl-N-dimethylaniline (2s).—Obtained as a yellow solid (120 mg, 64%). Mp: 45–47 °C. TLC 10% ethyl acetate–hexanes, R_f = 0.68 (UV, KMnO_4). ^1H NMR (400 MHz, CDCl_3): δ 7.27 (d, J = 9.3 Hz, 2H), 6.63 (d, J = 9.1 Hz, 2H), 3.49 (m, 1H), 2.73 (s, 3H), 1.87–1.80 (m, 2H), 1.79–1.67 (m, 3H), 1.47–1.31 (m, 4H), 1.12 (m, 1H). ^{13}C NMR (101 MHz, CDCl_3): δ 149.2, 131.8, 114.7, 107.9, 58.3, 31.3, 30.1, 26.3, 26.0. HRMS: $\text{ESI}^+ [\text{M} + \text{H}]^+$ calcd for $\text{C}_{13}\text{H}_{19}\text{NBr}$ 268.0701, found 268.0689.

7-Bromo-N-methyl-1,2,3,4-tetrahydroquinoline (2t).—Obtained as a yellow oil (110 mg, 69%). TLC 10% ethyl acetate–hexanes, R_f = 0.59 (UV, KMnO_4). ^1H NMR (400 MHz, CDCl_3): δ 7.13 (dd, J_1 = 8.7 Hz, J_2 = 2.5 Hz, 1H), 7.06–7.03 (m, 1H), 6.43 (d, J = 8.7 Hz, 1H), 3.23–3.17 (m, 2H), 2.85 (s, 3H), 2.73 (t, J = 6.5 Hz, 2H), 1.98–1.92 (m, 2H). ^{13}C NMR (101 MHz, CDCl_3): δ 145.8, 131.2, 129.7, 125.0, 112.5, 107.8, 51.2, 39.2, 27.8, 22.3. HRMS: $\text{ESI}^+ [\text{M} + \text{H}]^+$ calcd for $\text{C}_{10}\text{H}_{13}\text{NBr}$ 226.0231, found 226.0221.

General Procedure for the Reaction of *N,N*-Dialkylanilines *N*-Oxides with SOCl_2 .

A solution of thionyl chloride (52 μL , 0.70 mmol, 1.0 equiv) in tetrahydrofuran (1 mL) was added dropwise in two portions over a period of 2 h to a solution of the *N,N*-dimethylaniline *N*-oxide (96 mg, 0.703 mmol, 1 equiv) in tetrahydrofuran (3 mL) at –78 °C. The resultant mixture was stirred at –78 °C for 4 h, whereupon triethylamine (365 μL , 2.80 mmol, 4.00 equiv) was added. The cooling bath was removed, and the resultant mixture was allowed to warm to 23 °C and stirred at that temperature for 45 min. The resultant mixture was diluted

with saturated aqueous sodium bicarbonate solution (4 mL), the layers were separated, and the aqueous layer was extracted with dichloromethane (3 × 5 mL). The combined organic layers were dried over anhydrous sodium sulfate, and the dried solution was concentrated. The resultant oily residue was purified by flash column chromatography (silica gel, starting with hexanes, grading to 1% ethyl acetate–hexanes) to afford **3a** as a yellow oil (isolated as a mix of regioisomers, 4.9:1 (2-Cl:4-Cl), asterisk denotes minor peaks, 53 mg, 0.34 mmol, 49%).

2-Chloro-N,N-dimethylaniline (3a).—TLC 10% ethyl acetate–hexanes, R_f = 0.57 (UV, KMnO_4). ^1H NMR (400 MHz, CDCl_3) δ : 7.35 (dd, J_1 = 7.9, J_2 = 1.6 Hz, 1H), 7.22 (dt, J_1 = 8.1, J_2 = 1.6 Hz, 1H), 7.17* (d, J = 9.1 Hz, 2H), 7.08 (dd, J_1 = 8.1, J_2 = 1.6 Hz, 1H), 6.95 (dt, J_1 = 7.9, J_2 = 1.6 Hz, 1H), 6.64* (d, J = 9.1 Hz, 2H), 2.93* (s, 6H), 2.82 (s, 6H). ^{13}C NMR (101 MHz, CDCl_3) δ : 150.5, 149.3*, 130.8, 130.4, 128.9*, 128.4, 127.5, 123.3, 120.1*, 113.7*, 43.9, 40.8*. HRMS: ESI^+ $[\text{M} + \text{H}]^+$ calcd for $\text{C}_8\text{H}_{11}\text{NCl}$ 156.0580, found 156.0570.

2-Chloro-4-methoxy-N,N-dimethylaniline (3b).—Obtained as a yellow oil (79 mg, 61%). TLC 10% ethyl acetate–hexanes, R_f = 0.38 (UV, KMnO_4). ^1H NMR (400 MHz, CDCl_3) δ : 7.03 (d, J = 8.8 Hz, 1H), 6.95 (d, J = 2.9 Hz, 1H), 6.78 (dd, J_1 = 8.8, J_2 = 2.9 Hz, 1H), 3.77 (s, 3H), 2.74 (s, 6H). ^{13}C NMR (101 MHz, CDCl_3) δ : 155.6, 144.0, 129.5, 120.8, 116.1, 113.1, 55.8, 44.5. HRMS: ESI^+ $[\text{M} + \text{H}]^+$ calcd for $\text{C}_9\text{H}_{13}\text{ONCl}$ 186.0686, found 186.0685.

2-Chloro-4-methyl-N,N-dimethylaniline (3c).—Obtained as a yellow oil (49 mg, 41%). TLC 10% ethyl acetate–hexanes, R_f = 0.56 (UV, KMnO_4). ^1H NMR (400 MHz, CDCl_3) δ : 7.18 (d, J = 1.9 Hz, 1H), 7.02–6.95 (m, 2H), 2.78 (s, 6H), 2.27 (s, 3H). ^{13}C NMR (101 MHz, CDCl_3) δ : 148.0, 133.3, 131.2, 128.2, 128.1, 119.9, 44.2, 20.5. HRMS: ESI^+ $[\text{M} + \text{H}]^+$ calcd for $\text{C}_9\text{H}_{13}\text{NCl}$ 170.0737, found 170.0729.

2-Chloro-1-(N,N-dimethylamino)naphthalene (3d, 4-Cl Regioisomer also Isolated).—Obtained as a colorless oil (75 mg, 52%). TLC 10% ethyl acetate–hexanes, R_f = 0.87 (UV, KMnO_4). ^1H NMR (400 MHz, CDCl_3) δ : 8.35 (m, 1H), 7.80 (m, 1H), 7.58 (d, J = 8.8 Hz, 1H), 7.50 (m, 2H), 7.37 (d, J = 8.8 Hz, 1H), 3.02 (s, 6H). ^{13}C NMR (101 MHz, CDCl_3) δ : 145.4, 133.9, 133.3, 130.4, 128.6, 128.0, 126.6, 126.5, 126.1, 124.6, 42.9. HRMS: ESI^+ $[\text{M} + \text{H}]^+$ calcd for $\text{C}_{12}\text{H}_{13}\text{NCl}$ 206.0737, found 206.0727. *4-Chloro-1-(N,N-dimethylamino)naphthalene.* Obtained as a yellow oil (11 mg, 8%). TLC 10% ethyl acetate–hexanes, R_f = 0.47 (UV, KMnO_4). ^1H NMR (400 MHz, CDCl_3) δ : 8.26 (m, 2H), 7.62–7.51 (m, 2H), 7.47 (d, J = 8.0 Hz, 1H), 6.98 (d, J = 8.1 Hz, 1H), 2.88 (s, 6H). ^{13}C NMR (101 MHz, CDCl_3) δ : 150.3, 131.7, 130.0, 126.9, 126.0, 126.0, 126.0, 125.0, 124.7, 114.2, 45.4. HRMS: ESI^+ $[\text{M} + \text{H}]^+$ calcd for $\text{C}_{12}\text{H}_{13}\text{NCl}$ 206.0737, found 206.0728.

Methyl 3-Chloro-4-(N,N-dimethylamino)benzoate (3e).—Obtained as a yellow oil (61 mg, 40%). TLC 10% ethyl acetate–hexanes, R_f = 0.41 (UV, KMnO_4). ^1H NMR (400 MHz, CDCl_3) δ : 8.00 (d, J = 2.1 Hz, 1H), 7.86 (dd, J_1 = 8.5, J_2 = 2.1 Hz, 1H), 7.01 (d, J = 8.5 Hz, 1H), 3.89 (s, 3H), 2.91 (s, 6H). ^{13}C NMR (101 MHz, CDCl_3) δ : 166.2, 154.3, 132.5,

129.1, 126.5, 123.9, 118.8, 52.2, 43.3. HRMS: ESI⁺ [M + H]⁺ calcd for C₁₀H₁₃O₂NCl: 214.0635, found 214.0625.

3-Chloro-4-(N,N-dimethylamino)benzonitrile (3f).—Obtained as a yellow oil (42 mg, 33%). TLC 10% ethyl acetate–hexanes, *R*_f = 0.22 (UV, KMnO₄). ¹H NMR (400 MHz, CDCl₃) δ: 7.58 (d, *J* = 2.0 Hz, 1H), 7.46 (dd, *J*₁ = 8.5, *J*₂ = 2.0 Hz, 1H), 6.99 (d, *J* = 8.5 Hz, 1H), 2.92 (s, 6H). ¹³C NMR (101 MHz, CDCl₃) δ: 154.1, 134.5, 131.6, 126.5, 119.4, 118.5, 104.5, 43.1. HRMS: ESI⁺ [M + H]⁺ calcd for C₉H₁₀N₂Cl 181.0533, found 181.0530.

2,4-Dichloro-N,N-dimethylaniline (3g).—Obtained as a yellow oil (73 mg, 55%). TLC 10% ethyl acetate–hexanes, *R*_f = 0.54 (UV, KMnO₄). ¹H NMR (400 MHz, CDCl₃) δ: 7.35 (d, *J* = 2.4 Hz, 1H), 7.17 (dd, *J*₁ = 8.6, *J*₂ = 2.5 Hz, 1H), 6.98 (d, *J* = 8.7 Hz, 1H), 2.79 (s, 6H). ¹³C NMR (101 MHz, CDCl₃) δ: 149.3, 130.4, 128.9, 127.7, 127.5, 120.8, 43.9. HRMS: ESI⁺ [M + H]⁺ calcd for C₈H₁₀NCl₂ 190.0190, found 190.0183.

2-Chloro-4-fluoro-N,N-dimethylaniline (3h).—Obtained as a yellow oil (37 mg, 30%). TLC 10% ethyl acetate–hexanes, *R*_f = 0.57 (UV, KMnO₄). ¹H NMR (400 MHz, CDCl₃) δ: 7.12 (m, 1H), 7.03 (m, 1H), 6.93 (m, 1H), 2.76 (s, 6H). ¹³C NMR (101 MHz, CDCl₃) δ: 158.1 (d, *J* = 244.3 Hz), 147.0 (d, *J* = 3.1 Hz), 129.2 (d, *J* = 10.3 Hz), 120.8 (d, *J* = 8.7 Hz), 117.9 (d, *J* = 25.3 Hz), 114.1 (d, *J* = 21.5 Hz), 44.2. HRMS: ESI⁺ [M + H]⁺ calcd for C₈H₁₀NClF 174.0486, found 174.0478.

6-Chloro-3-methyl-N,N-dimethylaniline (3i).—Obtained as a yellow oil (isolated as a mixture of regioisomers, 5.3:3.7:1 (6-Cl:2-Cl:4-Cl), asterisk denotes 2-Cl, plus denotes 4-Cl, 57 mg, 47%). TLC 10% ethyl acetate–hexanes, *R*_f = 0.64–54 (UV, KMnO₄). ¹H NMR (400 MHz, CDCl₃) δ: 7.22 (d, *J* = 8.0 Hz, 1H), 7.16⁺ (d, *J* = 8.8 Hz, 1H), 7.11* (t, *J* = 7.7 Hz, 1H), 6.94* (m, 2H), 6.87 (d, *J* = 2.0 Hz, 1H), 6.76 (dd, *J*₁ = 8.0, *J*₂ = 1.7 Hz, 1H), 6.58⁺ (d, *J* = 3.1 Hz, 1H), 6.50⁺ (dd, *J*₁ = 8.8, *J*₂ = 3.1 Hz, 1H), 2.91⁺ (s, 6H), 2.80 (s, 6H), 2.79* (s, 6H), 2.39* (s, 3H), 2.33⁺ (s, 3H), 2.31 (s, 3H). ¹³C NMR (101 MHz, CDCl₃) δ: 150.9*, 150.1, 149.4⁺, 137.8*, 137.4, 136.2⁺, 130.4, 129.3⁺, 128.9*, 126.6*, 125.2, 125.1*, 124.1, 122.1⁺, 120.9, 117.7*, 115.0⁺, 111.6⁺, 44.2*, 44.0, 40.9⁺, 21.3, 21.1*, 20.7⁺. HRMS: ESI⁺ [M + H]⁺ calcd for C₉H₁₃NCl 170.0737, found 170.0733. *4-Chloro-3-methyl-N,N-dimethylaniline.* Obtained as a yellow oil (8 mg, 7%). TLC 10% ethyl acetate–hexanes, *R*_f = 0.54 (UV, KMnO₄). ¹H NMR (400 MHz, CDCl₃) δ: 7.16 (d, *J* = 8.8 Hz, 1H), 6.58 (d, *J* = 3.0 Hz, 1H), 6.50 (dd, *J*₁ = 8.8, *J*₂ = 3.1 Hz, 1H), 2.91 (s, 6H), 2.33 (s, 3H). ¹³C NMR (101 MHz, CDCl₃) δ: 149.5, 136.2, 129.3, 122.1, 115.0, 111.6, 40.9, 20.7. HRMS: ESI⁺ [M + H]⁺ calcd for C₉H₁₃NCl 170.0737, found 170.0733.

6-Chloro-3-methoxy-N,N-dimethylaniline (3j).—Obtained as a yellow oil (isolated as a mixture of regioisomers, 4.9:1 (6-Cl/4-Cl), asterisk denotes minor peaks, 55 mg, 43%). TLC 10% ethyl acetate–hexanes, *R*_f = 0.46 (UV, KMnO₄). ¹H NMR (400 MHz, CDCl₃) δ: 7.24 (d, *J* = 8.7 Hz, 1H), 7.17* (t, *J* = 8.2 Hz, 1H), 6.74* (dd, *J*₁ = 8.2, *J*₂ = 1.3 Hz, 1H), 6.65* (dd, *J*₁ = 8.3, *J*₂ = 1.3 Hz, 1H), 6.62 (d, *J* = 2.9 Hz, 1H), 6.49 (dd, *J*₁ = 8.7 Hz, *J*₂ = 2.9 Hz, 1H), 3.90* (s, 1H), 3.79 (s, 3H), 2.81* (s, 1H), 2.80 (s, 6H). ¹³C NMR (101 MHz, CDCl₃) δ: 159.1, 156.2*, 152.1*, 151.4, 131.0, 127.1*, 119.8, 116.5*, 112.4*, 107.5, 107.0,

106.3*, 56.4*, 55.6, 44.1*, 43.8. HRMS: ESI⁺ [M + H]⁺ calcd for C₉H₁₃ONCl 186.0686, found 186.0683.

2-Chloro-6-methyl-N,N-dimethylaniline (3k).—Obtained as a yellow oil (isolated as mixture of regioisomers, 1.3:1 (6-Cl:4-Cl), asterisk denotes minor peaks, 23 mg, 19%). TLC 10% ethyl acetate–hexanes, *R_f* = 0.89 (UV, KMnO₄). ¹H NMR (400 MHz, CDCl₃): δ 7.17 (d, *J* = 6.8 Hz, 1H), 7.13* (d, *J* = 2.0 Hz, 1H), 7.10* (dd, *J*₁ = 8.4, *J*₂ = 2.0 Hz, 1H), 7.03 (d, *J* = 8.4 Hz, 1H), 6.96 (t, *J* = 7.2 Hz, 1H), 6.93* (d, *J* = 8.4 Hz, 1H), 2.70 (s, 6H), 2.67* (s, 6H), 2.34 (s, 3H), 2.30* (s, 3H). ¹³C NMR (101 MHz, CDCl₃): δ 152.7*, 151.3, 134.0, 132.1, 131.2*, 130.8, 127.4*, 126.4*, 126.2, 122.5*, 119.6, 118.3*, 44.3*, 44.2, 18.4*, 18.3. HRMS: ESI⁺ [M + H]⁺ calcd for C₉H₁₃NCl 170.0737, found 170.0727.

2-Chloro-N-butyl-N-dimethylaniline (3l).—Obtained as a yellow oil (isolated as a mixture of regioisomers, 2.9:1 (6-Cl:2-Cl), asterisk denotes minor peaks, 77 mg, 55%). TLC 10% ethyl acetate–hexanes, *R_f* = 0.70 (UV, KMnO₄). ¹H NMR (400 MHz, CDCl₃): δ 7.37 (dd, *J*₁ = 7.9 Hz, *J*₂ = 1.6 Hz, 1H), 7.22 (td, *J*₁ = 7.7 Hz, *J*₂ = 1.6 Hz, 1H), 7.17* (d, *J* = 9.1 Hz, 2H), 7.09 (dd, *J*₁ = 8.1 Hz, *J*₂ = 1.5 Hz, 1H), 6.96 (td, *J*₁ = 7.6 Hz, *J*₂ = 1.6 Hz, 1H), 6.61* (d, *J* = 9.1 Hz, 2H), 3.33–3.27* (m, 2H), 3.07–3.00 (m, 2H), 2.92* (s, 3H), 2.80 (s, 3H), 1.62–1.51* (m, 2H), 1.62–1.51 (m, 2H), 1.41–1.28* (m, 2H), 1.41–1.28 (m, 2H), 0.95* (m, 3H), 0.95 (m, 3H). ¹³C NMR (101 MHz, CDCl₃): δ 150.2, 148.0*, 130.7, 129.0, 129.0*, 127.3, 123.2*, 121.3, 120.6*, 113.2*, 55.8, 52.7*, 40.9, 38.6*, 29.5, 28.8*, 20.5*, 20.4, 14.2, 14.1*. HRMS: ESI⁺ [M + H]⁺ calcd for C₁₁H₁₇NCl 198.1050, found 198.1040.

2-Chloro-N-isopentyl-N-dimethylaniline (3m).—Obtained as a yellow oil (isolated as a mixture of regioisomers, 2.8:1 (6-Cl:2-Cl), asterisk denotes minor peaks, 89 mg, 59%). TLC 10% ethyl acetate–hexanes, *R_f* = 0.75 (UV, KMnO₄). ¹H NMR (400 MHz, CDCl₃): δ 7.35 (dd, *J*₁ = 7.9 Hz, *J*₂ = 1.6 Hz, 1H), 7.22–7.17 (m, 1H), 7.15* (d, *J* = 9.1 Hz, 2H), 7.07 (dd, *J*₁ = 8.1 Hz, *J*₂ = 1.5 Hz, 1H), 6.93 (td, *J*₁ = 7.6 Hz, *J*₂ = 1.6 Hz, 1H), 6.59* (d, *J* = 9.1 Hz, 2H), 3.32–3.26* (m, 2H), 3.05–2.99 (m, 2H), 2.89* (s, 3H), 2.77 (s, 3H), 1.62–1.52* (m, 1H), 1.62–1.52 (m, 1H), 1.50–1.39* (m, 2H), 1.50–1.39 (m, 2H), 0.94* (d, *J* = 6.6 Hz, 6H), 0.90 (d, *J* = 6.5 Hz, 6H). ¹³C NMR (101 MHz, CDCl₃): δ 150.2, 147.9*, 130.7, 129.0, 129.0*, 127.3, 123.2, 121.3, 120.6*, 113.3*, 54.4, 51.2*, 41.0, 38.4*, 36.2, 35.0*, 26.4, 26.3*, 22.9, 22.8*. HRMS: ESI⁺ [M + H]⁺ calcd for C₁₂H₁₉NCl 212.1206, found 212.1196.

2-Chloro-N-cyclohexyl-N-dimethylaniline (3n).—Obtained as a yellow oil (isolated as a mixture of regioisomers, 4.9:1 (6-Cl:2-Cl), asterisk denotes minor peaks, 103 mg, 65%). TLC 10% ethyl acetate–hexanes, *R_f* = 0.82 (UV, KMnO₄). ¹H NMR (600 MHz, CDCl₃): δ 7.35 (dd, *J*₁ = 7.9 Hz, *J*₂ = 1.6 Hz, 1H), 7.19–7.16 (m, 1H), 7.15* (d, *J* = 9.0 Hz, 2H), 7.07 (dd, *J*₁ = 8.1 Hz, *J*₂ = 1.5 Hz, 1H), 6.91 (td, *J*₁ = 7.6 Hz, *J*₂ = 1.6 Hz, 1H), 6.68* (d, *J* = 9.1 Hz, 2H), 3.52–3.46* (m, 1H), 3.14 (m, 1H), 2.74* (s, 3H), 2.71 (s, 3H), 1.84 (m, 1H), 1.87–1.73 (m, 3H), 1.69* (m, 1H), 1.65–1.58* (m, 3H), 1.45 (m, 3H), 1.39–1.30* (m, 3H), 1.29–1.20 (m, 3H), 1.11* (m, 3H). ¹³C NMR (101 MHz, CDCl₃): δ 150.0, 148.8*, 130.7, 129.4, 129.2*, 128.9*, 127.0, 123.0, 122.7, 114.3*, 61.6, 58.5*, 33.8, 31.4*, 30.1*, 29.2, 26.3*, 26.2, 26.1, 26.0*. HRMS: ESI⁺ [M + H]⁺ calcd for C₁₃H₁₉NCl 224.1206, found 224.1195.

9-Chloro-N-methyl-1,2,3,4-tetrahydroquinoline (3o).—Obtained as a yellow oil (isolated as a mixture of regioisomers, 2.1:1 (9-Cl:7-Cl), asterisk denotes minor peaks, 78 mg, 61%). TLC 20% ethyl acetate- hexanes, R_f = 0.64 (UV, KMnO_4). ^1H NMR (400 MHz, CDCl_3): δ 7.18–7.15 (m, 1H), 7.01–6.98* (m, 1H), 6.97–6.94 (m, 1H), 6.91* (dt, J_1 = 2.4 Hz, J_2 = 1.0 Hz, 1H), 6.83 (t, J = 7.7 Hz, 1H), 6.48* (d, J = 8.7 Hz, 1H), 3.22–3.18* (m, 1H), 3.16–3.12 (m, 1H), 2.88* (s, 3H), 2.86 (s, 3H), 2.80 (t, J = 6.7 Hz, 2H), 2.73* (t, J = 6.6 Hz, 2H), 1.98–1.92* (m, 2H), 1.88–1.82 (m, 2H). ^{13}C NMR (101 MHz, CDCl_3) δ : 146.0*, 145.4*, 131.4, 128.4 (2 resonances), 128.3, 128.0, 127.7*, 126.7*, 124.5*, 122.2, 120.7*, 112.0, 52.1, 51.2*, 42.9, 39.3*, 28.0, 27.8*, 22.3*, 17.1. HRMS: ESI^+ $[\text{M} + \text{H}]^+$ calcd for $\text{C}_{10}\text{H}_{13}\text{NCl}$ 182.0737, found 182.0728.

ACKNOWLEDGMENTS

The University of Delaware (UD), the American Chemical Society Petroleum Research Fund (ACS PRF 54452-ND1), and the National Science Foundation (NSF CHE-1664954) are gratefully acknowledged for financial support. Spectral data were acquired at UD on instruments obtained with the assistance of NSF and NIH funding (NSF CHE0421224, CHE0840401, CHE1229234, CHE1048367; NIH S10 OD016267-01, S10 RR026962-01, P20GM104316, P30GM110758).

REFERENCES

- (1). (a) Henry MC; Mostafa MA; Sutherland A Recent Advances in Transition-Metal-Catalyzed, Directed Aryl C–H/N–H Cross-Coupling Reactions. *Synthesis* 2017, 49, 4586–4598. (b) Jana R; Pathak TP; Sigman MS Advances in Transition Metal (Pd, Ni, Fe)-Catalyzed Cross-Coupling Reactions Using Alkyl-organometallics as Reaction Partners. *Chem. Rev* 2011, 111, 1417–1492. [PubMed: 21319862] (c) Johansson Seechurn CCC; Kitching MO; Colacot TJ; Snieckus V Palladium-Catalyzed Cross-Coupling: A Historical Contextual Perspective to the 2010 Nobel Prize. *Angew. Chem., Int. Ed* 2012, 51, 5062–5085.
- (2). Seyferth D Alkyl and Aryl Derivatives of the Alkali Metals: Useful Synthetic Reagents as Strong Bases and Potent Nucleophiles. 1. Conversion of Organic Halides to Organoalkali-Metal Compounds. *Organometallics* 2006, 25, 2–24.
- (3). (a) Ashby EC Grignard reagents. Compositions and mechanisms of reaction. *Q. Rev., Chem. Soc* 1967, 21, 259–285. (b) Seyferth D The Grignard Reagents. *Organometallics* 2009, 28, 1598–1605. (c) Richey HG Grignard Reagents: New Developments; Wiley: New York, 2000.
- (4). (a) Gribble GW Naturally Occurring Organohalogen Compounds. *Acc. Chem. Res* 1998, 31, 141–152. (b) Wilcken R; Zimmermann MO; Lange A; Joerger AC; Boeckler FM Principles and Applications of Halogen Bonding in Medicinal Chemistry and Chemical Biology. *J. Med. Chem* 2013, 56, 1363–1388. [PubMed: 23145854]
- (5). (a) Carreno MC; Garcia Ruano JL; Sanz G; Toledo MA; Urbano A N-Bromosuccinimide in Acetonitrile: A Mild and Regiospecific Nuclear Brominating Reagent for Methoxybenzenes and Naphthalenes. *J. Org. Chem* 1995, 60, 5328–5331. (b) Goldberg Y; Alper H Biphasic electrophilic halogenation of activated aromatics and heteroaromatics with N-halosuccinimides catalyzed by perchloric acid. *J. Org. Chem* 1993, 58, 3072–3075. (c) Gervat S; Leonel E; Barraud J-Y; Ratovelomanana V High regioselectivity of bromination of anilines by tetraethylammonium chloride/methanol system as cocatalysts under mild conditions. *Tetrahedron Lett.* 1993, 34, 2115–2118. (d) Olah GA; Kuhn SJ; Hardie BA Aromatic Substitution. XVII. 1 Ferric Chloride and Aluminum Chloride Catalyzed Chlorination of Benzene, Alkylbenzenes, and Halobenzenes. *J. Am. Chem. Soc* 1964, 86, 1055–1060. (e) Hope E; Riley GF Chlorination of benzoyl chloride. Part I. *J. Chem. Soc., Trans* 1922, 121, 2510–2527.
- (6). For recent examples of metal-mediated halogenation of anilines, see: (a) Urones B; Martínez AM; Rodríguez N; Gómez Arrayas R; Carretero JC Copper-catalyzed ortho-halogenation of protected anilines. *Chem. Commun* 2013, 49, 11044–11046. (b) Kalyani D; Dick AR; Anani WQ; Sanford MS A Simple Catalytic Method for the Regioselective Halogenation of Arenes. *Org. Lett* 2006, 8, 2523–2526. [PubMed: 16737304]

- (7). (a)Arora A in Aromatic Organic Chemistry; Discovery Publishing House, India, 2008.
(b)Hepworth JD; Waring DR; Waring JM In Aromatic Chemistry; Royal Society of Chemistry: Cambridge, 2002.(c)Taylor R Electrophilic Aromatic Substitution; Wiley: New York, 1990.
(d)Katritzky AR; Taylor R Electrophilic Substitution of Heterocycles: Quantitative Aspects In Advances in Heterocyclic Chemistry; Academic Press: New York, 1990; Vol. 47.(e)Taylor R In Comprehensive Chemical Kinetics; Bamford CH, Tipper CFH, Eds.; Elsevier: New York, 1972; Vol. 13, pp 1–406.(f)De la Mare PB Electrophilic Halogenation; Cambridge University Press: Cambridge, 1976; Chapter 5.
- (8). (a)Lewis RS; Wisthoff MF; Grissmerson J; Chain WJ Metal-Free Functionalization of N,N-Dialkylanilines via Temporary Oxidation to *N,N*-Dialkylaniline *N*-Oxides and Group Transfer. Org. Lett 2014, 16, 3832–3835. [PubMed: 24992642] (b)Nakashige ML; Lewis RS; Chain WJ Transformation of N,N-dimethylaniline-*N*-oxides into *N*-methylindolines by a tandem Polonovski–Mannich reaction. Tetrahedron Lett. 2015, 56, 3531–3533.
- (9). N-Arylhydroxylamine rearrangement:(a)Bassoli A; Di Gregorio G; Galliani G; Riboldi M; Rindone B; Tollari S; Chioccare F The heterolytic and homolytic cleavage of the oxygen-nitrogen bond in O,N-diacylarylhydroxylamines. Bull. Chim. Soc. Fr 1988, 293–297.(b)Pereira M; Manuela A; Santos PP In Chemistry of Hydroxylamines, Oximes and Hydroxamic Acids, Part 1;Rappoport Z, Liebman JF, Eds.; John Wiley and Sons: Chichester, 2009.(c)Luo YR In Comprehensive Handbook of Chemical Bond Energies; CRC Press and Taylor & Francis Group: Boca Raton, 2007; p 353.For a recent review of the chemistry of hydroxylamines, see:(d)Tabolin AA; Ioffe SL Rearrangement of N-Oxyenamines and Related Reactions. Chem. Rev 2014, 114, 5426–5476. [PubMed: 24628517] (e)Horner L; Steppan H Studien Zum Ablauf Der Substitution XII. Umlagerung Und Thermischer Zerfall Acylierter Hydroxylamine. Justus Liebigs Ann. Chem 1957, 606, 24–47.(f)Oae S; Sakurai T; Kimura H; Kozuka S Oxygen-18 Tracer Study of The Rearrangement of o-Benzoyl-N-(p-Toluenesulfonyl)-arylhydroxylamines. Chem. Lett 1974, 3, 671–674.(g)Gutschke D; Heesing A; Heuschkel U Zur thermischen umlagerung von O-arylsulfonyl-N-phenylhydroxylaminen. Tetrahedron Lett. 1979, 20, 1363–1364.
- (10). N-Arylhydroxylamine halogenation:(a)Ayyangar NR; Kalkote UR Novel orthohalogenation reaction. Synthesis of orthochloroarylamines from nitroarenes. Tetrahedron Lett. 1982, 23, 1099–1102.(b)Marx L; Rassat A Hetero-Cope rearrangement for the synthesis of potassium 5-tert-butyl-2-(tert-butyl-aminoxyl)-benzoate, a highly water-soluble stable free radical. Tetrahedron Lett. 2002, 43, 2613–2614.(c)Kikugawa Y; Mitsui K Migration of the Hydroxyl Group of N-Hydroxy-N-phenylamides to the Phenyl Group with Tertiary Phosphines and Tetrachloromethane. A Novel Transhydroxylation Reaction. Chem. Lett 1993, 22, 1369–1372.
- (11). (a)Hill RK In Comprehensive Organic Synthesis; Trost BM, Ed.; Elsevier: New York, 1991; Vol. 5, pp 785–826.(b)Wipf P In Comprehensive Organic Synthesis; Trost BM, Ed.; Elsevier: New York, 1991; Vol. 5, pp 827–873.
- (12). Pangborn AB; Giardello MA; Grubbs RH; Rosen RK; Timmers FJ Safe and Convenient Procedure for Solvent Purification. Organometallics 1996, 15, 1518–1520.
- (13). Still WC; Kahn M; Mitra A Rapid chromatographic technique for preparative separations with moderate resolution. J. Org. Chem 1978, 43, 2923–2925.
- (14). Chandrasekharam M; Chiranjeevi B; Gupta KSV; Sridhar B Iron-Catalyzed Regioselective Direct Oxidative Aryl–Aryl Cross-Coupling. J. Org. Chem 2011, 76, 10229–10235. [PubMed: 22070185]

Previous Efforts **This Work**

Scheme 1.
Aniline Functionalization by Group Transfer

Table 1.

Optimization of the *N,N*-Dimethylaniline *N*-Oxide Halogenation Reactions^a

entry	solvent	temp (°C)	X = Br, yield (4-Br) (%)	X = Cl, yield (2-Cl/4-Cl) (%)
1	CH_2Cl_2	-78	31	25 (4.3:1)
2	CH_2Cl_2	-40	26	22 (4.0:1)
3	CH_2Cl_2	-85	13	15 (6.6:1)
4	Et_2O	-78	27	20 (3.6:1)
5	THF	-78	55	35 (4.0:1)
6 ^b	THF	-78	53	49 (4.9:1)

^aYields of isolated products; reactions were performed on a 0.7 mmol scale. Addition of SOX_2 was followed by stirring at the indicated temperature for 4 h.

^bAddition of SOX_2 was divided into two portions (0.5 equiv, followed by 0.5 equiv after 2 h).

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 2.

Selective 4-Bromination of Substituted *N,N*-Dialkylaniline *N*-Oxides^a

Yields of isolated products; reactions were performed on a 0.7 mmol scale.

η Reaction conducted in dichloromethane as solvent.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 3.

Yields of isolated products; reactions were performed on a 0.7 mmol scale.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript